Title	New stent exchange technique following endoscopic ultrasound-guided nasobiliary drainage (with video)
Author(s)	Kawakami, Hiroshi; Kuwatani, Masaki; Sakamoto, Naoya
Citation	Digestive Endoscopy, 25(3), 343-344 https://doi.org/10.1111/den.12051
Issue Date	2013-05
Doc URL	http://hdl.handle.net/2115/55278
Rights	The definitive version is available at www.wileyonlinelibrary.com
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	den12051.pdf

[Instructions for use](#)

**Internal biliary stent placement following endoscopic ultrasound-guided
nasobiliary drainage**

Running head: Stent exchange technique after EUS-BD

Hiroshi Kawakami,¹; Masaki Kuwatani,¹; Naoya Sakamoto,¹

¹ Department of Gastroenterology and Hepatology, Hokkaido University Graduate

School of Medicine, Sapporo, Japan

Address correspondence to: Hiroshi Kawakami, MD, PhD

Department of Gastroenterology and Hepatology, Hokkaido University Graduate School
of Medicine

Kita 15, Nishi 7, Kita-ku, Sapporo 060-8638, Japan

Fax: +81 11 706 7867

E-mail: hiropon@med.hokudai.ac.jp (H. Kawakami)

An 83-year-old man with obstructive jaundice was admitted to our department. He had undergone chemotherapy for malignant lymphoma. Seventeen months later, computed tomography revealed a distal bile duct mass. Endoscopic ultrasound (EUS)-guided fine-needle aspiration of the mass revealed malignant lymphoma. Endoscopic retrograde cholangiopancreatography (ERCP) was unsuccessful due to difficult cannulation, so EUS-guided choledochoduodenostomy (EUS-CDS) was performed. Although a 0.025-inch guidewire was advanced into the intrahepatic bile duct, a tapered ERCP/balloon/Soehendra biliary dilation catheter (SBDC-6, Cook-Japan, Tokyo, Japan) could not be passed across the tract. We attempted to dilate the tract using a guidewire-induced needle knife, but the knife was displaced from the axis of the bile duct between the duodenal bulb and bile duct. Then, a 5-Fr nasobiliary tube was put in place. Fourteen days after EUS-CDS, we tried to exchange the nasobiliary stent with a duodenoscope. First, the nasobiliary tube was cut using scissors (FS-3L-1, Olympus Medical Systems, Tokyo, Japan) and loop cutter forceps (FS-5L-1, Olympus). Next, the remnant tube was cannulated using a wire-guided technique via the ERCP catheter. The tube was trapped in the snare with the snare-over-the-wire technique [1] (**Fig. 1**). After removing the remnant tube via the working channel with the snare-over-the-wire technique, a 7-Fr plastic stent was put in place without complications (**Fig. 2**).

EUS-CDS is a new biliary drainage technique [2]. However, when the nasobiliary stent is first put in place, an internal stent is necessary for the second step via the first puncture tract. We describe a rare case of stent exchange after EUS-guided nasobiliary drainage that was performed using scissors, loop cutter forceps, and a snare-over-the-wire technique. This technique might be particularly useful after EUS-CDS is put in place with nasobiliary drainage.

REFERENCES

1. Fujita N, Sugawara T, Noda Y *et al.* Snare-over-the-wire technique for safe exchange of a stent following endosonography-guided biliary drainage. Dig Endosc 2009;21:48-52
2. Itoi T, Isayama H, Sofuni A *et al.* Stent selection and tips on placement technique of EUS-guided biliary drainage: transduodenal and transgastric stenting. J Hepatobiliary Pancreat Sci 2011;18:664-72

FIGURE LEGENDS

Fig. 1: Endoscopic image showing the snare-over-the-wire technique via the remnant nasobiliary tube following endoscopic ultrasound-guided nasobiliary drainage.

Fig. 2: A 7-Fr plastic stent was placed via the fistula without complications after endoscopic ultrasound-guided nasobiliary drainage.

Video

The nasobiliary tube was cut using scissors (FS-3L-1, Olympus Medical Systems, Tokyo, Japan) and loop cutter forceps (FS-5L-1, Olympus). The remnant tube was cannulated using a wire-guided technique via the ERCP catheter. The tube was trapped in the snare with the snare-over-the-wire technique. After removing the remnant tube via the working channel with the snare-over-the-wire technique, a 7-Fr plastic stent was put in place without complications.