



Title	PARTIALLY ORDERED ABELIAN SEMIGROUPS II. ON THE STRONGNESS OF THE LINEAR ORDER DEFINED ON ABELIAN SEMIGROUPS
Author(s)	Nakada, Osamu
Citation	Journal of the Faculty of Science Hokkaido University. Ser. 1 Mathematics, 12(2), 073-086
Issue Date	1952
Doc URL	http://hdl.handle.net/2115/55977
Type	bulletin (article)
File Information	JFSHIU_12_N2_073-086.pdf



[Instructions for use](#)

PARTIALLY ORDERED ABELIAN SEMIGROUPS II. ON THE STRONGNESS OF THE LINEAR ORDER DEFINED ON ABELIAN SEMIGROUPS

By

Osamu NAKADA

In my previous paper⁽¹⁾, I note that a strong linearly ordered abelian semigroup (strong l.o. semigroup) is always normal, its elements are of infinite order except the unit element (if there exists) and the product cancellation law is held in it. In this Part, I shall seek for the other conditions which are exchangeable for the cancellation law. But the some theorems in this note will be expressed in the form of a partially ordered abelian semigroup.

Definition 1. Let a be any element of an abelian semigroup S . We denote the set of all positive powers of a , in other words, the sub-semigroup of S generated by a , by $S(a)$, which is called the *sector* by a . And let $T(a)$ be the set of all elements of S whose some positive powers belong to $S(a)$, that is, the set of all elements x of S such that $x^m = a^n$ for some positive integers m and n . $T(a)$ is called the *complete sector* by a .

Let a and b be any two elements of S . Then the two complete sectors $T(a)$ and $T(b)$ are either disjoint or identical. Hence, S is the union of mutually disjoint complete sectors, and clearly $T(a)$ is a sub-semigroup of S .

Definition 2.⁽²⁾ Let a be any element of an abelian semigroup S , and we shall consider the sector $S(a)$ by a . There are two possible cases.

(1) Partially ordered abelian semigroups. I. On the extension of the strong partial order defined on abelian semigroups. Journal of the Faculty of Science, Hokkaido University, Series I, vol. XI, No. 4 (1951) pp. 181-189; this is referred to hereafter as "O.I."

A set S is said to be a *partially ordered abelian semigroup* (p.o. semigroup), when S is (I) an abelian semigroup (not necessarily contains the unit element), (II) a partially ordered set, and satisfies (III) the *homogeneity*: $a \geq b$ implies $ac \geq bc$ for any c of S . When a partial order P of an abelian semigroup S satisfies the condition (III), we call P a *partial order defined on an abelian semigroup* S . (Definition 1, O.I.)

(2) Cf. D. REES: On semi-groups, Proc. Cambridge Phil. Soc., vol. 36 (1940), pp. 387-400.

First, a may satisfy no equation of the form $a^m = a^n$ ($m \neq n$). In this case, the element a is said to be of *infinite order*.

Secondly, a may satisfy a relation of the form $a^m = a^n$ ($n > m$). If a satisfies such a relation, choose one for which n is a minimum. Then clearly a, a^2, \dots, a^{n-1} are distinct elements of $S(a)$. We write $n = m + r$, $r \geq 1$. Then, if q is any positive integer, by simple induction we have immediately $a^m = a^{m+qr}$. Now, if N is any integer not less than n , we may write it in the form $N = m + qr + s$ ($1 \leq q, 0 \leq s < r$). Then, by the above, we have $a^N = a^{m+s}$. Accordingly, it follows that the sector $S(a)$ just consists of the $n-1$ elements a, a^2, \dots, a^{n-1} . Further, let t be the least integer of the form $qr+1$ greater than m and put $d = a^t$. Then, since t is prime to r , the elements d, d^2, \dots, d^r are distinct from one another and coincide with the elements $a^m, a^{m+1}, \dots, a^{n-1}$, and it is easily seen that

$$\begin{aligned} d^r d^s &= d^s d^r = d^{r+s} = d^s & (s \leq r), \\ d^{r-s} d^s &= d^s d^{r-s} = d^r & (s < r). \end{aligned}$$

Then the above shows that $a^m, a^{m+1}, \dots, a^{n-1}$ form a cyclic group, generated by d , with the identity d^r . Such an element a is said to be of *finite order*. We shall term m the *length* and r the *period* of the element a of finite order.

An element a of finite order is called *cyclic, quasi-idempotent or idempotent*, when $m = 1, r = 1$ or $m = 1$ and $r = 1$ respectively. The immediate consequence of the above consideration is that any semigroup containing elements of finite order contains idempotent elements.

Moreover, from the definition of the complete sector we get readily the following: The complete sector by the element of finite order does not contain the element of infinite order and contains one and only one idempotent element, and the complete sector by the element of infinite order does not contain the element of finite order.

Definition 3. We consider the following conditions for an abelian semigroup S :

- (α): $a^n = b^n$ for some positive integer n implies $a = b$.
- (β): All elements of S are of infinite order except the unit element (if there exists).
- (γ): If a is not the unit element, then $ab \neq b$ for any b of S .
- (δ): If two complete sectors $T(a)$ and $T(b)$ are disjoint, then $c \cdot T(a)$ ⁽³⁾

(3) $c \cdot T(a)$ is the set of all elements $cx, x \in T(a)$.

and $c \cdot T(b)$ are disjoint for any c of S .

Moreover, we consider the following conditions for a p.o. semigroup S :

(A): If a is positive or negative⁽⁴⁾, then $ab \geq b$ or $b \geq ab$ for any b of S respectively.

(B): If a (\neq the unit element) is positive or negative, then $ab > b$ or $b > ab$ for any b of S respectively.

Theorem 1. If an abelian semigroup S satisfies the condition (A), then the element of S is idempotent or of infinite order.

Proof. Let a be an element of finite order, m and r be the length and the period of a respectively. And let $Z = \{a^m, \dots, a^{m+r-1}\}$, then Z is a cyclic group. Hence there exists a positive integer q such that $a^q = (a^2)^q = y$, where y is the identity of the cyclic group Z . Therefore, by the condition (A) we have $a = a^2$, that is, a is idempotent.

Corollary. An element of a normal⁽⁵⁾ p.o. semigroup S is idempotent or of infinite order.

Proof. For, S satisfies the condition (A) by Theorem 12, O.I.

Example 1. Let S_1' be a free abelian semigroup generated by one element a and $S_1'' = \{b, e = \text{the identity}\}$ be a group of order 2, which does not satisfy the conditions (A) and (B). And let S_1 be a direct product of S_1' and S_1'' in the usual sense. Then S_1 becomes the abelian semigroup which satisfies the conditions (B), (C) and (D), but, since $(xe)^2 = (xb)^2$ in spite of $xe \neq xb$ ($x \in S_1'$), does not satisfy the condition (A).

Clearly an abelian semigroup, all elements of which are idempotent, satisfies the condition (A). For instance, see Example 8, which does not satisfy the conditions (B), (C) and (D).

The conditions (A) and (B) are equivalent to each other in abelian groups but not in semigroups.

Theorem 2. If an abelian semigroup S satisfies the condition (A), then any complete sector $T(a)$ is isomorphic with a sub-semigroup of the additive semigroup of the non-negative rational numbers.

Proof. By Theorem 1, an element of S is idempotent or of infinite order.

(4) An element a of a p.o. semigroup S is called *positive* or *negative*, when $a^2 \geq a$ or $a \geq a^2$ respectively. (Definition 4, O.I.)

(5) A partial order defined on an abelian semigroup S is called *normal*, when the following condition is satisfied:

$a^n \geq b^n$ for some positive integer n implies $a \geq b$. (Definition 6, O.I.)

i) a is idempotent. Let x be any element of $T(a)$, then x is of finite order, and hence x is idempotent. Therefore, $x = a$. Then the complete sector $T(a)$ consists of only a . Consider the correspondence $a \longleftrightarrow 0$.

ii) a is of infinite order. Let x be any element of the complete sector $T(a)$. Then x is of infinite order and there exist positive integers n and m such that $x^m = a^n$. We consider the correspondence $x \rightarrow n/m$.

First, let $x^m = a^n$ and $x^{m'} = a^{n'}$, then $x^{mm'} = a^{nn'} = x^{nm'}$. Hence $mm' = nm'$ because x is of infinite order. Therefore we have $n/m = n'/m'$, i.e., the correspondence is unique.

Now, let $y \in T(a)$, $y \rightarrow j/i$, that is, $y^i = a^j$. If $n/m = j/i$, then $x^{mi} = a^{ni} = a^{mj} = y^{mi}$, and hence $x = y$ by the condition (α) , i.e., this correspondence is one-to-one.

Finally, since $x^{mi} = a^{ni}$ and $y^{mi} = a^{mj}$, we have $(xy)^{mi} = a^{ni+mj}$, therefore, $xy \rightarrow (ni+mj)/mi = n/m + j/i$.

Consequently, the correspondence $x \longleftrightarrow n/m$ is the isomorphism of $T(a)$ into the additive semigroup of the positive rational numbers.

Corollary 1. If an abelian semigroup S satisfies the condition (α) , then the product cancellation law is held in a (complete) sector by any element of S .

Corollary 2. If an abelian semigroup S satisfies the condition (α) , then a strong⁽⁶⁾ linear order may be defined on a (complete) sector by any element of $S^{(7)}$.

Definition 4. Let S be an abelian semigroup and P be a partial order defined on S . And let T be a subset of S . If any two elements of T are comparable in P , then T is called a *comparable subset* (in P) of S , on the contrary if any two distinct elements of T are non-comparable in P , then T is called a *non-comparable subset* (in P) of S .

Next, let U and V be two disjoint subsets of S . If any elements $u \in U$ and $v \in V$ are comparable in P , then U and V are called *comparable subsets to each other* (in P) of S , and if any elements u and v are non-comparable in P , then they are called *non-comparable subsets to each other* (in P) of S (or we say that U is comparable (or non-comparable) to V in P).

If U and V are comparable subsets to each other in P and $u > v$ in P for any $u \in U$ and $v \in V$, then we denote $U > V$ in P .

(6) A p.o. semigroup S is called *strong*, when the following condition is satisfied in S : $ac \geq bc$ implies $a \geq c$. (Definition 2, O.I.)

(7) Cf. Theorem 15, O.I.

If a complete sector by an element of *infinite order* contains a positive element, then it does not contain a negative element. Accordingly, we shall call such a complete sector *positive*. Similarly, we may define a *negative* complete sector.

Theorem 3. Let S be a normal p.o. semigroup. Then the positive or negative complete sector is the comparable subset of S .

Proof. Let $T(a)$ be a positive complete sector, where we assume that a is positive without loss of generality. Then clearly the sector $S(a)$ is the comparable subset. Now let u and v be any two elements of the complete sector $T(a)$, then there exist positive integers i, j and k such that $u^k = a^i$ and $v^k = a^j$. Since a^i and a^j are comparable, say that $a^i \geq a^j$, we have $u^k = a^i \geq a^j = v^k$, therefore by the normality $u \geq v$. Similarly, a negative complete sector is a comparable subset of S .

Example 2. Let S_2 be a free abelian semigroup generated by one element a with the order-relation

$$a^{2n+1} > a^{2m+1} \ (n > m \geq 0), \quad a^{2n} > a^{2m} \ (n > m > 0).$$

Then S_2 is a p.o. semigroup which satisfies the condition (α) and $T(a) = S_2$. But S_2 is not normal because a^2 and a^3 are non-comparable in spite of $(a^3)^2 = a^6 > a^4 = (a^2)^2$. Since $(a^2)^2 = a^4 > a^2$, a^2 is positive, that is, $T(a)$ is the positive complete sector but is not the comparable subset.

Or, by putting the other order-relation

$$a^{n+1} > a^n \text{ for any positive integer } n > N \geq 1,$$

S_2 becomes a p.o. semigroup which is not normal. And $T(a) = S_2$ is the positive complete sector but is not the comparable subset.

Or, we define the order-relation in S_1 (Example 1) as follows:

$$a^{n+1}e > a^ne, \quad a^{n+1}b > a^nb \text{ for any positive integer } n.$$

Then S_1 becomes the p.o. semigroup which is not normal because S_1 does not satisfy the condition (α) , and $T(ae) = S_1$ is the positive complete sector but is not the comparable subset.

Example 3. Let S_3' be a free abelian semigroup generated by one element a and S_3'' be an abelian semigroup generated by two elements b and 0 with the relations

$$b^2 = 0, \quad 0^2 = 0, \quad b0 = 0b = 0.$$

And let S_3 be a direct product of S_3' and S_3'' . Then S_3 satisfies the conditions (β) , (γ) and (δ) .

By putting the order-relation

$$a0 < ab < a^2 0 < \dots < a^n 0 < a^n b < a^{n+1} 0 < \dots,$$

S_3 becomes a l.o. semigroup. Clearly S_3 does not satisfy the condition (α), i.e., S_3 is not normal, but $T(a0) = S_3$ is a comparable positive complete sector.

We note that S_3'' satisfies the condition (δ) but does not the conditions (α), (β) and (γ).

Example 4. Let S_4' and S_4'' be free abelian semigroups generated by elements a and b respectively. And by defining the order-relations

$$a^{m+1} > a^m \quad (m \geq 1), \quad b^{n+1} > b^n \quad (n > 1),$$

S_4' and S_4'' becomes a l.o. semigroup and a p.o. semigroup respectively. Let S_4 be a direct product of S_4' and S_4'' . Then S_4 is an abelian semigroup which satisfies the conditions (α), (β), (γ) and (δ), and S_4 is the union of mutually disjoint complete sectors:

$$S_4 = \bigcup_{i,j} T(a^i b^j),$$

where i and j run over all relatively prime positive integers. For, let x be any element of S_4 , then $x = a^m b^n = (a^p b^q)^d$, where $(m, n) = d$ and $(p, q) = 1^{(8)}$, and hence $x \in T(a^p b^q)$. Next, suppose that $T(a^i b^j) = T(a^m b^n)$ and $(i, j) = (m, n) = 1$. Then we get easily $i = m$ and $j = n$.

Now, we define the order-relation in S_4 as follows:

$$a^i b^j > a^m b^n$$

if and only if $a^i > a^m$ or $a^i = a^m$ and $b^j > b^n$.

Then S_4 becomes a p.o. semigroup. Since ab^2 and ab are non-comparable in spite of $(ab^2)^2 = a^2 b^4 > a^2 b^2 = (ab)^2$, S_4 is not normal. On the other hand, any complete sector $T(a^i b^j)$, where without loss of generality $(i, j) = 1$, is a comparable subset and is positive. For, if x and y are any two elements of $T(a^i b^j)$, then they are expressed in the form $x = (a^i b^j)^p$ and $y = (a^i b^j)^q$. Hence, as can easily be seen x and y are comparable, and, since $(a^i b^j)^3 > a^i b^j$, $T(a^i b^j)$ is a positive complete sector.

Theorem 4. Let S be a normal p.o. semigroup. Then a complete sector $T(a)$ is order-isomorphic or anti-order-isomorphic with a sub-semigroup of the additive semigroup of the non-negative rational numbers, when a is positive or negative respectively.

(8) (i, j) is G.C.M. of i and j .

Proof. An element a of S is of finite order, i.e., a is idempotent by Theorem 1, if and only if a is positive and negative. In such a case, $T(a) = a$, consequently the theorem is trivial.

Suppose now that a is of infinite order and positive. Then $a^2 > a$, and hence $a^n > a^m$ if and only if $n > m$. Further, by Theorem 3 the complete sector $T(a)$ is a comparable subset of S . Let x be any element of $T(a)$, then there exist positive integers m and n such that $x^m = a^n$. By Theorem 2, the correspondence $x \longleftrightarrow n/m$ is the algebraic isomorphism of $T(a)$ into the additive semigroup of positive rational numbers.

Let $y \longleftrightarrow j/i$, $y \in T(a)$, that is, $y^i = a^j$.

i) Let $n/m > j/i$, i.e., $ni > mj$. Since $x^{mi} = a^{ni}$ and $y^{mi} = a^{mj}$, we have $x^{mi} = a^{ni} > a^{mj} = y^{mi}$. By the normality we have $x > y$.

ii) Conversely, let $x > y$. Then we have $x^{mi} > y^{mi}$ by the normality. And hence $a^{ni} > a^{mj}$. Therefore, $ni > mj$, i.e., $n/m > j/i$.

Similarly, a complete sector $T(a)$, where a is of infinite order and negative, is anti-order-isomorphic with a sub-semigroup of the additive semigroup of the positive rational numbers.

Corollary. If a l.o. semigroup S satisfies the condition (α), then any complete sector $T(a)$ is order-isomorphic or anti-order-isomorphic with a sub-semigroup of the additive semigroup of the non-negative rational numbers when a is positive or negative respectively.

Proof. By the linearity of S , any elements a of S is positive or negative. And, since the following property is held always in a l.o. semigroup: $a^n > b^n$ for some positive integer n implies $a > b$, then the condition (α) and the normality are equivalent to each other in a l.o. semigroup.

Theorem 5. Let S be a l.o. semigroup. And let $T(a)$ be a positive, $T(b)$ be a negative complete sector and g be an element of finite order. Then

- i) $T(a) > T(b)$ or $T(b) > T(a)$,
- ii) if $T(u)$ is any complete sector different from $T(g)$, then $T(u) > T(g)$ or $T(g) > T(u)$,
- iii) if $T(b) > T(a)$, then $b > ab > a$ and $ab = a^m b^n$ for any positive integers m and n .

Proof. i): Suppose that $a > b$. Let x be any element of $T(a)$ and y be any element of $T(b)$, then there exist positive integers i, j and k such that $x^i = a^i$ and $y^k = b^k$. But, since $a^i \geq a > b \geq b^k$, we have $x^i = a^i > b^k = y^k$ and hence by the linearity $x > y$, that is, $T(a) > T(b)$. Next,

let $b > a$. If there exist the elements $x \in T(a)$ and $y \in T(b)$ such that $x > y$, $x^k = a^i$ and $y^k = b^j$, then we have $x^k > y^k$ because $T(a)$ and $T(b)$ are disjoint, and hence $a^i > b^j$. If $i \geq j$, then $a^i > b^j \geq b^i$, hence we have $a > b$, which is absurd. Similarly, $j \geq i$ implies $a^j \geq a^i > b^j$, i.e., we have $a > b$, contrary to $b > a$. Therefore, $T(b) > T(a)$.

ii): Let f be the (only one) idempotent element belonging to $T(g)$. Clearly $T(g) = T(f)$. Suppose now that $u > f$. Since $T(u)$ and $T(f)$ are disjoint, we have $u^n > f$ for every positive integer n . If $x \in T(u)$ and $y \in T(f)$, then we get easily $x > y$, therefore $T(u) > T(g)$. Similarly, if $f > u$, then $T(g) > T(u)$.

iii): $b > a$ implies that $b > b^2 \geq ab \geq a^2 > a$, i.e., $b > ab > a$. Furthermore, we have

$$\begin{aligned} b &> b^2 > b^3 \geq ab^2 \geq a^2b \geq ab, \\ ab &\geq ab^2 \geq a^2b \geq a^3 > a^2 > a. \end{aligned}$$

Hence $ab^2 = ab = a^2b$. Now we assume that $a^m b^n = ab$, $m+n = N$. Then $a^{m+1}b^n = a^2b = ab = ab^2 = a^m b^{n+1}$. Since ab is idempotent and $b > ab > a$, we have $T(b) > T(ab) > T(a)$ by ii).

Let f be an idempotent element and $T(f) \not\supseteq f$. If g ($\neq f$) of $T(f)$ is positive and h ($\neq f$) of $T(f)$ is negative, then

$$\begin{aligned} h &> h^2 > \dots > h^t = f = g^s > \dots > g^2 > g, \\ h^m g^n &= f \text{ for any positive integers } m \text{ and } n, \end{aligned}$$

where s and t are the lengths of g and h respectively.

Corollary. If S has the unit element e , then the above theorem is somewhat brief:

- 1) $T(a) > e > T(b)$, $T(e) = e$,
- 2) if $T(g) > e$, then the element h of $T(g)$ differing from the idempotent element f belonging to $T(g)$, is positive, i.e., $f \geq h^2 > h$ and $fh = f$; and its dual.

The proof is obvious.

Theorem 6. Let S be a p. o. semigroup which satisfies the condition (A). Then the following properties are held in S :

- 1) If a is positive and $b \geq a$, then b is positive.
- 2) If a is negative and $a \geq b$, then b is negative.
- 3) If a is positive and b is negative, then $a \geq b$.

Proof. 1): $b \geq a$ implies $b^2 \geq ab$ by the homogeneity, and $ab \geq b$ by the condition (A). Therefore, we have $b^2 \geq b$, which says that b is posi-

tive. 2): Similarly to 1). 3): By the condition (A), we have $a \geq ab \geq b$.

Example 5. Let S_5 be an abelian semigroup generated by two elements a and b with the relation

$$ab = a^m b^n \quad \text{for any positive integers } m \text{ and } n.$$

Then S_5 satisfies the condition (α) but does not the conditions (β), (γ) and (δ).

And by putting the order-relation

$$a^{m+1} > a^m > ab > b^n > b^{n+1}$$

for any positive integers m and n ,

S_5 becomes a l.o. semigroup which satisfies the properties 1), 2) and 3) of Theorem 6 but does not the condition (A) because $a > ab = (ab)a$ in spite of $(ab)^2 \geq ab$.

Theorem 7. Let S be a p.o. semigroup which satisfies the condition (A). Then S may contain only one idempotent element, or what is the same, only one positive and negative element. Further, if there exists an idempotent element e , then it is necessarily the unit element and all positive or negative elements are comparable to e .

Proof. If S has a positive and negative element e , then by the condition (A) $eb \geq b$ and $b \geq eb$, i.e., $eb = b$ for any b of S , and hence e is the unit element. Accordingly, all elements of finite order of S belong to the complete sector $T(e)$. Let a be any positive (or negative) element. By the condition (A), we have $a = ae \geq e$ (or $e \geq ae = a$).

Corollary 1. If a p.o. semigroup S satisfies the conditions (α) and (A), then S satisfies the condition (β).

Proof. For, by Theorem 1 an element of S is idempotent or of infinite order.

Corollary 2. If a l.o. semigroup S satisfies the condition (A), then S satisfies the condition (β).

Proof. If there exists an element a of finite order, then S contains an idempotent element e . By the above theorem, e is the unit element of S and $a \in T(e)$. Moreover, by the property 1) of Corollary, Theorem 5, $T(e) = e$. Therefore we have $a = e$, that is S satisfies the condition (β).

Example 6. Let S_6 be an abelian group of finite order $n \geq 2$. And consider the vacuous order-relation of S_6 . Then S_6 satisfies the condition (A) but does not satisfy the conditions (α) and (β).

Example 7. Let S_7 be an abelian semigroup generated by two elements a and e with the relations

$$e^2 = e, \quad a^m e = e a^m = a^m \quad \text{for any positive integer } m, \\ a^n > a^m \quad \text{for } n > m.$$

Then S_7 is a p.o. semigroup with the unit element e and satisfies the conditions (α) and (β) . But, since $ae = a$ and e are non-comparable and $a^2 > a$, S_7 does not satisfy the condition (A) . (Or, see Examples 1 and 2.)

Theorem 8. Let S be a p.o. semigroup which satisfies the condition (A) . If S has not the unit element, then one can adjoin the new element e to S , so that $S^* = S \cup e$ becomes a p.o. semigroup in which e is the unit element, and S is order-embedded⁽⁹⁾ in S^* , moreover, all positive or negative elements are comparable to e in S^* .

Proof. We define the multiplication in S^* as follows:

$$e \cdot e = e, \quad e \cdot a = a \cdot e = a \quad \text{for any } a \text{ of } S,$$

and for any two elements a and b of S the product is the same as in S . Thus S^* becomes the abelian semigroup with the unit element e under the multiplication introduced above.

Let P be the partial order defined on S . Let us now define the order-relation Q in S^* as follows:

$$a > b \text{ in } Q (a, b \in S) \text{ if and only if } a > b \text{ in } P, \\ a > e \text{ in } Q (a \in S) \text{ if and only if } a \text{ is positive in } P, \\ e > a \text{ in } Q (a \in S) \text{ if and only if } a \text{ is negative in } P.$$

S has not a positive and negative element, because if there exists such an element a , then a is the unit element, by Theorem 7, which contradicts the hypothesis of the theorem. Therefore, $a > b$ and $b > a$ in Q are contradictory.

The transitivity of Q is verified by the properties 1), 2) and 3) of Theorem 6 and the homogeneity of Q is easily verified by the condition (A) itself.

Then S^* becomes a p.o. semigroup with the unit element e , and it is clear that S is order-embedded in S^* and also S^* satisfies the condition (A) .

Example 8. Let S_8 be a set of three elements a, b and c with the relations

(9) A p.o. semigroup S will be said to be *order-embedded* in a p.o. semigroup S' , if there exists an order-isomorphism of S into S' . (Definition 3, O.I.)

$$\begin{aligned} ab = ba = c, \quad ac = ca = c, \quad bc = cb = c, \\ a^2 = a, \quad b^2 = b, \quad c^2 = c, \quad a > c > b. \end{aligned}$$

Then S_s is a l.o. semigroup without the unit element. It is easy to verify that S_s does not satisfy the condition (A) and that one cannot adjoin the new element e to S_s , so that $S_s^* = S_s \cup e$ becomes a l.o. semigroup in which e is the unit element and S_s is order-embedded in S_s^* .

Example 9. Let S_9 be a set of two elements 0 and e with the relations

$$0^2 = 0, \quad e^2 = e, \quad 0e = e0 = 0, \quad 0 > e.$$

Then S_9 is a l.o. semigroup with the unit element e . Since $0 \geq 0^2$ and $0e = 0 > e$, S_9 does not satisfy the condition (A). (Or, see Example 7.)

Example 10. Let T be an abelian semigroup generated by two elements a and f with the relations

$$\begin{aligned} f^2 = f, \\ f > a > af > a^2 > \dots > a^n > a^n f > a^{n+1} > \dots \end{aligned}$$

Then T is a l.o. semigroup without the unit element and does not satisfy the condition (A) because $a > af$ in spite of $f^2 \geq f$. But one can adjoin the new element e to T , so that $T^* = T \cup e$ becomes a l.o. semigroup in which e is the unit element, by adding the following relations:

$$e^2 = e, \quad ex = xe = x, \quad e > x \quad \text{for all } x \text{ in } T.$$

Theorem 9. Let S be a p.o. semigroup which satisfies the condition (β). If S has the unit element e and all positive or negative elements are comparable to e , then S satisfies the condition (A).

Proof. Let a ($\neq e$) be positive. If $a^2 \geq a$ implies $a \succ e$, then we have $e > a$ by the hypothesis of the theorem, and hence $a \geq a^2$, therefore $a = a^2$, which contradicts the condition (β). Hence we have $a > e$, therefore $ab \geq b$ for any b of S . Similarly, if a ($\neq e$) is negative then $b \geq ab$ for any b of S .

Corollary 1. Let S be a p.o. semigroup with the unit element e which satisfies the condition (α). Then S satisfies the condition (β) and all positive or negative elements are comparable to e if and only if it satisfies the condition (A). (Cf. Corollary 1 of Theorem 7.)

Corollary 2. Let S be a l.o. semigroup with the unit element. Then S satisfies the condition (β) if and only if it satisfies the condition (A). (Cf. Corollary 2 of Theorem 7.)

Example 11. Let T_1 be an abelian semigroup generated by two

elements a and e with the relations

$$e^2 = e, \quad a^m e = e a^m = a^m \quad \text{for any positive integer } m,$$

$$a^{2n+1} > a^{2m+1} \quad (n > m \geq 0),$$

$$a^{2n} > a^{2m} > e \quad (n > m > 0).$$

Then T_1 is a p.o. semigroup with the unit element e and satisfies the conditions (α) and (β) , but is not normal. All positive elements (there exist no negative elements except e) are comparable to e , and clearly T_1 satisfies the condition (A).

Theorem 10. If a l.o. semigroup S satisfies the condition (β) , then S satisfies the properties 1), 2) and 3) of Theorem 6.

Proof. If S has the unit element e , then by Corollary 2 of Theorem 9 S satisfies the condition (A) and hence the properties 1), 2) and 3) by Theorem 6. Now we assume that S has not the unit element, that is, all elements of S are of infinite order. Let a be positive, b be negative, i.e., $a^2 > a$ and $b > b^2$. If $b > a$, then $T(b) > T(a)$ and hence $T(b) > ab > T(a)$ and ab is idempotent by Theorem 5, iii). This contradicts the hypothesis. Therefore we have $a > b$. Similarly, let $c > a$ ($b > c$), then c is positive (negative).

We note that a l.o. semigroup S , which does not satisfy the condition (β) , may satisfy the properties 1), 2) and 3); see Example 5.

Example 12. Let T_2 be an abelian semigroup generated by two elements a and b with the relation

$$a^m b^n = ab^n.$$

Then T_2 satisfies the conditions (α) and (β) . And by putting the order-relation

$$a^{m+1} > a^m > b^n > ab^n > b^{n+1} > ab^{n+1}$$

for any positive integers m and n ,

T_2 becomes a l.o. semigroup which does not satisfy the condition (A) because $b > ab$ in spite of $a^2 > a$,

Or, by putting the other-relation

$$b^{n+1} > ab^{n+1} > b^n > ab^n > a^{m+1} > a^m$$

for any positive integers m and n ,

T_2 becomes a l.o. semigroup which does not satisfy the condition (A) because $b > ab$ in spite of $a^2 > a$.

Theorem 11. In a l.o. semigroup the conditions (r) and (B) are equivalent to each other.

Proof. Let S be a l.o. semigroup which satisfies the condition (r) and let a (\neq the unit element) be a positive element of S . Then $a^2 > a$ and hence $a^2b = a(ab) > ab$ for any b of S . By the linearity, we have $ab > b$. Similarly, if a (\neq the unit element) is negative, then $b > ab$ for any b of S . The converse is obvious.

We note that if an abelian semigroup S satisfies the condition (r), then the element a of finite order is cyclic. For, let m be the length and r be the period of a , then $a^m = a^m a^r$ and hence a^r is the unit element, thereby $a = a \cdot a^r$, thus we have $m = 1$, i.e., a is cyclic.

Example 13. Let T_3 be an abelian semigroup generated by two elements a and b with the relations

$$a^m b^n = b^n, \quad a^{m+1} > a^m > b^n > b^{n+1}$$

for any positive integers m and n .

Then T_3 becomes a l.o. semigroup which satisfies the conditions (a), (b), (d) and (A) but does not the conditions (r) and (B).

Theorem 12. A l.o. semigroup S is strong if and only if S satisfies the conditions (a), (d) and (B).

Proof. Necessity: For the condition (a), see Theorem 10, O.I. Let $c \cdot T(a)$ and $c \cdot T(b)$ be intersect for some c of S , then there exist two elements $x \in T(a)$ and $y \in T(b)$ such that $cx = cy$. By the strongness, we have $x = y$. Therefore $T(a) = T(b)$. Since in the strong l.o. semigroup the element of finite order is the unit element (Theorems 9 and 10, O.I.), if a (\neq the unit element e) is positive, then we have $a^2 > a$. Hence by the strongness we have $a^2b > ab$, thereby $ab > b$ for any b of S . Similarly, if a ($\neq e$) is negative, then we have $b > ab$ for any b of S .

Sufficiency: Let $xz = yz$. i) $T(x) = T(y) = T(z)$. By Corollary 1 of Theorem 2, we have $x = y$. ii) $T(x) = T(y) \neq T(z)$. Since S satisfies the condition (B), S satisfies the condition (b). If x is of finite order, that is, x is the unit element, then $T(x) = T(y) = x$. Suppose that x is of infinite order. Then there exist positive integers n and m such that $x^n = y^m$. And hence $y^n z^n = x^n z^n = y^m z^n$. If $n \neq m$, say that $n > m$, then we have $y^m z^n = y^{n-m} (y^m z^n)$, which contradicts the condition (B), since y is not of finite order, that is, is not the unit element. Hence $n = m$, i.e., $x^n = y^n$, therefore we have $x = y$ by the condition (a). iii) $T(x) \neq T(y)$. By the condition (d), $xz \neq yz$ for all z of S . Therefore, the product

cancellation law is held in S . By Theorem 4, O.I., S is strong.

Corollary 1. A l.o. semigroup S is strong if and only if S satisfies the conditions (α) , (γ) and (δ) .

Corollary 2. A strong linear order may be defined on an abelian semigroup S if and only if S satisfies the conditions (α) , (γ) and (δ) .

Proof. Let x be an element of finite order, then by the condition (α) x is idempotent, furthermore by the condition (γ) x is the unit element. That is, S satisfies the condition (β) . Similarly to the proof of Theorem 12, we see that the product cancellation law is held in S . Therefore the proof is complete by Theorem 15, O.I.

Example 14. Let T_4' be a free abelian semigroup generated by one element a and T_4'' be an abelian semigroup generated by two elements 0 and e with the relations

$$0^2 = 0, \quad e^2 = e, \quad 0e = e0 = 0.$$

And let T_4 be a direct product of T_4' and T_4'' , then T_4 is an abelian semigroup which satisfies the conditions (α) , (β) and (γ) but does not the condition (δ) .

By putting the order-relation

$$a^n 0 < a^n e < a^{n+1} 0 < a^{n+1} e \quad \text{for any positive integer } n,$$

T_4 becomes a l.o. semigroup.

Examples 3, 13 and 14 show that the conditions (α) , (γ) and (δ) are independent in l.o. semigroups.

Theorem 13. A l.o. semigroup S with the unit element e is strong if and only if S satisfies the conditions (α) and (δ) .

Proof. Necessity: It is clear.

Sufficiency: Suppose that $ab = b$ for some b of S , that is, $ab = eb$. By the condition (δ) , $T(a) = T(e)$. Since $T(e) = e$ by the condition (α) , we have $a = e$, consequently S satisfies the condition (γ) .

Corollary 1. A strong linear order may be defined on an abelian semigroup S with the unit element if and only if S satisfies the conditions (α) and (δ) .

Corollary 2. The product cancellation law is held in an abelian semigroup with the unit element, when it has the following properties:

- 1) $a^n = b^n$ for some positive integer n implies $a = b$.
- 2) $ac = bc$ implies $a^m = b^m$ for some positive integers m and n .