



Title	POSITIVE LINEAR OPERATORS IN SEMI.ORDERED LINEAR SPACES
Author(s)	Andô, Tsuyoshi
Citation	Journal of the Faculty of Science Hokkaido University. Ser. 1 Mathematics, 13(3-4), 214-228
Issue Date	1957
Doc URL	<a href="http://hdl.handle.net/2115/55997">http://hdl.handle.net/2115/55997</a>
Type	bulletin (article)
File Information	JFSHIU_13_N3-4_214-228.pdf



[Instructions for use](#)

# POSITIVE LINEAR OPERATORS IN SEMI-ORDERED LINEAR SPACES

By

Tsuyoshi ANDÔ

Since in 1907 O. PERRON [9] discovered a remarkable spectral property of positive matrices and shortly later G. FROBENIUS [1], [2] and R. JENTZSCH [4] investigated and generalized it further, many authors have considered special properties of positive linear operators. Especially M. KREIN and M. A. RUTMAN [5]<sup>\*)</sup> considered with success a generalization to Banach spaces with a cone. They obtained particularly important results, when the space is lattice ordered or the cone has an interior point. In this paper, we consider spectral properties of positive compact (=completely continuous) linear operators on a universally continuous Banach space (= conditionally complete Banach lattice). Our main aim is to generalize the results of G. FROBENIUS [2] to infinite dimensional spaces.

In §1 preliminary definitions are summarized. In §2 the fundamental theorem on the maximum positive spectrum is proved (Theorem 2.1). In §3 we define *completely positive* linear operators. The operators of this class play a similar rôle as strongly positive operators in [5]. In §4 we obtain under some additional conditions a necessary and sufficient condition for that a positive compact linear operator is quasinilpotent (Theorem 4.7). In §5 the proper values with maximum modulus of a positive compact linear operator are determined (Theorem 5.2).

§ 1. Preliminaries. We recall briefly definitions from the theory of semi-ordered linear spaces and linear operators. A lattice ordered linear space (with real scalar)  $R$  is said to be *universally continuous*, if for any  $a_\lambda \geq 0$  ( $\lambda \in A$ ) there exists  $\bigcap_{\lambda \in A} a_\lambda$ . A linear manifold  $N$  of  $R$  is said to be *normal* if there exists a positive linear projection  $[N]$  of  $R$  onto  $N$  such that  $|x - [N]x| \wedge |y| = 0$  for  $x \in R$  and  $y \in N$ . The projection

---

<sup>\*)</sup> The author of the present paper expresses his thanks to Dr. S. YAMAMURO at Institute for Advanced Study, Princeton, who kindly communicated the main results of [5], which was not available to him.

onto the normal manifold generated by  $a$  is denoted by  $[a]$ . Normal manifolds and order-projections correspond to each other in one-to-one way.

When we consider spectral problems, it is convenient to define a *complex extension*  $\hat{R}$  of  $R$ , whose elements consist of all pair of elements of  $R$ ,  $(a, b) \equiv a + ib$ , the *absolute value* of  $a + ib$  is defined by

$$(1.1) \quad |a + ib| = \bigcup_{0 < \theta < 2\pi} |a \cos \theta + b \sin \theta|.$$

When  $R$  is normed, the norm, in this paper, satisfies the following additional condition:

$$(1.2) \quad |a| \leq |b| \quad \text{implies} \quad \|a\| \leq \|b\|,$$

The norm on  $\hat{R}$  is defined, when  $R$  is normed, by

$$(1.3) \quad \|a + ib\| = \||a + ib|\|.$$

A norm on  $R$  is said to be *continuous* if  $a_\lambda \downarrow_{\lambda \in A} 0$  implies  $\|a_\lambda\| \downarrow_{\lambda \in A} 0$ . A bounded linear functional  $\tilde{a}$  is said to be *universally continuous*, if  $a_\lambda \downarrow_{\lambda \in A} 0$  implies  $\inf_{\lambda \in A} |\tilde{a}(a_\lambda)| = 0$ .  $\bar{R}$  and  $\bar{R}$  denote the space of all bounded linear functionals on  $R$  and that of all universally continuous linear functionals respectively. For the other notations and definitions, we refer to [6].

For a bounded linear operator  $A$  on a complex Banach space  $R$  into itself,  $\sigma(A)$  denotes the set of all spectra of  $A$ , and  $\rho(A)$  the *resolvent set*. If  $(\lambda I - A)x = 0$  has a non-trivial solution in  $R$ ,  $\lambda$  is said to be a *proper value* and its solutions are *proper elements*. We put

$$(1.4) \quad r(A) = \sup_{\xi \in \sigma(A)} |\xi|$$

$$(1.5) \quad R(\lambda) = (\lambda I - A)^{-1} \quad \text{for } \lambda \in \rho(A).$$

It is well known (cf. [3]) that

$$(1.6) \quad r(A) = \lim_{\nu \rightarrow \infty} \|A^\nu\|^{\frac{1}{\nu}}$$

$A$  is said to be *quasi-nilpotent*, if  $r(A) = 0$ .

A bounded linear operator  $A$  is said to be *compact* if the unit sphere is mapped by  $A$  into a compact set. We assume in this paper the results of F. RIESZ ([10], chap. IV and V) concerning the spectral properties of compact linear operators. If  $A$  is a compact linear operator, every non-zero spectrum is a proper value and the corresponding proper manifold is each finite dimensional.  $\sigma(A)$  constitutes a totally discon-

nected set with the only possible limiting point 0. For a non-zero complex number  $\lambda$  we can define a bounded linear projection operator  $E(\lambda)$  relative to  $A$  by

$$(1.6) \quad E(\lambda) = \frac{1}{2\pi i} \int R(\zeta) d\zeta$$

where the integration is formed along a Jordan curve surrounding  $\lambda$ , whose boundary and interior intersect  $\sigma(A)$  in  $\lambda$  alone. The *index*  $\mu(\lambda)$  of  $\lambda$  is the smallest integer  $n$  satisfying  $(\lambda I - A)^n E(\lambda) = 0$ . For other definitions in operator theories, we refer to [3].

Examples of universally continuous Banach spaces are:  $L_p$ ,  $l_p$  ( $1 \leq p \leq \infty$ ), and more generally *modulated spaces* studied in [6].

As we develop in the following a theory of positive compact linear operators in a universally continuous Banach space, it has immediate applications to the theory of integral equations or linear equations with infinite unknowns in the spaces mentioned above.

**§ 2. The maximum positive spectrum.** Throughout the paper  $R$  denotes a universally continuous semi-ordered Banach space and  $A$  a linear operator on  $R$  into itself, if the contrary is not mentioned.

Though some of results are known under weaker conditions (see [5]), we prove them for the sake of completeness, since under our conditions proofs are sometimes simple.

$A$  is said to be *positive*, if  $a \geq 0$  implies  $Aa \geq 0$ .

*Lemma 2.1.* A positive linear operator is necessarily bounded.

*Proof.* For  $\tilde{a} \in \bar{R}$ , putting  $\tilde{b}(x) = \tilde{a}(Ax)$ ,  $\tilde{b}$  is an (o)-bounded linear functional on  $R$ , so by Theorem 31.3 in [6], is norm-bounded. This means that the image of the unit sphere by  $A$  is weakly bounded. The assertion follows from the known theorem on weakly bounded sets.

*Theorem 2.1.* If  $A$  is positive,  $r(A)$  is in  $\sigma(A)$ .

*Proof.* Suppose  $r = r(A)$  is not in  $\sigma(A)$ . Then  $R(\lambda) = \sum_{\nu=0}^{\infty} \frac{A^\nu}{\lambda^{\nu+1}}$  for  $\lambda > r$  and  $\sup_{r < \lambda} \|R(\lambda)\| < \infty$ . Since, considering  $\hat{R}$ , by (1.3)

$$\|R(\lambda e^{i\theta})x\| \leq \|R(\lambda)x\| \quad \text{for } \lambda > r, 0 \leq \theta \leq 2\pi \text{ and } x > 0,$$

hence  $\sup_{\lambda > r} \|R(\lambda e^{i\theta})\| < \infty$ , this implies  $re^{i\theta} \notin \sigma(A)$ ,  $0 \leq \theta \leq 2\pi$ , contradicting the assumption (cf. [3]).

*Corollary 2.1.1.* If  $A$  is positive,  $R(\lambda) \geq 0$  if and only if  $\lambda > r$ .

*Proof.* If  $R(\lambda) \geq 0$ ,  $\lambda$  is apparently real and by the resolvent equation

([3], p. 99)  $R(\lambda) - R(\mu) = \frac{R(\lambda)R(\mu)}{\mu - \lambda} \geq 0$  for  $\mu > \text{Max}(\lambda, r)$ . Hence as in Theorem 2.1  $\lambda > r(A)$ . The converse part is obvious.

*Corollary 2.1.2.* Let  $A$  be positive. If for a  $\lambda > 0$ , there exists  $x \neq 0$ , such that  $\lambda x \geq Ax$ , then  $\lambda \leq r(A)$ .

*Proof.* If  $\lambda \in \sigma(A)$ ,  $\lambda \leq r(A)$  by Theorem 2.1. If  $\lambda \in \rho(A)$ , from the hypothesis  $R(\lambda)$  is not positive, the assertion follows from Corollary 2.1.1.

Concerning the indice on the circle of radius  $r = r(A)$ , we obtain

*Theorem 2.2.* If  $A$  is positive compact with  $r = r(A) > 0$ ,

$$(2.1) \quad \mu(\lambda) \leq \mu(r) \quad \text{for } |\lambda| = r$$

*Proof.* By the Laurent resolution ([3], p. 109)  $\sup_{0 < \epsilon < \delta} \epsilon^{\mu(r)} \|R(r + \epsilon)\| < \infty$  for a small  $\delta$ . Since from (1.3)  $\epsilon^{\mu(r)} \|R(re^{i\theta} + \epsilon e^{i\theta})\| \leq \epsilon^{\mu(r)} \|R(r + \epsilon)\|$ , we obtain  $\mu(\lambda) \leq \mu(r)$ .

*Theorem 2.3.* If  $A$  is positive compact  $r = r(A) > 0$ ,  $A$  has a positive proper element corresponding to the proper value  $r$ .

*Proof.* We know that  $\lim_{\epsilon \rightarrow 0} \epsilon^{\mu(r)} R(r + \epsilon) = (A - rI)^{\mu(r)-1} E(r)$  ([3], p. 109). Since  $R(r + \epsilon) \geq 0$ ,  $(A - rI)^{\mu(r)-1} E(r) \geq 0$ , there exists  $x > 0$  with  $y = (A - rI)^{\mu(r)-1} E(r)x > 0$ . We obtain that  $Ay = ry$  and  $y > 0$ .

**§ 3. Completely positive linear operators.** A linear operator  $A$  is said to be *universally continuous*, if  $a_\lambda \downarrow_{\lambda \in A} 0$  implies  $\bigcap_{\lambda \in A} |Aa_\lambda| = 0$ .

*Lemma 3.1.* If  $A$  is positive, compact and universally continuous, the range of the conjugate operator  $A^*$  is contained in  $\bar{R}$ .

*Proof.* Since  $A$  is positive and compact, for any  $a_\lambda \downarrow_{\lambda \in A} 0$   $\{Aa_\lambda\}_{\lambda \in A}$  has a limiting point and it must be equal to 0. So  $\lim_{\lambda} Aa_\lambda = 0$ . This implies that  $A^* \bar{a}$  is universally continuous for every  $\bar{a} \in \bar{R}$ .

An element  $a$  of  $R$  is said to be *complete*, if  $[a] = I$ . A bounded linear functional  $\bar{a}$  is said to be *complete* if  $|\bar{a}|(|a|) = 0$  implies  $\bar{a} = 0$ . We remark that if  $a$  is complete,  $\bar{a} \in \bar{R}$   $|\bar{a}|(|a|) = 0$  implies  $\bar{a} = 0$ .

*Theorem 3.1.* Let  $A$  be universally continuous, positive and compact with  $r = r(A) > 0$ . If  $A$  has a positive complete proper element, then  $\mu(r) = 1$  and the proper element corresponds to  $r$ .

*Proof.* Let  $Aa = \lambda a > 0$  and  $[a] = I$ . If  $r \neq \lambda$ ,  $\bar{a}(a) = 0$  for the positive proper element of  $A^*$  corresponding to  $r$ , which exists by Theorem 2.3 and is universally continuous by Lemma 3.1. This implies  $\bar{a} = 0$ . So

$\lambda$  must be equal to  $r$ . Suppose that  $\mu(r) \geq 2$ . There exists a positive linear functional  $\tilde{b} = (A^* - rI)^{\mu(r)-1} E^*(r) \tilde{x}$  as in Theorem 2.3. But by Lemma 3.1  $\tilde{b}(a) = 0$  and  $\tilde{b} \in \bar{R}$ , this implies  $\tilde{b} = 0$ , contradicting the assumption.

The special property of the proper manifold corresponding to is contained in:

*Theorem 3.2.* *If  $A$  is positive compact and  $A^*$  has a positive complete proper element  $\tilde{a}$ , the proper manifold corresponding to  $r$  of  $A$  is a linear lattice manifold.*

*Proof.*  $\tilde{a}$  corresponds to  $r$  by Theorem 3.1. If  $Aa = ra$ ,  $Aa^+ \geq ra^+$ , so we obtain  $\tilde{a}(Aa^+ - ra^+) = 0$ . Since  $\tilde{a}$  is complete, this implies that  $Aa = ra$ . Hence the proper manifold corresponding to  $r$  is a linear lattice manifold.

*Corollary 3.2.* *Under the same assumption as Theorem 3.2, if  $a \in \hat{R}$  is a proper element corresponding to  $\xi$  with  $|\xi| = r$ ,  $A|a| = r|a|$ .*

*Proof.* By the definition (1.1),  $Aa = \xi a$  we have  $A \cdot |a| \geq r|a|$ . The assertion follows as above.

As a special class of positive linear operators, we define: a positive universally continuous linear operator  $A$  is said to be *completely positive* if

$$(3.1) \quad \bigcup_{\nu=1}^{\infty} [A^\nu x] = I \quad \text{for every } x > 0$$

This means that for a positive  $x > 0$ ,  $a \wedge A^\nu x = 0$  ( $\nu = 1, 2, \dots$ ) implies  $a = 0$ .

*Lemma 3.2.* *If  $A$  is positive and universally continuous, for  $a > 0$ , putting  $\bigcup_{\nu=1}^{\infty} [A^\nu a] = [N]$ , we obtain an invariant normal manifold, that is,*

$$(3.2) \quad A[N] = [N]A[N]$$

*Proof.* Since for  $x > 0$ ,  $[N]x = \bigcup_{\nu=1}^{\infty} (x \wedge_\nu (Aa + \dots + A^\nu a))$  and  $A$  is universally continuous,  $A[N]x = \bigcup_{\nu=1}^{\infty} A(x \wedge_\nu (Aa + \dots + A^\nu a))$  so we obtaine  $[A[N]x] \leq \bigcup_{\nu=1}^{\infty} [Aa + \dots + A^\nu a] \leq [N]$ .

Complete positiveness corresponds to "Unzerlegbarkeit" in [2], as is seen in the following:

*Theorem 3.3.* *A positive universally continuous linear operator is completely positive if and only if it has no non-trivial invariant normal manifold.*

This is an immediate consequence of Lemma 3.2.

**Lemma 3.3.** *If  $A$  is compact and completely positive, every positive proper element of  $A$  (and  $A^*$ ) is complete.*

*Proof.* If  $a$  is a proper element of  $A$ ,  $[a] = [A^\nu a]$  ( $\nu = 1, 2, \dots$ ) and so  $[a] = I$ . If  $\bar{a}$  is a proper element of  $A^*$ ,  $\bar{a}(a) = 0$  implies  $\bar{a}(A^\nu a) = 0$  ( $\nu = 1, 2, \dots$ ). Since  $\bar{a}$  is in  $\bar{R}$  by Lemma 3.1,  $a = 0$ .

**Theorem 3.4.** *If  $A$  is compact and completely positive with  $r = r(A) > 0$ , the multiplicity of the proper value  $r$  is equal to 1 (cf. §5 later).*

*Proof.* By Theorem 3.2 and Lemma 3.3, the proper space corresponding to  $r$  is a linear lattice manifold. Since  $A$  is completely positive, all positive proper elements are complete, so the multiplicity must be equal to 1.

Next we consider the distribution of proper values corresponding to positive proper elements.

For a bounded linear operator  $A$  on a complex Banach space  $R$  into itself, the *spectral radius* of  $x$ ,  $r(x, A)$  or  $r(x)$  (if there is no confusion), is defined by

$$(3.2) \quad r(x, A) \equiv r(x) = \overline{\lim}_{\nu \rightarrow \infty} \|A^\nu x\|^{\frac{1}{\nu}}$$

This functional satisfies the following properties:

- 1)  $0 \leq r(x) \leq r(A)$
- 2)  $\sup_{x \in R} r(x) = r(A)$
- 3)  $r(\alpha x) = r(x)$  for  $\alpha \neq 0$
- 4)  $r(x+y) \leq \text{Max} \{r(x), r(y)\}$
- 5)  $r(Ax) = r(x)$

We remark that  $r(x)$  is nothing but the maximum modulus of singularities of analytic continuation of  $R(\lambda)x$  ( $\lambda \in \rho(A)$ ).

Since the set of all spectra of a compact operator is a totally disconnected set with the only possible limiting point 0, the functional  $r(x)$  is rather convenient.

**Lemma 3.4.** *If  $A$  is compact, the functional  $r(x)$  satisfies the following:*

- a)  $r(x) = \text{Max} \{|\lambda| : E(\lambda)x \neq 0\}$  for  $x \neq 0$
- b)  $r(x)$  is lower semi-continuous,
- c) the range of  $r(x)$  coincides with the set  $\{|\lambda|; \lambda \in \sigma(A)\}$ .

*Proof.* a) is an immediate consequence of remarks stated above. b) follows from a). c) is evident, since every non-zero spectrum is a proper value.

For a positive compact linear operator, we obtain:

*Theorem 3.5.* *If  $A$  is positive, universally continuous and compact, the set  $\{r(x); r(x) > 0, x > 0\}$  coincides with the set of all non-zero proper values corresponding to positive proper elements.*

*Proof.* For  $\lambda > 0$ , the set  $S_\lambda = \{x; r(|x|) \leq \lambda\}$  is a closed linear manifold by Lemma 3.4. If  $0 \leq x_\rho \uparrow_\rho x (x_\rho \in S_\lambda), x \in S_\lambda$ , because  $E(\xi)$  is universally continuous, hence  $S_\lambda$  is normal. If there exists  $x \in S_\lambda$  such that  $r(|x|) = \lambda$ , by formula (5)  $A[S_\lambda] = [S_\lambda]A[S_\lambda]$  and  $r(A[S_\lambda]) = \lambda$ . Hence by Theorem 2.1 there exists  $0 \leq a_\lambda \in S_\lambda$  such that  $Aa_\lambda = \lambda a_\lambda$ . Conversely if  $Aa = \rho a > 0, r(a) = \rho$ .

*Theorem 3.6.* *Let  $A$  be positive, universally continuous and compact, with  $r = r(A) > 0, A$  is completely positive if and only if*

- a)  *$A$  has a unique positive proper element (up to scalar) which is complete,*
- b)  $r(x) > 0$  *for every  $x > 0$ .*

*Proof.* If  $A$  is completely positive, then by Theorems 3.2 and 3.4 the positive proper element is unique and complete. Since  $A^*$  has a positive complete proper element  $\tilde{a}$  corresponding to  $r$ , for any positive  $a > 0, r(a) \geq \lim_{\nu \rightarrow \infty} |\tilde{a}(A^\nu a)|^{\frac{1}{\nu}} = r$ . Conversely suppose that  $A$  satisfies a) and b). If a normal manifold  $N$  is invariant relative to  $A$ , by b)  $r(A[N]) > 0$  and by Theorem 2.3 there exists a positive proper element in  $N$  which is not complete.

We may replace the condition b) by the condition

- b')  *$A^*$  has a complete positive proper element.*

*Lemma 3.5.* *For a compact linear operator  $A$  on a Banach space  $R$*

$$\sup_{|\lambda|=r} \mu(\lambda) \leq 1 \text{ if and only if } \|A^\nu\| \leq M \cdot r^\nu \quad (\nu = 1, 2, \dots)$$

*for some  $M$ , where  $r = r(A)$ . And in this case*

$$(3.3) \quad E(r) = \lim_{\nu \rightarrow \infty} \frac{1}{\nu} \sum_{k=1}^{\nu} \left(\frac{A}{r}\right)^k$$

The proof is well known (cf. [10] and [5]).

If the maximum spectrum is not simple, the following decomposition holds:

*Theorem 3.7.* Let  $A$  be positive, universally continuous and compact. If  $A$  and  $A^*$  both have positive complete proper elements, there exists a decomposition of identity such that

$$(3.4) \quad \begin{aligned} I &= \sum_{\nu=1}^n [a_\nu], & [a_\nu][a_\mu] &= 0 & (\nu \neq \mu) \\ A[a_\nu] &= [a_\nu]A & & & (\nu=1, 2, \dots, n) \end{aligned}$$

and  $A$  is completely positive on  $[a_\nu]R$  ( $\nu=1, 2, \dots, n$ ).

*Proof.* By Lemma 3.5 and Theorem 3.2 the positive projection  $E(r)$  is written in a form  $E(r)x = \sum_{\nu=1}^n \tilde{a}_\nu(x)a_\nu$  such that

$$Aa_\nu = ra_\nu > 0, \quad A^*\tilde{a}_\nu = r\tilde{a}_\nu > 0, \quad \tilde{a}_\nu(a_\mu) = \delta_{\nu\mu} \quad (\nu, \mu=1, 2, \dots, n).$$

Since  $\bigcup_{\nu=1}^n [a_\nu] = \bigcup_{\nu=1}^n [\tilde{a}_\nu]^R = I$  and  $A[a_\nu] = [a_\nu]A[a_\nu]$ ,  $A^*[\tilde{a}_\nu] = [\tilde{a}_\nu]A^*[\tilde{a}_\nu]$  ( $\nu=1, 2, \dots, n$ ),  $A$  is completely positive on  $[a_\nu]R$  ( $\nu=1, 2, \dots, n$ ).

We define a somewhat weaker condition than complete positiveness: a positive linear operator  $A$  is said to be *naturally decomposable*, if there exists a decomposition of identity  $[N_\rho]$  ( $\rho \in \Lambda$ ) and  $[M]$  such that

$$(3.5) \quad \begin{aligned} I &= \sum_{\rho \in \Lambda} [N_\rho] + [M], & [N_{\rho_1}][N_{\rho_2}] &= 0 & (\rho_1 \neq \rho_2), & [N_\rho][M] &= 0 \\ A[N_\rho] &= [N_\rho]A & \text{and} & & A[M] &= [M]A, \\ r(A[N_\rho]) &> 0 & (\rho \in \Lambda) & & \text{and} & r(A[M]) &= 0, \end{aligned}$$

and  $A$  acts as a completely positive operator on  $[N_\rho]R$  ( $\rho \in \Lambda$ ).

*Theorem 3.8* Let  $A$  be positive, universally continuous and compact.  $A$  is naturally decomposable if and only if

$$\bigcup [a] = \bigcup [\tilde{a}]^R$$

where  $a$  (and  $\tilde{a}$ ) varies in all positive proper elements of  $A$  (and of  $A^*$  respectively).

*Proof.* If  $A$  is naturally decomposable with the decomposition (3.5), then by Lemma 3.3.  $\bigcup [a] = \bigcup_{\rho \in \Lambda} [\tilde{a}]^R = \bigcup [N_\rho]$ . Conversely, let  $\bigcup [a] = \bigcup [\tilde{a}]^R$ . We arrange the proper values of  $A$  corresponding to positive proper elements in the descending order  $\lambda_1 > \lambda_2 > \dots$ . Since each corresponding proper manifold is finite dimensional, there exist maximum projectors  $[p_\nu]$  corresponding to  $\lambda_\nu$  ( $\nu=1, 2, \dots$ ). Analogously we obtain  $\tilde{q}_\nu$  relative to  $A^*$  corresponding to  $\mu_1 > \mu_2 > \dots$ . By hypothesis  $\bigcup_{\nu=1}^\infty [p_\nu] = \bigcup_{\nu=1}^\infty [\tilde{q}_\nu]^R$  and  $\tilde{q}_k(p_\nu) = 0$  if  $\lambda_\nu \neq \mu_k$ . We obtain  $[p_\nu] = [\tilde{q}_\nu]^R$  ( $\nu=1, 2, \dots$ ) and  $[p_\nu]A = A[p_\nu]$ .

Since  $(1 - \bigcup_{\nu=1}^{\infty} [p_{\nu}])A$  is quassi-nilpotent by Theorem 2.3, we obtain the assertion by Theorem 3.7.

Next we consider a characterization of natural decomposability analogous to Theorem 3.6. For a projection  $[N]$ , we put

$$r([N]) = \sup_{[N]x=x} r(x) \quad \text{and} \quad r^*([N]) = \sup_{\bar{x}[N]=\bar{x}} r^*(\bar{x})$$

where

$$r^*(\bar{x}) = \overline{\lim}_{\nu \rightarrow \infty} \|A^{*\nu} \bar{x}\|^{1/\nu}$$

*Theorem 3.9.* Let  $A$  be positive, universally continuous and compact, and  $R$  semi-regular.  $A$  is naturally decomposable, if and only if it satisfies

- a)  $r([N]) = r^*([N])$  for every projection  $[N]$ ,
- b)  $\sup_{\nu=1,2,\dots} \frac{\|A^{\nu} x\|}{r(x)^{\nu}} < \infty$  for  $x > 0$  with  $r(x) > 0$ .

*Proof.* If  $A$  is naturally decomposable with the decomposition (3.5), it is easy to see that for a projection  $[N]$  and  $x > 0$

$$r([N]) = \sup_{[N][N_{\rho}] \neq 0} r(A[N_{\rho}]) = r^*([N])$$

$$r(x) = \sup_{[N_{\rho}]x \neq 0} r(A[N_{\rho}])$$

b) follows from Lemma 3.5. Conversely, suppose that a) and b) are satisfied. Put  $S_{\lambda} = \{x; r(|x|) \leq \lambda\}$  and  $\bar{S}_{\lambda} = \{\bar{a}; \bar{a} \in \bar{R}, r^*(|\bar{a}|) \leq \lambda\}$ . By Lemma 3.4 and a) we have  $[S_{\lambda}] = [\bar{S}_{\lambda}]^R$ , hence  $A[S_{\lambda}] = [S_{\lambda}]A$ . If  $a$  and  $b$  are positive proper elements corresponding to different positive proper values  $\lambda_1$  and  $\lambda_2$  respectively. If  $[p] = [a][b] \neq 0$ , as in the proof of Theorem 3.6,

$$r([p]) \leq \text{Min} \{r([a]), r([b])\} = \text{Min} \{\lambda_1, \lambda_2\}$$

and

$$r^*([p]) \geq \text{Max} \{r^*([a]), r^*([b])\} = \text{Max} \{\lambda_1, \lambda_2\}$$

contradicting a). So  $a \wedge b = 0$ . Let  $\lambda_1 > \lambda_2 > \dots$  be proper values of  $A$  corresponding to positive proper elements in the descending order. Putting  $[S_{\nu}] = [S_{\lambda_{\nu}}] - [S_{\lambda_{\nu+1}}]$ , we have  $A[S_{\nu}] = [S_{\nu}]A$  ( $\nu = 1, 2, 3, \dots$ ). By Lemma 3.5 and b) we can prove that  $E(\lambda_{\nu})x > 0$  and  $r(x) = \lambda_{\nu}$  for  $0 < x \in [S_{\nu}]R$ . Hence there exists  $0 < \bar{a} \in \bar{R}$  such that

$$A^* \bar{a} = \lambda_{\nu} \bar{a} \quad \text{and} \quad [\bar{a}]^R = [S_{\nu}]$$

Again using a), we obtain that there exists  $0 < a \in R$  such that  $Aa = \lambda_{\nu} a$

and  $[a]=[S_\nu]$ . Now Theorem 3.7 is applicable.

§ 4. **Quasi-nilpotent operators.** In this § we consider relations between  $A$  and its restriction to normal manifolds.

*Lemma 4.1.* Let  $A$  be a compact linear operator on a Banach space  $R$ . If  $F$  is a bounded linear operator such that  $F^2=F$  and  $AF=FAF$ , then

$$(4.1) \quad \sigma(A) = \sigma(AF) \cup \sigma((1-F)A(1-F)),$$

$$(4.2) \quad r(A) = \text{Max} \{r(AF), r((1-F)A(1-F))\}$$

*Proof.* Let  $0 \neq \lambda \in \rho(AF) \cup \rho((1-F)A(1-F))$ . If  $(\lambda I - A)x = 0$ ,  $(1-F)(\lambda I - A)x = (1-F)(\lambda - (1-F)A(1-F))x = 0$  hence  $(1-F)x = 0$  and similarly  $Fx = 0$ , so,  $x = 0$ , hence  $\lambda \in \rho(A)$ . Conversely, if  $\lambda \in \rho(A)$ ,  $(\lambda I - A)F$  is one-to-one on  $FR$ , hence by RIESZ's theorem  $\lambda \in \rho(AF)$ . Similarly  $\lambda \in \rho(A^*(1-F^*)) = \rho((1-F)A(1-F))$ .

For positive linear operators  $A$  and  $B$  such that  $A \geq B$ , it is evident that  $r(A) \geq r(B)$ . In particular, for any projection  $[N]$ ,  $r([N]A[N]) \leq r(A)$ .

*Theorem 4.1.* Let  $A$  be positive, universally continuous and compact with  $r=r(A) > 0$ .  $A$  is completely positive, if and only if for a projection  $[N]$   $r([N]A[N]) = r(A)$  implies  $[N] = I$ .

*Proof.* Suppose first that  $A$  is completely positive. If  $r([N]A[N]) = r$ , by Theorem 2.3 there exists  $a > 0$  such that  $[N]A[N]a = ra$ . Since  $Aa \geq ra$ , as in the proof of Theorem 3.2, we obtain  $Aa = ra$ . Complete positiveness implies  $[N] = [a] = I$ . Next suppose that  $A$  is not completely positive. There exists by Theorem 3.3 a non-trivial normal manifold  $[N]$  such that  $A[N] = [N]A[N]$ . Lemma 4.1 shows that  $\text{Max} \{r(A[N]), r((I-[N])A(I-[N]))\} = r$ .

Between a positive proper value distinct from  $r=r(A)$  and  $r([N]A[N])$  the following relation holds:

*Theorem 4.2.* Let  $A$  be positive and compact, and  $\lambda$  a positive proper value distinct from  $r=r(A)$ . For any non-zero projection  $[N]$  there exists a non-zero projection  $[M]$  such that  $[N] \geq [M]$  and  $r((I-[M])A(1-[M])) \geq \lambda$ .

*Proof.* There exists  $a \neq 0$  such that  $Aa = \lambda a$ , so  $Aa^+ \geq \lambda a^+$  and  $Aa^- \geq \lambda a^-$ . By Corollary 2.1.2,  $r([a^+]A[a^+]) \geq \lambda$  (and  $r([a^-]A[a^-]) \geq \lambda$ ) if  $[a^+] \neq 0$  (and  $[a^-] \neq 0$ ). If  $[N][a] = 0$ , we put  $[M] = [N]$ . If  $[N][a] \neq 0$  and  $a^- = 0$ , we have  $Aa = \lambda a > 0$  and  $r([a]A[a]) = \lambda$ . Since by Lemma 4.1  $\text{Max} \{r(A[a]), r((I-[a])A(1-[a]))\} = r$ , we put  $[M] = [N][a]$ . If  $[N][a^+] \neq 0$  and  $[a^-] \neq 0$ , we put  $[M] = [N][a^+]$ . The other case is treated similarly.

*Corollary 4.2.* Under the same conditions as in Theorem 4.2, if  $p$  is

a discrete element of  $R$ ,  $r((1-[p])A(1-[p])) \geq \lambda$ .

*Proof.* Since  $p$  is a discrete element, putting  $[N]=[p]$ ,  $[M]$  must coincide with  $[N]$ .

*Lemma 4.2.* If  $A_\lambda$  ( $\lambda \in \Lambda$ ), ( $\Lambda$  being a directed set), are compact linear operators defined on a Banach space, such that  $\lim_{\lambda} \|A_\lambda - A\| = 0$ , then

$$(4.3) \quad \lim_{\lambda} \sigma(A_\lambda) = \sigma(A) \quad \text{in the sense of metric,}$$

$$(4.4) \quad \lim_{\lambda} r(A_\lambda) = r(A).$$

The proof is found in [8].

*Lemma 4.3.* Let  $F_\lambda$  ( $\lambda \in \Lambda$ ), ( $\Lambda$  being a directed set), be bounded linear operators on a Banach space  $R$  such that  $F_\lambda^2 = F_\lambda$  ( $\lambda \in \Lambda$ ),  $F^2 = F \sup_{\lambda \in \Lambda} \|F_\lambda\| < \infty$  and  $\lim_{\lambda \in \Lambda} F_\lambda x = Fx$  ( $x \in R$ ). If  $A$  is a compact linear operator on  $R$ , then  $\lim_{\lambda \in \Lambda} r(F_\lambda A F_\lambda) = r(FAF)$ .

*Proof.* Since the image of the unit sphere by a compact linear operator is relatively compact, it is easy to see that  $\lim_{\lambda} \|F_\lambda A F_\lambda A - F A F A\| = 0$ . By Lemma 4.2,  $\lim_{\lambda} r(F_\lambda A F_\lambda A) = r(FAFA)$ . But since  $r(FAFA) = \lim_{\nu \rightarrow \infty} \|(FAFA)^\nu\|^{\frac{1}{\nu}} = \lim_{\nu \rightarrow \infty} \|(FAF)^{2\nu-1} A\|^{\frac{1}{\nu}} \leq r(FAF)^2$  and  $r(FAFA) = \lim_{\nu \rightarrow \infty} \|(FAFA)^\nu\|^{\frac{1}{\nu}} \geq \lim_{\nu \rightarrow \infty} \left\{ \frac{1}{\|F\|} \|(FAF)^{2\nu}\| \right\}^{\frac{1}{\nu}} = r(FAF)^2$ , we obtain  $\lim_{\lambda} r(F_\lambda A F_\lambda A) = r(FAF)^2$ . Hence  $\lim_{\lambda} r(F_\lambda A F_\lambda) = r(FAF)$ .

*Theorem 4.3.* Let  $R$  have no discrete element and be of continuous norm. If  $A$  is positive compact, there exist  $p_\lambda \in R$  ( $0 \leq \lambda \leq r(A)$ ), such that

$$(4.5) \quad [p_\lambda] \leq [p_\mu] \quad (0 \leq \lambda \leq \mu)$$

and 
$$r([p_\lambda]A[p_\lambda]) = \lambda \quad (0 \leq \lambda \leq r(A))$$

*Proof.* We choose, ZORN's lemma, a maximal linearly ordered family of projections  $[q_\rho]$ . The assumptions on  $R$  and Lemma 4.3 imply that  $\sup_{\xi < \rho} r([q_\xi]A[q_\xi]) = \inf_{\xi > \rho} r([q_\xi]A[q_\xi])$ . We can choose  $[p_\lambda]$  from  $[q_\rho]$  with  $r([q_\rho]A[q_\rho]) = \lambda$ , the remaining part is easily proved.

In the operator theory, it is important to study conditions assuring non-quasi-nilpotentness. Naturally a problem arises whether complete positiveness implies non-quasi-nilpotentness. We have been able to solve this problem only under some additional conditions.

A bounded linear operator  $A$  is said to be *totally continuous*, if for

any positive  $a \in R$  and positive  $\bar{a} \in \bar{R}$

$$(4.6) \quad [\bar{p}] \bar{x}(A[p]x) = \int_{[p] \times [\bar{p}]^R} \psi(\eta, \eta) \bar{a}(d\eta x) \bar{x}(d\eta a) \quad \text{for } [p] \leq [a] \text{ and } [\bar{p}] \leq [\bar{a}],$$

for a fixed Borel function  $\psi(\eta, \eta)$  on the product space  $\mathfrak{G} \times \mathfrak{G}$  of the proper space of  $R$  (see [7] §5). H. NAKANO [7] proved that  $A$  is totally continuous if and only if  $|a_\nu| \leq a$  ( $\nu = 1, 2, \dots$ )  $s\text{-}\lim_{\nu \rightarrow \infty} a_\nu = 0$  (star-convergence) implies  $(o)\text{-}\lim_{\nu \rightarrow \infty} Aa_\nu = 0$ . (It is easy to prove that here star-convergence may be replaced by weak-convergence).

*Lemma 4.4* For a bounded linear operator  $A$  on a Banach space, put  $B = \sum_{\nu=1}^{\infty} \frac{A^\nu}{\lambda^{\nu+1}}$  for some  $\lambda > r(A)$ . If  $B$  has a non-zero spectrum,  $A$  has one also. If  $A$  is positive and compact,  $B$  is so.

*Proof.* Since  $R(\lambda) = \frac{1}{\lambda} I + B$ , by the spectral mapping theorem (cf. [3] p. 122)  $\sigma(A) = \left\{ \lambda - \frac{1}{\frac{1}{\lambda} + \xi} ; \xi \in \sigma(B) \right\}$ , if  $\sigma(B)$  contains non-zero

number,  $\sigma(A)$  does also.

*Lemma 4.5.* Let  $R$  be reflexive as a Banach space. If  $A_\nu, A$  ( $\nu = 1, 2, \dots$ ) are positive compact such that

$$A_1 \leq A_2 \leq A_3 \leq \dots \quad \text{and} \quad \lim_{\nu \rightarrow \infty} A_\nu a = Aa \quad (a \in R),$$

then  $\lim_{\nu \rightarrow \infty} \|A_\nu - A\| = 0$

*Proof.* Considering  $A$  as a continuous function  $\bar{a}(Aa)$  on  $S \times \bar{S}$ , where  $S$  and  $\bar{S}$  are the positive unit spheres of  $R$  and  $\bar{R}$  respectively, topologized by weak topologies. Since  $S \times \bar{S}$  is compact, the assertion follows from the well-known theorem of Dini and the definition of norms of operators.

*Theorem 4.5.* Let  $R$  be reflexive as a Banach space. If  $A$  is compact, totally continuous and completely positive, then  $A$  is not quasi-nilpotent.

*Proof.* By Lemma 4.4 considering the compact positive operator  $B$ , we may assume that  $[Ax] = I$  for every  $x > 0$ . Further we may assume that for some  $a > 0$  and  $\bar{a} > 0$   $[a] = [\bar{a}]^R = I$  and  $\bar{a}(a) = 1$ . Suppose that  $A$  is represented in a form (4.6).  $A^2$  satisfies the same conditions as  $A$  and the corresponding function may be given by

$$\psi(\eta, \eta) = \int \psi(\eta, \xi) \psi(\xi, \eta) \bar{a}(d\xi a)$$

If  $\Psi(q, p)$  vanishes on a set of positive measure, from the measure theory, there exists a measurable subset  $\mathfrak{N} \subset \mathfrak{G}$  such that

a) the measure of  $\mathfrak{N}$  is positive, (the measure being defined by  $\bar{a}([p]a)$ ).

b)  $\text{meas}(\mathfrak{B}_p) > 0$  for  $p \in \mathfrak{N}$ , where  

$$\mathfrak{B}_p = \{q; \int \varphi(q, \xi) \varphi(\xi, p) \bar{a}(d\xi a) = 0\}$$

c)  $\varphi(q, p)$  is measurable with respect to  $q$  for  $p \in \mathfrak{N}$ .

If  $\varphi(\xi, p) > 0$  on a set of positive measure  $\mathfrak{G}$  for some  $p \in \mathfrak{N}$   $\varphi(q, \xi) = 0$  almost everywhere on  $\mathfrak{B}_p \times \mathfrak{G}$ , contradicting the assumption that  $[Ax] = I$  for every  $x > 0$ . Thus  $\varphi(p, q) = 0$  almost everywhere on  $\mathfrak{G} \times \mathfrak{N}$ , also contradicting the assumption. Hence  $\Psi(q, p) > 0$  almost everywhere. It is known [7] that  $\bar{a} \otimes a = \bigcap_{\nu=1}^{\infty} (\nu A^2 \wedge \bar{a} \otimes a)$  and  $r(\bar{a} \otimes a) = 1$ . Lemmas 4.5 and 4.2 imply  $r(A^2) = r(A)^2 > 0$ .

Next theorem is proved in [11], but we give a somewhat different proof.

*Theorem 4.6.* Let  $F_\lambda$  ( $0 \leq \lambda \leq 1$ ) be bounded linear operators defined on a Banach space  $R$  such that  $F_0 = 0, F_1 = I, F_\lambda F_\mu = F_{\min(\lambda, \mu)}$  and  $\lim_{\rho \uparrow \lambda} F_\rho x = \lim_{\rho \downarrow \lambda} F_\rho x = F_\lambda x$  ( $x \in R$ ). If  $A$  is a compact linear operator on  $R$  such that  $AF_\lambda = F_\lambda AF_\lambda$  ( $0 \leq \lambda \leq 1$ ), then  $A$  is quasi-nilpotent.

*Proof.* Since by Lemma 4.1  $\text{Max}\{r(F_\lambda AF_\lambda), r((I - F_\lambda)A(I - F_\lambda))\} = r(A)$ , there exists, by induction, sequences  $\lambda_\nu \uparrow_{\nu=1}^{\infty} a$  and  $\mu_\nu \downarrow_{\nu=1}^{\infty} \beta$  such that  $r((F_{\mu_\nu} - F_{\lambda_\nu})A(F_{\mu_\nu} - F_{\lambda_\nu})) = r(A)$ , ( $\nu = 1, 2, \dots$ ). But by Lemma 4.3  $r(A) = r((F_\beta - F_\alpha)A(F_\beta - F_\alpha))$ . Continuing this method, we obtain  $r(A) = 0$ .

Combining Theorems 4.5 and 4.6, we obtain a generalization of criteria of Volterra-type ([10] p. 147).

*Theorem 4.7.* Let  $R$  be reflexive as a Banach space and have no discrete element, and  $A$  be positive, compact and totally continuous. Then  $A$  is quasi-nilpotent if and only if there exist projections  $[N_\lambda]$  ( $0 \leq \lambda \leq 1$ ), such that

$$(4.7) \quad [N_0] = 0, [N_1] = \bigcup_{x \in R} [Ax], \bigcap_{\rho > \lambda} [N_\rho] = \bigcup_{\rho < \lambda} [N_\rho] = [N_\lambda]$$

$$A[N_\lambda] = [N_\lambda]A[N_\lambda] \quad (0 \leq \lambda \leq 1).$$

*Proof.* Since by Theorem 4.5 any non-trivial invariant normal manifold contains the other non-trivial one, the proof proceeds as in Theorem 4.3.

§ 5. Proper values with maximum modulus. Here the distribution

of proper values with maximum modulus of a positive compact linear operator is considered.

*Lemma 5.1.* *If  $A$  is compact and completely positive with  $r=r(A)>0$ , the proper values with maximum modulus are all simple and are the solutions of the equation*

$$(5.1) \quad \xi^k - r^k = 0 \quad \text{for some } k.$$

*Proof.* Since by Theorem 3.1  $\mu(r)=1$ , the assertion follows from Theorem 8.1 of [5].

*Lemma 5.2.* *For a compact linear operator  $A$  on a Banach space  $R$ , the following conditions are equivalent to each other*

- 1)  $w\text{-}\lim_{\nu \rightarrow \infty} A^\nu x = E(1)x \quad (x \in R)$
- 2)  $\lim_{\nu \rightarrow \infty} A^\nu x = E(1)x \quad (x \in R)$
- 3)  $\lim_{\nu \rightarrow \infty} \|A^\nu - E(1)\| = 0$
- 4)  $r(A) \leq 1$  and 1 is the only possible proper value with modulus 1.

The proof is similar to that of Lemma 3.5.

*Theorem 5.1.* *Let  $A$  be compact and completely positive with  $r(A)=1$ . The following conditions are equivalent:*

- 1) 1 is the unique proper value with maximum modulus,
- 2)  $\lim_{\nu \rightarrow \infty} A^\nu x$  exists and is complete for every  $x > 0$ ,
- 3)  $A^\nu (\nu=1, 2, \dots)$  are all completely positive.

*Proof.* By Theorem 2.1 and Lemma 5.2, 1) and 2) are equivalent. Let 2) be satisfied. Since  $\lim_{n \rightarrow \infty} A^{\nu n} x = E(1)x$ , 3) follows from 2) by the definition. Finally if  $A^\nu (\nu=1, 2, \dots)$  are all completely positive and  $\lambda$  is a proper value with  $|\lambda|=1$  distinct from 1, then by Lemma 5.1 there exists a positive integer  $k$  such that  $\lambda^k=1$ , so 1 is a proper value of  $A^k$  of multiplicity greater than 1, contradicting the assumption by Theorem 3.4.

*Lemma 5.3.* *Let  $R$  be reflexive as a Banach space and  $A$  be compact. If  $[N_\lambda] \downarrow_{\lambda \in A} [N]$  and  $\sigma([N_\lambda] A [N_\lambda]) \subseteq \sigma(A)$ , then  $\sigma([N] A [N]) \subseteq \sigma(A)$ .*

*Proof.* The reflexivity implies that that the norms of  $R$  and  $\bar{R}$  are both continuous. Since  $\lim_{\lambda \in A} \|[N_\lambda] A [N_\lambda] - [N] A [N]\| = 0$ , by Lemma 4.2  $\sigma([N] A [N]) = \lim_{\lambda \in A} \sigma([N_\lambda] A [N_\lambda]) \subseteq \sigma(A)$ .

*Theorem 5.2.* *Let  $R$  be reflexive as a Banach space. If  $A$  is positive*

compact, the proper values with maximum modulus coincides with the solutions of the equation

$$(5.2) \quad \prod_{i=1}^n (\xi^{k_i} - r^{k_i}) = 0$$

where  $k_i (i=1, 2, \dots, n)$  are some positive integers.

*Proof.* Reflexivity of  $R$  implies universal continuity of  $A$ . Let  $\lambda$  be a proper value with maximum modulus. Considering, by ZORN'S lemma, a maximal linearly ordered family  $[N_\rho]_{\rho \in A}$  such that  $\lambda \in \sigma([N_\rho] A [N_\rho]) \subseteq \sigma(A)$ . Putting  $[N] = \bigcap [N_\lambda]$ , by Lemma 5.3, we obtain  $\lambda \in \sigma([N] A [N]) \subseteq \sigma(A)$ . Since  $r([N] A [N]) \leq r(A)$ ,  $\lambda$  is a proper value of  $[N] A [N]$  with maximum modulus. By Lemma 4.1, the maximal hypothesis implies that there is no  $0 < [M] < [N]$  with  $[N] A [M] = [M] A [M]$ , that is,  $[N] A [N]$  is completely positive. Hence by Lemma 5.1  $\lambda$  is a solution of an equation  $\xi^k = r^k$  for some  $k$  and all its solutions are contained in  $\sigma([N] A [N]) \subseteq \sigma(A)$ . Since  $A$  has only a finite number of proper values on the circle with radius  $r$ , this completes the proof.

### References

- [1] G. FROBENIUS: Über Matrizen aus positiven Elementen I. II. Sitz. Ber. Preuss. Akad. (1908) 471-476, (1909) 514-518.
- [2] G. FROBENIUS: Über Matrizen aus nichtnegativen Elementen, Sitz. Ber. Preuss. Akad. (1912) 456-477.
- [3] E. HILLE: Functional analysis and semi-groups, New York 1948.
- [4] R. JENTZSCH: Über Integralgleichungen mit positiven Kern, Crelles Jour. Bd. 141 (1912) 235-244.
- [5] M. KREIN and M. A. RUTMAN: Linear operators leaving a cone invariant in a BANACH space, Uspehi Math. 3 (1948) 3-95.
- [6] H. NAKANO: Modulated semi-ordered linear spaces, Tokyo 1950.
- [7] H. NAKANO: Product spaces of semi-ordered linear spaces, Jour. Fac. Sci. Hokkaido Univ. Ser. I Vol. 12 (1953) 163-210.
- [8] J. D. NEWBURGH: The variation of spectra, Duke Math. Jour. Vol. 18 (1951) 165-176.
- [9] O. PERRON: Grundlagen für eine Theorie des Jacobischer Kettenbruchalgorithmus, Math. Ann. Bd. 64 (1907) 1-76.
- [10] F. RIESZ and B. SZ-NAGY: Leçon d'analyse fonctionelle, 1952.
- [11] J. R. RINGROSE: Compact linear operators of Volterra type, Proc. Cambridge Phil. Soc. Vol. 51 (1954) 44-55.