



Title	ON THE DISTRIBUTION OF INTEGERS REPRESENTABLE AS A SUM OF TWO $n$ -TH POWERS
Author(s)	Uchiyama, Saburô
Citation	Journal of the Faculty of Science Hokkaido University. Ser. 1 Mathematics, 18(3-4), 124-127
Issue Date	1965
Doc URL	<a href="http://hdl.handle.net/2115/56058">http://hdl.handle.net/2115/56058</a>
Type	bulletin (article)
File Information	JFSHIU_18_N3-4_124-127.pdf



[Instructions for use](#)

# ON THE DISTRIBUTION OF INTEGERS REPRESENTABLE AS A SUM OF TWO $h$ -TH POWERS

By

Saburô UCHIYAMA

Our aim in this note is to present some elementary results concerning the distribution of integers which can be expressed as a sum of two  $h$ -th powers, where  $h \geq 2$  is a fixed integer.

1. According to P. Erdős [2], R. P. Bambah and S. Chowla [1] have proved that for some *sufficiently large* constant  $C$  the interval  $(n, n + Cn^{\frac{1}{4}})$  always contains an integer of the form  $x^2 + y^2$ ,  $n, x$  and  $y$  being integral, and Erdős [2] conjectures (among others) that this holds for every  $C$  if  $n \geq n_0(C)$ . We cannot, at present, prove this conjecture of Erdős, but it is possible to refine the result of Bambah and Chowla in the following form:

**Theorem 1.** *For every  $n \geq 1$  there are integers  $x, y$  with  $xy \neq 0$  satisfying*

$$n < x^2 + y^2 < n + 2^{\frac{3}{2}} n^{\frac{1}{4}}.$$

*Proof.* For  $n=1$  and  $n=2$  the result is obvious. Assume now that  $n \geq 3$ . Let  $\delta, 0 < \delta < 1$ , be a fixed real number: the exact value of  $\delta$  (which may depend on  $n$ ) will be determined in a moment later.

Write

$$[n^{\frac{1}{2}}] = n^{\frac{1}{2}} - (1 - \varepsilon) \quad (0 < \varepsilon \leq 1).$$

Here, and in what follows,  $[t]$  denotes, as usual, the greatest integer not exceeding  $t$ .

We distinguish two cases.

*Case 1:*  $0 < \varepsilon \leq \delta$ . We take

$$x = [n^{\frac{1}{2}}] + 1, \quad y = 1.$$

Then we have

$$n < x^2 + y^2 = n + 2\varepsilon n^{\frac{1}{2}} + \varepsilon^2 + 1 < n + 2^{\frac{3}{2}} n^{\frac{1}{4}},$$

if

$$2\epsilon n^{\frac{1}{2}} + \epsilon^2 + 1 < 2^{\frac{3}{2}} n^{\frac{1}{4}}$$

or

$$(1) \quad \delta^2 + 2\delta n^{\frac{1}{2}} - (2^{\frac{3}{2}} n^{\frac{1}{4}} - 1) < 0.$$

Case 2:  $\delta < \epsilon \leq 1$ . We put

$$x = [n], \quad y = [(n - [n^{\frac{1}{2}}]^2)^{\frac{1}{2}}] + 1.$$

Then we have

$$n < x^2 + y^2 \leq n + 2 \left( 2(1 - \epsilon) n^{\frac{1}{2}} - (1 - \epsilon)^2 \right)^{\frac{1}{2}} + 1 < n + 2^{\frac{3}{2}} n^{\frac{1}{4}},$$

if

$$2 \left( 2(1 - \epsilon) n^{\frac{1}{2}} - (1 - \epsilon)^2 \right)^{\frac{1}{2}} + 1 < 2^{\frac{3}{2}} n^{\frac{1}{4}}$$

or

$$(2) \quad \delta^2 + 2\delta(n^{\frac{1}{2}} - 1) - \left( 2^{\frac{1}{2}} n^{\frac{1}{4}} - \frac{5}{4} \right) > 0.$$

Now, let  $\delta_0$  be the (unique) positive zero of the quadratic equation

$$\delta_0^2 + 2\delta_0(n^{\frac{1}{2}} - 1) - \left( 2^{\frac{1}{2}} n^{\frac{1}{4}} - \frac{5}{4} \right) = 0.$$

It is easy to see that  $\delta_0 < 1$  and that

$$\delta_0^2 + 2\delta_0 n^{\frac{1}{2}} - (2^{\frac{3}{2}} n^{\frac{1}{4}} - 1) = 2\delta_0 - \left( 2^{\frac{1}{2}} n^{\frac{1}{4}} + \frac{1}{4} \right) < 0$$

for  $n \geq 3$ . Thus, we may take any  $\delta$  less than 1 and slightly greater than  $\delta_0$ , so that the inequalities (1) and (2) hold true simultaneously. This proves the theorem.

**Corollary 1.** *For every  $\epsilon > 0$  the set of integers  $n$  for which the interval  $(n, n + \epsilon n^{\frac{1}{4}})$  contains an integer of the form  $x^2 + y^2$  has a positive density.*

*Proof.* For every  $\delta, 0 < \delta \leq 1$ , the set of integers  $n$  satisfying  $n^{\frac{1}{2}} - \delta < [n^{\frac{1}{2}}] \leq n^{\frac{1}{2}}$  is of positive density. It suffices to take  $\delta = \delta(\epsilon)$  small enough.

2. Here we wish to state two conjectures related to Theorem 1. They are:

**Conjecture 1.** *Let  $C_1$  be a constant  $> 2^{-\frac{1}{4}} \cdot 3$ . Then for all  $n \geq 1$  there are integers  $x, y$  with  $xy \neq 0$  satisfying*

$$n < x^2 + y^2 < n + C_1 n^{\frac{1}{4}};$$

and

**Conjecture 2.** Let  $C_2$  be a constant  $> 2^{-\frac{1}{2}} \cdot 5^{\frac{3}{4}}$ . Then for all  $n \geq 1$  there are integers  $x, y$  satisfying

$$n < x^2 + y^2 < n + C_2 n^{\frac{1}{4}}.$$

Either of these conjectures, if true, is the best possible in the sense that if  $C_1 = 2^{-\frac{1}{4}} \cdot 3$  or  $C_2 = 2^{-\frac{1}{2}} \cdot 5^{\frac{3}{4}}$  then the corresponding result cannot be correct any longer (to see this we put, for instance,  $n=2$  or  $n=20$ ). We note also that our Conjectures 1 and 2 have been verified by M. Uchiyama up to  $n=1000$ .

3. For a general  $h \geq 2$  we shall mention the following rather trivial

**Theorem 2.** Let  $g(n)$  be a function of  $n$  satisfying the inequality

$$g(n) > \sum_{j=1}^h \binom{h}{j} (n - [n^{1/h}]^h)^{(h-j)/h}$$

for  $n \geq n_0$ . Then there exist integers  $x, y$  with  $xy \neq 0$  such that

$$n < x^h + y^h < n + g(n)$$

for all  $n \geq n_0$ .

*Proof.* Put

$$x = [n^{1/h}], \quad y = [(n - [n^{1/h}]^h)^{1/h}] + 1.$$

**Corollary 2.** For any  $\varepsilon > 0$  there is an  $n_0 = n_0(\varepsilon)$  such that for all  $n \geq n_0$  there exist integers  $x, y$  with  $xy \neq 0$  satisfying

$$n < x^h + y^h < n + (c + \varepsilon)n^a,$$

where

$$a = \left(1 - \frac{1}{h}\right)^2, \quad c = h^{(2h-1)/h}.$$

For  $h=2$  this is of course weaker than Theorem 1.

**Corollary 3.** For every  $\varepsilon > 0$  the set of integers  $n$  for which the interval  $(n, n + \varepsilon n^a)$ , with  $a = \left(1 - \frac{1}{h}\right)^2$ , contains an integer of the form  $x^h + y^h$  has a positive density.

Proof is similar to that of Corollary 1.

### References

- [1]\* R. P. BAMBAH and S. CHOWLA: On numbers which can be expressed as a sum of two squares, *Proc. Nat. Inst. Sci. India*, vol. 13 (1947), pp. 101-103.
- [2] P. ERDÖS: Some unsolved problems, *Publ. Math. Inst. Hungar. Acad. Sci.*, vol. 6, ser. A (1961), pp. 221-254.

Department of Mathematics,  
Hokkaidō University

(Received September 10, 1964)

---

\* The writer has been unable to consult this paper.