Title	Prenatal exposure to perfluoroalkyl acids and allergic diseases in early childhood
Author(s)	Okada, Emiko; Sasaki, Seiko; Kashino, Ikuko; Matsuura, Hideyuki; Miyashita, Chihiro; Kobayashi, Sumitaka; Itoh, Kumiko; Ikeno, Tamiko; Tamakoshi, Akiko; Kishi, Reiko
Citation	Environment International, 65, 127-134 https://doi.org/10.1016/j.envint.2014.01.007
Issue Date	2014-04
Doc URL	http://hdl.handle.net/2115/56344
Type	article (author version)
File Information	Environ Int_65_127-134.pdf

[Instructions for use](#)

Prenatal exposure to perfluoroalkyl acids and allergic diseases in early childhood

Emiko Okada^a, Seiko Sasaki^a, Ikuko Kashino^a, Hideyuki Matsuura^b, Chihiro Miyashita^c,
Sumitaka Kobayashi^a, Kumiko Itoh^a, Tamiko Ikeno^c, Akiko Tamakoshi^a, Reiko Kishi^{c,*}

^aDepartment of Public Health Sciences, Hokkaido University Graduate School of Medicine,
North 15 West 7 Kita-ku, Sapporo, 060-8638, Japan.

^bLaboratory of Bioorganic Chemistry, Division of Applied Bioscience, Research Faculty of
Agriculture, Hokkaido University, North 9 West 9 Kita-ku, Sapporo, 060-8589, Japan.

^cCenter for Environmental and Health Sciences, Hokkaido University, North 12 West 7
Kita-ku, Sapporo, 060-0812, Japan.

Email address of each author: Emiko Okada^a <e-okada@med.hokudai.ac.jp>, Seiko Sasaki^a,
Ikuko Kashino^a, Hideyuki Matsuura^b, Chihiro Miyashita^c, Sumitaka Kobayashi^a, Kumiko
Itoh^a, Tamiko Ikeno^c, Akiko Tamakoshi^a, Reiko Kishi <rkishi@med.hokudai.ac.jp>

*Corresponding author: Center for Environmental and Health Sciences, Hokkaido
University, North 12 West 7 Kita-ku, Sapporo, 060-0812, Japan. Tel: +81-11-706-4746;
Fax: +81-11-706-4725; E-mail: rkishi@med.hokudai.ac.jp (R. Kishi).

ABSTRACT

Perfluoroalkyl acids (PFAAs) are persistent organic pollutants that are detected in humans worldwide. Laboratory animal studies have shown that PFAAs are associated with immunotoxic effects. However, epidemiological studies investigating the role of PFAAs, in particular PFAAs with longer chains than perfluorooctanoic acid, are scarce. We investigated associations between prenatal exposure to PFAAs, including long-chain compounds, and infant allergic diseases at 12 and 24 months in a large study population. The participants included mothers and their infants who enrolled in the Hokkaido Study on Environment and Children's Health 2003–2009. Eleven PFAAs were measured in maternal plasma taken at 28–32 weeks of gestation using ultra-performance liquid chromatography coupled to triple quadrupole tandem mass spectrometry. Characteristics of participants and information on infant allergic diseases were obtained from self-administered questionnaires and medical records. At 24 months, the adjusted odds ratio (OR) (first vs. fourth quartiles) for eczema in association with higher maternal perfluorotridecanoic acid (PFTrDA) levels was 0.62 (95% confidence interval (CI) 0.45, 0.86). After stratification by gender, the adjusted ORs in female infants from mothers with higher maternal perfluoroundecanoic acid (PFUnDA) and PFTrDA levels were also statistically significant (PFUnDA: OR = 0.50; 95% CI, 0.30, 0.81; PFTrDA: OR = 0.39; 95% CI, 0.23, 0.64). Our findings suggest that lower prenatal exposure to PFTrDA may decrease the risk of developing eczema in early childhood, only in female infants.

1

2 **Key words:** Perfluoroalkyl acids, perfluorotridecanoic acid, prenatal exposure, allergic
3 diseases, eczema

4

5 **Funding:** This study was funded by a Grant-in-Aid for Scientific Research from the
6 Japanese Ministry of Health, Labor, and Welfare; the Ministry of Education, Culture,
7 Sports, Science, and Technology; and the Japan Society for the Promotion of Science.

8 The authors declare that there are no conflicts of interest.

9

10 **Ethics approval:** This study was conducted with written informed consent from all
11 participants and was approved by the institutional ethical board for epidemiological studies at
12 the Hokkaido University Graduate School of Medicine.

13

14 **Abbreviations:**

15 PFAAs, perfluoroalkyl acids

16 PFCAs, perfluorinated carboxylic acids

17 PFHxA, perfluorohexanoic acid

18 PFHpA, perfluoroheptanoic acid

19 PFOA, perfluorooctanoic acid

20 PFNA, perfluorononanoic acid

- 1 PFDA, perfluorodecanoic acid
- 2 PFUnDA, perfluoroundecanoic acid
- 3 PFDoDA, perfluorododecanoic acid
- 4 PFTrDA, perfluorotridecanoic acid
- 5 PFTeDA, perfluorotetradecanoic acid
- 6 PFHxS, perfluorohexane sulfonate
- 7 PFOS, perfluorooctane sulfonate
- 8 MDL, method detection limits
- 9 CI, confidence interval
- 10 OR, odds ratio
- 11 Ig, immunoglobulin
- 12 ETS, environmental tobacco smoke
- 13 ISAAC, International Study of Asthma and Allergies in Childhood
- 14
- 15
- 16

1. Introduction

Perfluoroalkyl acids (PFAAs) are used in a broad range of consumer products because of their surface properties, which include insulation and water resistance. These compounds are persistent organic pollutants that are widespread within the environment, wildlife, and humans (Lau et al., 2007). Contamination of drinking water and foodstuffs such as seafood, leaching from food packaging and non-stick cookware, and household dust are major known routes of human exposure (Fromme et al., 2009). Potential health effects associated with PFAA exposure in humans are worsened by both bioaccumulation and persistence.

PFAA exposure has been suggested to have immunotoxic effects in laboratory animals including altered inflammatory responses, production of cytokines, and adaptive and innate immune responses (Dewitt et al., 2009). Cytokine expression and signaling related to inflammation and T-helper cell responses are altered in PFAA-exposed animals (Dewitt et al., 2012). PFAAs cross the placental barrier and are transferred to the fetus in humans (Midasch et al., 2007; Monroy et al., 2008). Previous epidemiological studies have shown a positive or negative association between perfluorooctane sulfonic acid (PFOS) and perfluorooctanoic acid (PFOA) and levels of cord blood immunoglobulin (Ig) E (Okada et al., 2012; Wang et al., 2011). Moreover, these studies have reported no association between prenatal PFOS, PFOA, or perfluorononanoic acid (PFNA) exposure and allergic and infectious diseases as health outcomes in children (Fei et al., 2010; Okada et al., 2012;

Wang JJ et al., 2011). In the C8 Health Project, which was a cross-sectional, immune biomarker study that investigated residents in the vicinity of a PFOA plant, IgA, IgE, and C-reactive protein levels significantly decreased with increasing PFOA levels in blood samples (Fletcher et al., 2009).

In 2002, after 50 years of production, the 3M Company phased out the manufacture and distribution of PFOS (Renner 2001). PFOS was also included in Annex B of the 2009 Stockholm Convention on Persistent Organic Pollutants (UNEP 2007; Wang et al., 2009). The Environmental Protection Agency of the United States (2006) launched a 2010/2015 PFOA Stewardship Program to voluntarily reduce PFOA emissions. Recent studies indicate that concentrations of PFOS and PFOA are declining in the general human population (Olsen et al., 2012; Sundström et al., 2011; Wang M et al., 2011). In contrast, concentrations of PFNA and perfluorodecanoic acid (PFDA), which are long-chain perfluorinated carboxylic acids (PFCAs), are increasing in the general human population (Wang M et al., 2011). However, the effects of prenatal exposure to other PFAAs, particularly PFCAs, which generally have longer chains than PFOA with a carbon chain length of eight (e.g., PFDA, perfluoroundecanoic acid (PFUnDA), perfluorododecanoic acid (PFDoDA), and perfluorotridecanoic acid (PFTrDA)), have not been characterized. PFCAs with chains longer than those of PFOA have high bioconcentration factors, suggesting that they are environmentally persistent (Martin et al., 2003). Furthermore, between 2003 and 2011, we reported increased PFNA and PFDA in maternal plasma levels

in Japanese, whereas levels of PFOS and PFOA decreased (Okada et al., 2013).

Epidemiological determination of whether exposure to long-chain PFCAAs affects immunity and allergic responses in humans is critical.

In this study, we explored associations between maternal PFAA levels, including long-chain compounds, and allergic diseases in early childhood using a prospective birth cohort study.

2. Methods

2.1. Study population

This prospective ongoing birth cohort study (Hokkaido Study on Environment and Children's Health) includes mothers who gave birth at hospitals in Hokkaido, Japan and their infants. The study was initiated in February 2003, and details have been described elsewhere (Kishi et al., 2011; Kishi et al., 2013). Briefly, participants were considered eligible if they were indigenous Japanese women who had received antenatal care at one of 37 participating hospitals within Hokkaido during their first trimester of pregnancy. Of the 33,500 eligible women invited to participate in the study from 2003 to 2009, 17,869 agreed to join (participation rate 53.3%). These participants signed informed consent forms, completed a baseline questionnaire, and also mailed follow-up questionnaires. From all participants (n = 17,869), we selected 12,847 who had submitted a baseline questionnaire and from whom we had obtained a third trimester blood sample and hospital birth records.

From these, we excluded cases of miscarriage and stillbirth ($n = 19$), congenital malformation ($n = 143$), and multiple births ($n = 162$), because these are common exclusion criteria for studies investigating allergies, infectious diseases, mental development, and endocrine metabolic disorders. From the selected 12,523 participants, we then extracted 6,335 participants who had completed all three self-administered questionnaires (at 4, 12, and 24 months after birth) for long-term follow-up of child development. Finally, from these 6,335 participants, we randomly extracted 300 participants per year from 2003 to 2008 and 295 participants in 2009 to give a total of 2,095 participants selected for the PFAA analysis of maternal plasma. Of these participants, we excluded cases of congenital malformations that became apparent from the follow-up questionnaire at 12 months ($n = 17$) and those whose maternal blood samples were taken before 26 weeks of gestation ($n = 15$) because the time of blood sampling during pregnancy may have affected concentrations due to increased maternal blood volume during gestation. Thus, a final total of 2,063 study participants met the specific exclusion and inclusion criteria for this study (Fig. 1). The protocol used in this study was approved by the institutional ethical board for epidemiological studies at the Hokkaido University Graduate School of Medicine.

2.2. Data collection

Participants completed a self-administered baseline questionnaire during the first trimester of pregnancy. The baseline questionnaire included maternal and paternal

information related to age, pre-pregnancy height and weight, previous medical history, educational level, household income, alcohol intake during pregnancy, and parity. Medical birth records from hospitals included the gestational age, infant gender, and birth weight, as well as miscarriage and stillbirth, multiple births, and congenital malformations. At 4 months post-delivery, participants completed a self-administered questionnaire including information about birth size, maternal complications during pregnancy, and maternal smoking status in the third trimester. At 12 and 24 months post-delivery, participants completed another self-administered questionnaire, which included information related to breast feeding, infant weight, length, head and chest circumferences, smoking status of parents, environmental tobacco smoke (ETS) exposure, pets in the home, day care attendance, infant vaccination, and previous or current medical history of infant allergic diseases (eczema, wheezing, and allergic rhinoconjunctivitis symptoms), infectious diseases, and other diseases. ETS exposure was defined as a self-reported positive response of whether a smoker was in the place where children lived their daily life at both 12 and 24 months of age.

2.3. Assessment of infant allergic diseases

Infant allergies that developed during the first 12 months of life and from months 12–24 were assessed based on the mothers' self-administered questionnaires that were obtained twice, at 12 and 24 months post-delivery. Allergic diseases were defined using a

modified part of the Japanese version of the International Study of Asthma and Allergies in Childhood (ISAAC) Phase Three questionnaire. In this study, we estimated eczema based on positive answers to all three of these questions: “Have you (has your child) had this itchy rash at any time in the past 12 months?”, “Have you (has your child) ever had a skin rash which was coming and going for at least 6 months?”, and “Has this itchy rash at any time affected any of the following places: the folds of the elbows; behind the knees; in front of the ankles; under the buttocks; or around the neck, ears, or eyes?” Wheezing was based on a positive answer to the question: “Have you (has your child) had wheezing or whistling in the chest in the past 12 months?” Current allergic rhinoconjunctivitis symptoms were based on all positive answers to both of these questions: “In the past 12 months, have you (has your child) had a problem with sneezing or a runny or blocked nose when you (he/she) did not have a cold or the flu?” and if yes, “In the past 12 months, has this nose problem been accompanied by itchy watery eyes?” (Asher et al., 2006). We also defined total allergic diseases as cases with at least one of the following symptoms: eczema, wheezing, allergic rhinoconjunctivitis symptoms.

2.4. Measurement of PFAA concentrations in maternal plasma

Detailed sampling and laboratory methods for analysis of PFAAs have been previously described (Okada et al., 2013). In brief, a 10-mL blood sample was taken from the maternal peripheral vein between 28 and 32 weeks of pregnancy. Maternal plasma was

analyzed using ultra-performance liquid chromatography coupled to triple quadrupole tandem mass spectrometry instrumentation (Waters, Tokyo, Japan). The concentrations of 11 PFAAs (perfluorohexanoic acid (PFHxA), perfluoroheptanoic acid (PFHpA), PFOA, PFNA, PFDA, PFUnDA, PFDoDA, PFTrDA, perfluorotetradecanoic acid (PFTeDA), perfluorohexane sulfonate (PFHxS), and PFOS) were measured in 2,095 maternal plasma samples. The method detection limits (MDLs) were: PFHxA, PFHpA, PFDA, PFUnDA, PFDoDA, PFTrDA, and PFTeDA (0.1 ng/mL), PFOA and PFHxS (0.2 ng/mL), and PFNA and PFOS (0.3 ng/mL).

2.5. Statistical analysis

For participants with PFAA concentrations below the MDL, a value equal to half of the MDL was assigned for statistical analyses. Participants were divided into four categories based on quartiles of maternal PFAA concentrations. Crude and adjusted logistic regression analyses were performed to evaluate associations between maternal PFAA concentrations and the risk of allergic diseases. In logistic models, odds ratios (ORs) for the risk of allergic diseases were evaluated with PFAA concentrations in the second through fourth quartiles and compared to those in the first quartiles. First, to see the risk of developing at least one of the symptoms (eczema, wheezing, and allergic rhinoconjunctivitis symptoms), we examined the relationship with total allergic diseases. Second, we examined the effects on each allergic disease. Potential confounding variables

considered in the analysis were: maternal age, educational levels, parental allergic history, infant gender, gestational age, birth season, breast feeding, siblings, ETS exposure, pets in the home, and day care attendance. Covariates in analysis were selected based on a review of the literature and on the change in estimate criteria, which were set to more than 10%.

The fully adjusted model used logistic regression analysis of total allergic diseases and was adjusted for maternal age, maternal educational level (≤ 9 years, 10–12 years, 13–16 years, and ≥ 17 years), parental allergic history (yes/no), infant gender, breast-feeding period (< 6 months or ≥ 6 months), number of older siblings, day care attendance (yes/no), and ETS exposure (yes/no). The number of older siblings was obtained from the parity information.

Logistic regression analysis of eczema was adjusted for maternal age, maternal educational level, parental allergic history, infant gender, breast-feeding period, and ETS exposure.

Logistic regression analysis of wheezing was adjusted for maternal age, maternal educational level, parental allergic history, infant gender, number of older siblings, day care attendance, and ETS exposure. In previous studies, gender differences were observed between prenatal exposure to PFAAs and birth weight or cord blood IgE levels (Okada et al., 2012; Washino et al., 2009), and therefore, we further analyzed the models including the multiplicative interaction term and stratified models to assess potential effect modification by gender. All statistical analyses were performed using JMP 10 Statistical Discovery Software for Windows (S.A.S. Institute Inc., Cary, NC). Differences were considered statistically significant at $p < 0.05$.

3. Results

Demographic characteristics of the parents and infants are shown in Table 1. The mean maternal age was 30.4 ± 4.5 years. The proportion with a maternal allergic history was 31.6%. Our population consisted of 1,044 (50.6%) male infants and 1,018 (49.4%) female infants.

Concentrations of maternal plasma PFAAs were measured (Table 2). Nearly all study participants had detectable plasma concentrations of PFOS, PFOA, PFNA, PFDA, PFUnDA, PFDoDA, and PFTTrDA (>90%), whereas PFHxS was detected in 81.9% of samples. PFHxA, PFHpA, and PFTeDA were detected in <50% of samples, and therefore, we did not include these compounds in the statistical analysis. PFOS was found at the highest median concentration (5.02 ng/mL), followed by PFOA (2.01 ng/mL), PFUnDA (1.40 ng/mL), and PFNA (1.15 ng/mL). We excluded a participant with an exceptionally high maternal PFOS concentration (464.8 ng/mL) as an outlier; this value was 93 times higher than the median concentration in our study and 51 times higher than the median concentration in the highest PFOS concentration area in Japan (Harada et al., 2010). Therefore, data from a final total of 2,062 participants were included in this study.

The incidences of infant allergic diseases during the first 24 months were determined for our study group (Table 3). The numbers of infants who developed allergic diseases up to age 24 months were as follows: eczema, 367 (17.8%); wheezing, 397 (19.3%); allergic

rhinoconjunctivitis symptoms, 91 (4.4%). The number of cases with at least one of these diseases was 714 (34.6%). We found significant gender differences in eczema and total allergic diseases, but no differences in wheezing or allergic rhinoconjunctivitis symptoms were observed between male and female infants.

Table 4 shows the results of logistic regression analyses between maternal PFAA concentrations in quartiles and total infant allergic diseases during the first 24 months. Crude and adjusted ORs ranged from 0.74 (95% confidence interval (CI): 0.57, 0.95) to 0.73 (95% CI: 0.56, 0.94) and from 0.71 (95% CI: 0.55, 0.92) to 0.73 (95% CI: 0.56, 0.94) for the three highest quartiles of PFTrDA compared with the lowest. After stratified analysis by infant gender, crude and adjusted ORs of the highest quartiles of PFAAs decreased compared with the lowest except for PFHxS, PFDA, and PFOS among female infants. The adjusted OR of PFTrDA for the highest quartiles versus the lowest quartile was 0.51 (95% CI: 0.35, 0.75). However, among male infants, the risk of allergic disease was not associated with maternal levels of the eight PFAAs. In interaction models based on quartiles of maternal PFAA levels, the adjusted ORs (95% CI) of total allergic diseases at 24 months for female infants compared to male infants were: 0.92 (0.85, 1.00) times lower for PFNA ($p = 0.050$), 0.90 (0.83, 0.98) times lower for PFUnDA ($p = 0.018$), and 0.90 (0.83, 0.98) times lower for PFTrDA ($p = 0.014$).

Crude and adjusted ORs for eczema and PFTrDA significantly decreased for the three highest quartiles compared with the lowest and ranged from 0.71 (95% CI: 0.52, 0.97)

to 0.64 (95% CI: 0.47, 0.88) and from 0.69 (95% CI: 0.50, 0.94) to 0.62 (95% CI: 0.45, 0.86), with a dose-response relationship (p for trend = 0.008 and 0.005, respectively; Table 5).

Among female infants, adjusted ORs of PFTrDA decreased for the three highest quartiles compared with the lowest and ranged from 0.60 (95% CI: 0.37, 0.95) to 0.39 (95% CI: 0.23, 0.64), with a dose-response relationship (p for trend <0.001). Also, the adjusted OR of PFUnDA for the highest quartiles versus lowest quartile was 0.50 (95% CI: 0.30, 0.81) among female infants (p for trend = 0.016). However, the risk of eczema was not associated with maternal levels of the eight PFAAs in male infants. In interaction models, the adjusted ORs (95% CI) of eczema at 24 months for female infants compared to male infants were: 0.89 (0.80, 0.99) times lower for PFUnDA (p = 0.033) and 0.88 (0.79, 0.97) times lower for PFTrDA (p = 0.014). Fig. 2 shows the adjusted ORs for eczema stratified by gender in association with the three highest quartiles for PFTrDA compared with the lowest.

At 12 months, no significant association was observed between eczema and PFAAs, although similar patterns of a decreased risk of eczema were seen (data not shown).

Regarding wheezing, no significant associations were observed with any maternal PFAA levels at 12 or 24 months (data not shown). We did not analyze allergic rhinoconjunctivitis symptoms because the number of cases with this type of allergic disease was very low, and sufficient statistical power could not be ensured in the multivariate analysis.

4. Discussion

Only in female infants during the first 24 months did we observe an association between high maternal PFAA levels (except for PFHxS, PFDA, and PFOS) and a decline in the risk of developing total allergic diseases as seen in cases with at least one of the following: eczema, wheezing, and allergic rhinoconjunctivitis symptoms. In the Sapporo study, which was a cohort study conducted of mothers and their infants who attended one local maternity hospital from 2002 to 2005 in Sapporo, Hokkaido, Japan, we previously reported that cord blood IgE levels decrease significantly with high maternal PFOA concentrations among female infants (Okada et al., 2012). The results of the C8 Health Project, which was a cross-sectional study that investigated residents exposed to PFOA-contaminated drinking water, showed a significant decreasing trend in IgE levels with increasing PFOA levels in blood samples among females (Fletcher et al., 2009). Two cohort studies showed that maternal PFOS, PFOA, PFNA, and PFHxS levels negatively correlate with antibody concentrations in children (Grandjean et al., 2012; Granum et al., 2013). These studies are supported by experimental studies showing adverse effects of PFOS and PFOA exposure on humoral immune function. Our results are consistent with laboratory animal experiments in which immunosuppression and reduced IgM antibody production were observed following PFAA exposure (Keil et al., 2008; Peden-Adams et al., 2007). However, the immunotoxic effect varies depending on the type of PFAA and the endpoint being evaluated (Dewitt et al., 2012). Reduced antibody concentrations in children exposed to PFAAs may lead to immunosuppression in childhood (Granum et al., 2013). In this study,

1 therefore, prenatal PFCA exposure may have suppressed the developing immune system in
2 infants and thereby indirectly reduced the risk of developing immune
3 hyperactivity/hypersensitivity diseases, such as eczema, wheezing, and allergic
4 rhinoconjunctivitis. However, despite the reduction in allergic diseases, general immune
5 suppression is not necessarily beneficial because this decrease may be linked to an immune
6 system deficit.

7 Our findings showed gender differences in the association of allergic diseases with
8 prenatal PFAA exposure. In the same study population (the Hokkaido Study on
9 Environment and Children's Health), we found a negative association between maternal
10 PFUnDA and PFTrDA levels and birth weight only in female infants (Kashino et al., 2013).
11 A previous study in China showed that compared to other PFAAs, PFTrDA in cord blood is
12 higher than in maternal blood, especially among female infants (Liu et al., 2011). In a
13 Korean study, levels of PFTrDA were negatively correlated with total thyroxine and
14 positively correlated with thyroid stimulating hormone levels, especially among females (Ji
15 et al., 2012). The transport of PFTrDA across the placental barrier from mothers to infants
16 suggests a gender difference. Therefore, prenatal PFTrDA exposure may have a potential
17 impact predominantly on female infants. Our finding also suggests that prenatal PFAAs
18 may differentially affect the development of allergic diseases in female infants. However,
19 the reason for the gender-specific association with PFTrDA is not clear. Very few studies
20 have reported the effects of long-chain PFCAs, particularly PFCAs that have longer chains

1 than PFDA such as PFTrDA. Further investigations into the effects of long-chain PFCAs in
2 different human populations are needed.

3 Levels of long-chain PFCAs such as PFUnDA, PFDoDA, and PFTrDA in plasma
4 in the present study were higher than those seen in many countries but lower than levels
5 reported in other areas of Japan (Harada et al., 2011). PFNA, PFUnDA, and PFTrDA are
6 manufactured primarily in Japan *via* the oxidation of a mixture of linear fluorotelomer
7 olefins (Prevedouros et al., 2006). Industrial application of these PFCAs may have
8 contributed to the observed accumulation of longer-chain PFCAs in East Asian populations.
9 Because longer-chain PFCAs have higher environmental persistence (Martin et al., 2003)
10 and longer half-lives (Ohmori et al., 2003), prenatal PFTrDA exposure may have led to a
11 reduction in the risk of developing eczema symptoms in infants in the current study. The
12 toxicity of PFCAs is correlated with the length of the carbon chain and the nature of the
13 functional group (Liao et al., 2009; Wolf et al., 2008).

14 We found no association between maternal PFOS and PFOA levels and eczema
15 and wheezing during the first 12 and 24 months. Our results are consistent with a previous
16 cohort study that examined prenatal exposure to PFOS and PFOA and the relationship with
17 atopic dermatitis, eczema, and wheezing (Granum et al., 2013; Okada et al., 2012; Wang JJ
18 et al., 2011). The case-control Genetic and Biomarkers study for Childhood Asthma
19 reported positive associations between serum PFAAs and asthma and positive associations
20 between PFAAs and IgE, absolute eosinophil counts, eosinophilic cationic protein levels,

and (to a lesser extent) asthma severity scores in asthmatic children (Dong et al., 2013). They investigated 10 types of PFAAs, but did not include PFTrDA and PFUnDA. The difference in the results between the present study and the Genetic and Biomarkers study for Childhood Asthma may be due to the differences in a prospective cohort study and a case-control study.

The present study has some limitations. First, we assessed allergic diseases in infants based on self-administered questionnaires by the mother. We did not investigate any biomarkers that indicate immunotoxicity. However, because we defined the development of allergic diseases with ISAAC questionnaires, which are internationally standardized procedures, these facts provided validity for the criteria for developing illness. Second, postnatal exposure to PFAAs from intake of food and drinking water or from indoor dust from birth to 24 months of age was not investigated in our study. Sources of postnatal exposure in infants also include breast milk and products in which PFAAs are used. Therefore, postnatal exposure to PFAAs may very well have affected the results. A strength of the present study is that it examined a large cohort of the general population from a relatively wide area (the entire Hokkaido region of Japan).

In conclusion, prenatal exposure to PFTrDA was associated with a decrease in the risk of developing eczema in early childhood in female infants. Prenatal PFCA exposure may have gender-specific effects on allergic diseases in infants. The immunotoxic potential of long-chain PFCAs, including PFTrDA, and the mechanisms of the gender-specific

differences warrant further studies.

Acknowledgments: We are grateful to all the participating mothers, the medical staff, and all persons involved in the Hokkaido study on Environment and Children's Health. We thank Toru Matsumura and Jun Yamamoto at the Institute of Environmental Ecology, IDEA Consultants, Inc. for assistance with blood analysis; and Yoichi M. Ito at the Department of Biostatistics, Division of Advanced Medical Sciences, Hokkaido University Graduate School of Medicine for assistance with the statistical analysis.

References

- Asher MI, Montefort S, Björkstén B, Lai CK, Strachan DP, Weiland SK. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet* 2006;368:733–43.
- DeWitt JC, Peden-Adams MM, Keller JM, Germolec DR. Immunotoxicity of perfluorinated compounds: recent developments. *Toxicol Pathol* 2012;40:300–11.
- DeWitt JC, Shnyra A, Badr MZ, Loveless SE, Hoban D, Frame SR, et al. Immunotoxicity of perfluorooctanoic acid and perfluorooctane sulfonate and the role of peroxisome proliferator-activated receptor alpha. *Crit Rev Toxicol* 2009;39:76–94.
- Dong GH, Tung KY, Tsai CH, Liu MM, Wang D, Liu W, et al. Serum polyfluoroalkyl

1 concentrations, asthma outcomes, and immunological markers in a case-control
2 study of Taiwanese children. *Environ Health Perspect* 2013;121:507–13.

3 Fei C, McLaughlin JK, Lipworth L, Olsen J. Prenatal exposure to PFOA and PFOS and risk
4 of hospitalization for infectious diseases in early childhood. *Environ Res*
5 2010;110:773–7.

6 Fletcher T, Steenland K, Savitz D. Status Report: PFOA and immune biomarkers in adults
7 exposed to PFOA in drinking water in the mid Ohio valley 2009:
8 <[http://www.c8sciencepanel.org/pdfs/Status_Report_C8_and_Immune_markers_Ma](http://www.c8sciencepanel.org/pdfs/Status_Report_C8_and_Immune_markers_March2009.pdf)
9 [rch2009.pdf](http://www.c8sciencepanel.org/pdfs/Status_Report_C8_and_Immune_markers_March2009.pdf)> (accessed 7 October 2013)

10 Fromme H, Tittlemier SA, Völkel W, Wilhelm M, Twardella D. Perfluorinated
11 compounds—exposure assessment for the general population in Western countries. *Int*
12 *J Hyg Environ Health* 2009; 212:239–70.

13 Grandjean P, Andersen EW, Budtz-Jørgensen E, Nielsen F, Mølbak K, Weihe P, et al. Serum
14 vaccine antibody concentrations in children exposed to perfluorinated compounds.
15 *JAMA* 2012;307:391–7.

16 Granum B, Haug LS, Namork E, Stølevik SB, Thomsen C, Aaberge IS, et al. Pre-natal
17 exposure to perfluoroalkyl substances may be associated with altered vaccine
18 antibody levels and immune-related health outcomes in early childhood. *J*
19 *Immunotoxicol* 2013; doi:10.3109/1547691X.2012.755580 [Online 25 January
20 2013].

1 Harada KH, Hitomi T, Niisoe T, Takanaka K, Kamiyama S, Watanabe T, et al. Odd-numbered
2 perfluorocarboxylates predominate over perfluorooctanoic acid in serum samples
3 from Japan, Korea and Vietnam. *Environ Int* 2011;37:1183–9.

4 Harada KH, Yang HR, Moon CS, Hung NN, Hitomi T, Inoue K, et al. Levels of
5 perfluorooctane sulfonate and perfluorooctanoic acid in female serum samples from
6 Japan in 2008, Korea in 1994-2008 and Vietnam in 2007-2008. *Chemosphere*.
7 2010;79:314–9.

8 Ji K, Kim S, Kho Y, Paek D, Sakong J, Ha J, et al. Serum concentrations of major
9 perfluorinated compounds among the general population in Korea: dietary sources
10 and potential impact on thyroid hormones. *Environ Int* 2012;45:78–85.

11 Kashino I, Okada E, Sasaki S, Miyashita C, Ikeno T, Araki A, et al. Prenatal Exposure to 11
12 Perfluorinated Compounds (PFCs) and infant weight in the Hokkaido Study on
13 Environmental and Children's Health. *Environment and Health - Bridging South,*
14 *North, East and West Conference of ISEE, ISES and ISIAQ. 19–23 August 2013,*
15 *Basel, Switzerland. 151.*
16 [http://ehp.niehs.nih.gov/ehbasel13/wp-content/uploads/2013/09/EHB13-Abstracts.](http://ehp.niehs.nih.gov/ehbasel13/wp-content/uploads/2013/09/EHB13-Abstracts.pdf)
17 [pdf](http://ehp.niehs.nih.gov/ehbasel13/wp-content/uploads/2013/09/EHB13-Abstracts.pdf)> (accessed 7 October 2013)

18 Keil DE, Mehlmann T, Butterworth L, Peden-Adams MM. Gestational exposure to
19 perfluorooctane sulfonate suppresses immune function in B6C3F1 mice. *Toxicol Sci*
20 2008;103:77–85.

- 1 Kishi R, Kobayashi S, Ikeno T, Araki A, Miyashita C, Itoh S, et al. Ten years of progress in
2 the Hokkaido birth cohort study on environment and children's health: cohort
3 profile-updated 2013. *Environ Health Prev Med* 2013;18:429–450.
- 4 Kishi R, Sasaki S, Yoshioka E, Yuasa M, Sata F, Saijo Y, et al. Cohort Profile: The
5 Hokkaido Study on Environment and Children's Health in Japan. *Int J Epidemiol*
6 2011;40:611–18.
- 7 Lau C, Anitole K, Hodes C, Lai D, Pfahles-Hutchens A, Seed J. Perfluoroalkyl acids: a
8 review of monitoring and toxicological findings. *Toxicol Sci* 2007;99:366–94.
- 9 Liao C, Wang T, Cui L, Zhou Q, Duan S, Jiang G. Changes in synaptic transmission, calcium
10 current, and neurite growth by perfluorinated compounds are dependent on the chain
11 length and functional group. *Environ Sci Technol* 2009;43:2099–104.
- 12 Liu J, Li J, Liu Y, Chan HM, Zhao Y, Cai Z, et al. Comparison on gestation and lactation
13 exposure of perfluorinated compounds for newborns. *Environ Int* 2011;37:1206–12.
- 14 Martin JW, Mabury SA, Solomon KR, Muir DCG. Bioconcentration and tissue distribution of
15 perfluorinated acids in rainbow trout (*Oncorhynchus mykiss*). *Environ Toxicol*
16 *Chem* 2003;22:196–204.
- 17 Midasch O, Drexler H, Hart N, Beckmann MW, Angerer J. Transplacental exposure of
18 neonates to perfluorooctanesulfonate and perfluorooctanoate: a pilot study. *Int Arch*
19 *Occup Environ Health* 2007;80:643–8.
- 20 Monroy R, Morrison K, Teo K, Atkinson S, Kubwabo C, Stewart B, et al. Serum levels of

perfluoroalkyl compounds in human maternal and umbilical cord blood samples.

Environ Res 2008;108:56–62.

Ohmori K, Kudo N, Katayama K, Kawashima Y. Comparison of the toxicokinetics between perfluorocarboxylic acids with different carbon chain length. Toxicology

2003;184:135–40.

Okada E, Kashino I, Matsuura H, Sasaki S, Miyashita C, Yamamoto J, et al. Temporal trends of perfluoroalkyl acids in plasma samples of pregnant women in Hokkaido, Japan,

2003-2011. Environ Int 2013;60:89–96.

Okada E, Sasaki S, Saijo Y, Washino N, Miyashita C, Kobayashi S, et al. Prenatal exposure to perfluorinated chemicals and relationship with allergies and infectious diseases in

infants. Environ Res 2012;112:118–25.

Olsen GW, Lange CC, Ellefson ME, Mair DC, Church TR, Goldberg CL, et al. Temporal trends of perfluoroalkyl concentrations in American Red Cross adult blood donors,

2000–2010. Environ Sci Technol 2012;46:6330–8.

Peden-Adams MM, EuDaly JG, Dabra S, EuDaly A, Heesemann L, Smythe J, et al.

Suppression of humoral immunity following exposure to the perfluorinated

insecticide sulfluramid. Toxicol Environ Health A 2007;70:1130–41.

Prevedouros K, Cousins IT, Buck RC, Korzeniowski SH. Sources, fate and transport of perfluorocarboxylates. Environ Sci Technol 2006;40:32–44.

Renner R. Scotchgard scotched—following the fabric protector's slippery trail to a new class

1 of pollutant. Sci Am 2001;284:18.

2 Sundström M, Ehresman DJ, Binert A, Butenhoff JL, Olsen GW, Chang SC. A temporal trend
3 study (1972–2008) of perfluorooctanesulfonate, perfluorohexanesulfonate, and
4 perfluorooctanoate in pooled human milk samples from Stockholm, Sweden.
5 Environ Int 2011;37:178–83.

6 UNEP, 2007. POPRC3: Development of Risk Management Evaluation;
7 UNEP/POPS/POPRC.3/20
8 <[http://chm.pops.int/Portals/0/Repository/poprc3/UNEP-POPS/POPRC.3-POPRC-3-](http://chm.pops.int/Portals/0/Repository/poprc3/UNEP-POPS/POPRC.3-POPRC-3-5.English.PDF)
9 5.English.PDF> (accessed 7 October 2013).

10 United States Environmental Protection Agency, 2006. 2010/15 PFOA Stewardship Program.
11 <<http://www.epa.gov/oppt/pfoa/pubs/stewardship/index.html>> (accessed 7 October
12 2013).

13 Wang IJ, Hsieh WS, Chen CY, Fletcher T, Lien GW, Chiang HL, et al. The effect of prenatal
14 perfluorinated chemicals exposures on pediatric atopy. Environ Res
15 2011;111:785–91.

16 Wang M, Park JS, Petreas M. Temporal changes in the levels of perfluorinated compounds in
17 California women's serum over the past 50 years. Environ Sci Technol
18 2011;45:7510–6.

19 Wang T, Wang YW, Liao CY, Cai YQ, Jiang GB. Perspectives on the inclusion of
20 perfluorooctane sulfonate into the Stockholm Convention on Persistent Organic

- 1 Pollutants. Environ. Sci. Technol 2009;43:5171–5.
- 2 Washino N, Saijo Y, Sasaki S, Kato S, Ban S, Konishi K, et al. Correlations between prenatal
- 3 exposure to perfluorinated chemicals and reduced fetal growth. Environ Health
- 4 Perspect 2009;117:660–7.
- 5 Wolf CJ, Takacs ML, Schmid JE, Lau C, Abbott BD. Activation of mouse and human
- 6 peroxisome proliferator-activated receptor alpha by perfluoroalkyl acids of different
- 7 functional groups and chain lengths. Toxicol Sci 2008;106:162–71.
- 8

Fig. 1. Flow chart of study participant selection.

A; All infants

B; Male infants

C; Female infants

Fig. 2.

ORs for eczema in association with the three highest quartiles for PFTrDA compared with the lowest. The data were adjusted for maternal age, maternal educational level, parental allergic history, infant gender, breast-feeding period, and ETS exposure in infancy at 24 months. (A) Among all infants. (B) Among male infants. (C) Among female infants.

Table 1

Characteristics of 2,062 parents and infants of the Hokkaido Study on Environment and Children's Health, Japan, 2003–2009.

		No.	(%)
Parental characteristics			
Maternal age (years) (mean $\pm$ SD)		30.4 $\pm$ 4.5	
Annual household income (million yen) ^a	<5	1,155	(61.8)
	$\geq$ 5	640	(34.5)
Maternal educational level (years)	<13	911	(44.2)
	$\geq$ 13	1,151	(55.8)
Parity (times)	0	944	(45.8)
	$\geq$ 1	1,118	(54.2)
Maternal smoking status during pregnancy	Nonsmoker	1,912	(92.7)
	Smoker	150	(7.3)
Maternal allergic history	Yes	652	(31.6)
Paternal allergic history	Yes	385	(18.7)
Infant characteristics			
Gender	Male	1,044	(50.6)
	Female	1,018	(49.4)
Older siblings (number)	0	944	(45.8)
	$\geq$ 1	1,118	(54.2)
Breast-feeding period (months) ^a	<6	420	(20.4)
	$\geq$ 6	1,640	(79.6)
Day care attendance at 24 months	Yes	583	(28.3)
ETS exposure at 24 months ^b	Yes	947	(45.9)

^aMissing data: annual household income (267), breast-feeding period (2).

^bETS: environmental tobacco smoke.

Table 2
Concentrations of 11 PFAs in 2,062 maternal plasma samples from the Hokkaido Study on Environment and Children's Health, Japan, 2003–2009.

Compound (carbon chain length)	MDL ^a	Detection		Concentration (ng/mL)					
		No.	(%)	Geometric mean	Mean	Minimum	25 th	50 th	75 th Maximum
PFHxA (C6)	0.2	1,688	(81.9)	0.275	0.324	<0.2	0.222	0.296	0.395 3.39
PFHxA (C6)	0.1	970	(47.0)	<0.1	0.104	<0.1	<0.1	<0.1	0.146 0.694
PFHpA (C7)	0.1	719	(34.9)	<0.1	0.096	<0.1	<0.1	<0.1	0.125 1.02
PFOS (C8)	0.3	2,062	(100)	5.01	5.56	1.00	3.71	5.02	6.83 30.3
PFOA (C8)	0.2	2,061	(100.0)	2.08	2.67	<0.2	1.31	2.01	3.26 24.9
PFNA (C9)	0.3	2,059	(99.9)	1.19	1.36	<0.3	0.873	1.15	1.57 13.2
PFDA (C10)	0.1	2,049	(99.4)	0.501	0.563	<0.1	0.382	0.510	0.684 2.43
PFUnDA (C11)	0.1	2,055	(99.7)	1.34	1.50	<0.1	1.02	1.40	1.87 5.89
PFDoDA (C12)	0.1	1,857	(90.1)	0.168	0.188	<0.1	0.138	0.182	0.230 0.729
PFTtDA (C13)	0.1	2,012	(97.6)	0.312	0.347	<0.1	0.244	0.329	0.424 1.33
PFTeDA (C14)	0.1	271	(13.1)	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1 0.303

^aMDL: method detection limit.

Table 3
Number and proportion of infants who developed allergic diseases during the first 24 months in the Hokkaido Study on Environment and Children's Health, Japan, 2003–2009 (n = 2,062).

Symptoms	Total (n = 2,062)		Male infants (n = 1,044)		Female infants (n = 1,018)		<i>p</i> ^a
	n	(%)	n	(%)	n	(%)	
Allergic diseases ^b	714	(34.6)	391	(37.5)	323	(31.7)	0.006
Eczema	367	(17.8)	212	(20.3)	155	(15.2)	0.003
Wheezing	397	(19.3)	212	(20.3)	185	(18.2)	0.241
Allergic rhinoconjunctivitis symptoms	91	(4.4)	51	(4.9)	40	(3.9)	0.335

^aFisher's exact test.

^b“Allergic diseases” indicates cases with at least one of the listed symptoms.

Table 4
Odds ratio (95% CI) between PFAA concentrations in maternal plasma and total allergic diseases during the first 24 months in the Hokkaido Study on Environment and Children's Health, Japan, 2003–2009 (n = 2,062).

		Total (n = 2,062)				Male infants (n = 1,044)				Female infants (n = 1,018)						
Compound (carbon chain length)	n	Crude		Adjusted ^a		n	Crude		Adjusted ^b		n	Crude		Adjusted ^b		
		OR ^c	(95% CI) ^d	OR ^c	(95% CI) ^d		OR ^c	(95% CI) ^d	OR ^c	(95% CI) ^d		OR ^c	(95% CI) ^d	OR ^c	(95% CI) ^d	
PFHxS (C6)	Quantile 1	189	1.00			112	1.00			77	1.00					
	Quantile 2	193	1.05	(0.81, 1.35)	1.04	(0.80, 1.35)	104	0.95	(0.67, 1.34)	0.93	(0.65, 1.33)	89	1.19	(0.82, 1.73)	1.18	(0.80, 1.74)
	Quantile 3	163	0.80	(0.62, 1.04)	0.82	(0.63, 1.07)	83	0.66	(0.46, 0.94)	0.65	(0.45, 0.94)	80	1.01	(0.69, 1.47)	1.10	(0.75, 1.64)
	Quantile 4	169	0.86	(0.66, 1.11)	0.93	(0.71, 1.21)	92	0.78	(0.55, 1.11)	0.81	(0.56, 1.16)	77	0.96	(0.66, 1.40)	1.13	(0.75, 1.69)
	<i>p</i> for trend			0.078				0.052		0.085			0.002		0.657	
PFOS (C8)	Quantile 1	195	1.00			102	1.00			93	1.00					
	Quantile 2	185	0.91	(0.71, 1.17)	0.97	(0.75, 1.26)	96	1.03	(0.72, 1.46)	1.12	(0.78, 1.61)	89	0.81	(0.56, 1.17)	0.81	(0.55, 1.18)
	Quantile 3	171	0.76	(0.59, 0.98)	0.80	(0.61, 1.04)	106	0.89	(0.63, 1.26)	0.91	(0.64, 1.30)	65	0.63	(0.43, 0.93)	0.68	(0.46, 1.01)
	Quantile 4	163	0.77	(0.59, 0.99)	0.86	(0.66, 1.13)	87	0.90	(0.63, 1.29)	0.95	(0.65, 1.37)	76	0.66	(0.45, 0.95)	0.79	(0.53, 1.17)
	<i>p</i> for trend			0.018				0.432		0.535			0.629		0.183	
PFOA (C8)	Quantile 1	197	1.00			102	1.00			95	1.00					
	Quantile 2	199	1.01	(0.79, 1.30)	1.05	(0.81, 1.37)	110	1.09	(0.77, 1.54)	1.11	(0.77, 1.60)	89	0.94	(0.65, 1.35)	1.01	(0.69, 1.47)
	Quantile 3	162	0.74	(0.57, 0.95)	0.80	(0.61, 1.06)	87	0.77	(0.54, 1.10)	0.82	(0.56, 1.20)	75	0.71	(0.49, 1.02)	0.77	(0.52, 1.15)
	Quantile 4	156	0.70	(0.54, 0.91)	0.79	(0.59, 1.04)	92	0.85	(0.60, 1.22)	0.93	(0.63, 1.37)	64	0.56	(0.38, 0.82)	0.64	(0.42, 0.97)
	<i>p</i> for trend			0.001				0.151		0.402			0.001		0.017	
PFNA (C9)	Quantile 1	203	1.00			99	1.00			104	1.00					
	Quantile 2	152	0.79	(0.61, 1.03)	0.81	(0.62, 1.06)	89	1.10	(0.76, 1.59)	1.10	(0.75, 1.60)	63	0.55	(0.38, 0.81)	0.59	(0.40, 0.88)
	Quantile 3	201	0.81	(0.63, 1.03)	0.82	(0.63, 1.05)	107	0.94	(0.67, 1.33)	0.96	(0.67, 1.37)	94	0.69	(0.48, 0.98)	0.71	(0.49, 1.02)
	Quantile 4	158	0.68	(0.53, 0.88)	0.73	(0.55, 0.95)	96	0.93	(0.65, 1.32)	0.95	(0.66, 1.38)	62	0.48	(0.32, 0.69)	0.55	(0.36, 0.82)
	<i>p</i> for trend			0.006				0.522		0.658			0.001		0.010	
PFDA (C10)	Quantile 1	187	1.00			96	1.00			91	1.00					
	Quantile 2	197	1.10	(0.85, 1.42)	1.12	(0.86, 1.44)	107	1.30	(0.91, 1.85)	1.31	(0.92, 1.89)	90	0.92	(0.64, 1.33)	0.94	(0.65, 1.37)
	Quantile 3	166	0.84	(0.65, 1.09)	0.84	(0.64, 1.09)	92	1.01	(0.71, 1.44)	0.98	(0.68, 1.41)	74	0.69	(0.48, 1.01)	0.73	(0.50, 1.08)
	Quantile 4	164	0.83	(0.64, 1.07)	0.89	(0.69, 1.17)	96	1.11	(0.78, 1.58)	1.13	(0.78, 1.64)	68	0.60	(0.41, 0.88)	0.70	(0.47, 1.04)
	<i>p</i> for trend			0.044				0.915		0.894			0.003		0.039	
PFUnDA (C11)	Quantile 1	190	1.00			93	1.00			97	1.00					
	Quantile 2	184	0.95	(0.74, 1.22)	0.92	(0.71, 1.19)	108	1.23	(0.87, 1.75)	1.22	(0.85, 1.74)	76	0.70	(0.48, 1.02)	0.65	(0.44, 0.96)

Quartile 3	169	0.83	(0.64, 1.08)	0.80	(0.62, 1.04)	90	1.11	(0.77, 1.59)	1.11	(0.76, 1.62)	79	0.63	(0.44, 0.91)	0.57	(0.39, 0.83)
Quartile 4	171	0.85	(0.66, 1.10)	0.82	(0.63, 1.07)	100	1.15	(0.81, 1.64)	1.13	(0.79, 1.63)	71	0.61	(0.42, 0.89)	0.58	(0.39, 0.86)
<i>p</i> for trend		0.136		0.092			0.594		0.642				0.008		0.004
PFDODA (C12)															
Quartile 1	202	1.00		1.00		100	1.00		1.00		102	1.00		1.00	
Quartile 2	158	0.69	(0.53, 0.89)	0.66	(0.51, 0.86)	99	0.88	(0.62, 1.25)	0.89	(0.62, 1.27)	59	0.51	(0.34, 0.74)	0.45	(0.30, 0.67)
Quartile 3	191	0.91	(0.71, 1.17)	0.87	(0.67, 1.13)	100	1.08	(0.76, 1.54)	1.10	(0.76, 1.58)	91	0.77	(0.54, 1.10)	0.67	(0.46, 0.97)
Quartile 4	163	0.73	(0.56, 0.94)	0.74	(0.57, 0.96)	92	0.89	(0.62, 1.27)	0.93	(0.65, 1.34)	71	0.59	(0.41, 0.85)	0.58	(0.39, 0.85)
<i>p</i> for trend		0.101		0.132			0.818		0.996				0.038		0.030
PFTrDA (C13)															
Quartile 1	205	1.00		1.00		98	1.00		1.00		107	1.00		1.00	
Quartile 2	169	0.74	(0.57, 0.95)	0.71	(0.55, 0.92)	98	0.91	(0.64, 1.29)	0.93	(0.64, 1.33)	71	0.58	(0.40, 0.84)	0.51	(0.34, 0.75)
Quartile 3	174	0.77	(0.60, 0.99)	0.75	(0.58, 0.98)	100	0.95	(0.67, 1.35)	1.01	(0.70, 1.45)	74	0.61	(0.42, 0.88)	0.54	(0.37, 0.79)
Quartile 4	166	0.73	(0.56, 0.94)	0.73	(0.56, 0.94)	95	0.99	(0.69, 1.42)	1.01	(0.70, 1.46)	71	0.54	(0.37, 0.77)	0.51	(0.35, 0.75)
<i>p</i> for trend		0.026		0.032			0.989		0.834				0.002		0.001

^a Adjusted for maternal age, maternal educational level, parental allergic history, infant gender, breast-feeding period, number of siblings, day care attendance, and ETS exposure in infancy at 24 months.

^b Adjusted for all the covariates except infant gender.

^cOR: odds ratio.

^dCI: confidence interval.

Table 5
Odds ratio (95% CI) between PFPA concentrations in maternal plasma and eczema during the first 24 months in the Hokkaido Study on Environment and Children's Health, Japan, 2003–2009 (n = 2,062)

Total (n = 2,062)										Male infants (n = 1,044)										Female infants (n = 1,018)									
Compound (carbon chain length)	n	Crude			Adjusted ^a			n	Crude			Adjusted ^b			n	Crude			Adjusted ^b										
		OR ^c	(95% CI) ^d		OR ^c	(95% CI) ^d			OR ^c	(95% CI) ^d		OR ^c	(95% CI) ^d			OR ^c	(95% CI) ^d		OR ^c	(95% CI) ^d									
PFHxS (C6)																													
Quantile 1	107	1.00			1.00			57	1.00			1.00			40	1.00			1.00										
Quantile 2	92	0.84	(0.61, 1.14)		0.82	(0.60, 1.13)		53	0.76	(0.50, 1.15)		0.75	(0.49, 1.13)		40	0.97	(0.60, 1.56)		0.90	(0.55, 1.47)									
Quantile 3	80	0.70	(0.51, 0.97)		0.69	(0.50, 0.95)		50	0.55	(0.35, 0.84)		0.55	(0.35, 0.85)		40	0.96	(0.60, 1.55)		0.93	(0.57, 1.52)									
Quantile 4	88	0.80	(0.58, 1.09)		0.79	(0.57, 1.08)		52	0.78	(0.52, 1.18)		0.78	(0.51, 1.19)		35	0.83	(0.50, 1.35)		0.82	(0.49, 1.36)									
<i>p</i> for trend			0.085			0.080				0.107			0.118				0.473			0.497									
PFOS (C8)																													
Quantile 1	94	1.00			1.00			59	1.00			1.00			41	1.00			1.00										
Quantile 2	99	1.06	(0.77, 1.45)		1.06	(0.77, 1.46)		53	0.96	(0.62, 1.48)		1.00	(0.64, 1.55)		50	1.18	(0.75, 1.88)		1.13	(0.71, 1.81)									
Quantile 3	90	0.94	(0.69, 1.30)		0.93	(0.67, 1.29)		56	1.32	(0.87, 1.99)		1.33	(0.87, 2.04)		27	0.58	(0.34, 0.97)		0.56	(0.32, 0.94)									
Quantile 4	84	0.87	(0.63, 1.20)		0.89	(0.64, 1.24)		44	0.93	(0.60, 1.44)		0.98	(0.63, 1.53)		37	0.81	(0.50, 1.32)		0.84	(0.51, 1.38)									
<i>p</i> for trend			0.311			0.372				0.833			0.706				0.093			0.124									
PFOA (C8)																													
Quantile 1	100	1.00			1.00			50	1.00			1.00			42	1.00			1.00										
Quantile 2	102	1.02	(0.75, 1.39)		1.03	(0.75, 1.41)		58	1.05	(0.70, 1.59)		1.11	(0.73, 1.69)		40	0.97	(0.60, 1.56)		0.98	(0.60, 1.58)									
Quantile 3	90	0.88	(0.64, 1.20)		0.86	(0.62, 1.19)		50	0.76	(0.49, 1.17)		0.76	(0.49, 1.19)		43	1.03	(0.65, 1.65)		0.99	(0.61, 1.61)									
Quantile 4	75	0.71	(0.51, 0.98)		0.72	(0.51, 1.00)		54	0.73	(0.47, 1.13)		0.75	(0.48, 1.18)		30	0.67	(0.40, 1.11)		0.65	(0.39, 1.09)									
<i>p</i> for trend			0.025			0.032				0.071			0.092				0.184			0.144									
PFNA (C9)																													
Quantile 1	97	1.00			1.00			54	1.00			1.00			50	1.00			1.00										
Quantile 2	82	0.97	(0.70, 1.34)		0.97	(0.70, 1.35)		44	1.54	(1.00, 2.40)		1.56	(1.00, 2.44)		26	0.53	(0.31, 0.87)		0.55	(0.32, 0.91)									
Quantile 3	107	0.97	(0.71, 1.31)		0.94	(0.69, 1.29)		56	1.20	(0.79, 1.83)		1.24	(0.81, 1.93)		46	0.77	(0.49, 1.19)		0.74	(0.47, 1.16)									
Quantile 4	81	0.80	(0.58, 1.11)		0.77	(0.55, 1.08)		58	1.01	(0.65, 1.57)		0.96	(0.61, 1.52)		33	0.62	(0.38, 1.00)		0.63	(0.38, 1.02)									
<i>p</i> for trend			0.224			0.145				0.756			0.643				0.127			0.122									
PFDA (C10)																													
Quantile 1	104	1.00			1.00			43	1.00			1.00			47	1.00			1.00										
Quantile 2	86	0.80	(0.58, 1.10)		0.80	(0.58, 1.10)		63	1.00	(0.66, 1.52)		1.03	(0.67, 1.59)		32	0.60	(0.37, 0.98)		0.60	(0.36, 0.97)									
Quantile 3	86	0.80	(0.58, 1.09)		0.78	(0.57, 1.08)		49	0.90	(0.59, 1.38)		0.92	(0.59, 1.43)		36	0.69	(0.43, 1.11)		0.68	(0.41, 1.09)									
Quantile 4	91	0.85	(0.62, 1.17)		0.85	(0.62, 1.17)		57	0.94	(0.62, 1.44)		0.93	(0.60, 1.44)		40	0.77	(0.48, 1.22)		0.78	(0.49, 1.25)									
<i>p</i> for trend			0.334			0.331				0.685			0.640				0.360			0.398									
PFUnDA (C11)																													
Quantile 1	100	1.00			1.00			47	1.00			1.00			51	1.00			1.00										
Quantile 2	88	0.85	(0.62, 1.17)		0.82	(0.60, 1.14)		57	1.21	(0.79, 1.85)		1.20	(0.78, 1.86)		30	0.54	(0.33, 0.88)		0.52	(0.31, 0.85)									
Quantile 3	93	0.91	(0.67, 1.25)		0.87	(0.63, 1.19)		51	1.16	(0.75, 1.80)		1.12	(0.71, 1.77)		43	0.72	(0.46, 1.12)		0.67	(0.42, 1.06)									
Quantile 4	86	0.83	(0.60, 1.14)		0.78	(0.56, 1.08)		57	1.18	(0.77, 1.81)		1.16	(0.75, 1.81)		31	0.54	(0.33, 0.88)		0.50	(0.30, 0.81)									
<i>p</i> for trend			0.341			0.185				0.526			0.607				0.036			0.016									
PFDoDA (C12)																													

^bAdjusted for all the covariates except infant gender.^dCI: confidence interval.^dCI: confidence interval.