Title	Variability of K-40 isotopic composition in forest soils under different environmental conditions
Author(s)	Fujiyoshi, Ryoko; Takekoshi, Naoki; Okamoto, Kazumasa
Citation	Journal of Radioanalytical and Nuclear Chemistry, 299(3), 1365-1371 https://doi.org/10.1007/s10967-013-2797-z
Issue Date	2014-03
Doc URL	http://hdl.handle.net/2115/56346
Type	article (author version)
File Information	10967_2013_2797_Author(changed by Fujiyoshi).pdf

[Instructions for use](#)

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.springerlink.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

Please note: Images will appear in color online but will be printed in black and white.

ArticleTitle	Variability of ^{40}K isotopic composition in forest soils under different environmental conditions	
Article Sub-Title		
Article CopyRight	Akadémiai Kiadó, Budapest, Hungary (This will be the copyright line in the final PDF)	
Journal Name	Journal of Radioanalytical and Nuclear Chemistry	
Corresponding Author	Family Name	Fujiyoshi
	Particle	
	Given Name	Ryoko
	Suffix	
	Division	Faculty of Engineering
	Organization	Hokkaido University
	Address	060-8628, Sapporo, Japan
	Email	fuji@eng.hokudai.ac.jp
Author	Family Name	Takekoshi
	Particle	
	Given Name	Naoki
	Suffix	
	Division	Faculty of Engineering
	Organization	Hokkaido University
	Address	060-8628, Sapporo, Japan
	Email	
Author	Family Name	Okamoto
	Particle	
	Given Name	Kazumasa
	Suffix	
	Division	Faculty of Engineering
	Organization	Hokkaido University
	Address	060-8628, Sapporo, Japan
	Email	
Schedule	Received	1 August 2013
	Revised	
	Accepted	
Abstract	Radiopotassium isotopic composition ($^{40}\text{K}/\text{K}$, %) of several forest soils did not show a constant value of generally known 1.17×10^{-2} %, but they were varied significantly from 0.4×10^{-2} to 1.3×10^{-2} % at different locations under different environmental conditions. Surface portion of a soil (2–4 cm in depth) gave always lower $^{40}\text{K}/\text{K}$ values compared with those of deeper soil layer (35–40 cm in depth). Ion exchange of K^+ with NH_4^+ did not affects the $^{40}\text{K}/\text{K}$ value in any soils, which revealed with chemical leaching experiments in the laboratory. Some plant species showed much lower $^{40}\text{K}/\text{K}$ values than those in the surface soil. Possible reasons for varying $^{40}\text{K}/\text{K}$ values obtained in this study may result from a dynamic behavior of potassium in soil, probably due to biological activity including root uptake and decomposing soil organic matter by microorganisms in the forest floor.	

Keywords (separated by '-') ⁴⁰K isotopic composition - Depth distribution - Carbonaceous soil - Volcanic soil - Soil organic matter

Footnote Information

Variability of ^{40}K isotopic composition in forest soils under different environmental conditions

Ryoko Fujiyoshi · Naoki Takekoshi ·
Kazumasa Okamoto

Received: 1 August 2013
© Akadémiai Kiadó, Budapest, Hungary 2013

Abstract Radiopotassium isotopic composition ($^{40}\text{K}/\text{K}$, %) of several forest soils did not show a constant value of generally known 1.17×10^{-2} %, but they were varied significantly from 0.4×10^{-2} to 1.3×10^{-2} % at different locations under different environmental conditions. Surface portion of a soil (2–4 cm in depth) gave always lower $^{40}\text{K}/\text{K}$ values compared with those of deeper soil layer (35–40 cm in depth). Ion exchange of K^+ with NH_4^+ did not affect the $^{40}\text{K}/\text{K}$ value in any soils, which revealed with chemical leaching experiments in the laboratory. Some plant species showed much lower $^{40}\text{K}/\text{K}$ values than those in the surface soil. Possible reasons for varying $^{40}\text{K}/\text{K}$ values obtained in this study may result from a dynamic behavior of potassium in soil, probably due to biological activity including root uptake and decomposing soil organic matter by microorganisms in the forest floor.

Keywords ^{40}K isotopic composition · Depth distribution · Carbonaceous soil · Volcanic soil · Soil organic matter

Introduction

Potassium (K) is an essential nutrient for plant growth and development. An extreme nutrition state was observed in a hilly region of tropical montane forest in northern Peru, where plants utilize a very small amount of K present only in the uppermost portion of the soil underlying acidic sand rock [1]. Potassium was supposed to circulate between

plants and topsoil with high (micro) biological activity. Likens et al. [2] summarized their long-term investigation of potassium and the environment as the biogeochemistry of potassium at Hubbard Brook in New Hampshire, USA. According to them, turnover rate for K (exchangeable K) in forest floor and mineral soil are relatively fast, 0.40 and 3.4 year, respectively. In spite of large quantities of K in abiotic and biotic pools and short residence time, K is the slowest to recover to pre-disturbance conditions such as clear-cutting and removal of harvest products compared with other major elements.

Potassium is believed to distribute rather homogeneously in soil within a horizon of interest [3]. Chemical speciation of K in soil revealed labile potassium (K^+) to be predominant at a depth of 10 cm under temperate coniferous stands with different tree ages in Tharndt, Germany [4]. The result is reasonably explained for the plant roots to uptake nutrients easily from the surface portion abundant in bio-available K (K^+ and ion exchangeable K). With respect to the behavior of K in plants, Szczesba et al. [5] reviewed K^+ transport in plants in view of plant molecular biology and physiology. They showed a wide variety of K^+ transport systems and catalyzing K^+ uptake across a wide spectrum of external concentrations, mediating K^+ movement within the plants as well as its efflux into the environment. There are many factors affecting K^+ transport in plants, such as external K^+ supply, the presence of other ions in the root environment and plant stresses. It is therefore important to elucidate dynamic properties of potassium in soil, which is closely related to the transportation in plants.

Naturally occurring potassium consists of stable isotopes of ^{39}K and ^{41}K , and a radioactive isotope ^{40}K with their isotopic composition (atom %) of 93.2581, 6.7302 and 0.0117 %, respectively. With a half-life of 1.25×10^9 year,

R. Fujiyoshi (✉) · N. Takekoshi · K. Okamoto
Faculty of Engineering, Hokkaido University,
Sapporo 060-8628, Japan
e-mail: fuji@eng.hokudai.ac.jp

^{40}K is changing by beta emission to ^{40}Ca (89 %) and by orbital electron capture to ^{40}Ar (11 %). In our previous investigation using ^{40}K as a natural radiotracer in soil, its distribution showed no smooth profile with depth within the same soil horizon. Radiopotassium isotopic composition ($^{40}\text{K}/\text{K}$) calculated from sets of data on ^{40}K activity concentration and total K content in soil were not constant (0.0117 %), but varied to a great extent depending on environmental factors including geology of the sites [1]. Considering dynamic behavior of potassium in soil–plant systems, $^{40}\text{K}/\text{K}$ values in soil may change during uptake and transport processes of the nutrients. Isotope fractionation of light elements like H, C, N and O occurs under various environmental conditions. Several reports have recently been published on isotope fractionation of heavy metals in soils, in which stable isotope fractionation of chromium, copper, zinc, cadmium and mercury in contaminated soils are useful tracers for evaluating hydrological and biogeochemical processes and sources of the contaminants [6–9]. However, there is little data on $^{40}\text{K}/\text{K}$ values in forest soils in the literature in spite that only ^{40}K activity has been measured extensively for monitoring and assessing external natural radiation dose, and for determining potassium content in soil so far [3, 10, 11].

This study aims at evaluating ^{40}K isotopic composition ($^{40}\text{K}/\text{K}$) in forest soils under different environmental conditions in Tomakomai (Hokkaido, Japan) and in Slovenia with non-destructive techniques of gamma spectrometry and of energy-dispersive X-ray fluorescence spectrometry (EDXRF), and at elucidating possible reasons for its inhomogeneous distribution in soil, and hopefully proposing $^{40}\text{K}/\text{K}$ values as a potential tracer to elucidate biogeochemical cycling of potassium in soil–plant systems.

Materials and methods

Description of the sites and soil sampling

Location map of the Tomakomai Experimental Forest of Hokkaido University in Tomakomai, Japan ($42^{\circ}40'\text{N}$, $141^{\circ}36'\text{N}$, 27 km^2 in total area) was shown in our previous paper [12]. Annual mean temperature and precipitation in the site are 6.4°C and $1,450\text{ mm}$, respectively. The ground surface of hilly site with altitude of $5\text{--}90\text{ m a.s.l.}$ was covered with volcanic ash derived from an active Mt. Tarumae which erupted 330 years ago. As shown in Fig. 1, a thin organic layer exists only on the uppermost part of the soil down to a depth of 6 cm from the ground surface, which then appears several laminar structures of volcanic ash down to a depth of 1 m. Each layer corresponds to past eruption of the volcano. About 100 tree species and 300 herbaceous species were recorded in the Tomakomai

Fig. 1 Cross sectional view of the volcanic soil in Tomakomai Experimental Forest, Tomakomai, Japan. Thin organic layer appeared only on the top 5 cm in depth. Several laminar layers of coarse volcanic ash were inter-located in the profile. The soil showed high permeability and low clay content (8 % in top soil and 0.5 % at a depth of 30 cm). The depth of the pit is 100 cm in depth

Experimental Forest in 1916 [13]. Predominant vegetation of the study site is deciduous broad-leaved stands like Mongolian oak (*Quercus crispula*), Painted maple (*Acer mono Maxim*) and Linden (*Tilia japonica*). The site was categorized to be high tree density ($>70\%$ canopy occupancy) and high tree height ($>16\text{ m}$ in mean height). The forest floor was covered with various plant species, with a lot of fallen leaves (twigs) and with snow depending on the season of the year.

Soil sampling was carried out in June 2009 and in November 2010. Each sample ($\sim 500\text{ g}$ in weight) was collected carefully at $2\text{--}5\text{ cm}$ intervals from the uppermost (organic layer) to a depth of about 40 cm (volcanic ash layer of different grain sizes). The samples dried at 110°C for 24 h were sieved with a stainless filter of 2 mm in pore size, and stored in plastic bottles for further analyses.

Other sites investigated in this study were located in Slovenia, where three major landscape regions and four provinces are lying; Alpine-High Nordic (Alpine province), Euro-Siberian–North American (Central European and Illyrian provinces) and Mediterranean (Adriatic province) regions [14]. The predominant forest vegetation cover at each site was oak (*Quercus pubescens*, *Quercus petraea*) and beech (*Fagus sylvatica*) trees and some pine trees (*Pinus sylvestris*) in Žirovski vrh (P1), beech trees (*Fagus sylvatica*) in Idrija (P2), beech (*Fagus sylvatica*) and pine (*Pinus sylvestris*) trees in Kočevski rog (P3), pine trees (*Pinus sylvestris*) in Pohorje (P4), beech (*Fagus sylvatica*)

148 and oak (*Quercus pubescens*, *Quercus petraea*) trees in
149 Gorišnica (P5) and beech (*Fagus sylvatica*) and pine (*Pinus*
150 *syvestris*) trees in Rakitna (P6). All the sites were cate-
151 gorized to be high tree density (>70 % canopy occupancy)
152 and high tree height (>16 m in mean height). Undergrowth
153 was not dense in any sites except for that in Žirovski vrh
154 where several types of shrubs grew. All the soils in
155 Slovenia investigated in this study were collected in Sep-
156 tember 2009.

157 Soil properties

158 Table 1 shows the results of some soil properties (density,
159 humidity, pH and soil organic matter) in Tomakomai and
160 in Slovenian forest sites [15, 16]. Details of the experi-
161 mental procedures were described previously [12].

162 Activity measurements of ^{40}K with a HPGe detection 163 system

164 Gamma spectrometry with two types of HPGe detectors
165 (GEM-25185P, GMX10P, AMETEC, USA) were applied
166 to evaluate activity concentration of ^{40}K in soil. Standard
167 reference materials (IAEA 327 and IAEA 444) were used
168 for the detector efficiency calibration with the same
169 geometry under identical operating conditions. Energy and
170 efficiency calibrations were periodically carried out as well
171 as background check. In order to evaluate reliability of the
172 ^{40}K activity measurement, a standard addition procedure
173 was applied, in which aliquots of 2 M KCl aqueous solu-
174 tion was added to a definite amount of reference soils to
175 evaluate increase in counting.

176 Determination of potassium in soil

177 Energy dispersive X-ray fluorescence spectrometry (JSX-
178 3220, JEOL, Japan) was used for determining total K
179 content of the soil. After energy calibration of the instru-
180 ment with a Rh checking source, calibration curve of K
181 was prepared with several standard soil samples of known
182 K content. X-ray energy spectra of soil samples show some
183 overlapping interference from the Ca ($\text{K}\alpha$) peak on that of
184 K ($\text{K}\alpha$) from 3.4 to 3.5 keV. The energy range for counting
185 was set from 3.14 to 3.32 keV in this study. Potassium in
186 soil was determined with a calibration technique using
187 standard soils of known K content (JSO-1, National Insti-
188 tute of Advanced Industrial Science and Technology,
189 Japan).

190 Chemical leaching of potassium in selected samples

191 Change in $^{40}\text{K}/\text{K}$ value was investigated in selected sam-
192 ples including reagent grade clay minerals using a chemical

leaching technique with 1 M ammonium acetate aqueous
solution or 1 M HCl. An aliquot of each chemical reagent
was added to a soil suspension (5 cm^3 per 1 g soil sample
in dry weight), which was stirred for 24 h at room tem-
perature. The suspension was filtered to collect the residual
portion, which was then used for ^{40}K and total K deter-
mination. The samples used for this treatment were both
surface and deeper portions from Slovenia (P1, P2, P4, P5
and P6), and those from Tomakomai (P1). The leaching
experiment was also conducted on several reagent grade
clay minerals (bentonite, kaolinite and vermiculite) pur-
chased from the Wako Pure Chemical Industries Ltd.,
Japan as references.

Results and discussion

Figure 2 shows depth distribution profiles of $^{40}\text{K}/\text{K}$ ratio in
forest soils collected from Slovenia (Fig. 2a) and from
Tomakomai (Fig. 2b). Vertical lines in the figures denote
the $^{40}\text{K}/\text{K}$ value of $1.17 \times 10^{-2} \%$ with its uncertainty
including systematic error determined in our laboratory
[17]. As shown in Fig. 2a, $^{40}\text{K}/\text{K}$ values were not constant
among soils in different locations and at different depths
within the same horizon of interest in Slovenia. Lower $^{40}\text{K}/\text{K}$
values (0.75×10^{-2} to $0.78 \times 10^{-2} \%$) appeared in the
surface portion (down to a depth of about 5 cm) compared
with those (0.86×10^{-2} to $1.07 \times 10^{-2} \%$) in deeper part
(lower than 30 cm in depth) of the soil independent of the
location. Similar $^{40}\text{K}/\text{K}$ depth distribution profiles were
obtained in other locations of different geology [17].
Results in the carbonaceous soil in Rakitna (P6) showed
high and variable $^{40}\text{K}/\text{K}$ values with two peaks appearing
about at 4–6 and 20–22 cm in depth, which may reflect
different soil properties such as density and pH [15].

In Tomakomai two types of soil horizon with different
 $^{40}\text{K}/\text{K}$ ratio appeared on the profile, as shown in Fig. 2b.
They are about 0.50×10^{-2} to 0.66×10^{-2} and $0.7 \times$
 10^{-2} to $1.0 \times 10^{-2} \%$ in the upper (<10 cm) and lower
(>20 cm) portions of the soil, respectively, which reflects a
boundary of distinct structures existing at a depth of about
15 cm as shown in Fig. 1. Depth distribution profiles of
soil organic matter (SOM) support the presence of this
boundary layer below which SOM becomes quite low
proportion (SOM <5 %) as shown in Table 1. The soil in
Tomakomai is categorized to immature volcanic ash
resulting in a specific $^{40}\text{K}/\text{K}$ value shown in Fig. 2b.

In order to find possible factors (and processes) affecting
 $^{40}\text{K}/\text{K}$ values in soil, they were compared with some soil
properties. As summarized in Table 1, humidity and SOM
correlate to each other depending on the soil depth
regardless of the sites, since highest values on both prop-
erties generally appear at the surface portion, and they

Table 1 Properties of soils (dry density, humidity, pH and the amount of soil organic matter) investigated in Tomakomai (Japan) and Slovenian forest sites

Soil depth (cm)	Tomakomai experimental forest site (coarse-textured volcanic soil)							
	Point 1				Point 3			
	pH	Humidity (%)	SOM (%)	Density (g/cm ³)	pH	Humidity (%)	SOM (%)	Density (g/cm ³)
0–2	4.88	56.8	20.0	1.27	–	55.0	18.9	1.39
2–4	4.81	49.0	15.4	1.38	–	46.8	13.7	1.12
4–6	4.93	33.1	6.2	1.51	–	39.9	10.5	1.53
6–8	5.12	25.1	3.1	1.76	–	33.4	8.8	–
8–10	5.27	19.0	1.8	1.67	–	27.5	5.5	–
10–12	5.52	18.7	1.3	1.76	–	18.0	2.5	–
12–15	5.42	21.5	1.6	1.98	–	16.7	1.4	–
15–20	5.36	22.3	0.7	1.66	–	21.0	1.6	–
20–25	5.58	19.9	0.5	1.86	–	22.4	–	–
25–30	–	19.7	0.6	1.76	–	23.6	–	–
30–35	–	21.6	0.6	1.67	–	21.8	–	–
35–40	5.4	23.1	1.1	1.68	5.04	20.6	0.8	1.67

Depth (cm)	Point 2				Point 4			
	pH	Humidity (%)	SOM (%)	Density (g/cm ³)	pH	Humidity (%)	SOM (%)	Density (g/cm ³)
0–2	5.2	57.3	18.2	1.36	5.3	43.8	10.5	1.38
2–4	5.33	45.4	10.0	1.40	5.23	–	4.4	1.39
4–6	5.32	35.4	7.0	1.67	5.16	18.0	2.2	1.40
6–8	5.47	25.7	3.8	1.76	4.95	15.1	1.1	1.44
8–10	5.64	18.9	2.3	1.67	5.04	17.1	1.2	1.51
10–12	–	17.4	1.3	1.86	5.04	22.7	1.2	1.51
12–15	5.61	21.1	1.5	–	4.94	22.2	1.8	1.58
15–20	–	24.4	1.5	1.75	5.14	22.6	0.9	1.58
20–25	–	25.5	1.0	1.62	–	–	1.0	1.30
25–30	–	21.2	0.7	1.74	–	–	0.7	1.67
30–35	–	20.1	0.6	1.76	–	–	–	–
35–40	5.78	21.4	0.6	1.86	–	–	0.6	1.64

Soil Depth (cm)	Slovenian forest sites ^a							
	Point 1 (Žirovski vrh) siliceous soil				Point 2 (Idria) carbonaceous soil			
	pH	Humidity (%)	SOM (%)	Density (g/cm ³)	pH	Humidity (%)	SOM (%)	Density (g/cm ³)
0–2	3.96	23.1	15.0	0.98	6.62	34.9	26.6	0.91
2–4	3.81	23.9	12.5	0.97	6.85	34.7	30.0	0.95
4–6	3.67	27.1	12.5	0.87	6.67	32.2	29.2	0.95
6–8	3.71	27.3	12.7	0.91	6.86	33.3	29.1	0.98
8–10	3.77	27.2	12.9	0.86	6.88	31.5	29.0	1.05
10–12	3.51	26.8	11.5	0.93	6.87	30.3	27.3	1.06
12–15	3.56	26.0	11.6	0.94	6.26	27.3	21.8	1.17
15–20	3.68	26.0	11.0	0.95	6.47	24.8	17.7	1.24
20–25	3.68	24.8	9.5	1.03	7.00	20.6	13.4	1.24
25–30	3.71	23.5	10.2	1.01	6.92	18.8	13.8	1.25
30–35	3.63	23.4	8.5	1.03	6.67	17.0	10.4	1.26
35–40	3.65	13.1	7.8	1.07	6.85	13.1	8.5	1.29

Table 1 continued

Depth (cm)	Point 3 (Kočevski rog) carbonaceous soil				Point 4 (Pohorje) siliceous soil			
	pH	Humidity (%)	SOM (%)	Density (g/cm ³)	pH	Humidity (%)	SOM (%)	Density (g/cm ³)
0–2	5.86	22.9	22.5	1.02	3.48	21.3	19.2	0.96
2–4	–	–	–	–	3.62	21.4	13.6	1.06
4–6	6.09	21.5	17.7	1.17	3.69	21.1	12.4	0.94
6–8	–	–	–	–	3.70	21.4	11.9	1.03
8–10	–	–	–	–	3.75	21.2	11.7	1.01
10–12	6.02	21.8	16.2	1.24	3.69	20.7	10.3	0.92
12–15	–	–	–	–	3.76	19.8	9.5	1.07
15–20	6.13	21.5	15.9	1.13	3.85	19.8	9.2	1.06
20–25	6.41	20.1	14.2	1.19	3.91	19.6	8.5	1.11
25–30	6.70	20.0	14.8	1.21	3.92	20.6	8.9	1.09
30–35	–	–	–	–	3.96	28.4	8.3	1.11
35–40	–	–	–	–	4.02	17.9	8.3	1.07

Depth (cm)	Point 5 (Gorišnica) fluvial sediments				Point 6 (Rakitna) carbonaceous soil			
	pH	Humidity (%)	SOM (%)	Density (g/cm ³)	pH	Humidity (%)	SOM (%)	Density (g/cm ³)
0–2	3.13	14.6	8.6	1.04	6.44	39.3	37.7	1.07
2–4	3.33	14.6	8.1	1.02	6.46	39.0	34.1	1.07
4–6	3.40	10.1	7.1	1.14	6.77	38.5	29.6	1.11
6–8	3.44	10.1	7.1	1.16	6.81	36.9	26.8	1.04
8–10	3.52	10.2	6.6	1.14	6.94	36.0	26.2	1.11
10–12	3.55	6.1	7.2	1.07	6.96	33.1	24.5	1.07
12–15	3.82	11.2	6.7	1.18	6.56	29.4	22.6	1.11
15–20	3.92	15.2	7.2	1.07	6.85	30.8	21.2	1.04
20–25	4.07	17.2	7.5	1.20	6.94	29.6	25.4	1.07
25–30	4.09	16.7	7.9	1.18	6.99	29.4	25.3	1.07
30–35	4.23	10.4	9.09	1.18	–	–	–	–
35–40	4.31	8.4	10.9	1.25	–	–	–	–

– No data

^a Cited from the reference [16]

decrease with increasing soil depth. Also in the table, carbonaceous surface soils in Slovenia (Points 2, 3 and 6) contain higher SOM (30–40 %) than those (<20 %) in siliceous soils (Points 1, 4 and 5). Depth distribution of ⁴⁰K/K values (Fig. 2a) showed similar profiles to those of SOM, which gives positive correlations between them in all the Slovenian soils investigated in this study.

With respect to immature volcanic soil in Tomakomai, good correlations also exist between humidity and SOM. However, different patterns of ⁴⁰K/K–SOM relations to those of Slovenian soils were obtained, in which lower ⁴⁰K/K values appeared only at the uppermost portions with low SOM. These findings suggest that the presence of SOM in early diagenetic stage (or possible biological activity) may affect ⁴⁰K/K values to a great extent. The presence of soil organic matter and biological activity were suggested to play an important role on ⁴⁰K/K distribution profiles in

Peruvian surface soil [17]. Further investigation is necessary to elucidate biological effects on soil ⁴⁰K/K dynamics by monitoring various soil components in the forest floor for a long time.

Figure 3 shows the relations between ⁴⁰K/K values and soil pH in Slovenian soils. Positive correlations (correlation coefficient $r > 0.9$) exist in the case of siliceous soil types (P4 and P5) in which a difference in 1 pH unit corresponds to an increase of 0.002 to 0.003 % in ⁴⁰K/K value within a small pH ranged from 3 to 4.5. In carbonaceous soils at Points 2, 3 and 6, ⁴⁰K/K values vary to a great extent within a narrow pH range ($6 < \text{pH} < 7$) attributing to pH buffer effect of carbonate minerals in soil–water systems. These results indicate that K (and ⁴⁰K) in soil may be labile in hydrogen exchange reactions in soil–water systems.

Ion exchange properties of potassium (⁴⁰K and two other stable isotopes of K) with hydrogen ions (H⁺) were

Fig. 2 a Depth distribution profiles of $^{40}\text{K}/\text{K}$ value in several forest soils from Slovenia. Žirovski VRH (Point 1), Idria (Point 2), Kočevski ROG (Point 3), Pohorje (Point 4), Gorišnica (Point 5), Rakitna (Point 6). A vertical line in the figure indicates the $^{40}\text{K}/\text{K}$ value of $1.17 \times 10^{-2} \%$ with uncertainty evaluated in this study. b Depth distribution profiles of $^{40}\text{K}/\text{K}$ value from Tomakomai Experimental Forest in Tomakomai, Japan. Four points (Points 1, 2, 3 and 4) in north eastern part of the flat zone in the forest (27.15 m^2 in area) were selected in this study. A vertical line in the figure indicates the $^{40}\text{K}/\text{K}$ value of $1.17 \times 10^{-2} \%$ with uncertainty evaluated in this study

Fig. 3 Plots of $^{40}\text{K}/\text{K}$ values against soil pH in Slovenian forest soils investigated in this study. Siliceous soils at Points 1, 4 and 5 showed high positive correlation with pH, whereas scattered $^{40}\text{K}/\text{K}$ values appeared in carbonaceous soils at Points 2, 3 and 6

Fig. 4 Results of chemical leaching experiments with 1 M HCl on the surface and deeper portions of several Slovenian soils at Points 1, 2, 4, 5 and 6. Notation after each point in the figure indicates a range of depth. For example, a notation (P1 0–2) denotes to a depth range from 0 to 2 cm at Point 1. A bar on each plot shows an estimated uncertainty obtained by triplicate experimental results. All the soils except for that at deeper portion of the Point 1 showed increasing $^{40}\text{K}/\text{K}$ values with acid treatments. No consistent trends appeared for carbonaceous soils at Points 2 and 6

leaching. In contrast, small amount of ^{40}K in the soil at Point 6 disappeared resulting in lower values of the $^{40}\text{K}/\text{K}$ isotope ratio. Dissolution of carbonate minerals (calcite and/or dolomite) in the carbonaceous soils at both points appeared after the acid treatment by power X-ray diffraction analyses [16]. The results suggest that potassium is dissolved partly into the acid solution by dissolving carbonate minerals (also some other soil components) rather randomly without keeping its isotope ratio to be constant. Dissolution of carbonate minerals (calcite) and subsequent potassium leaching by the acid treatment result in ^{40}K deficiency to a greater extent in carbonaceous soils at Point 6. It should be noted that dolomite contained in carbonaceous soils in deeper portion at Points 2 and 6 was little dissolved with the acid treatment. Results of powder X-ray diffraction analyses support the findings [16].

Jalali and Kolahchi [18] investigated short-term potassium release and fixation in some calcareous soils in Hamadan province, Iran. They used a modified quantity-to-intensity experiment to evaluate exchangeable K^+ and nonexchangeable K^+ in the laboratory. They proposed a set of parameters characterizing the exchange equilibrium and the release or fixation of potassium in agricultural soils. Although their study is only concerned with agricultural soils, it is likely to compare the findings on $^{40}\text{K}/\text{K}$ values in this study to theirs.

Another factor probably affecting $^{40}\text{K}/\text{K}$ value is included in various organic matter (SOM) present in forest soils (or possibly biological activity). As shown in Fig. 2a, the $^{40}\text{K}/\text{K}$ values were higher at organic-rich surface portion (30–40 % in SOM) in carbonaceous soils at P2 and P6 in Slovenia. The result indicates that ^{40}K is enriched in the soils abundant in SOM. Our previous investigation in Sapporo gave a similar result [17]. As a possible evidence of changing $^{40}\text{K}/\text{K}$ ratios, a plant species (*Equisetum hyemale* L.) and a litter sample gave quite different $^{40}\text{K}/\text{K}$ values to those in the soil (0.9×10^{-2} to 1.3×10^{-2} %) collected in November 2010 from semi-natural woods on the campus of Hokkaido University in Sapporo, Japan. They were $0.42 (\pm 0.01) \times 10^{-2}$ and $0.59 (\pm 0.02) \times 10^{-2}$ % for the plant and litter, respectively indicating reluctant uptake of radiopotassium (^{40}K) by the plant in late autumn.

The root-fungal association in forest ecosystems is known to have a unique soil microenvironment different from the bulk soil. Arocena et al. [19] studied the spatial variations in the composition of soil solution and mineralogy of the rhizosphere as influenced by the root-fungal association of Norway spruce (*Piloderma croceum*) in the cultivated experiment. They found significantly lower amounts of K^+ and other nutrients in the *Piloderma*-colonized treatment culture (closer to 1 cm from the roots) compared to the control. They referred the result to an evidence of *Piloderma*

modified soil solution and mineralogy through acquisition of essential elements for its own survival and/or for the uptake by plant roots. Monitoring $^{40}\text{K}/\text{K}$ value in soil environment may be another indicator to evaluate bioavailability of potassium in root-fungal association in forest ecosystems.

Dynamic behavior of potassium in undisturbed forest soil environment influences its isotopic composition ($^{40}\text{K}/\text{K}$) to a great extent. Biological activity and subsequent properties of soil organic matter may play an important role on changing $^{40}\text{K}/\text{K}$ values in soil components. Further investigation should be necessary to establish $^{40}\text{K}/\text{K}$ as a potential parameter to elucidate isotope fractionation of potassium in forest soil environment.

Acknowledgments We would greatly thank Prof. Janja Vaupotič and Prof. Emeritus Ivan Kobal of Jozef Stefan Institute in Slovenia for helping field works in the forest sites in Slovenia and giving valuable background information on the geology, climate and topology etc. The authors would also appreciate Prof. Hiura and Dr. Nakaji at Tomakomai Research Station, Field Science Center for Northern Biosphere, Hokkaido University for supporting their research work in Tomakomai Experimental Forest.

References

- Fujiyoshi R, Satake Y, Sumiyoshi S, Dietz J, Zimmermann R (2009) J Radioanal Nucl Chem 279(2):509
- Likens GE, Driscoll CT, Buso DC, Siccama TG, Johnson CE, Lovett GM, Ryan DF, Fahey T, Reiners WW (1994) Biogeochem 25:61
- Albers BP, Rackwitz R, Kleinschroth S, Bunzl K (2000) In: Markert B, Friese K (eds) Trace elements-their distribution and effect in the environment. Elsevier, New York
- Fujiyoshi R, Sawamura S (2004) Sci Total Environ 320:177
- Szczerba MW, Britto DT, Kronzucker HJ (2009) J Plant Physiol 166:447
- Bergquist BA, Blum JD (2007) Sci 318(5849):417
- Chen J, Gaillardet J, Louvat P (2008) Environ Sci Tech 42(17):6494
- Foucher D, Ogrinc N, Bullen T (2009) Environ Sci Tech 43(1):33
- Caldelas C, Dong S, Araus JL, Weiss DJ (2011) J Exp Bot 62(6):2169
- Zikovskiy L, Blagoeva R (1994) J Radioanal Nucl Chem 187(6):423
- Petersen H, Wunderlich T, al Hagrey SA, Rabbel W (2012) J. Plant Nutr Soil Sci 175:651
- Fujiyoshi R, Okabayashi M, Sakuta Y, Okamoto K, Sumiyoshi T, Kobal I, Vaupotič J (2013) Environ Earth Sci. doi:10.1007/s12665-012-2203-0
- Hiura T (2005) Ecol Res 20:271
- Repe B (2004) Slovenia Str. 57
- Fujiyoshi R, Yamaguchi T, Okamoto K, Sumiyoshi T (2011) Cent Eur J Geosci 3(3):291
- Fujiyoshi R, Amano H, Sakuta Y, Okamoto K, Sumiyoshi T, Kobal I, Vaupotič J (2012) Environ Earth Sci 67(1):133
- Fujiyoshi R, Satake Y, Sumiyoshi T (2009) J Radioanal Nucl Chem 281(3):553
- Jalali M, Lolahchi Z (2007) J Plant Nutr Soil Sci 170:530
- Arocena JM, Göttlein A, Raidl S (2004) J Plant Nutr Soil Sci 167:479