Title	Regioselective aromatic C–H silylation of five-membered heteroarenes with fluorodisilanes catalyzed by iridium(i) complexes
Author(s)	Ishiyama, Tatsuo; Sato, Kazuaki; Nishio, Yukihiro; Saiki, Takeaki; Miyaura, Norio
Citation	Chemical Communications, 40, 5065-5067 https://doi.org/10.1039/b511171d
Issue Date	2005
Doc URL	http://hdl.handle.net/2115/56376
Type	article (author version)
File Information	(44) Si-Si + H-HetAr (Com).pdf

[Instructions for use](#)

Regioselective aromatic C-H silylation of five-membered heteroarenes with fluorodisilanes catalyzed by iridium(I) complexes†

Tatsuo Ishiyama,* Kazuaki Sato, Yukihiro Nishio, Takeaki Saiki and Norio Miyaoura*

Receipt/Acceptance Data [DO NOT ALTER/DELETE THIS TEXT]

Publication data [DO NOT ALTER/DELETE THIS TEXT]

DOI: 10.1039/b000000x [DO NOT ALTER/DELETE THIS TEXT]

The aromatic C-H silylation of five-membered heteroarenes with 1,2-di-*tert*-butyl-1,1,2,2-tetrafluorodisilane regioselectively proceeded at 120 °C in octane in the presence of a catalytic amount of iridium(I) complexes generated from 1/2[Ir(OMe)(COD)]₂ and 2-*tert*-butyl-1,10-phenanthroline.

Transition metal-catalyzed aromatic C-H silylation of aromatic compounds is highly attractive as a convenient, economical, and environmentally benign process for preparing synthetically useful aromatic silicon compounds.¹ Although several studies on this type of transformation with disilanes² or hydrosilanes^{2a,3} have been carried out, the protocol has been limitedly applied to the synthesis of less-reactive aromatic triorganosilicon derivatives. Recently, we found that iridium(I) complexes comprised of an air-stable, easily available 1/2[Ir(OMe)(COD)]₂ precursor and a simple, commercially available 2,2'-bipyridine (bpy) or 4,4'-di-*tert*-butyl-2,2'-bipyridine (dtbpy) ligand effectively catalyzed aromatic C-H silylation of neat arenes with 1,2-di-*tert*-butyl-1,1,2,2-tetrafluorodisilane (*t*-BuF₂Si)₂ at 120 °C to give the corresponding aryl fluorosilanes in high yields with high regioselectivities.^{4,5} Also, synthetic utility of the aryl fluorosilanes was demonstrated by their palladium-catalyzed cross-coupling⁶ with aromatic electrophiles and rhodium-catalyzed 1,4-addition⁷ to enones.⁴ As an extension of our methodology to other aromatic substrates, we describe here the iridium(I)-catalyzed aromatic C-H silylation of five-membered heteroarenes with (*t*-BuF₂Si)₂ in octane at 120 °C (Scheme 1). Thiophene and furan derivatives were exclusively silylated at the α-position by using 1/2[Ir(OMe)(COD)]₂-(2-*tert*-butyl-1,10-phenanthroline)⁸ (tbphen) catalysts. On the other hand, the reactions of 1-triisopropylsilylpyrrole and -indole catalyzed by 1/2[Ir(OMe)(COD)]₂-tbphen or -dtbpy complexes selectively yielded β-silyl products.

Scheme 1 C-H silylation of five-membered heteroarenes.

† Electronic Supplementary Information (ESI) available: experimental procedures and spectral analyses of products. See <http://www.rsc.org/suppdata/xx/b0/b000000x/>

* xxxxx@aaa.bbb.ccc

Table 1 Reaction conditions for 3-methylthiophene^a

Entry	R ¹	R ²	Yield (%) ^b	5-:4-:2- (%) ^c
1	H	H	99	66:4:30
2	Me	Me	85 ^d	81:0:19
3	<i>n</i> -Bu	<i>n</i> -Bu	65	91:0:9
4	<i>i</i> -Pr	<i>i</i> -Pr	65	93:0:7
5	<i>t</i> -Bu	<i>t</i> -Bu	0	—
6	<i>i</i> -Pr	H	91	72:0:28
7	<i>t</i> -Bu	H	40	98:0:2
8	<i>t</i> -Bu	H	96 ^e	99:0:1

^a Reactions were carried out at 120 °C for 16 h by using (*t*-BuF₂Si)₂ (1.0 mmol), 3-methylthiophene (5.0 mmol), [Ir(OMe)(COD)]₂ (0.015 mmol), ligand (0.03 mmol), and octane (6 mL). ^b GC yields based on (*t*-BuF₂Si)₂. ^c Isomer ratios were determined by ¹H NMR. ^d 8 h. ^e 10 mmol of 3-methylthiophene, 32 h.

The silylation of 3-methylthiophene (60 mmol) with (*t*-BuF₂Si)₂ (1.0 mmol) in the presence of catalysts generated from [Ir(OMe)(COD)]₂ (0.015 mmol) and dtbpy (0.03 mmol) resulted in only a 38% yield of the silylated product after 16 h at 120 °C,⁴ probably due to the high coordinating ability of sulfur atom which retards the formation of a coordinatively unsaturated iridium species active for C-H activation.⁹ This result prompted us to use an appropriate solvent inactive toward the C-H silylation. Indeed, a silylated product was obtained in 90% yield when 5.0 mmol of the substrate was diluted in 6 mL of octane. At that time, we expected formation of regioisomerically pure 5-silyl-3-methylthiophene, because previous studies on C-H silylation of alkylated arenes demonstrated that the reactions did not occur at sterically hindered *ortho* positions.⁴ However, the present reaction gave a regioisomeric mixture of 5-, 4-, and 2-silylated products in a ratio of 69:3:28, indicating that the dtbpy-ligated iridium species was insufficient to recognize the steric environment of the substrate. Thus, we examined a sterically hindered 6,6'-di-Me-bpy ligand, but the reaction resulted in only an 8% yield. Presumably, twisting between two pyridyl units arising from steric hindrance around the iridium center leads to the low activity.⁴ Based on this hypothesis,

Table 2 C-H silylation of thiophene derivatives^a

Entry	Product	Yield (%) ^b	Isomer ratio (%) ^c
1		96 ^d	5-:4-:2- = 99:0:1
2		97	5-:4-:2- = 87:2:11
3		92	5-:4-:2- = 98:0:2
4		93	5-:4-:2- = 98:0:2
5		99 ^d	5-:4-:3- = 95:5:0
6		88	5-:4-:3- = 95:5:0
7		98	2-:3- = >99:<1
8		96 ^d	2-:3- = 93:7

^a Reactions were carried out at 120 °C for 48 h by using (*t*-BuF₂Si)₂ (1.0 mmol), thiophenes (10 mmol), [Ir(OMe)(COD)]₂ (0.015 mmol), *tbphen* (0.03 mmol), and octane (6 mL). ^b GC yields based on (*t*-BuF₂Si)₂. ^c Isomer ratios were determined by ¹H NMR. ^d 32 h.

we investigated the reaction by using structurally rigid 1,10-phenanthroline (phen) derivatives as ligands for iridium (Table 1).

As we expected, phen ligands even having substituents at 2 and 9 positions displayed high reactivity. Although unsubstituted phen produced the desired 5-silylated products with 66% selectivity (Entry 1), the selectivity was improved to 81%, 91% and 93% when using 2,9-di-Me-phen, 2,9-di-*n*-Bu-phen and 2,9-di-*i*-Pr-phen, respectively (Entries 2-4). However, the use of sterically more hindered 2,9-di-*t*-Bu-phen led to no reaction (Entry 5). The result prompted us to examine monosubstituted phen derivatives. 2-*i*-Pr-phen resulted in a 72% selectivity (Entry 6), but 2-*t*-Bu-phen (*tbphen*) improved the selectivity to 98%, while its reactivity was low (Entry 7). Finally, we obtained the desired 5-silyl-3-methylthiophene in 96% yield with 99% regioselectivity when the reaction was carried out for 32 h by using 10 mmol of 3-methylthiophene (Entry 8).

Reactions of thiophene derivatives with (*t*-BuF₂Si)₂ catalyzed by the combination of 1/2[Ir(OMe)(COD)]₂ and *tbphen* at 120 °C in octane are summarized in Table 2. 3-Substituted (Entries 1-4), 2-substituted (Entries 5 and 6), benzo-fused (Entry 7), and unsubstituted (Entry 8) thiophenes were all viable substrates to produce the corresponding silylated products in excellent yields with high regioselectivities. The reactions were suitable for

substrates possessing various functional groups, such as OMe, Cl, and CO₂Me or benzylic C-H bonds¹⁰ (Entries 1-6). The regiochemistry was controlled by both electronic and steric effects. The electronegative sulfur atom caused the α-C-H bonds to be active,¹¹ and the substituents on thiophene blocked activation of the neighboring C-H bonds.⁴ The low selectivity observed in the reaction of 3-methoxythiophene may be ascribed to small steric hindrance of the MeO group (Entry 2).

Although furan derivatives also underwent smooth C-H silylation, the regiochemistry was perplexing. Reactions of 3-methylfuran and benzofuran using *dtbpy* as a ligand predominantly yielded unexpected 4-silylated and 3-silylated products, respectively. In such cases, use of a *tbphen* ligand resulted in the expected 5-silylated and 2-silylated products with 88% and 83% selectivities (Scheme 2).

Scheme 2 C-H silylation of furan derivatives.

The C-H silylation of unsubstituted pyrrole and indole by using either *dtbpy* or *tbphen* as a ligand produced complex mixtures probably due to the presence of an acidic hydrogen in the substrates, but the reactions of 1-triisopropylsilylpyrrole and -indole successfully gave silylated products. The silylation occurred only at the β-positions when using either *dtbpy* or *tbphen* because of large steric hindrance of the TIPS group (Scheme 3).

Scheme 3 C-H silylation of pyrrole derivatives.

In summary, iridium complexes comprised of 1/2[Ir(OMe)(COD)]₂ and 2-*tert*-butyl-1,10-phenanthroline were found to be efficient catalysts for the regioselective aromatic C-H

silylation of five-membered heteroarenes with 1,2-di-*tert*-butyl-1,1,2,2-tetrafluorodisilane. Further investigations to survey the scope and limitation of this C-H silylation, including that of other heteroarenes such as pyridines, as well as to elucidate the reaction mechanisms are in progress.

This work was partially supported by Grant-in-Aid for Scientific Research on Priority Areas (No. 14078101, "Reaction Control of Dynamic Complexes") from Ministry of Education, Culture, Sports, Science and Technology, Japan. T.I. thanks The Akiyama Foundation and Takeda Science Foundation for support of a part of his work.

Tatsuo Ishiyama,* Kazuaki Sato, Yukihiko Nishio, Takeaki Saiki and Norio Miyaoura*

Division of Chemical Process Engineering, Graduate School of Engineering, Hokkaido University, Sapporo 060-8628, Japan. Fax: +81 11 706 6562; Tel: +81 11 706 6562; E-mail: ishiyama@org-mc.eng.hokudai.ac.jp

Notes and references

- For a review, see: F. Kakiuchi and N. Chatani, *Adv. Synth. Catal.*, 2003, **345**, 1077.
- (a) T. Sakakura, Y. Tokunaga, T. Sodeyama and M. Tanaka, *Chem. Lett.*, 1987, 2375; (b) M. Ishikawa, S. Okazaki, A. Naka and H. Sakamoto, *Organometallics*, 1992, **11**, 4135; (c) M. Ishikawa, A. Naka and J. Ohshita, *Organometallics*, 1993, **12**, 4987; (d) N. A. Williams, Y. Uchimarui and M. Tanaka, *Chem. Commun.*, 1995, 1129; (e) A. Naka, K. K. Lee, K. Yoshizawa, T. Yamabe and M. Ishikawa, *Organometallics*, 1999, **18**, 4524; (f) N. A. Williams, Y. Uchimarui and M. Tanaka, *Dalton Trans.*, 2003, 236.
- (a) W. A. Gustavson, P. S. Epstein and M. D. Curtis, *Organometallics*, 1982, **1**, 884; (b) Y. Uchimarui, A. M. M. El Sayed and M. Tanaka, *Organometallics*, 1993, **12**, 2065; (c) K. Ezbiarsky, P. I. Djurovich, M. LaForest, D. J. Sinning, R. Zayes and D. H. Berry, *Organometallics*, 1998, **17**, 1455; (d) F. Kakiuchi, K. Igi, M. Matsumoto, N. Chatani and S. Murai, *Chem. Lett.*, 2001, 422; (e) F. Kakiuchi, K. Igi, M. Matsumoto, T. Hayamizu, N. Chatani and S. Murai, *Chem. Lett.*, 2002, 396; (f) F. Kakiuchi, M. Matsumoto, K. Tsuchiya, K. Igi, T. Hayamizu, N. Chatani and S. Murai, *J. Organomet. Chem.*, 2003, **686**, 134.
- T. Ishiyama, K. Sato, Y. Nishio and N. Miyaoura, *Angew. Chem., Int. Ed.*, 2003, **42**, 5346.
- The catalysts were also effective for aromatic C-H borylation, see: (a) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaoura, N. R. Anastasi and J. F. Hartwig, *J. Am. Chem. Soc.*, 2002, **124**, 390; (b) J. Takagi, K. Sato, J. F. Hartwig, T. Ishiyama and N. Miyaoura, *Tetrahedron Lett.*, 2002, **43**, 5649; (c) T. Ishiyama, J. Takagi, J. F. Hartwig and N. Miyaoura, *Angew. Chem., Int. Ed.*, 2002, **41**, 3056; (d) T. Ishiyama, J. Takagi, Y. Yonekawa, J. F. Hartwig and N. Miyaoura, *Adv. Synth. Catal.*, 2003, **345**, 1103; (e) T. Ishiyama, Y. Nobuta, J. F. Hartwig and N. Miyaoura, *Chem. Commun.*, 2003, 2924; (f) T. Ishiyama and N. Miyaoura, *J. Organomet. Chem.*, 2003, **680**, 3; (g) T. Ishiyama and N. Miyaoura, *Chem. Rec.*, 2004, **3**, 271; (h) K. Kurotobi, M. Miyauchi, K. Takakura, T. Marufuji and Y. Sugihara, *Eur. J. Org. Chem.*, 2003, 3663; (i) A. Datta, A. Köllhofer and H. Plenio, *Chem. Commun.*, 2004, 1508; (j) D. N. Coventry, A. S. Batsanov, A. E. Goeta, J. A. K. Howard, T. B. Marder and R. N. Perutz, *Chem. Commun.*, 2005, 2172; (k) G. A. Chotana, M. A. Rak and M. R. Smith, III, *J. Am. Chem. Soc.*, 2005, **127**, 10539; (l) For a theoretical study, see: H. Tamura, H. Yamazaki, H. Sato and S. Sakaki, *J. Am. Chem. Soc.*, 2003, **125**, 16114.
- (a) T. Hiyama and E. Shirakawa, *Top. Curr. Chem.*, 2002, **219**, 61; (b) S. E. Denmark and R. F. Sweis, *Acc. Chem. Res.*, 2002, **35**, 835.
- (a) S. Oi, M. Moro and Y. Inoue, *Organometallics*, 2001, **20**, 1036; (b) T.-S. Huang and C.-J. Li, *Chem. Commun.*, 2001, 2348; (c) K. Fagnou and M. Lautens, *Chem. Rev.*, 2003, **103**, 169.
- M. P. T. Sjögren, S. Hansson, B. Åkermarck and A. Vitagliano, *Organometallics*, 1994, **13**, 1963.
- J. J. Li and G. W. Gribble in *Tetrahedron Organic Chemistry Series*, ed. J. E. Baldwin and R. M. Williams, Pergamon, Amsterdam, 2000, vol. 20, p. 233.
- D. P. Curran in *Comprehensive Organic Synthesis*, ed. B. M. Trost, Pergamon, Oxford, 1991, vol. 4, p. 715.
- (a) L. Dong, S. B. Duckett, K. F. Ohman and W. D. Jones, *J. Am. Chem. Soc.*, 1992, **114**, 151; (b) T. Morikita, M. Hirano, A. Sasaki and S. Komiya, *Inorg. Chim. Acta*, 1999, **291**, 341; (c) S. N. Ringelberg, A. Meetsma, B. Hessen and J. H. Teuben, *J. Am. Chem. Soc.*, 1999, **121**, 6082.

Regioselective aromatic C-H silylation of five-membered heteroarenes with
fluorodisilanes catalyzed by iridium(I) complexes

Tatsuo Ishiyama,* Kazuaki Sato, Yukihiro Nishio, Takeaki Saiki and Norio Miyaoura*

Division of Chemical Process Engineering, Graduate School of Engineering, Hokkaido University, Sapporo 060-8628, Japan

Electronic Supplementary Information (ESI)

General Methods. All the experiments were carried out under a nitrogen atmosphere. ^1H and ^{13}C NMR spectra were recorded in CDCl_3 solutions using a JEOL JNM-A400II spectrometer (400 or 100 MHz) and Me_4Si or residual protiated solvent as an internal standard. High-resolution mass spectra were obtained on a JEOL JMS-DX303. GC analyses were performed on a Hitachi G-3500 instrument equipped with a glass column (OV-101 on Uniport B, 2 m). 1,2-Di-*tert*-butyl-1,1,2,2-tetrafluorodisilane,¹ methyl 3-thiophenecarboxylate,² 1-triisopropylsilylpyrrole,³ 1-triisopropylsilylindole,⁴ $[\text{Ir}(\text{OMe})(\text{COD})]_2$,⁵ 2,9-dibutyl-1,10-phenanthroline,⁶ 2,9-di-*tert*-butyl-1,10-phenanthroline,⁶ and 2-*tert*-butyl-1,10-phenanthroline⁷ were synthesized by the reported procedures. 2,9-Diisopropyl-1,10-phenanthroline and 2-isopropyl-1,10-phenanthroline were prepared by the methods similar to those for 2,9-di-*sec*-butyl-1,10-phenanthroline⁸ and 2-*sec*-butyl-1,10-phenanthroline,⁹ respectively. Octane and hetroarenes were purified by distillation from appropriate drying agents. All of other compounds were used as received.

2,9-Diisopropyl-1,10-phenanthroline. The purity determined by NMR analyses: > 95%; ^1H NMR δ 1.47 (d, 12 H, $J = 7.1$ Hz), 3.50-3.61 (m, 2 H), 7.54 (d, 2 H, $J = 8.3$ Hz), 7.68 (s, 2 H), 8.14 (d, 2 H, $J = 8.3$ Hz); ^{13}C NMR δ 22.89, 37.29, 120.07, 125.43, 127.25, 136.37, 145.24, 167.84; exact mass calcd for $\text{C}_{18}\text{H}_{20}\text{N}_2$ 264.1626, found 264.1640.

2-Isopropyl-1,10-phenanthroline. The purity determined by NMR analyses: > 95%; ^1H NMR δ 1.46 (d, 6 H, $J = 7.1$ Hz), 3.63-3.73 (m, 1 H), 7.59 (d, 1 H, $J = 4.4$ Hz), 7.61 (d, 1 H, $J = 4.6$ Hz), 7.72 (d, 1 H, $J = 8.8$ Hz), 7.76 (d, 1 H, $J = 8.8$ Hz), 8.20 (d, 1

H, $J = 8.3$ Hz), 8.23 (dd, 1 H, $J = 8.1$ and 1.7 Hz), 9.23 (dd, 1 H, $J = 4.3$ and 1.8 Hz); ^{13}C NMR δ 23.10, 37.63, 119.96, 122.54, 125.45, 126.36, 127.05, 128.72, 135.96, 136.53, 145.21, 146.07, 150.20, 168.50; exact mass calcd for $\text{C}_{15}\text{H}_{14}\text{N}_2$ 222.1157, found 222.1155.

General Procedure for C-H Silylation with 1,2-Di-*tert*-butyl-1,1,2,2-tetrafluorodisilane. A resealable Schlenk tube containing $[\text{Ir}(\text{OMe})(\text{COD})]_2$ (0.015 mmol) and 2-*tert*-butyl-1,10-phenanthroline (0.03 mmol) was flushed with nitrogen, and then charged with octane (6 mL), 1,2-di-*tert*-butyl-1,1,2,2-tetrafluorodisilane (1.0 mmol), and a heteroarene (10 mmol). The tube was sealed with a Teflon screwcap, and the mixture was stirred at 120 °C for 16-48 h. The product was isolated by Kugelrohr distillation to give an analytically pure sample.

(*tert*-Butyldifluorosilyl)-3-methylthiophene (Table 1 and Entry 1 in Table 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (5-silyl isomer) 1.08 (s, 9 H), 2.31 (s, 3 H), 7.30 (s, 1 H), 7.34 (s, 1 H), (4-silyl isomer) 1.05 (s, 9 H), 2.38 (s, 3 H), 7.01 (m, 1 H), 7.72 (d, 1 H, $J = 2.7$ Hz), (2-silyl isomer) 1.08 (s, 9 H), 2.42 (s, 3 H), 7.03 (d, 1 H, $J = 4.4$ Hz), 7.59 (d, 1 H, $J = 4.4$ Hz); ^{13}C NMR δ (5-silyl isomer) 14.87, 17.61 (t, $J = 15.3$ Hz), 24.60, 124.59 (t, $J = 21.1$ Hz), 129.13, 139.27, 140.49, (4-silyl and 2-silyl isomers) not assigned; exact mass calcd for $\text{C}_9\text{H}_{14}\text{F}_2\text{SiS}$ 220.0553, found 220.0559.

(*tert*-Butyldifluorosilyl)-3-methoxythiophene (Entry 2 in Table 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (5-silyl isomer) 1.10 (s, 9 H), 3.84 (s, 3 H), 6.69 (d, 1 H, $J = 1.0$ Hz), 7.17 (d, 1 H, $J = 1.2$ Hz), (4-silyl isomer) 1.06 (s, 9 H), 3.83 (s, 3 H), 6.30 (m, 1 H), 7.65 (d, 1 H, $J = 2.9$ Hz), (2-silyl isomer) 1.09 (s, 9 H), 3.88 (s, 3 H), 6.96 (d, 1 H, $J = 4.9$ Hz), 7.60 (d, 1 H, $J = 4.9$ Hz); ^{13}C NMR δ (5-silyl isomer) 17.59 (t, $J = 15.3$ Hz), 24.52, 57.86, 105.38, 124.13 (t, $J = 22.3$ Hz), 129.33, 160.26, (4-silyl and 2-silyl isomers) not assigned; exact mass calcd for $\text{C}_9\text{H}_{14}\text{OF}_2\text{SiS}$ 236.0503, found 236.0505.

(*tert*-Butyldifluorosilyl)-3-chlorothiophene (Entry 3 in Table 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (5-silyl isomer) 1.11 (s, 9 H), 7.39 (d, 1 H, $J = 0.5$ Hz), 7.52 (d, 1 H, $J = 0.5$ Hz), (2-silyl isomer, aromatic region) 7.12 (d, 1 H, $J = 4.9$ Hz), 7.68 (d, 1 H, $J = 4.9$ Hz); ^{13}C NMR δ (5-silyl isomer) 17.59 (t, $J = 14.9$ Hz), 24.45, 126.26 (t, $J = 21.9$ Hz), 127.39, 127.90, 137.96, (2-silyl isomer) not assigned; exact mass calcd for $\text{C}_8\text{H}_{11}\text{ClF}_2\text{SiS}$ 240.0007, found 240.0015.

Methyl (*tert*-butyldifluorosilyl)-3-thiophenecarboxylate (Entry 4 in Table 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (5-silyl isomer) 1.11 (t, 9 H, $J = 1.0$ Hz), 3.90 (s, 3 H), 7.98 (d, 1 H, $J = 1.0$ Hz), 8.46 (s, 1 H), (2-silyl isomer, aromatic region) 7.10 (d, 1 H, $J = 5.4$ Hz), 7.47 (d, 1 H, $J = 5.1$ Hz); ^{13}C NMR δ (5-silyl isomer) 17.53 (t, $J = 14.9$ Hz), 24.42, 51.97, 126.43 (t, $J = 21.5$ Hz), 135.29, 139.04, 139.87, 162.69, (2-silyl isomer) not assigned; exact mass calcd for $\text{C}_{10}\text{H}_{14}\text{O}_2\text{F}_2\text{SiS}$ 264.0452, found 264.0447.

(*tert*-Butyldifluorosilyl)-2-methylthiophene (Entry 5 in Table 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (5-silyl isomer) 1.07 (t, 9 H, $J = 1.0$ Hz), 2.55 (s, 3 H), 6.90-6.91 (m, 1 H), 7.37 (d, 1 H, $J = 3.4$ Hz), (4-silyl isomer) 1.05 (t, 9 H, $J = 1.0$ Hz), 2.51 (d, 3 H, $J = 1.0$ Hz), 6.90-6.91 (m, 1 H), 7.57 (d, 1 H, $J = 1.2$ Hz); ^{13}C NMR δ (5-silyl isomer) 14.98, 17.60 (t, $J = 15.7$ Hz), 24.61, 122.45 (t, $J = 21.5$ Hz), 127.12, 138.79 (t, $J = 1.7$ Hz), 148.71, (4-silyl isomer) not assigned; exact mass calcd for $\text{C}_9\text{H}_{14}\text{F}_2\text{SiS}$ 220.0553, found 220.0558.

(*tert*-Butyldifluorosilyl)-2-chlorothiophene (Entry 6 in Table 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (5-silyl isomer) 1.10 (s, 9 H), 7.07 (d, 1 H, $J = 3.7$ Hz), 7.35 (d, 1 H, $J = 3.7$ Hz), (4-silyl isomer) 1.07 (s, 9 H), 7.07 (m, 1 H), 7.59 (d, 1 H, $J = 1.2$ Hz); ^{13}C NMR δ (5-silyl isomer) 17.56 (t, $J = 15.3$ Hz), 24.44, 124.59 (t, $J = 21.9$ Hz), 127.79, 138.00, 138.29, (4-silyl isomer) not assigned; exact mass calcd for $\text{C}_8\text{H}_{11}\text{ClF}_2\text{SiS}$ 240.0007, found 240.0003.

2-(*tert*-Butyldifluorosilyl)benzo[*b*]thiophene (Entry 7 in Table 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.15 (s, 9 H), 7.39-7.45 (m, 2 H), 7.83 (s, 1 H), 7.91-7.95 (m, 2 H); ^{13}C NMR δ 17.67 (t, $J = 14.9$ Hz), 24.54, 122.28, 124.45, 124.59, 125.81, 126.26 (t, $J = 20.7$ Hz), 135.72, 139.91, 143.85; exact mass calcd for $\text{C}_{12}\text{H}_{14}\text{F}_2\text{SiS}$ 256.0554, found 256.0565.

(*tert*-Butyldifluorosilyl)thiophene (Entry 8 in Table 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (2-silyl isomer) 1.09 (s, 9 H), 7.27 (t, 1 H, $J = 4.0$ Hz), 7.59 (d, 1 H, $J = 3.4$ Hz), 7.76 (d, 1 H, $J = 4.6$ Hz), (3-silyl isomer) 1.06 (s, 9 H), 7.30 (d, 1 H, $J = 4.9$ Hz), 7.47 (m, 1 H), 7.83 (d, 1 H, $J = 1.7$ Hz); ^{13}C NMR δ (2-silyl isomer) 17.62 (t, $J = 15.3$ Hz), 24.57, 124.57 (t, $J = 21.1$ Hz), 128.38, 133.40, 138.33, (3-silyl isomer) not assigned; exact mass calcd for $\text{C}_8\text{H}_{12}\text{F}_2\text{SiS}$ 206.0397, found 206.0396.

(*tert*-Butyldifluorosilyl)-3-methylfuran (Scheme 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (5-silyl isomer) 1.10 (t, 9 H, $J = 1.0$ Hz),

2.06 (s, 3 H), 6.90 (s, 1 H), 7.48 (d, 1 H, $J = 0.7$ Hz), (4-silyl isomer) 1.07 (t, 9 H, $J = 1.0$ Hz), 2.11 (s, 3 H), 7.30 (s, 1 H), 7.58 (s, 1 H), (2-silyl isomer) 1.10 (t, 9 H, $J = 1.0$ Hz), 2.20 (s, 3 H), 6.31 (d, 1 H, $J = 1.7$ Hz), 7.60 (d, 1 H, $J = 1.7$ Hz); ^{13}C NMR δ (5-silyl isomer) 9.12, 17.29 (t, $J = 15.3$ Hz), 24.39, 119.98, 127.66 (t, $J = 1.7$ Hz), 145.62 (t, $J = 1.7$ Hz), 148.05 (t, $J = 26.5$ Hz), (4-silyl isomer) 10.24, 17.69 (t, $J = 15.7$ Hz), 24.57, 109.70 (t, $J = 20.7$ Hz), 123.29, 140.42 (t, $J = 2.5$ Hz), 151.31 (t, $J = 4.5$ Hz); exact mass calcd for $\text{C}_9\text{H}_{14}\text{OF}_2\text{Si}$ 204.0782, found 204.0783.

(*tert*-Butyldifluorosilyl)benzo[*b*]furan (Scheme 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ (2-silyl isomer) 1.16 (t, 9 H, $J = 1.0$ Hz), 7.28 (t, 1 H, $J = 7.8$ Hz), 7.39 (t, 1 H, $J = 8.3$ Hz), 7.41 (s, 1 H), 7.57 (d, 1 H, $J = 8.3$ Hz), 7.67 (d, 1 H, $J = 7.8$ Hz), (3-silyl isomer) 1.12 (t, 9 H, $J = 0.9$ Hz), 7.30 (t, 1 H, $J = 7.3$ Hz), 7.36 (t, 1 H, $J = 7.8$ Hz), 7.57 (d, 1 H, $J = 8.1$ Hz), 7.71 (d, 1 H, $J = 7.6$ Hz), 7.85 (s, 1 H); ^{13}C NMR δ (2-silyl isomer) 17.34 (t, $J = 14.5$ Hz), 24.32, 111.86, 121.49, 121.99, 123.13, 126.22, 126.49, 150.16 (t, $J = 26.5$ Hz), 158.22, (3-silyl isomer) 17.69 (t, $J = 15.3$ Hz), 24.45, 104.68 (t, $J = 21.9$ Hz), 111.51, 122.45, 123.52, 125.04, 129.47, 153.08 (t, $J = 4.1$ Hz), 155.36; exact mass calcd for $\text{C}_{12}\text{H}_{14}\text{OF}_2\text{Si}$ 240.0782, found 240.0787.

3-(*tert*-Butyldifluorosilyl)-1-triisopropylsilylpyrrole (Scheme 3). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.05 (t, 9 H, $J = 0.5$ Hz), 1.09 (d, 18 H, $J = 7.6$ Hz), 1.41-1.53 (m, 3 H), 6.49-6.50 (m, 1 H), 6.89 (s, 1 H), 7.10 (s, 1 H); ^{13}C NMR δ 11.66, 17.71, 17.71 (t, $J = 16.5$ Hz), 24.80, 106.67 (t, $J = 21.1$ Hz), 115.22 (t, $J = 2.1$ Hz), 125.53, 133.20 (t, $J = 2.9$ Hz); exact mass calcd for $\text{C}_{17}\text{H}_{33}\text{NF}_2\text{Si}_2$ 345.2120, found 345.2126.

3-(*tert*-Butyldifluorosilyl)-1-triisopropylsilylindole (Scheme 3). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.09 (t, 9 H, $J = 0.7$ Hz), 1.15 (d, 18 H, $J = 7.6$ Hz), 1.67-1.79 (m, 3 H), 7.16-7.22 (m, 2 H), 7.55 (s, 1 H), 7.56 (d, 1 H, $J = 6.8$ Hz), 7.76 (d, 1 H, $J = 7.6$ Hz); ^{13}C NMR δ 12.68, 18.00, 18.03 (t, $J = 16.5$ Hz), 24.84, 101.34 (t, $J = 21.5$ Hz), 114.01, 120.82, 121.96, 122.18, 134.67, 140.93 (t, $J = 4.1$ Hz), 141.79; exact mass calcd for $\text{C}_{21}\text{H}_{35}\text{NF}_2\text{Si}_2$ 395.2276, found 395.2277.

References

1. B. Reiter and K. Hassler, *J. Organomet. Chem.*, 1994, **467**, 21.
2. M. Pomerantz, A. S. Amarasekara and H. V. R. Dias, *J. Org. Chem.*, 2002, **67**, 6937.

3. B. L. Bray, P. H. Mathies, R. Naef, D. R. Solas, T. T. Tidwell, D. R. Artis and J. M. Muchowski, *J. Org. Chem.*, 1990, **55**, 6317.
4. P. J. Beswick, C. S. Greenwood, T. J. Mowlem, G. Nechvatal and D. A. Widdowson, *Tetrahedron*, 1988, **44**, 7325.
5. R. Uson, L. A. Oro and J. A. Cabeza, *Inorg. Synth.*, 1985, **23**, 126.
6. C. O. Dietrich-Buchecker, P. A. Marnot and J. P. Sauvage, *Tetrahedron Lett.*, 1982, **23**, 5291.
7. M. P. T. Sjögren, S. Hansson, B. Åkermark and A. Vitagliano, *Organometallics*, 1994, **13**, 1963.
8. A. J. Pallenberg, K. S. Koenig and D. M. Barnhart, *Inorg. Chem.*, 1995, **34**, 2833.
9. G.-J. ten Brink, I. W. C. E. Arends, M. Hoogenraad, G. Verspui and R. A. Sheldon, *Adv. Synth. Catal.*, 2003, **345**, 1341.

RSC Author Templates - ChemDraw (CDX) - Graphical Abstracts

The aromatic C-H silylation of five-membered heteroarenes with 1,2-di-*tert*-butyl-1,1,2,2-tetrafluorodisilane regioselectively proceeded at 120 °C in octane in the presence of a catalytic amount of iridium(I) complexes generated from 1/2[Ir(OMe)(COD)]₂ and 2-*tert*-butyl-1,10-phenanthroline.