Title	Room temperature borylation of arenes and heteroarenes using stoichiometric amounts of pinacolborane catalyzed by iridium complexes in an inert solvent
Author(s)	Ishiyama, Tatsuo; Nobuta, Yusuke; Hartwig, John F.; Miyaura, Norio
Citation	Chemical Communications, 23, 2924-2925 https://doi.org/10.1039/b311103b
Issue Date	2003
Doc URL	http://hdl.handle.net/2115/56377
Type	article (author version)
File Information	(37) B-H + H-Ar (rt) (Com).pdf

[Instructions for use](#)

Room temperature borylation of arenes and heteroarenes by stoichiometric amounts of pinacolborane catalyzed by iridium complexes in an inert solvent†

Tatsuo Ishiyama,^{a*} Yusuke Nobuta,^a John F. Hartwig^b and Norio Miyaura^{a*}

^a Division of Molecular Chemistry, Graduate School of Engineering, Hokkaido University, Sapporo 060-8628, Japan. Fax: +81-11-706-6562; Tel: +81-11-706-6562; E-mail: ishiyama@org-mc.eng.hokudai.ac.jp

^b Department of Chemistry, Yale University, P.O. Box 208107, New Haven, Connecticut 06520-8107, USA.

*This submission was created using the RSC ChemComm Template (DO NOT DELETE THIS TEXT)
(LINE INCLUDED FOR SPACING ONLY - DO NOT DELETE THIS TEXT)*

Aromatic C-H borylation of arenes and heteroarenes by stoichiometric amounts of pinacolborane was catalyzed by an iridium complex generated from 1/2[Ir(OMe)(COD)]₂ and 4,4'-di-*tert*-butyl-2,2'-bipyridine at room temperature in hexane and afforded the corresponding aryl- and heteroarylboronates in high yields with excellent regioselectivities.

Aryl- and heteroarylboron derivatives have been applied to various fields of chemistry.¹ Traditional methods for their synthesis are based on the reactions of trialkylborates with arylmagnesium or -lithium reagents derived from haloarenes, which are most common and convenient for large-scale preparations.² However, two milder catalytic methods have been developed recently. First, catalytic coupling of aryl and heteroaryl halides with tetra(alkoxo)diboron³ or di(alkoxo)borane⁴ reagents forms arylboronates without magnesium or lithium intermediates. Second, catalytic borylation of arenes and heteroarenes forms arylboronates without any halogenated arene.

Previous work has demonstrated the reaction of arenes and heteroarenes with bis(pinacolato)diboron (pin₂B₂, pin = Me₄C₂O₂) or pinacolborane (pinBH) in the presence of various transition metal complexes to form arylboronates.⁵⁻¹¹ Most of these studies have been conducted with excess of substrate, and the resulting methods have required the use of pin₂B₂ instead of pinBH to obtain high yields. No reactions of a 1:1 ratio of substrate and the readily accessible and inexpensive pinBH, instead of the expensive pin₂B₂, have been reported. We disclose the culmination of several steps toward the development of such an aromatic C-H borylation process. With modest loadings of 1/2[Ir(OMe)(COD)]₂ (COD = 1,5-cyclooctadiene) and 4,4'-di-*tert*-butyl-2,2'-bipyridine (dtbpy), the reaction of pinBH with many arenes and heteroarenes in a 1:1 ratio occurs in an inert solvent at room temperature (Scheme 1).

Scheme 1

To achieve the borylation of arenes and heteroarenes at room temperature with equimolar amounts of pinBH and substrate, several combinations of Ir(I) precursors (0.03 mmol of Ir) and ligands (0.03 mmol) were investigated as catalysts for the reaction of pinBH (1.1 mmol) with 1,3-dichlorobenzene (1.0

mmol) in hexane (6 mL) at 25 °C for 8 h. Of the precursors and ligands examined, the combination of 1/2[Ir(OMe)(COD)]₂ and dtbpy¹² efficiently catalyzed the borylation to form isomerically pure 5-boryl-1,3-dichlorobenzene in 86% yield.

The choice of catalyst precursor was crucial to observe room temperature reactions. Although the combination of 1/2[Ir(OAc)(COD)]₂ and dtbpy produced the borylated product in 42% yield after 8 h, the combination of dtbpy and either 1/2[IrCl(COD)]₂ or [Ir(COD)₂]BF₄ formed no borylated product.

The effects of steric and electronic properties of bipyridine ligands were evaluated with 1/2[Ir(OMe)(COD)]₂ as a catalyst precursor. Catalysts bearing 2,2'-bipyridine (bpy), 4,4'-di-Me-bpy, and 5,5'-di-Me-bpy displayed moderate or good reactivity, but catalysts bearing 3,3'-di-Me-bpy or 6,6'-di-Me-bpy displayed little activity. These results indicated the importance of a parallel arrangement of two pyridine rings and a relatively unhindered coordination sphere at iridium. Reactions catalyzed by complexes containing electron-rich derivatives of bpy generated more active catalysts than those containing electron-poor derivatives. Catalysts containing 4,4'-di-Me₂N-bpy produced the highest yields (88%). We evaluated reactions catalyzed by 1/2[Ir(OMe)(COD)]₂ and dtbpy for studies on reaction scope because of the high solubility of the catalyst and the commercial availability of the ligand.

Proper choice of inert solvent was also important to observe efficient borylation. The reactions were faster in non-polar solvents, such as hexane, than in more polar and coordinating solvents. The order of reactivity in different solvents was hexane > mesitylene > DME > DMF.

Reactions of equimolar amounts of pinBH with arenes and heteroarenes catalyzed by the combination of 1/2[Ir(OMe)(COD)]₂ and dtbpy at room temperature in hexane are summarized in Table 1. In contrast to the control of regioselectivity of electrophilic and nucleophilic substitution of arene by the electronic properties of substituents, the regiochemistry of C-H borylation of arene is primarily controlled by the steric effects of these substituents. Reactions occurred at C-H bonds located meta or para to a substituent in preference to those located ortho. Thus, 1,2-, 1,4-, and 1,3-dichlorobenzenes gave a single product (Entries 1-3), but the 1,4-isomer reacted slowly (Entry 2). In addition, the borylation of 1,3-disubstituted arenes containing two different substituents at the 1 and 3 positions produced isomerically pure arylboronates in excellent yields. (Entries 4-9). In the case of five-membered heteroarenes, the electronegative heteroatom causes the C-H bonds at the α-positions to be active¹³ so that the borylation of indole, benzo[b]furan, and benzo[b]thiophene selectively occurred at the α-positions to form single isomers in high yields (Entries 10-12).

Functional group tolerance of the borylation is higher than that of boronate syntheses through magnesium or lithium reagents. The reaction occurred with substrates possessing Cl, Br, I, CF₃, and OMe groups, but also the more reactive CO₂Me and

† Electronic Supplementary Information (ESI) available: experimental procedures and spectral analyses of products. See <http://www.rsc.org/suppdata/cc/b0/b000000a/>

Table 1 C-H borylation of arenes and heteroarenes^a

Entry	Product	Yield (%) ^b	Entry	Product	Yield (%) ^b
1		73% (8 h)	7		74% (2 h)
2		22% (24 h)	8		83% (1 h)
3		86% (8 h)	9		73% (24 h)
4		67% (8 h)	10		99% (0.5 h)
5		70% (24 h)	11		90% (1 h)
6		80% (8 h)	12		99% (2 h)

^a All reactions were carried out at 25 °C with pinacolborane (1.1 mmol), arene or heteroarene (1.0 mmol), [Ir(OMe)(COD)]₂ (0.015 mmol), dtbpy (0.03 mmol), and hexane (6 mL). ^b GC yields based on arenes or heteroarenes and reaction times are in parentheses.

CN. The aryl iodide and bromide underwent borylation at the C-H bond (Entries 4, 6, and 7) without carbon-halogen bond cleavage.¹⁴ Electron-withdrawing substituents activated the arene for the borylation process.

The mechanism of the catalytic borylation of arenes and heteroarenes by pinBH may proceed through an Ir(III)-Ir(V) cycle.^{10-12,15} Generation of a (pinB)₃Ir(III) intermediate^{10,11a,16} by the reaction of an Ir(I) complex with pinBH, oxidative addition of a C-H bond to the (pinB)₃Ir(III) intermediate to yield an (Ar)(H)(pinB)₃Ir(V) species, and reductive elimination of pinB-Ar from the (Ar)(H)(pinB)₃Ir(V) species to give a (H)(pinB)₂Ir(III) complex would be followed by oxidative addition of pinBH and reductive elimination of H₂ to regenerate the (pinB)₃Ir(III) intermediate.

The direct preparation of aryl- and heteroarylboronates from pinBH and the corresponding arenes or heteroarenes creates an efficient, one-pot, two-step procedure for the synthesis of unsymmetrical biaryls. The biaryl shown in Scheme 2 was produced in 91% yield by sequential generation of 5-boryl-1,3-dichlorobenzene from pinBH (1.43 mmol) and 1,3-dichlorobenzene (1.3 mmol) in hexane (2 mL), followed by reaction of the resulting arylboronate with added methyl 4-bromobenzoate (1.0 mmol), PdCl₂(dppf) (0.03 mmol), K₃PO₄ (3.0 mmol), and DMF (4 mL) at 60 °C.¹⁷

In summary, aromatic C-H borylation of arenes and heteroarenes by pinBH without excess of substrate or reagent provides a convenient, economical, and environmentally benign route to regiodefined aryl- and heteroarylboronates. Further investigations to survey the scope and limitation of this C-H borylation, including that of monosubstituted arenes, other heteroarenes, alkenes, and alkanes, as well as the application of

Scheme 2

this catalyst system to other types of C-H functionalizations are in progress.

This work was partially supported by Grant-in-Aid for Scientific Research on Priority Areas (No. 14078101, "Reaction Control of Dynamic Complexes") from Ministry of Education, Culture, Sports, Science and Technology, Japan and by the National Science Foundation (JFH). T.I. thanks Itoh Science Foundation and The Akiyama Foundation for support of a part of his work.

Notes and references

- (a) M. Vaultier and B. Carboni in *Comprehensive Organometallic Chemistry II*, ed. E. W. Abel, F. G. A. Stone and G. Wilkinson, Pergamon Press, Oxford, 1995, vol. 11, p. 191; (b) K. Ishihara and H. Yamamoto, *Eur. J. Org. Chem.*, 1999, 527; (c) S. Shinkai, M. Ikeda, A. Sugasaki and M. Takeuchi, *Acc. Chem. Res.*, 2001, **34**, 494; (d) C. D. Entwistle and T. B. Marder, *Angew. Chem. Int. Ed.*, 2002, **41**, 2927; (e) A. H. Soloway, W. Tjarks, B. A. Barnum, F.-G. Rong, R. F. Barth, I. M. Codogni and J. G. Wilson, *Chem. Rev.*, 1998, **98**, 1515; (f) W. Yang, X. Gao and B. Wang, *Med. Res. Rev.*, 2003, **23**, 346.
- A. N. Nesmeyanov and R. A. Sokolik, *Methods of Elemento-Organic Chemistry*, North-Holland, Amsterdam, 1967, vol. 1.
- (a) T. Ishiyama and N. Miayura, *J. Synth. Org. Chem., Jpn.*, 1999, **57**, 503; (b) T. Ishiyama and N. Miayura, *J. Organomet. Chem.*, 2000, **611**, 392.
- M. Murata, T. Oyama, S. Watanabe and Y. Masuda, *J. Org. Chem.*, 2000, **65**, 164.
- A review, see: T. Ishiyama and N. Miayura, *J. Organomet. Chem.*, 2003, **680**, 3.
- C. N. Iverson and M. R. Smith, III, *J. Am. Chem. Soc.*, 1999, **121**, 7696.
- H. Chen and J. F. Hartwig, *Angew. Chem. Int. Ed.*, 1999, **38**, 3391.
- (a) H. Chen, S. Schlecht, T. C. Semple and J. F. Hartwig, *Science*, 2000, **287**, 1995; (b) J.-Y. Cho, C. N. Iverson and M. R. Smith, III, *J. Am. Chem. Soc.*, 2000, **122**, 12868; (c) M. K. Tse, J.-Y. Cho and M. R. Smith, III, *Org. Lett.*, 2001, **3**, 2831.
- S. Shimada, A. S. Batsanov, J. A. K. Howard and T. B. Marder, *Angew. Chem. Int. Ed.*, 2001, **40**, 2168.
- J.-Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka, Jr. and M. R. Smith, III, *Science*, 2002, **295**, 305.
- (a) T. Ishiyama, J. Takagi, K. Ishida, N. Miayura, N. R. Anastasi and J. F. Hartwig, *J. Am. Chem. Soc.*, 2002, **124**, 390; (b) J. Takagi, K. Sato, J. F. Hartwig, T. Ishiyama and N. Miayura, *Tetrahedron Lett.*, 2002, **43**, 5649.
- T. Ishiyama, J. Takagi, J. F. Hartwig and N. Miayura, *Angew. Chem. Int. Ed.*, 2002, **41**, 3056.
- (a) L. Dong, S. B. Duckett, K. F. Ohman and W. D. Jones, *J. Am. Chem. Soc.*, 1992, **114**, 151; (b) T. Morikita, M. Hirano, A. Sasaki and S. Komiyama, *Inorg. Chim. Acta*, 1999, **291**, 341; (c) S. N. Ringelberg, A. Meetsma, B. Hessen and J. H. Teuben, *J. Am. Chem. Soc.*, 1999, **121**, 6082.
- J. D. Atwood in *Comprehensive Organometallic Chemistry II*, ed. E. W. Abel, F. G. A. Stone and G. Wilkinson, Pergamon Press, Oxford, 1995, vol. 8, p. 303.
- Theoretical studies, see: H. Yamazaki, H. Tamura, M. Sugimoto, H. Sato and S. Sakaki, *49th Symposium on Organometallic Chemistry, Japan*, 2002, PA119.
- P. Nguyen, H. P. Blom, S. A. Westcott, N. J. Taylor and T. B. Marder, *J. Am. Chem. Soc.*, 1993, **115**, 9329.
- (a) N. Miayura and A. Suzuki, *Chem. Rev.*, 1995, **95**, 2457; (b) N. Miayura, *Top. Curr. Chem.*, 2002, **219**, 11; (c) A. Suzuki and H. C. Brown, *Organic Syntheses Via Boranes*, Aldrich Chemical Company, Inc., Milwaukee, 2003, vol. 3.

Room temperature borylation of arenes and heteroarenes by stoichiometric amounts of pinacolborane catalyzed by iridium complexes in an inert solvent

Tatsuo Ishiyama,* Yusuke Nobuta and Norio Miyaoura*

Division of Molecular Chemistry, Graduate School of Engineering, Hokkaido University, Sapporo 060-8628, Japan

John F. Hartwig

Department of Chemistry, Yale University, P.O. Box 208107, New Haven, Connecticut 06520-8107, USA

Electronic Supplementary Information (ESI)

General Methods. All the experiments were carried out under a nitrogen atmosphere. ^1H and ^{13}C NMR spectra were recorded in CDCl_3 solutions using a JEOL JNM-A400II spectrometer (400 or 100 MHz) and Me_4Si or residual protiated solvent as an internal standard. High-resolution mass spectra were obtained on a JEOL JMS-DX303. GC analyses were performed on a Hitachi G-3500 instrument equipped with a glass column (OV-101 on Uniport B, 2 m). Solvents, arenes, and heteroarenes were purified by distillation from appropriate drying agents. Pinacolborane was prepared by Knochel's method¹ and purified by distillation through a Widmer column. $[\text{Ir}(\text{OMe})(\text{COD})]_2$,² $[\text{Ir}(\text{OAc})(\text{COD})]_2$,³ 4,4'-bis(*N,N*-dimethylamino)-2,2'-bipyridine,⁴ 4,4'-dimethoxy-2,2'-bipyridine,⁴ 4,4'-dichloro-2,2'-bipyridine,⁴ and 4,4'-dinitro-2,2'-bipyridine⁵ were synthesized by the reported procedures. All of other compounds were used as received.

General Procedure for Aromatic C-H Borylation by Pinacolborane (Table 1). A 25-mL flask assembled a magnetic stirring bar, a septum inlet, a condenser, and a bubbler was charged with $[\text{Ir}(\text{OMe})(\text{COD})]_2$ (0.015 mmol) and 4,4'-di-*tert*-butyl-2,2'-bipyridine (0.03 mmol), and then flushed with nitrogen. Dry hexane (6 mL), pinacolborane (1.1 mmol), and an arene or a heteroarene (1.0 mmol) were added, and the mixture was stirred at 25 °C for the period shown in Table 1. The reaction mixture was

treated with water at room temperature, extracted with benzene, washed with brine, and dried over MgSO_4 . Kugelrohr distillation in vacuo gave an analytically pure sample.

1,2-Dichloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzene (Entry 1). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.34 (s, 12 H), 7.44 (d, 1 H, $J = 7.8$ Hz), 7.60 (d, 1 H, $J = 8.1$ Hz), 7.87 (s, 1 H); ^{13}C NMR δ 24.82, 84.31, 129.98, 132.23, 133.73, 135.46, 136.53; exact mass calcd for $\text{C}_{12}\text{H}_{15}\text{BCl}_2\text{O}_2$ 272.0542, found 272.0534.

1,4-Dichloro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzene (Entry 2). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.37 (s, 12 H), 7.27 (d, 1 H, $J = 8.1$ Hz), 7.30 (dd, 1 H, $J = 8.5$ and 2.2 Hz), 7.65 (d, 1 H, $J = 2.0$ Hz); ^{13}C NMR δ 24.75, 84.46, 130.68, 131.69, 132.08, 136.02, 137.7; exact mass calcd for $\text{C}_{12}\text{H}_{15}\text{BCl}_2\text{O}_2$ 272.0542, found 272.0549.

1,3-Dichloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzene (Entry 3). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.34 (s, 12 H), 7.43 (t, 1 H, $J = 2.0$ Hz), 7.65 (d, 2 H, $J = 2.0$ Hz); ^{13}C NMR δ 24.81, 84.48, 131.05, 132.67, 134.69; exact mass calcd for $\text{C}_{12}\text{H}_{15}\text{BCl}_2\text{O}_2$ 272.0542, found 272.0566.

1-Chloro-3-iodo-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzene (Entry 4). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.34 (s, 12 H), 7.72 (d, 1 H, $J = 1.5$ Hz), 7.78 (t, 1 H, $J = 1.2$ Hz), 8.00 (s, 1 H); ^{13}C NMR δ 24.81, 84.46, 94.18, 133.71, 134.70, 139.42, 141.42; exact mass calcd for $\text{C}_{12}\text{H}_{15}\text{BClIO}_2$ 363.9899, found 363.9880.

Methyl 3-chloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (Entry 5). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.35 (s, 12 H), 3.92 (s, 3 H), 7.95 (d, 1 H, $J = 1.5$ Hz), 8.09 (t, 1 H, $J = 1.2$ Hz), 8.33 (s, 1 H); ^{13}C NMR δ 24.82, 52.30, 84.42, 131.32, 132.10, 133.70, 134.28, 138.79, 165.93; exact mass calcd for $\text{C}_{14}\text{H}_{18}\text{BClO}_4$ 296.0987, found 296.0993.

3-Bromo-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzotrifluoride (Entry 6). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.36 (s, 12 H), 7.83 (s, 1 H), 7.97 (s, 1 H), 8.10 (s, 1 H); ^{13}C NMR δ 24.82, 84.66, 122.49, 123.26 (q, $J = 272.9$ Hz), 129.83 (q, $J = 3.3$ Hz), 130.80 (q, $J = 4.1$ Hz), 131.88 (q, $J = 32.8$ Hz), 140.82; exact mass calcd for $\text{C}_{13}\text{H}_{15}\text{BBrF}_3\text{O}_2$ 350.0300, found 350.0309.

3-Bromo-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzonitrile (Entry 7). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.35 (s, 12 H), 7.85 (s, 1 H), 8.01 (s, 1 H), 8.13 (s, 1 H); ^{13}C NMR δ 24.81, 84.85, 113.77, 117.32, 122.60, 136.70, 136.74, 141.73; exact mass calcd for $\text{C}_{13}\text{H}_{15}\text{BBrNO}_2$ 307.0379, found 307.0387.

3-Trifluoromethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzonitrile (Entry 8). The purity determined by NMR and GC analyses: > 95%; ^1H NMR δ 1.37 (s, 12 H), 7.98 (s, 1 H), 8.26 (s, 2 H); ^{13}C NMR δ 24.81, 85.03, 112.96, 117.39, 123.02 (q, $J = 272.6$ Hz), 131.01 (q, $J = 4.1$ Hz), 131.30

(q, $J = 33.6$ Hz), 135.22 (q, $J = 3.3$ Hz), 141.33; exact mass calcd for $C_{14}H_{15}BF_3NO_2$ 297.1148, found 297.1153.

3-Trifluoromethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)anisole (Entry 9). The purity determined by NMR and GC analyses: > 95%; 1H NMR δ 1.35 (s, 12 H), 3.86 (s, 3 H), 7.21 (s, 1 H), 7.48 (d, 1 H, $J = 2.2$ Hz), 7.65 (s, 1 H); ^{13}C NMR δ 24.81, 55.52, 84.29, 114.10 (q, $J = 3.3$ Hz), 122.50, 123.43 (q, $J = 3.3$ Hz), 124.01 (q, $J = 272.6$ Hz), 131.34 (q, $J = 32.0$ Hz), 159.16; exact mass calcd for $C_{14}H_{18}BF_3O_3$ 302.1301, found 302.1314.

2-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)indole (Entry 10). The purity determined by NMR and GC analyses: > 95%; 1H NMR δ 1.36 (s, 12 H), 7.09 (t, 1 H, $J = 7.7$ Hz), 7.11 (s, 1 H), 7.23 (t, 1 H, $J = 8.3$ Hz), 7.38 (d, 1 H, $J = 8.3$ Hz), 7.67 (d, 1 H, $J = 7.8$ Hz), 8.56 (br s, 1 H); ^{13}C NMR δ 24.78, 84.13, 111.25, 113.83, 119.75, 121.57, 123.59, 128.24, 138.18; exact mass calcd for $C_{14}H_{18}BNO_2$ 243.1431, found 243.1433.

2-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[b]furan (Entry 11). The purity determined by NMR and GC analyses: > 95%; 1H NMR δ 1.39 (s, 12 H), 7.23 (t, 1 H, $J = 7.4$ Hz), 7.34 (dt, 1 H, $J = 1.2$ and 7.8 Hz), 7.40 (s, 1 H), 7.57 (d, 1 H, $J = 8.5$ Hz), 7.63 (d, 1 H, $J = 7.8$ Hz); ^{13}C NMR δ 24.73, 84.64, 111.92, 119.50, 121.84, 122.68, 125.89, 127.43, 157.45; exact mass calcd for $C_{14}H_{17}BO_3$ 244.1271, found 244.1278.

2-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[b]thiophene (Entry 12). The purity determined by NMR and GC analyses: > 95%; 1H NMR δ 1.38 (s, 12 H), 7.35 (ddd, 1 H, $J = 1.7, 7.3$, and 8.8 Hz), 7.37 (ddd, 1 H, $J = 1.8, 7.1$, and 9.0 Hz), 7.85 (dd, 1 H, $J = 2.2$ and 9.0 Hz), 7.89 (s, 1 H), 7.91 (dd, 1 H, $J = 1.5$ and 9.0 Hz); ^{13}C NMR δ 24.79, 84.42, 122.50, 124.08, 124.35, 125.29, 134.47, 140.41, 143.67; exact mass calcd for $C_{14}H_{17}BO_2S$ 260.1042, found 260.1019.

One-Pot Synthesis of methyl 4-(3,5-dichlorophenyl)benzoate via Borylation-Coupling Sequence (Scheme 2). To a solution of 1,3-dichloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzene resulted by the reaction of pinacolborane (1.43 mmol) with 1,3-dichlorobenzene (1.3 mmol) in dry hexane (2 mL) were added methyl 4-bromobenzoate (1.0 mmol), $PdCl_2(dppf)$ (0.03 mmol), K_3PO_4 (3.0 mmol), and DMF (4 mL), and the mixture was stirred at 60 °C for 1 h. The product was extracted with benzene, washed with water, and dried over $MgSO_4$. Column chromatography over silica gel provided analytically pure methyl 4-(3,5-dichlorophenyl)benzoate: The purity determined by NMR and GC analyses: > 95%; 1H NMR δ 3.95 (s, 3 H), 7.39 (t, 1 H, $J = 1.8$ Hz), 7.49 (d, 2 H, $J = 1.7$ Hz), 7.61 (dt, 2 H, $J = 8.5$ and 1.8 Hz), 8.12 (dt, 2 H, $J = 8.3$ and 1.7 Hz); ^{13}C NMR δ 52.25, 125.77, 127.03, 127.95, 130.04, 130.29, 135.49, 142.76, 142.95, 166.59; exact mass calcd for $C_{14}H_{10}Cl_2O_2$ 280.0058, found 280.0047.

References

1. C. E. Tucker, J. Davidson and P. Knochel, *J. Org. Chem.*, 1992, **57**, 3482.
2. R. Uson, L. A. Oro and J. A. Cabeza, *Inorg. Synth.*, 1985, **23**, 126.
3. R. N. Haszeldine, R. J. Lunt and R. V. Parish, *J. Chem. Soc. A*, 1971, 3696.
4. P. Wehman, G. C. Dol, E. R. Moorman, P. C. J. Kamer, P. W. N. M. van Leeuwen, J. Fraanje and K. Goubitz, *Organometallics*, 1994, **13**, 4856.
5. D. Wenkert and R. B. Woodward, *J. Org. Chem.*, 1983, **48**, 283.

RSC Author Templates - ChemDraw (CDX) - Graphical Abstracts

Aromatic C-H borylation of arenes and heteroarenes by stoichiometric amounts of pinacolborane was catalyzed by an iridium complex generated from $1/2[\text{Ir}(\text{OMe})(\text{COD})]_2$ and 4,4'-di-*tert*-butyl-2,2'-bipyridine at room temperature in hexane, and afforded the corresponding aryl- and heteroarylboronates in high yields with excellent regioselectivities.