Title	Concentrations and human health risk assessment of organochlorine pesticides in edible fish species from a Rift Valley lake-Lake Ziway, Ethiopia
Author(s)	Yohannes, Yared B.; Ikenaka, Yoshinori; Saengtienchai, Aksorn; Watanabe, Kensuke P.; Nakayam, Shouta M. M.; Ishizuka, Mayumi
Citation	Ecotoxicology and Environmental Safety, 106, 95-101 https://doi.org/10.1016/j.ecoenv.2014.04.014
Issue Date	2014-08
Doc URL	http://hdl.handle.net/2115/56870
Type	article (author version)
File Information	EES106_p.95-101.pdf

[Instructions for use](#)

Concentrations and human health risk assessment of organochlorine pesticides in edible fish species from a Rift Valley Lake – Lake Ziway, Ethiopia

Yared B. Yohannes ^{a,b}, Yoshinori Ikenaka ^a, Aksorn Saengtienchai ^a, Kensuke P. Watanabe ^a, Shouta M.M. Nakayama ^a and Mayumi Ishizuka ^{a,*}

^a Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Kita 18, Nishi 9, Kita-ku, Sapporo 060–0818, Japan

^b University of Gondar, Faculty of Natural and Computational Science, Department of chemistry, P.O. Box 196, Gondar, Ethiopia

* Corresponding author

Tel.: +81 11 706 6949

E-mail address: ishizum@vetmed.hokudai.ac.jp

Abstract

Fish consumption is known to have several health benefits for humans. However, the accumulation of organic pollutants, like organochlorine pesticides (OCPs) could pose health hazards. Thus, OCPs in edible fish species (*Oreochromis niloticus*, *Tilapia zillii*, *Carassius* spp., and *Clarias gariepinus*) from Lake Ziway, an Ethiopian Rift Valley Lake were investigated to assess the potential human health hazards of these contaminants. Dichlorodiphenyltrichloroethanes (DDTs), hexachlorocyclohexanes (HCHs), chlordanes, and heptachlors were observed with Σ OCPs concentration ranging from 1.41 to 63.8 ng g⁻¹ ww. DDTs were the predominant contaminants (0.9 to 61.9 ng g⁻¹ ww), followed by HCHs. The predominance of DDTs may be attributed to their current use in vector control and contamination from past usage. The estimated daily intakes (EDIs) of OCPs from all fish species were much lower than the acceptable daily intakes (ADIs), indicating that consumption of fish is at little risk to human health at present. However, the cancer risk estimates in the area of concern and the hazard ratios (HRs) of HCHs, DDTs, and heptachlors exceeded the threshold value of one, indicating daily exposure to these compounds is a potential concern. This may result in a lifetime cancer risk greater than of 1 in 10⁶.

Key words: Organochlorine pesticides, Fish, Lake Ziway, Risk assessment

1. Introduction

Organochlorine pesticides (OCPs) have been widely used and become a worldwide concern due to their persistence, bioaccumulative potential, chronic toxicity, and potential negative impacts on humans and wildlife (UNEP, 2001). It is known that most of the total intake of pesticide residues by human beings is through the food chain (Martinez et al., 1997). Fish are known to biomagnify pesticides from the surrounding environment (Mackay and Fraser, 2000), and transfer the pesticides to humans when consumed. Epidemiological studies indicate that some of these compounds may be associated with cancers in humans (Snedeker, 2001; Beard, 2006; IARC, 2008), and also influence the concentration of thyroid hormones (Meeker et al., 2007). Eskenazi et al. (2006) reported delays in neurodevelopment during early childhood due to the impacts of prenatal exposure to dichlorodiphenyltrichloroethanes (DDTs).

Although the use of OCPs has been banned or restricted, developing countries like Ethiopia still use them for agricultural and health purposes, and as a consequence they can be found in aquatic (Deribe et al., 2011; Yohannes et al., 2013a,b) and terrestrial ecosystems, for example in cow's milk (Gebremichael et al., 2013). Because it is landlocked, Ethiopia is highly dependent on lake aquatic environments for its economic development. The Ethiopian Rift Valley region, encompassing seven principal lakes, is a densely populated area confined with various agricultural activities where there is still an increasing trend of pesticide usage (Amera and Abate, 2008). Moreover, Ethiopia has implemented indoor residual spraying (IRS) with DDT for malaria control in the past few decades (WHO, 2007). Approximately 400 metric tons of active-ingredient DDT per year is used for IRS in many parts of the country including the Rift Valley, a malaria epidemic prone region (Biscoe et al.,

2005; Van den Berg, 2009). In addition, Ethiopia is one of the many African countries burdened with the problem of obsolete pesticides, which have been accumulated since the first imports in the 1960s (Haylamicheal and Dalvie, 2009). These were mostly organochlorine compounds such as chlordane, DDT, dieldrin and lindane that are banned or restricted in most countries. In this view, there is great likelihood that the Ethiopian Rift Valley ecosystem is exposed to large amounts of pesticides.

Lake Ziway, one of the Ethiopian Rift Valley lakes, is located in an area with many agricultural activities but few soil conservation efforts in its catchment area. Intensive agriculture in the proximity of the lake and municipal waste discharges are sources of pollution into this fresh water ecosystem (Hengsdijk and Jansen, 2006). It is therefore necessary to evaluate the current status of the OCPs in different fish species from Lake Ziway. A recent study on the lake examined only the levels and biomagnification of DDTs (Deribe et al., 2013). No other studies have been carried out on the levels and risk assessment of other OCPs in the lake.

Therefore, objectives of this study are to assess the accumulation levels of OCPs in edible fish species collected from Lake Ziway and to evaluate the potential risks to human health posed through dietary consumption of these fish. This study gives a comprehensive overview of OCPs' status in the fish species of different trophic levels in Lake Ziway and provides a basis for decision-makers to take effective measures aimed at mitigating potential health and ecological risks.

2. Materials and methods

2.1. Study area

The study area, Lake Ziway (surface area: 434 km²) is a shallow freshwater lake located in the northern section of the Rift Valley (Fig. 1). It is fed by two inflowing rivers, the Meki River from the north-west and the Katar River from the east, and drains towards the Lake Abijata, through the Bulbula River. The lake has a large littoral zone containing emergent and submergent vegetation, which provides feeding, breeding and nursery habitats for fish (Admassu and Ahlgren, 2000; Erko et al., 2006). Lake Ziway contains different fish species including Nile tilapia (*Oreochromis niloticus*), Redbelly tilapia (*Tilapia zillii*), African big barb (*Barbus intermedius*), African sharptooth catfish (*Clarias gariepinus*), and Carp spp. (*Carassius carassius* and *Carassius auratus*) (Lemma, 2005). Fisheries on Lake Ziway are an open and easily accessible source of income and have always been a source of food and income for the people living on the shores of the lake. The landings of Lake Ziway used to be dominated by *O. niloticus*, but species of *C. gariepinus*, *T. zillii*, and *Carassius* spp. (*C. carassius* and *C. auratus*) are increasingly becoming a part of the catch. The potential yield of all the species of the lake is estimated to range between 2,500 and 6,680 tons/yr (Spliethoff et al., 2009).

2.2. Sampling

A total of 100 individual fish belonging to *O. niloticus*, *T. zillii*, *Carassius* spp., and *C. gariepinus* fish species of Lake Ziway were purchased from the local fishermen in January 2011. Samples were transported to Ziway fisheries research laboratory where the body size and body weight were recorded. General information about the fish is given in Table 1.

Fishes were dissected to obtain dorsal muscles and stored at $-20\text{ }^{\circ}\text{C}$. The frozen samples were then transported to Japan for analysis. Each individual sample was lyophilized, homogenized separately and used for chemical analysis.

2.3. OCPs analysis

Samples were processed and analyzed using a method described by Yohannes et al. (2013a) with slight modifications. Approximately 10 g of muscle tissue from each fish was taken and mixed with anhydrous sodium sulfate. After spiking with the surrogate standard of 2,4,5,6-tetrachloro-*m*-xylene (TCmX), each sample was extracted using Soxtherm apparatus (S306AK Automatic Extractor, Gerhardt, Germany) with *n*-hexane:acetone (3:1, *v/v*) for 4 h. An aliquot of the extract (20%) was used for lipid measurement using gravimetric method. The remaining extract was applied to a column filled with 6 g florisil (activated at $150\text{ }^{\circ}\text{C}$ overnight) for clean-up and eluted with a mixture of *n*-hexane:dichloromethane (7:3, *v/v*). The eluate was concentrated to 2 mL on rotary evaporator, and further to near dryness under gentle nitrogen flow. Finally, the extract was redissolved in 100 μL *n*-decane, and the internal standard pentachloronitrobenzene was added before instrumental analysis.

OCPs including DDTs (*o,p'*-DDT, *p,p'*-DDT, *o,p'*-DDE, *p,p'*-DDE, *o,p'*-DDD and *p,p'*-DDD), hexachlorocyclohexanes (HCHs; α -, β -, γ - and δ -HCH), heptachlors (HPTs; heptachlor, *cis*- and *trans*-heptachlor epoxide), chlordanes (CHLs; *cis*- and *trans*-chlordanes, *cis*- and *trans*-nonachlor and oxychlordanes), drins (aldrin, dieldrin and endrine) and hexachlorobenzene (HCB) were analyzed by gas chromatography equipped with an electron capture detector (Shimadzu GC-2014, Kyoto, Japan). An ENV-8MS capillary column (30 m $\times$ 0.25 mm i.d., 0.25 μm film thickness) with splitless injection was

used to separate OCPs. One μL of each sample was injected. The column oven temperature was initially set at $100\text{ }^{\circ}\text{C}$ for 1 min, increased to $180\text{ }^{\circ}\text{C}$ at $20\text{ }^{\circ}\text{C min}^{-1}$ and then to $260\text{ }^{\circ}\text{C}$ at $4\text{ }^{\circ}\text{C min}^{-1}$, which was held for 5 min. The injector and detector temperatures were $250\text{ }^{\circ}\text{C}$ and $310\text{ }^{\circ}\text{C}$, respectively. Helium at a flow rate of 1.0 mL min^{-1} and nitrogen at 45 mL min^{-1} were used as carrier gas and make-up gas, respectively.

2.4. Quality control and quality assurance

OCPs were identified by comparing their retention time with the reference to the corresponding standards. Multi-level calibration curves were created for the quantification and linearity ($R^2 \geq 0.995$) was achieved. Quality control was performed by analysis of procedural blanks and spiked blanks. Results showed that no target analysts were detected in blank samples and recoveries for spiked blanks ranged from 90% to 105%. The recovery rate of the surrogate, TCmX was $85 \pm 11\%$. To check for the validity of the method used for the extraction and analysis of the samples, the standard reference material SRM 1947 (Lake Michigan Fish Tissue) was analyzed during the analysis of samples, and the recoveries ranged from 85% to 105% with $\text{RSD} < 10\%$. The values reported here were not corrected for recoveries. Detection limits based on 3:1 signal to noise ratio (S/N) were between 0.05 and 0.1 ng/g for all OCPs. Concentrations were expressed on a wet weight (ww) basis.

2.5. Risk assessment

Various international organizations have subsequently established a series of standards and instructions to estimate the risks to human health from environmental pollutants in fish (USEPA, 2013). A straight forward risk assessment is performed through comparison with the levels set by laws and guidelines. However, this comparison was made without the

consideration of factors like different eating habits and consumption rates. Thus, in this study, we investigated the risk assessment by two approaches. To comprehensively evaluate the health risk assessment, the 50th and 95th percentile measured concentrations were used.

2.5.1. Estimated daily intake (EDI)

Estimated dietary intakes of OCPs were calculated as follows:

$$EDI = \frac{C \times DR}{BW} \quad (1)$$

where C is the measured concentration of OCPs (ng/g ww), DR is average daily consumption rate of fish (g/day) and BW is body weight (kg), which was set at 60 kg (WHO, 2010). The average daily consumption rate was derived from FAO (2011). Though Ethiopians are traditionally meat eaters, eating habits have been shifting in favor of fish in areas and communities where there is regular and sufficient supply. In those communities, annual fish consumption can exceed 10 kg/person (FAO, 2011). Thus, the DR was estimated at 30 g/day per person.

2.5.2. Potential carcinogenic risks

To assess public health risks posed through fish consumption, the cancer risk estimates and hazard ratios (HRs) were assessed on the basis of the guidelines of the United States Environmental Protection Agency (USEPA). Cancer risks associated with OCPs were estimated by combining the exposure dose and slope factor (USEPA, 2005). A public screening criteria for carcinogens is set at a carcinogenic risk level of 10⁻⁶. Carcinogenic risks below 10⁻⁶ are considered acceptable, while carcinogenic risks above 10⁻⁴ are considered unacceptable. An area of concern is present between 10⁻⁶ and 10⁻⁴ (USEPA, 2005).

HR for cancer risks was assessed by comparing the EDI with the benchmark concentration (BMC) (Solomon et al., 2000; Jiang et al., 2005) using the following equation:

$$HR = \frac{EDI}{BMC} \quad (2)$$

The BMC for carcinogenic effects was derived from the cancer slope factor (CSF), which was obtained from the USEPA (USEPA, 2012). The BMC for carcinogenic effects represents the exposure concentration at which lifetime cancer risk is one in a million for lifetime exposure. A hazard ratio that is greater than one indicates that there is potential risk to human health (Dougherty et al., 2000).

2.6. *Statistical analysis*

Statistical analysis was performed using JMP 9 (SAS Institute, Cary, NC, USA). Descriptive statistics using one-way analysis of variance (ANOVA) were used to characterize the levels of OCPs in the studied fish species. Concentrations below the limit of detections were given a value of zero. Multiple comparisons among the fish species were tested using Tukey's HSD post hoc test. A significant level of $p < 0.05$ was used.

3. Results and discussion

The length and weight of the studied fish species varied from 120 to 560 mm and from 111 to 1910 g, respectively (Table 1). A continuous increase in length and weight was observed for all individuals with a significant and positive correlation ($R^2 = 0.70$, $p < 0.001$). The mean lipid content was in the range $0.75 \pm 0.68\%$ to 1.34 ± 2.52 , and there was no significant difference ($p > 0.05$) among the studied fish species (Table 1). There was no significant correlation between the biometric data and lipid content ($p > 0.05$).

3.1. Levels of OCPs

OCPs were detected in muscle samples of all fish species, indicating their widespread contamination in Lake Ziway. DDTs, HCHs, HPTs, and CHLs were detected with varying concentrations (Table 2). The total concentrations of OCPs ranged from 1.41 to 63.8 ng g^{-1} ww, with a mean concentration of $7.72 \pm 6.90 \text{ ng g}^{-1}$ ww. The highest concentrations of OCPs were found in *C. gariepinus* ($p < 0.05$), which is a carnivorous fish and found at top trophic position. Among the OCPs analyzed, DDTs were the most commonly detected and were dominant in all samples. It accounted for $64.5 \pm 10\%$ (SD) (ranging from 52 to 78%) of the total OCPs. In general, the contamination pattern of OCPs in fish samples detected in this study was in the order of DDTs > HCHs > CHLs $\cong$ HPTs. This result indicates the high degree of exposure to DDTs in biota from the Ethiopian Rift Valley region, which is most likely due to recent use of DDT for malaria control through IRS (Biscoe et al., 2005; Van den Berg, 2009) as well as from past usage and spills from obsolete pesticides (Haylamicheal and Dalvie, 2009). It is also reported that DDT is still ongoing use by farmers in the region (Amera and Abate, 2008). Log transformed OCPs show a positive correlation

with total length for all fish species ($R^2 = 0.18$; $p < 0.001$), whereas no significant correlation was found between lipid content and concentration of OCPs ($R^2 = 0.00$; $p = 0.140$).

3.1.1. DDTs

DDT and its metabolites were detected in all fish species (Table 2). Concentrations of DDTs in the muscle tissue are found at large variations ranging from 0.77–61.9 ng g⁻¹ ww (mean concentration of 5.27 ± 6.73 ng g⁻¹ ww). *C. gariepinus* with 9.0 ± 11.7 ng g⁻¹ ww and *O. niloticus* with 2.33 ± 1.09 ng g⁻¹ ww had the highest and lowest concentrations, respectively. This may be attributed to their different feeding habits because *C. gariepinus* is a carnivorous and *O. niloticus* is almost herbivorous fish species (Table 1). Overall, the concentrations of DDTs were higher than those of other OCPs. The possible reasons for the presence of high level of DDTs in the region may be its current use in vector control, illegal usage and contamination from obsolete pesticides (Haylamicheal and Dalvie, 2009; Van den Berg, 2009). Reports from other African lakes also indicate much higher levels of DDT in aquatic organisms compared to other OCPs. In Lake Koka, Ethiopia DDT ranged from 0.05–72.53 ng g⁻¹ ww and it was the predominant pesticide by a factor of 10 when compared to the other OCPs (Deribe et al., 2011) and in Lake Malawi DDT concentrations were up to 60 times higher than other OCPs (Kidd et al., 2001). Concentrations of DDTs found in this study (mean concentration of 2.33 to 9.0 ng g⁻¹ ww) are higher than those found in Lake Victoria, Uganda (mean 1.39 to 1.67 ng g⁻¹ ww) (Kasozi et al., 2006). However, they are lower than those in fish from Southern Lake Victoria, Tanzania (mean 15 and 20 ng g⁻¹ ww) (Henry and Kishimba, 2006) and fish from Lake Burullus, Egypt (mean 2.76 to 45.13 ng g⁻¹ ww) (Said et al., 2008), and comparable to fish from Lake Awassa, Ethiopia with Σ DDTs

mean concentration of 1.80 and 9.0 ng g⁻¹ ww for *O. niloticus* and *C. gariepinus*, respectively (Yohannes et al., 2013a). Direct comparisons should be made with caution since these studies were conducted on different species. With all the data pooled together, the concentration of DDTs (log transformed) was significantly correlated ($R^2 = 0.18$; $p < 0.001$) to total length of the fish, but not with % lipid content ($R^2 = 0.02$; $p = 0.139$).

The composition profiles of DDTs in the muscle tissue of the four fish species are shown in Fig. 2. Among the metabolites, *p,p'*-DDE was the predominant congener, accounting for 55% ± 15.72 (from 41 to 77%), followed by *p,p'*-DDT (15% ± 6.42), and *p,p'*-DDD (13% ± 4.80). The proportion of *p,p'*-DDE was higher in *C. gariepinus* than in the others, comprising 77% of the mean DDT concentrations, showing that *C. gariepinus* found at high trophic level is more likely feeding on prey (both fish and invertebrates) and accumulates DDE, a more degraded form of DDT. In addition, this may be attributed to the more persistent nature of *p,p'*-DDE, and to its rate of biomagnification along the food chain in freshwater ecosystems (Rognerud et al., 2002). In contrast, the proportions of parent compounds (*o,p'*- and *p,p'*-DDT) in *O. niloticus* (29.8%), *T. zillii* (32.6%), and *Carassius* spp. (34.4%) were higher than in *C. gariepinus* (11.7%). This may be probably as a result of more efficient transfer of DDT to phytoplankton and macrophyte consuming herbivorous fish (Zhou et al., 2007). Technical DDT generally contains 75% *p,p'*-DDT, 15% *o,p'*-DDT, 5% *p,p'*-DDE, and <5% others. DDT can be metabolized into DDE under aerobic conditions or into DDD in anaerobic environments (Hitch and Day, 1992). Thus, the ratio of (*p,p'*-DDE + *p,p'*-DDD)/ΣDDTs can indicate past or recent usage of technical DDT. A ratio greater than 0.5 generally indicates long term biotransformation of DDT, whereas a ratio of less than 0.5 may indicate recent input of DDT. In the present study, the ratio ranged from

0.55 to 0.87, suggesting that DDTs in fish from Lake Ziway were mainly due to historical use, and to its current use for vector control in the region since the Ethiopian government decided to continue using DDT because of the high incidence of malaria and the corresponding fatalities (Biscoe et al., 2005; WHO, 2007).

3.1.2. HCHs

HCHs were the second most prevalent OCP contaminants in the studied fish species and accounted for 17% (from 10% to 25%) of the total OCPs measured. The levels of HCHs in *T. zillii* ($1.45 \pm 0.61 \text{ ng g}^{-1} \text{ ww}$) and *O. niloticus* ($1.26 \pm 1.04 \text{ ng g}^{-1} \text{ ww}$) were significantly higher ($p < 0.05$) than that of *Carassius* spp. ($0.61 \pm 0.31 \text{ ng g}^{-1} \text{ ww}$), and *C. gariepinus* ($0.72 \pm 0.47 \text{ ng g}^{-1} \text{ ww}$) (Table 2). A negative relationship ($R^2 = 0.07$; slope = -0.02 ; $p < 0.01$) between log transformed ΣHCH and length of fish was found, whereas no significant relationship ($p > 0.05$) was found between lipid content and concentration of HCHs. Of the HCHs measured, α and γ -HCHs were frequently detected and the γ -isomer (lindane) was the predominant, accounting for 60% on an average of ΣHCHs in the muscle tissue. The higher γ -HCH concentrations in the samples indicate current usage of lindane around the lake. A recent study in the Ethiopian rift valley region also showed high concentrations of lindane in tissues taken from cattle with the highest level of $0.14 \text{ mg kg}^{-1} \text{ ww}$ in liver samples obtained from Holeta, Ethiopia (Letta and Attah, 2012). In general, the concentrations of HCHs in this study are lower than those in fish from Lake Taabo, Cote d'Ivoire (Roche et al., 2007), and fish from Lake Burullus, Egypt (Said et al., 2008).

3.1.3. CHLs and HPTs

With regard to the residual levels of CHLs, *trans*-chlordane, *cis*-chlordane and *trans*-nonachlor were detected in most of the samples as they are the dominant constituents in technical chlordane (Xu et al., 2004) whereas oxy-chlordane was rarely encountered. The presence of these compounds in the environment at relatively high concentrations as compared to oxy-chlordane likely indicates recent inputs of chlordane to the environment. The mean residual levels of CHLs in the muscle tissues in the present study ranged from 0.40-0.91 ng g⁻¹ ww (Table 2). The use of chlordane is permitted in Ethiopia as a general insecticide (Ritter et al., 1995). Chlordane is imported to Ethiopia under the regulation of Ministry of Agriculture for termiticide usage only (EICDCR, 2004).

It was found that HPTs (*cis*-heptachlor epoxide and *trans*-heptachlor epoxide) were also present in most of the fish collected. The *cis* and *trans*-heptachlor epoxides predominated with a mean concentration of 0.28 ± 0.27 ng g⁻¹ ww and 0.31 ± 0.21 ng g⁻¹ ww, respectively (Table 2). The highest residual levels of HPTs (0.90 ± 0.35 ng g⁻¹ ww) were found in *O. niloticus*, the herbivorous fish species.

3.2. Human health risk assessment

Fish consumption has been proven to be one of the major routes of human exposure to organic contaminants. To better understand the concentration levels, the concentrations of OCPs in the present study were evaluated against international existing limits. The EDI was calculated and compared with the acceptable daily intake (ADI) recommended by the Food and Agriculture Organization and the World Health Organization (FAO/WHO) Joint Meeting on Pesticide Residue (WHO, 2010). To comprehensively evaluate risk exposure,

the 50th and 95th percentile EDIs of OCPs for each fish species were calculated. The EDIs of OCPs expressed as nanogram per kilogram body weight per day (ng/kg bw/d) through consumption of fish for the population are presented in Table 3. EDI of HCHs, HPTs, CHLs, and DDTs at both exposure levels were far below the ADI, indicating that consumption of fish at present would not pose a human health risk.

A carcinogenic risk assessment for OCPs was conducted using cancer risk estimates and HRs at the 50th and 95th percentile measured concentrations. As shown in Table 4, heptachlors showed much higher carcinogenic risk than other OCPs in all fish species. Regard to DDTs, the cancer risk for the 50th exposure level ranged from 3.7 in *O. niloticus* to 8.4×10^{-4} in *C. gariepinus* suggested that a person would have a chances of about 4 and 8 in 10000 to develop cancer from DDTs, respectively. This carcinogenic risk increased from 7.6 to 36×10^{-4} on 95th exposure level, which was unacceptable for human health. In general, the overall cancer risk estimates for all OCPs ranged from 0.7×10^{-4} to 36×10^{-4} on both the 50th and 95th exposure levels, and when compared to a target risk of $>1 \times 10^{-4}$, are considered unacceptable. Thus, the carcinogenic risk of HCHs, HPTs, CHLs and DDTs among humans at present should be of concern.

HRs based on the 50th and 95th percentile exposure levels were assessed in each fish species and the results are shown in Fig. 3. HRs for cancer risk based on the 95th percentile concentrations of HCHs, HPTs, and DDTs were greater than one. The HRs for the OCPs followed almost the following sequence: HPTs > DDT ≥ HCHs > CHLs. For all fish species, the HRs for HPTs were greater than one, showing that consuming fish is harmful to humans. Based on landings, *O. niloticus* is the most caught fish in Lake Ziway. The carcinogenic risk due to HCHs for this fish species in also greater than one while for DDTs is less than one.

However, for *T. zillii*, *Carassius* spp. and *C. gariepinus* the HRs for DDTs were greater than one. In general, cumulative daily exposure to OCPs because of fish consumption would yield a lifetime cancer risk of greater than one in a million. The results indicate that these compounds may be of particular concern because they are still in use.

4. Conclusion

This is the first study reporting on the levels and risk assessment of some OCPs in the most commonly caught fish species from the Ethiopian Rift Valley lake – Lake Ziway. The rift valley region is a populated area that is influenced by heavy pollution stemming from urban, agricultural and industrial activities. Our results indicated the presence of HCHs, HPTs, CHLs and DDTs with varying concentrations among the fish species. The overall conclusion of the evaluation is that DDTs were the main abundant pollutants, attributed to its current use in vector control and contamination from past usage. Dietary intakes estimated from the 50th and 95th percentile exposure level were far below ADIs. In contrast, the calculated cancer risk estimates and HRs of the studied fish species indicated that the consumption of most of the fish species could cause cancer as HR for cancer risk based on the 95th percentile concentrations of HCHs, HPTs and DDTs was greater than one. In this study, only fish and some OCPs were investigated to assess the risk. The consumption of water, vegetables, and animal meat, and the levels of other environmental pollutants were not considered. Therefore, the actual health risk for local people through dietary intake could be higher.

Acknowledgments

This study was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT), and from the Japan Society for the Promotion of Science (JSPS). We would also like to acknowledge the financial support provided by the Mitsui & Co., Ltd. Environment Fund, The Akiyama Life Science Foundation, and The Nihon Seimei Foundation. The authors are most thankful to Mr. Lemma Abera, Director of the Institute of Ziway Fisheries Resources Research Center laboratory for his kind help in every aspect during sampling and for allowing us to perform dissections in the laboratory. The support offered by Mr. Abraham G/Tsadik and Mr. Kasim during sampling and dissection is highly appreciated.

Conflict of interest

The authors declare no conflicts of interest.

367 **References**

- 368 Admassu, D., Ahlgren, I., 2000. Growth of juvenile catfish, *Oreochromis niloticus* L. from
 369 Lakes Zwai, Langeno and Chamo (Ethiopian rift valley) based on otolith micro-
 370 increment analysis. Ecol. Freshw. Fish 9, 127–137.
- 371 Amera, T., Abate, A., 2008. An assessment of the pesticide use, practice and hazards in the
 372 Ethiopian Rift Valley. Published by: Institute for Sustainable Development (ISD) and
 373 Pesticide Action Network (PAN-UK).
- 374 Beard, J., 2006. DDT and human health. Sci. Total Environ. 355, 78–89.
- 375 Biscoe, M.L., Mutero, C.M., Kramer, R.A., 2005. Current policy and status of DDT use for
 376 malaria control in Ethiopia, Uganda, Kenya and South Africa. Colombo, Sri Lanka:
 377 International Water Management Institute. [Working paper 95]
 378 doi: <http://dx.doi.org/10.3910/2009.268>
- 379 Deribe, E., Rosseland, B.O., Borgstrøm, R., Salbu, B., Gebremariam, Z., Dadebo, E., Norli,
 380 H.R., Eklo, O.M., 2011. Bioaccumulation of persistent organic pollutants (POPs) in fish
 381 species from Lake Koka, Ethiopia: the influence of lipid content and trophic position. Sci.
 382 Total Environ. 410-411, 136–145.
- 383 Deribe, E., Rosseland, B.O., Borgstrøm, R., Salbu, B., Gebremariam, Z., Dadebo, E.,
 384 Skipperud, L., Eklo, O.M., 2013 Biomagnification of DDT and its metabolites in four
 385 fish species of a tropical lake. Ecotoxicol. Environ. Saf. 95, 10–18.
- 386 Dougherty, C.P., Holtz, S.H., Reinert, J.C., Panyacosit, L., Axelrad, D.A., Woodruff, T.J.,
 387 2000. Dietary exposures to food contaminants across the United States. Environ. Res. 84,
 388 170–185.
- 389 EICDCR (Export and Imports of Certain Dangerous Chemicals Regulations), 2004.
 390 Subsidiary legislation 504.59 pp 19, report
 391 <http://www.justiceservices.gov.mt/DownloadDocument.aspx?app=lom&itemid=11536&l=1>
 392 (Accessed April, 2014).
- 393 Erko, B., Balcha, F., Kifle, D., 2006. The ecology of *Biomphalaria sudanica* in Lake Ziway,
 394 Ethiopia. Afr. J. Ecol. 44, 347–352.
- 395 Eskenazi, B., Marks, A.R., Bradman, A., Fenster, L., Johnson, C., Barr, D.B., Jewell, N.P.,
 396 2006. In utero exposure to dichlorodiphenyltrichloroethane (DDT) and

397 dichlorodiphenyldichloroethylene (DDE) and neurodevelopment among young Mexican
 398 American children. *Pediatrics*, 118, 233–241.

399 FAO (Food and Agriculture Organization), 2011. Fishery and aquaculture country profiles
 400 Ethiopia. http://www.fao.org/fishery/countrysector/FI-CP_ET/en (Accessed September, 2013).

401 Gebremichael, S., Birhanu, T., Tessema, D.A., 2013. Analysis of organochlorine pesticide
 402 residues in human and cow's milk in the towns of Asendabo, Serbo and Jimma in South-
 403 Western Ethiopia. *Chemosphere* 90, 1652–1657.

404 Haylamicheal, I., Dalvie, M., 2009. Disposal of obsolete pesticides, the case of Ethiopia.
 405 *Environ. Int.* 35, 667–673.

406 Hengsdijk, H., Jansen, H., 2006. Agricultural development in the Central Ethiopian Rift
 407 valley: a desk-study on water-related issues and knowledge to support a policy dialogue.
 408 Plant Research International B.V., Wageningen. <http://library.wur.nl/WebQuery/wurpubs/347623>

409 Henry, L., Kishimba, M.A., 2006. Pesticide residues in Nile Tilapia (*Oreochromis niloticus*)
 410 and Nile perch (*Lates niloticus*) from Southern Lake Victoria, Tanzania. *Environ. Pollut.*
 411 140, 348–354.

412 Hitch, R.K., Day, H.R., 1992. Unusual persistence of DDT in some Western USA soils. *Bull*
 413 *Environ. Contam. Toxicol.* 48, 259–264.

414 IARC (International Agency for Research on Cancer), 2008. Overall evaluations of
 415 carcinogenicity to humans. <http://monographs.iarc.fr/ENG/Classification/index.php> (accessed
 416 September, 2013).

417 Jiang, Q.T., Lee, T.K., Chen, K., Wong, H.L., Zheng, J.S., Giesy, J.P., Lo, K.K., Yamashita,
 418 N., Lam, P.K., 2005. Human health risk assessment of organochlorines associated with
 419 fish consumption in a coastal city in China. *Environ. Pollut.* 136, 155–165.

420 Kasozi, G.N., Kiremire, B.T., Bugenyi, F.W.B., Kirsch, N.H., Nkedi-Kizza, P., 2006.
 421 Organochlorine residues in fish and water samples from Lake Victoria, Uganda. *J.*
 422 *Environ. Qual.* 35, 584–589.

423 Kidd, K.A., Bootsma, H.A., Hesslein, R.H., Muir, D.C.G., Hecky, R.E., 2001.
 424 Biomagnification of DDT through the benthic and pelagic food webs of Lake Malawi,
 425 East Africa: importance of trophic level and carbon sources. *Environ. Sci. Technol.* 35,
 426 14–20.

427 Lemma, A., 2005. Site action plan for the conservation and sustainable use of the Lake Ziway
 428 biodiversity (Rift Valley Lakes Project). Institute of Biodiversity, Addis Ababa, Ethiopia.

429 Letta, B.D., Attah, L.E., 2013. Residue levels of organochlorine pesticides in cattle meat and
 430 organs slaughtered in selected towns in West Shoa Zone, Ethiopia. J. Environ. Sci. Heal.
 431 B 48, 23–32.

432 Mackay, D., Fraser, A., 2000. Bioaccumulation of persistent organic chemicals: mechanisms
 433 and models. Environ. Pollut. 110, 375–391.

434 Martinez, M.P., Angulo, R., Pozo, R., Jodral, M., 1997. Organochlorine pesticides in
 435 pasteurized milk and associated health risks. Food Chem. Toxicol. 35, 621–624.

436 Meeker, J.D., Altshul, L., Hauser, R., 2007. Serum PCBs, *p,p'*-DDE and HCB predict thyroid
 437 hormone levels in men. Environ. Res. 104, 296–304.

438 Ritter, L., Solomon, K.R., Forget, J., 1995. A review of selected persistent organic pollutants.
 439 http://www.who.int/ipcs/assessment/en/pes_95_39_2004_05_13.pdf (Accessed September, 2013).

440 Roche, H., Tidou, A., Persic, A., 2007. Organochlorine pesticides and biomarker response in
 441 two fishes *Oreochromis niloticus* (Linnaeus, 1758) and *Chrysichthys nigrodigitatus*
 442 (Lacepede, 1803) and an Invertebrate, *Macrobrachium vollenhovenii* (Herklot, 1857),
 443 from the Lake Taabo (Cote d'Ivoire), J. Applied Sci. 7, 3860–3869.

444 Rognerud, S., Grimalt, J.O., Rosseland, B.O., Fernandez, P., Hofer, R., Lackner, R., et al.,
 445 2002. Mercury and organochlorine contamination in brown trout (*Salmo trutta*) and
 446 Arctic charr (*Salvelinus alpinus*) from high mountain lakes in Europe and the Svalbard
 447 Archipelago. Water Air Soil Pollut: Focus 2, 209–232.

448 Said, T.O., El Moselhy, K.M., Rashad, A.A.M., Shreadah, M.A., 2008. Organochlorine
 449 contaminants in water, sediment and fish of Lake Burullus, Egyptian Mediterranean Sea.
 450 Bull Environ. Contam. Toxicol. 81, 136–146.

451 Snedeker, S.M., 2001. Pesticides and breast cancer risk: a review of DDT, DDE, and dieldrin.
 452 Environ. Health Perspect. 109, 35–47.

453 Solomon, K.R., Giesy, J., Jones, P., 2000. Probabilistic risk assessment of agrochemicals in
 454 the environment. Crop Protection 19, 649–655.

455 Spliethoff, P., Wudneh, T., Tariku, E., Senbeta, G., 2009. Past, current and potential
 456 production of fish in lake Ziway: Central Rift Valley in Ethiopia. Capacity development
 457 and institutional change programme Wageningen International, the Netherlands.

458 UNEP (United Nations Environmental Protection), 2001. Final act of the conference of
 459 plenipotentiaries on the Stockholm convention on persistent organic pollutants,
 460 Stockholm, Sweden.

461 USEPA (United States Environmental Protection Agency), 2005. Guidelines for carcinogen
 462 risk assessment. <http://www.epa.gov/cancerguidelines/2005> (Accessed September, 2013).

463 USEPA (United States Environmental Protection Agency), 2012. Human health risk
 464 assessment; risk based screening table.
 465 http://www.epa.gov/reg3hwmd/risk/human/pdf/NOV_2012_FISH.pdf (Accessed September, 2013).

466 USEPA (United States Environmental Protection Agency), 2013. Food safety.
 467 <http://www.epa.gov/oecaagct/tfsy.html> (Accessed September, 2013)

468 Van den Berg, H., 2009. Global status of DDT and its alternatives for use in vector control to
 469 prevent disease. *Environ. Health Perspect.* 117, 1656–1663.

470 WHO (World Health Organization), 2007. Implementation of indoor residual spraying of
 471 insecticides for malaria control in the WHO African Region Report: Irs report 2007 Draft,
 472 pp. 20–22.

473 WHO (World Health Organization), 2010. Inventory of IPCS and other WHO pesticide
 474 evaluations and summary of toxicological evaluations performed by the Joint Meeting on
 475 Pesticide Residues (JMPRs) through 2010.
 476 http://www.who.int/foodsafety/chem/jmpr/publications/jmpr_pesticide/en/index.html (Accessed September,
 477 2013).

478 Xu, D., Zhong, W., Deng, L., Chai, Z., Mao, X., 2004. Regional distribution of
 479 organochlorinated pesticides in pine needles and its indication for socioeconomic
 480 development. *Chemosphere* 54, 743–752.

481 Yohannes, Y.B., Ikenaka, Y., Nakayama, S.M.M., Saengtienchai, A., Watanabe, K., Ishizuka,
 482 M., 2013a. Organochlorine pesticides and heavy metals in fish from Lake Awassa,
 483 Ethiopia: Insights from stable isotope analysis. *Chemosphere* 91, 857–863.

484 Yohannes, Y.B., Ikenaka, Y., Saengtienchai, A., Watanabe, K.P, Nakayama, S.M.M.,
 485 Ishizuka, M., 2013b. Occurrence, distribution, and ecological risk assessment of DDTs
 486 and heavy metals in surface sediments from Lake Awassa-Ethiopian Rift Valley Lake.
 487 *Environ. Sci. Pollut. Res.* 20, 8663–8671.

488 Zhou, R., Zhu, L., Kong, Q., 2007. Persistent chlorinated pesticides in fish species from
 489 Qiantang River in East China. *Chemosphere* 68, 838–847.

Table 1
Biometry data of fish species in this study from Lake Ziway.

Fish species	<i>N</i>	Length (mm)	Weight (g)	Lipid content (%)	Main food*
		Mean $\pm$ SD	Mean $\pm$ SD	Mean $\pm$ SD	
		Min–max	Min–max	Min–max	
<i>O. niloticus</i>	27	213 $\pm$ 28	315 $\pm$ 111	^a 0.75 $\pm$ 0.68	Blue green algae, detritus, macrophytes
		167–270	178–554	0.10–3.60	
<i>T. zillii</i>	19	174 $\pm$ 21	199 $\pm$ 59	^a 0.90 $\pm$ 0.48	Macrophytes
		120–205	111–312	0.18–2.13	
<i>Carassius</i> spp.	27	267 $\pm$ 39	585 $\pm$ 230	^a 0.87 $\pm$ 0.59	Macrophytes, detritus, green algae
		160–332	231–1199	0.15–2.14	
<i>C. gariepinus</i>	27	353 $\pm$ 88	559 $\pm$ 454	^a 1.34 $\pm$ 2.52	Insect, fish eggs, fish, gastropods
		235–560	154–1910	0.23–5.3	

N = number of samples.

Mean $\pm$ standard deviation (SD).

^a Means with different letter superscript are significantly different (Tukey test is applied; $p < 0.05$).

* Reference: Deribe et al., 2013.

Table 2Levels of OCPs (ng g⁻¹ ww) in muscle of four fish species from Lake Ziway.

	<i>O. niloticus</i>	<i>T. zillii</i>	<i>Carassius</i> spp.	<i>C. gariepinus</i>
α-HCH	0.22 ± 0.06	0.19 ± 0.03	0.25 ± 0.09	0.27 ± 0.12
β-HCH	ND	0.31 ± 0.09	0.03 ± 0.11	ND
γ-HCH	0.67 ± 0.33	0.68 ± 0.52	0.22 ± 0.18	0.47 ± 0.42
δ-HCH	ND	0.27 ± 0.04	0.11 ± 0.21	ND
ΣHCHs	^a 1.26 ± 1.04	^a 1.45 ± 0.61	^b 0.61 ± 0.31	^b 0.72 ± 0.47
*	0.29–5.10	0.91–3.54	0.16–1.85	0.27–2.01
Heptachlor	ND	ND	ND	ND
cis-heptachlor-epoxide	0.57 ± 0.27	0.08 ± 0.10	0.24 ± 0.11	0.23 ± 0.11
trans-heptachlor-epoxide	0.32 ± 0.09	0.20 ± 0.02	0.31 ± 0.23	0.42 ± 0.29
ΣHPTs	^a 0.90 ± 0.35	^c 0.42 ± 0.11	^{b,c} 0.59 ± 0.27	^b 0.65 ± 0.28
*	0.44–2.27	0.19–0.69	0.20–1.52	0.34–1.56
oxy-chlordane	0.04 ± 0.04	0.16 ± 0.07	0.11 ± 0.06	0.10 ± 0.07
cis-chlordane	0.18 ± 0.04	0.26 ± 0.11	0.12 ± 0.07	0.16 ± 0.05
trans-chlordane	0.16 ± 0.05	0.20 ± 0.03	0.26 ± 0.21	0.29 ± 0.12
trans-nonachlor	0.03 ± 0.07	0.29 ± 0.10	0.37 ± 0.59	0.35 ± 0.23
ΣCHLs	^b 0.40 ± 0.10	^a 0.91 ± 0.22	^a 0.87 ± 0.82	^a 0.90 ± 0.25
*	0.17–0.61	0.65–1.32	0.19–4.00	0.58–1.50
p,p'-DDE	1.32 ± 0.81	1.89 ± 2.02	2.42 ± 1.60	6.92 ± 11.47
o,p'-DDE	0.10 ± 0.08	0.35 ± 0.12	0.26 ± 0.36	0.12 ± 0.10
p,p'-DDD	0.40 ± 0.21	0.85 ± 0.41	0.58 ± 0.35	0.79 ± 0.68
o,p'-DDT	0.43 ± 0.17	0.53 ± 0.21	0.68 ± 0.74	0.43 ± 0.11
p,p'-DDT	0.31 ± 0.18	0.77 ± 0.66	0.57 ± 0.73	0.62 ± 0.40
ΣDDTs	^b 2.33 ± 1.09	^{a,b} 4.38 ± 2.67	^{a,b} 4.55 ± 2.80	^a 9.0 ± 11.7
*	0.90–5.12	1.35–13.2	0.77–10.6	2.36–61.9
ΣOCPs	^b 4.89 ± 1.85	^{a,b} 7.16 ± 2.63	^{a,b} 6.62 ± 3.71	^a 11.2 ± 11.7
*	2.46–10.9	3.59–15.2	1.41–15.0	4.00–63.8

ND = below detection limit.

Mean ± standard deviation (SD).

* Min–max.

Values with different letters (a, b, c) within a row are significantly different at $p < 0.05$ level (Tukey test is applied).

Table 3
Estimated daily intake values (ng/kg bw/d) of OCPs through the studied fish species by human.

	ADI	50 th (95 th) percentile measured concentrations (ng/g ww)				50 th (95 th) estimated daily intakes			
		<i>O. niloticus</i>	<i>T. zillii</i>	<i>Carassius</i> spp.	<i>C. gariepinus</i>	<i>O. niloticus</i>	<i>T. zillii</i>	<i>Carassius</i> spp.	<i>C. gariepinus</i>
HCHs	5000 ^a	1.08 (3.39)	1.32 (2.60)	0.52 (1.20)	0.54 (1.53)	0.54 (1.70)	0.66 (1.30)	0.26 (0.66)	0.27 (0.77)
HPTs	100	0.86 (1.34)	0.42 (0.56)	0.53 (0.93)	0.60 (1.20)	0.43 (0.67)	0.21 (0.28)	0.27 (0.49)	0.30 (0.60)
CHLs	500	0.39 (0.60)	0.86 (1.26)	0.65 (2.41)	0.85 (1.41)	0.19 (0.30)	0.43 (0.63)	0.33 (1.20)	0.42 (0.72)
DDTs	10000	2.20 (4.49)	3.97 (8.23)	4.61 (9.21)	4.91 (21.21)	1.10 (2.24)	1.99 (4.12)	2.30 (4.61)	2.46 (10.61)

ADI = Acceptable daily intake (ng/kg bw/d).

^a for γ -HCH. (WHO, 2010).

Table 4
Cancer risk estimates for HCHs, HPTs, CHLs and DDTs.

OCPs	Cancer slope factor * [per (mg/kg day)]	50 th (95 th) percentile cancer risks (x 10 ⁻⁴)					^a for γ - HC H. *Ca ncer slop e fact
		<i>O. niloticus</i>	<i>T. zillii</i>	<i>Carassius</i> spp.	<i>C. gariepinus</i>		
HCHs	1.1 ^a	5.9 (18.7)	7.2 (14.3)	2.8 (6.6)	2.9 (8.4)		
HPTs	4.5	19 (30)	0.9 (13)	12 (22)	13 (27)		
CHLs	0.35	0.7 (1.0)	1.5 (2.2)	1.1 (4.2)	1.5 (2.5)		
DDTs	0.34	3.7 (7.6)	6.7 (14.0)	7.8 (15.7)	8.4 (36)		

ors were from the United States Environmental Protection Agency (USEPA, 2012).

Figure captions

Fig. 1. The map of Lake Ziway. (Deribe et al., 2013).

Fig. 2. Relative abundance of individual DDT components in four fish species from Lake Ziway.

Fig. 3. Carcinogenic hazard ratios for daily consumption of fish from Lake Ziway, Ethiopia. MEC, measured concentration. (The horizontal line represents the hazard ratio of > 1 , and any ratio higher than that indicates a risk.)

Fig. 1.

Fig. 2.

Fig. 3.