Title	Recovery of forest-floor vegetation after a wildfire in a Picea mariana forest
Author(s)	Tsuyuzaki, Shiro; Narita, Kenji; Sawada, Yuki; Harada, Koichiro
Citation	Ecological research, 28(6), 1061-1068 https://doi.org/10.1007/s11284-013-1087-0
Issue Date	2013-11
Doc URL	http://hdl.handle.net/2115/57303
Rights	The final publication is available at link.springer.com .
Type	article (author version)
File Information	Tsuyuzaki_14EcRs.pdf

[Instructions for use](#)

Ecol Res (201__) __: __-__

DOI: __

ORIGINAL ARTICLE

Shiro Tsuyuzaki · Kenji Narita · Yuki Sawada · Koichiro Harada

Recovery of forest-floor vegetation after a wildfire in a *Picea mariana* forest

S. Tsuyuzaki (✉)

Graduate School of Environmental Earth Science, Hokkaido University, Sapporo 060-0810
Japan

E-mail: tsuyu@ees.hokudai.ac.jp

K. Narita

Faculty of Education and Human Studies, Akita University, Akita 010-8502, Japan

E-mail: knarita@ed.akita-u.ac.jp

Y. Sawada

Faculty of Urban Management, Fukuyama City University, Fukuyama, 721-0964, Japan

E-mail: y-sawada@fcu.ac.jp

K. Harada

School of Food, Agricultural and Environmental Sciences, Miyagi University, 982-0215,
Japan

E-mail: haradak@myu.ac.jp

Abstract We aimed to detect the trajectories of forest-floor vegetation recovery in a *Picea mariana* forest after wildfire. Since fire severity in boreal forests is expected to increase due to changes in climate, we investigated the effects of ground-surface burned severity, a surrogate to the overall fire severity, on the revegetation. We annually monitored vegetation less than 1.3 m high in 80 1 m × 1 m quadrats at Poker Flat Research Range (65°12'N, 147°46'W, 650 m a.s.l.) near Fairbanks, interior Alaska, where a large wildfire occurred in the summer of 2004, from 2005 to 2009. *Sphagnum* mosses were predominant on the unburned ground surface. In total, 66% of the ground surface was burned completely by the wildfire. Total plant cover increased from 48% in 2005 to 83% in 2009. The increase was derived mostly by the vegetative reproduction of shrubs on the unburned surface, and by the immigration of non-*Sphagnum* mosses and deciduous trees on the burned surface. Deciduous trees, which had not been established before the wildfire, colonized only on the burned surface and grew faster than *P. mariana*. Although species richness decreased with increasing slope gradient, these deciduous trees established even on steep slopes. The wildfire that completely burned the ground surface distorted the revegetation, particularly, on steep slopes. Therefore, restoration of the *Sphagnum* surface was prerequisite when the severe wildfire occurred, because *Sphagnum* cover seemed to be difficult to return to predominance in the short term.

Key words: Burned ground surface · Deciduous trees · Mosses · Revegetation · Slope gradient

1 Introduction

2 Lightning-caused wildfire is a key to maintain the regeneration of taiga dominated by *Picea*
 3 *mariana* (P. Mill.) B.S.P. (black spruce) in boreal regions including interior Alaska
 4 (Engelmark 1999). Lightning has led to crown fire, which moderately burned the forest
 5 floor providing safe sites for *P. mariana* seedlings (Greene et al. 2004; Jayen et al. 2006).
 6 *P. mariana* produces a semi-serotinous cone that releases most of its seeds after wildfire, and
 7 the seedlings establish themselves steadily on incompletely-burned ground surface (Ilisson
 8 and Chen 2009). In recent years, however, wildfires have been increasing fuel
 9 consumption above and below the ground surface in the boreal forest zones of the Northern
 10 Hemisphere, because of dry lightning (Kashischke and Turetsky 2006, Johnstone et al. 2011).
 11 Climate projections suggest that these extreme disturbances will increase during this century
 12 (Flannigan et al. 2000; Anisimov et al. 2007).

13 Species recovering after wildfire are often divided into two types: seeders that recover
 14 by sexual reproduction and sprouters that recover by vegetative reproduction. Revegetation
 15 patterns after wildfire differ with fire severity, due to changes in the contributions of seeders
 16 and sprouters (Santana et al. 2012). When fire severity is low, i.e., the ground surface is
 17 incompletely burned, sprouters contribute more to revegetation, and *vice versa* (Schimmel
 18 and Granstrom 1996, Gurvich et al. 2005). Consequently, vegetation recovery is delayed or
 19 altered when disturbance exceeds the threshold of the resilience of sprouters (Dale et al.
 20 2001; Johnstone and Chapin 2006). Delayed recovery induces increase in active layer
 21 depth in permafrost zones (Burn 1998; Tsuyuzaki et al. 2009), and accelerates positive
 22 feedback on global warming (Kaplan and New 2006). Severe wildfire burning of ground
 23 surfaces may accelerate establishment of species that did not exist pre-fire and reorganize the
 24 flora.

25 Physical site characteristics, such as aspect, elevation and slope gradient, affect
 26 regeneration patterns after disturbances (Chapin et al. 2006). Permafrost distribution is
 27 related with the presence of *P. mariana*-dominated forests in discontinuous permafrost zones

(Davis 2001; Hollingsworth et al. 2006), and is decreased in post-fire vegetation due to decreasing albedo and increasing soil temperature until the vegetation recovers well (Chambers et al. 2005; Tsuyuzaki et al. 2009). Therefore, the paces and patterns of revegetation are a key to sustain permafrost and to conserve the ecosystems. Since large-scale disturbances, including burn-out wildfire, delay or alter the revegetation more than small-scale ones such as crown fire (Dale et al. 2001; Rydgren et al. 2004), we hypothesized that burned-out wildfire was associated with shifts in species composition by removing the ground-surface plant cover and by species immigrating from the external environments. We had a chance to obtain revegetation patterns after various-scale wildfires, including both crown and burn-out fires, coded as “Boundary Fire” that occurred in interior Alaska in the summer of 2004, and reported the first five years of monitoring of forest-floor vegetation after the wildfire.

Materials and methods

Study area

The study site is located at Poker Flat Research Range, approximately 50 km north of Fairbanks, Alaska. It is a scientific rocket launching facility owned by the University of Alaska Fairbanks. The region is located in the discontinuous permafrost zone. Annual precipitation averaged 297.4 mm at Fairbanks from 1971 to 2000, the maximum monthly mean air temperature was 23.0°C in June, and the minimum was -25.0°C in January (ACRC 2007). The Boundary Fire occurred in the region including Poker Flat from mid-June to late August in 2004, and burned ca 217,000 ha of forests with various severities (Betts and Jones 2009; Johnstone et al. 2011). Before the wildfire, the tree layer was dominated by *P. mariana* (Tsuyuzaki et al. 2009) and the ground surface was covered with thick mosses dominated by *Sphagnum* spp. (*S. capillifolium* (Ehrh.) Hedw., *S. subsecundum* Nees, and others) and with *Hylocomium splendens* (Hedw.) Schimp. Since *Sphagnum* spp. often establishes on wetter sites than *Hylocomium* in pre-burned forests (Heijmans et al. 2004;

Nichole and Yves 2006), *Sphagnum* was likely to remain more abundant than *Hylocomium* on the un-burned surface after the wildfire.

Sampling

Sixteen 10 m × 10 m plots were established on a north slope of different fire severity, ranging from 0% to 100%, before leaf flushing began in the spring of 2005, to detect relationships between burned area and revegetation (Tsuyuzaki et al. 2009). Five 1 m × 1 m quadrats were randomly set up in each plot, and the percentage of burned area was estimated visually in each quadrat. Burned ground surface was divided into two types in each quadrat; complete burning that removed the organic layer and exposed soil, and incompletely burning that left blackened organic matter such as peat. The measurements of burned areas were conducted over the whole burned area, including complete and incomplete burning, and completely-burned area. Monitoring continued annually from 2005 to 2009.

Percent cover was recorded for each plant taxa in a layer less than 1.3 m high on each quadrat every summer. The vascular plant species were identified by the first author, and the vouchers have been stored in SAPS. The mosses were sampled, identified by M. Higuchi, NMNS, and the vouchers have been kept in TNS. The lichens were sampled, and the vouchers have been kept in GSES, HU. The layers more than 1.3 m high had few vascular plants except *P. mariana*, the density of which averaged 24 stems per plot.

Photos were taken toward the canopy on each quadrat at 1.3 m above the ground surface by a fish-eye lens in the summers of 2005 and 2008. The photos were used to evaluate canopy openness, using a freeware Gap Light Analyzer ver. 2.0 (Frazer et al. 1999). Longitude and latitude on each quadrat and on the center of each plot were measured by differential GPS receivers (StarBox SSII-51CPN-19, Amtechs, Tokyo) with antenna (GPS-701, NovAtel, Calgary). Location, aspect and slope gradient on each quadrat were measured by a laser level meter. Based on these measurements, the elevation on each

1 quadrat was calculated. Thermometers were established from 2 cm to the surface of
2 permafrost of which maximum was 150 cm in three sites in August 2007 at three sites of
3 which fire severities differed between none and completely burned. Thaw depth was
4 inspected by temperature profiles recorded at 1 hour intervals.

5 6 *Data analysis*

7 Diversity and evenness in each quadrat were calculated by Shannon-Wiener indices.
8 Generalized linear mixed-effects models (GLMM) were used to detect significant
9 environmental factors on species richness, diversity, evenness, and total plant cover. Four
10 response variables—total plant cover, species richness, diversity, and evenness—were
11 examined in the models. The distributions were assumed as Poisson for richness because
12 of discontinuous function, gamma distribution for diversity because of continuous function
13 without zero, and binomial for evenness and cover because of the ratio. The examined
14 explanatory variables were burned ratio, canopy openness, elevation, slope, aspect, and years
15 after wildfire on each plot. Plot locations, i.e., longitude and latitude, were used as random
16 effect, to reduce the effects of pseudo-replicated sampling designs.

17 Non-metric multi-dimensional scaling (NMDS) was applied to investigate relationships
18 between environmental factors and plant cover on each species, using all taxa. NMDS is an
19 ordination technique well suited for data that are nonnormal, or occur along arbitrary or
20 discontinuous scales and is considered the most effective ordination method for ecological
21 community data (McCune and Grace 2002). NMDS does not assume a unimodal model of
22 species responses to the environment, and allows for the possibility that the community
23 variations are related to unmeasured environmental variables. Therefore, community
24 patterns are retained, regardless of what environmental variables were measured. In
25 addition, multi-response permutation procedures (MRPP) of 999 permutations were used to
26 examine significant differences between groups of sampling units (Reich et al. 2001).

27 Since the presence or absence *Sphagnum* mat greatly influence revegetation and/or

carbon cycle on and in the ground after wildfires in boreal ecosystems (Greene et al. 2004; Shetler et al. 2008; Whinam et al. 2010), two similarity analyses were compared using Jaccard similarity index to inspect the habitat preferences, in particular, to *Sphagnum* mat, of species that showed high cover and/or frequency (hereafter, i.e., dominant species) in relation to *Sphagnum*. In the first analysis, percentage similarities between *Sphagnum fuscum* (Schimp.) Klinggr. and each dominant species or each lifeform were calculated. In the second analysis, the similarities were calculated between non-*Sphagnum* cover and the species or lifeform to investigate the preferences of species to ground surface not occupied by *Sphagnum*. On the similarity between *Sphagnum* mat and moss lifeform, *Sphagnum* species were excluded from the lifeform. The non-*Sphagnum* surface was mostly created by burning in the first year (i.e., 2005). Therefore, non-*Sphagnum* cover was a surrogate for the burned area in 2005. All statistical analyses were conducted with the statistical program R (ver. 2.10.1) (R Development Core Team, 2010).

Results

Initial vegetation patterns and the environments

There were 29 vascular plant species (27 seed plants and 2 ferns), and over 13 non-vascular plant species, i.e., mosses and lichens, recorded in the 80 quadrats. The two ferns were *Equisetum silvaticum* L. and *Lycopodium annotinum* L. Of the mosses and lichens, *S. fuscum*, which occupied 29.3% of the unburned surface in 2005. The vascular plants consisted of 13 herbs (forbs, grasses and ferns), 12 shrubs, and 4 trees. Visual observation confirmed that the shrubs survived throughout the wildfire and recovered mostly by vegetative reproduction, i.e., sprouters. Of the four tree taxa, *P. mariana* was evergreen, and the others (*Betula neoalaskana* Sarg., *Populus tremuloides* Michx., and *Salix* spp.) were deciduous. *P. mariana* survived through the fire when they were rooted in *Sphagnum*. Therefore, the cover was high in 2005 (0.6%) and was derived mostly from surviving saplings. In contrast, the most of all broad-leaved trees were regenerated by seedlings.

Canopy openness ranged from 57% to 94%. The burned area averaged 66%, ranging from 0% to 100%. *Sphagnum* cover decreased annually, probably due to the delayed effects of burning, such as desiccation. Thaw depth increased with increasing burned surface on plot level from 50 cm to 330 cm in August 2007. The total number of species in all the quadrats ranged from 39 to 41 for five years, and did not differ largely, indicating that the species composition was fixed soon after the wildfire. Plant cover was negatively correlated to burned area (GLMM, $P < 0.01$).

Slope gradient was negatively correlated to richness ($P < 0.01$), showing that steep slopes restricted species richness. Species diversity and evenness ranged from 1.24 to 1.61 and from 0.56 to 0.68, respectively, and were not correlated to any examined environmental variables. Canopy openness and elevation were not related to these four parameters on plant community structure ($P > 0.01$).

Temporal changes in vegetation

Total plant cover increased from $60\% \pm 50$ (mean with standard deviation) in 2005 to $108\% \pm 38$ in 2009 (GLMM, $P < 0.01$). Similarly, species richness ranging from 9.8 to 11.3 was negatively correlated to burned area ($P < 0.01$) and positively correlated to year ($P < 0.01$). A sedge, *Carex bigelowii* Torr., and a grass, *Calamagrostis canadensis* (Michx.) Beauv., gradually increased their cover from 0.9–1.7% to 3.0–7.7% over the five years. A forb, *Epilobium angustifolium* L., had a peak of cover ($3.0\% \pm 9.0$) two years after the wildfire and then gradually decreased to $1.9\% \pm 3.4$. Ferns were infrequent in the quadrats. All of the dominant shrub species gradually increased their cover from 0.3–4.3% in 2005 to 0.9–11.3% in 2009. Shrubs, such as *Ledum groenlandicum* L., *Vaccinium vitis-idaea* L., *Vaccinium uliginosum* L., *Betula nana* L., and *Oxycoccus microcarpus* Turcz., had high cover even soon after the wildfire.

All of the trees gradually increased in cover across time. The cover of *P. mariana* increased slowly, and was 0.9% even in 2009. In contrast, *B. neoalaskana* and *P.*

1 *tremuloides* established from seeds, i.e., seeders, and thus the initial cover was low (less than
 2 0.1%). Increase in cover was, however, faster for the two deciduous trees than for *P.*
 3 *mariana*. In 2009, cover of *P. tremuloides* and *B. neoalaskana* reached 1.0% and 0.7%,
 4 respectively. Of the mosses and lichens, *S. fuscum* was dominant but decreased in cover
 5 annually to 21.1%. *Polytrichum commune* Hedw. and *Ceratodon purpureus* Hedw.
 6 increased in cover annually reaching 6% and 15%, respectively, in 2009. *Ceratodon*
 7 *purpureus* was established in 80% of quadrats in 2009. These two mosses, *P. commune* and
 8 *C. purpureus* were uncommon on unburned surfaces.

9 Coefficients of determination on NMDS indicated that all examined variables
 10 significantly explained the ordination patterns (test of random data permutations, $P < 0.01$).
 11 As axis I explained 52% of variance, axis II 26%, and axis III 14%, the first two axes were
 12 examined. The stress was 0.191. Of these, fire severity expressed by burned surface was
 13 the prime factor determining vegetation structures (Fig. 1). Burned area and canopy
 14 openness were both negatively correlated with axis I, showing that fire severity is a strong
 15 factor in determining post-fire species composition. The other three site factors—elevation,
 16 aspect and slope gradient—were related more to axis II, showing that burn severity was not
 17 greatly related to the site characteristics. Slope gradient was related to axis II more than
 18 aspect and elevation. Year had the weakest effects of the examined variables on vegetation
 19 structures, showing that revegetation pathways were not unique. MRPP also supported
 20 these results, i.e., these examined variables explained significantly different between the
 21 sampled groups, including continuous variables, at $P < 0.01$ (A ranging from 0.022 to 0.718,
 22 $\delta < 61.2$ that were less than expected $\delta = 62.58$, and $n = 400$).

23 Species scores on NMDS showed distinct patterns between the four life forms—herbs,
 24 shrubs, trees, and mosses (Fig. 1). Shrubs had high scores on axis I close to *Sphagnum*
 25 *fuscum*, while non-*Sphagnum* mosses, *P. commune* and *C. purpureus*, had low scores far
 26 from *S. fuscum* on axis I. These patterns indicated that the shrubs established less with the
 27 non-*Sphagnum* mosses. Deciduous trees scored low and clustered to each other on axis I

and axis II but *P. mariana* scored high on axis I, showing that the two deciduous trees established least with *P. mariana*. The scores of two non-*Sphagnum* mosses came close to those of deciduous trees but were slightly higher on axis I, showing that the mosses established primarily on burned surface but less so on the unburned surface. Although species richness decreased with increasing slope gradient, two deciduous trees (*P. tremuloides* and *B. neoalaskana*) and a moss (*P. commune*) established more on steep slopes, as shown by the low scores on axis II correlated to slope gradient.

High quadrat scores on axis I meant that quadrats were burned less, and *vice versa* (Fig. 2). The annual fluctuations of quadrat scores on NMDS became larger on more burned areas indicating that vegetation structures changed faster on burned surfaces with increasing fire severity. In addition, the scores on severely burned surfaces fluctuated along axis II rather than axis I, showing that the vegetation changes did not go toward less-burned vegetation. The major causes of the fluctuation patterns were derived from the colonization of two deciduous trees (*P. tremuloides* and *B. neoalaskana*, a herb, *E. angustifolium*) and two mosses (*P. commune* and *C. purpureus*), all of which established more on burned surfaces (Fig. 1).

Habitat preferences

The patterns of yearly fluctuations on the two similarities were different between burned and unburned surfaces (Fig. 3). The similarity of each lifeform were stable on unburned *Sphagnum* surface for the five years, as compared with the similarity on burned surface. These implied that the vegetation structures temporally changed more on burned surfaces than on unburned surfaces. The similarity of trees increased gradually on both unburned and burned surfaces, indicating that the trees increased their cover with time on both the surfaces.

The patterns were not synchronized between the three herbs (Fig. 3). The similarity of *E. angustifolium* to *Sphagnum* cover was extremely low and to non-*Sphagnum* cover was

high, showing that *E. angustifolium* established least on the unburned surface. In contrast, the similarity indicated *C. canadensis* recovered rapidly not only on *Sphagnum* surface but also on non-*Sphagnum* surface. *C. bigelowii* established on the *Sphagnum* surface more than on the non-*Sphagnum* surface, and the annual changes in the two similarities were small.

For all the dominant shrubs, the similarities to *Sphagnum* cover were high but to non-*Sphagnum* cover were low, showing that shrubs established well with *Sphagnum* as indicated by NMDS (Fig. 1).

P. mariana showed contrast patterns on the similarities on the *Sphagnum* and non-*Sphagnum* surfaces to the two deciduous species (Fig. 3). *P. mariana* showed higher similarity to the *Sphagnum* mat than any other tree species, and did not show low similarity to the non-*Sphagnum* mat. Although *P. mariana* established more on unburned surfaces, that did not mean *P. mariana* did not establish on burned surfaces. In contrast, two deciduous trees established least with *Sphagnum* and increased annually the similarities to non-*Sphagnum*, showing that these species gradually in cover on the non-*Sphagnum* surface.

Discussion

Vegetation patterns along gradients of fire severity

NMDS on quadrats (Fig. 1) indicated that revegetation patterns on the floor differed greatly between burned and unburned surfaces since the recovery on unburned surfaces was dependent on regeneration and that on burned surfaces was on colonization. Species producing wind-dispersed seeds contribute more to revegetation after severer wildfire (Johnson and Miyanishi 2007). Fire severity determines community composition derived from differences in regeneration strategy and root depth among species (Hollingworth et al. 2013).

The similarities between ground-surface combustion and plants indicated as follows (Fig. 3): *P. mariana* of the tree species produced semi-serotinous wind-dispersed seeds, but

established more on the unburned surface probably because of a habitat preference that is adapted to crown fire. Deciduous trees needed high severity sites where most *Sphagnum* carpets were removed for their establishments, while *P. mariana* trees colonized in both high and low severity sites because they were either able to survive fire on thick *Sphagnum* and germinated on burned and unburned surfaces. When post-fire residual organic matter is more than 2 cm in thickness, vegetation composition diverges in *P. mariana* forest of eastern Canada (Siegwart Collier and Mullik 2010).

Species producing wind-dispersed seeds or spores established steadily when suitable habitats were provided by burning the *Sphagnum* surface; viz. all the deciduous trees produced wind-dispersed seeds, and a perennial herb, *Epilobium angustifolium*, did, too. Mosses produced spores that should be dispersed by wind. Deciduous trees do not recruit seedlings in thick organic mats but outcompete with conifers on mineral soils after wildfire (Johnstone and Chapin 2006). There should be no safe sites for deciduous trees before *P. mariana* forests were burned.

Vegetation patterns along gradients of topography

Species richness decreased with increasing slope gradients. In addition, NMDS on species showed that revegetation on steep slopes was altered by the establishment of deciduous trees and non-*Sphagnum* mosses (Fig. 2). Wind-dispersed seeds often accumulate in rough microtopography provided by live plants, litter and coarse-textured soil, all of which function as seed traps (Koyama and Tsuyuzaki 2010). Deciduous trees disperse numerous seeds to treeless areas after wildfire (Landhausser and Wein 1993). The wind-dispersed seeds of broad-leaf trees should be captured by coarse and/or concave ground surfaces, even on steep slopes. Revegetation after wildfire is determined not only by fire severity but also pre-disturbance conditions, including topography, and is not interpreted by a vegetation gradient related to time after disturbance (Rydgren et al. 2004).

Lifeforms in relation to seeders vs sprouters

All the deciduous trees were seeders at Poker Flat after the wildfire. In contrast, shrubs, most of which were sprouters, increased in cover annually, depending on the unburned surface. The clonal shrubs (e.g., *Vaccinium uliginosum* and *V. myrtilloides*) rarely produced seedlings on the *Sphagnum* mat (Eriksson 1989; Moola and Mallik 1998). Perennial grasses and sedges also reproduced vegetatively on unburned surface. Sprouters, such as perennial sedges, grasses, and shrubs are common when wildfire is not severe in a mixed-evergreen forest, Oregon (Donato et al. 2009). Revegetation is progressed by species in soil bud-bank (sprouters) and seed-bank after weak fire that consumes relatively-shallow moss layer in a boreal Swedish forest, while the revegetation is conducted by seeders after deep-burning fire (Schimmel and Granstrom 1996).

Such differences in regeneration strategies between species or lifeforms determine the plant community structures and should be used for predicting successional sere with different fire severities.

Temporal changes in vegetation with reference to habitat preferences

Sphagnum gradually decreased in cover for five years and did not re-colonize on the burned surface, due probably to post-fire stresses such as by desiccation. Albedo was reduced when the plant cover burned and did not return to the pre-fire status until plant cover became high (Tsuyuzaki et al. 2009). Low albedo leads to high ground-surface temperature and deep thaw depth (Jorgenson et al. 2001). These changes induce drought stresses for plants, particularly *Sphagnum* mosses, which require more water than vascular plants for the short term. This is because they develop shallow belowground organs for water acquisition and have no special organs for water transportation (Schouwenaars and Gosen 2007; Thompson and Waddington 2008). *Sphagnum* is likely to recover slower than vascular plants, particularly when ground surface is completely consumed by wildfire.

The increase in moss cover was mostly derived from the pioneer mosses, represented by

Polytrichum strictum and *Ceratodon purpureus*, on burned surfaces. Mineral soils exposed after disturbances, including severe wildfires, were covered with the pioneer mosses, and the vegetation was totally dissimilar to the pre-fire vegetation (Bernhardt et al. 2011). The aboveground biomass and productivity of non-vascular plants are lower on drier sites after a wildfire in interior Alaska, in particular, where *Ceratodon* and/or *Polytrichum* were dominant (Mack et al. 2008). Deciduous trees established with *P. commune* and *C. purpureus* on the burned surface. Since these two mosses established well on burned surfaces, they should induce the alteration of tree regeneration. The revegetation did not go towards *P. mariana* forest directly when wildfire burns out the ground surface.

Forest revegetation seems to be delayed by severe burn (Dale et al. 2001). Furthermore, the complete removal of moss and organic matter promoted the colonization of non-*Sphagnum* mosses and deciduous trees. In conclusion, seeders represented by broad-leaved trees have a great role on revegetation after severe wildfire in *P. mariana* forests that completely burns out not only *Sphagnum* mat but also organic layer.

Acknowledgements The authors thank Miyoshi, C., Takeuchi, F., and Hirata, A.K.B. for field assistance. We also thank the staff members of Poker Flat Research Range for research permission. M. Higuchi, NMNS, identified the mosses. T. Alton, IARC, improved English. This work is partly supported by IARC-JAXA (Terrestrial Team Leader, M. Fukuda) and JSPS.

References

- ACRC (The Alaska Climate Research Center) (2007) Climate Data: climatological data and information for locations across Alaska. Division normals (Interior). <http://climate.gi.alaska.edu/Climate/Normals/interior.html>
- Anisimov OA, Vaughan DG, Callaghan TV, Furgal C, Marchant H, Prowse DT, Vilhjalmsen H, Walsh JE (2007) Polar regions (Arctic and Antarctic). Climate Change 2007: Impacts,

Adaptation and Vulnerability. In: the 4th Assessment Report of IPCC (Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE, eds.). Cambridge University Press, Cambridge. 653-685 pp

Bernhardt EL, Hollingsworth TN, Chapin III FS (2011) Fire severity mediates climate-driven shifts in understory community composition of black spruce stands of interior Alaska. *J Veg Sci* 22: 32-44

Betts EF, Jones Jr JB (2009) Impact of wildfire on stream nutrient chemistry and ecosystem metabolism in boreal forest catchments of interior Alaska. *Arc Antarc Alpine Res* 41: 407-417

Burn CR (1998) The response (1958-1997) of permafrost and near-surface ground temperatures to forest fire, Takhini River Valley, southern Yukon Territory. *Canad J Earth Sci* 35: 184-199

Chambers SD, Beringer J, Randerson JT, Chapin III FS (2005) Fire effects on net radiation and energy partitioning: contrasting responses of tundra and boreal forest ecosystems. *J Geophys Res* 110: D09106. doi:10.1029/2004JD005299

Chapin III FS, Oswood MW, Cleve KV, Verbyla DL (2006) Alaska's changing boreal forest. Oxford University Press, New York

Dale VH, Joyce LA, McNulty S, Neilson RP, Ayres MP, Flannigan MD, Hanson PJ, Irland LC, Lugo AE, Peterson CJ, Simberloff D, Swanson FJ, Stocks BJ, Wotton BM (2001) Climate change and forest disturbances. *BioSci* 51: 723-734

Davis N (2001) Permafrost. A guide to frozen ground in transition. University of Alaska Press, Fairbanks

Donato DC, Fontaine JB, Robinson WD, Kauffman JB, Law BE (2009) Vegetation response to a short interval between high-severity wildfires in a mixed-evergreen forest. *J Ecol* 97: 142-154

Engelmark O (1999) Boreal forest disturbances. In: Walker LR (ed.). *Ecosystems of the world 16: Ecosystems of disturbed ground*. Elsevier, Amsterdam. pp. 161-186

- 1 Eriksson O (1989) Seedling dynamics and life histories in clonal plants. *Oikos* 55: 231-238
- 2 Flannigan MD, Stocks BJ, Wotton BM (2000) Climate change and forest fires. *Sci Total*
3 *Environ* 262: 221-229
- 4 Frazer GW, Canham CD, Lertzman KP (1999) Gap light analyzer (GLA), version 2.0:
5 imaging software to extract canopy structure and gap light transmission indices from
6 true-colour fisheye photographs, users manual and program documentation. Simon
7 Fraser University, Burnaby, British Columbia, and the Institute of Ecosystem Studies,
8 Millbrook
- 9 Greene DF, Noel J, Bergeron Y, Rousseau M, Gauthier S (2004) Recruitment of *Picea*
10 *mariana*, *Pinus banksiana*, and *Populus tremuloides* across a burn severity gradient
11 following wildfire in the southern boreal forest of Quebec. *Canad J For Res* 34:
12 1845-1857
- 13 Gurvich DE, Enrico L, Cingolani AM (2005) Linking plant functional traits with post-fire
14 sprouting vigour in woody species in central Argentina. *Austral Ecol* 30: 789-796
- 15 Heijmans MMPD, Arp WJ, Chapin III FS (2004) Carbon dioxide and water vapour exchange
16 from understory species in boreal forest. *Agr For Meteorol* 123: 135-147
- 17 Hollingworth TN, Walker MD, Chapin III FS, Parsons AL (2006) Scale-dependent
18 environmental controls over species composition in Alaskan black spruce communities.
19 *Canad J For Res* 36: 1781-1796
- 20 Hollingworth TN, Johnstone F, Bernhardt EL, Chapin FS (2013) Fire severity filters
21 regeneration traits to shape community assembly in Alaska's boreal forest. *PloS One* 8:
22 e56033. doi:10.1371/journal.pone.0056033
- 23 Ilisson T, Chen HYH (2009) The direct regeneration hypothesis in northern forests. *J Veg Sci*
24 20: 735-744
- 25 Jayen K, Leduc A, Bergeron Y (2006) Effect of fire severity on regeneration success in the
26 boreal forest of northwest Quebec, Canada. *EcoSci* 13: 143-151
- 27 Johnson EA, Miyanishi K (2007) Plant disturbance ecology. Academic Press, Amsterdam

- 1 Johnstone JF, Chapin III FS (2006) Effects of soil burn severity on post-fire tree recruitment
2 in boreal forest. *Ecosystems* 9: 14-331
- 3 Johnstone JF, Rupp TS, Olson M, Verbyla D (2011) Modeling impacts of fire severity on
4 successional trajectories and future fire behavior in Alaskan boreal forests. *Landscape*
5 *Ecol* 26: 487-500
- 6 Jorgenson MT, Rachine CH, Walters JC, Osterkamp TE (2001) Permafrost degradation and
7 ecological changes associated with a warming climate in central Alaska. *Climatic*
8 *Change* 48: 551-579
- 9 Kaplan JO, New M (2006) Arctic climate change with a 2°C global warming: timing,
10 climate patterns and vegetation change. *Climatic Change* 79: 213-241
- 11 Kasischke ES, Turetsky Mr (2006) Recent changes in the fire regime across the North
12 American boreal region. *Geophys Res Lett* 33: doi.10.1029/2006GL025677
- 13 Koyama A, Tsuyuzaki S (2010) Effects of sedge and cottongrass tussocks on plant
14 establishment patterns in a post-mined peatland, northern Japan. *Wetlands Ecol Manag*
15 18: 135-148
- 16 Landhausser SM, Wein RW (1993) Postfire vegetation recovery and tree establishment at
17 the arctic treeline: climate-change-vegetation-response hypotheses. *J Ecol* 81: 665-672
- 18 Mack MC, Treseder KK, Manies KL, Harden JW, Schuur EAG, Vogel JG, Randerson JT,
19 Chapin III FS (2008) Recovery of aboveground plant biomass and productivity after fire
20 in mesic and dry black spruce forests of interior Alaska. *Ecosystems* 11: 209-225
- 21 McCune, B., Grace, J.B. 2002. Analysis of ecological communities. MjM Software Design,
22 Gleneden Beach, Oregon
- 23 Moola FM, Mallik AU (1998) Morphological plasticity and regeneration strategies of velvet
24 leaf blueberry (*Vaccinium myrtilloides* Michx.) following canopy disturbance in boreal
25 mixedwood forests. *For Ecol Manag* 111: 35-50
- 26 Nicole J, Yves B (2006) Facilitative succession in a boreal bryophyte community driven by
27 changes in available moisture and light. *J Veg Sci* 17: 65-76

- R Development Core Team (2010) R: A language and environment for statistical computing.
R Foundation for Statistical Computing, Vienna, Austria
- Reich PB, Bakken P, Carlson D, Frelich LE, Friedman SK, Grigal DF. 2001. Influence of
logging, fire, and forest type on biodiversity and productivity in southern boreal forests.
Ecology 82: 2731-2748
- Rydgren K, Okland RH, Hestmark G (2004) Disturbance severity and community resilience
in a boreal forest. Ecology 85: 1906-1915.
- Santana VM, Baeza MJ, Maestre FT (2012) Seedling establishment along post-fire
succession in Mediterranean shrublands dominated by obligate seeders. Acta Oecol 39:
51-60
- Schimmel J, Granstrom A (1996) Fire severity and vegetation response in the boreal Swedish
forest. Ecology 77: 1436-1450
- Schouwenaars JM, Gosen AM (2007) The sensitivity of *Sphagnum* to surface layer
conditions in a re-wetted bog: a simulation study of water stress. Mires Peat 2: (Article
02) 1-19.
- Shetler GS, Turetsky MR, Kane E, Kasischke E (2008) Sphagnum mosses limit total carbon
consumption during fire in Alaskan black spruce forests. Canadian Journal of Forest
Research 38: 2328-2336
- Siegwart Collier LC, Mallik AU (2010) Does post-fire abiotic habitat filtering create
divergent plant communities in black spruce forests of eastern Canada? Oecologia 164:
465-477.
- Thompson DK, Waddington JM (2008) *Sphagnum* under pressure: towards an
ecohydrological approach to examining *Sphagnum* productivity. Ecohydrol 1: 299-308
- Tsuyuzaki S, Kushida K, Kodama Y (2009) Recovery of surface albedo and plant cover after
wildfire in a *Picea mariana* forest in interior Alaska. Climatic Change 93: 517-525
- Whinam J, Hope G, Good, R, Wright G (2010) Post-fire experimental trials of vegetation
restoration techniques in the peatlands of Namadgi (ACT) and Kosciuszko National

1 Parks (NSW), Australia. In: Haberle SG, Stevenson J, Prebble M (ed.). *Altered*
2 *Ecologies: Fire, climate and human influence on terrestrial landscapes* (Terra Australis
3 32), ANU ePress, Canberra Australia, pp. 363-379

Fig. 1 NMDS ordination diagram on species scores in 400 quadrats (80 × 5 years) surveyed from 2005 to 2009 after the 2004 wildfire. Life forms: closed circles = herbs and ferns (13 taxa), open circles = shrubs (12), closed triangles = trees (4), and open triangles = mosses and lichens (13). Species codes explained in the text: Car big = *Carex bigelowii*, Cal can = *Calamagrostis canadensis*, Epi ang = *Epilobium angustifolium*, Led gro = *Ledum groenlandicum*, Vac vit = *Vaccinium vitis-idaea*, Vac uli = *Vaccinium uliginosum*, Bet nan = *Betula nana*, Oxy mic = *Oxycoccus microcarpus*, Pic mar = *Picea mariana*, Bet neo = *Betula neoalaskana*, Pop tre = *Populus tremuloides*, Pol for = *Polytrichum commune*, Sph fus = *Sphagnum fuscum*, and Cer pur = *Ceratodon purpureus*. Environmental factors: Yr = years after wildfire, Brn: burned area (%), Cbn: completely burned area (%) where the soils were exposed. Cnp = canopy openness (%), Elv = elevation (m), Slp = slope gradient (°), and Asp = aspect (°).

Fig. 2 NMDS ordination diagram on quadrat scores in 400 quadrats (80×5 years) surveyed for five years after the 2004 wildfire. The connected lines show the same quadrats surveyed from 2005 to 2009. NMDS scores at the first survey, i.e., in 2005, are shown by solid circles, and the others are by open circles. The arrows of environmental factors are not shown.

Fig. 3 Percentage similarities between *Sphagnum fuscum* surface and species or lifeform (shown by solid lines with closed symbols), and between non-*Sphagnum* surface and species (by interrupted lines with open symbols). Gray symbols and lines indicate the lifeforms. Herb: ●/○ = *Carex bigelowii*, ■/□ = *Calamagrostis canadensis*, ▲/△ = *Epilobium angustifolium*, Shrub: ●/○ = *Ledum groenlandicum*, ■/□ = *Vaccinium vitis-idaea*, ▲/△ = *Vaccinium uliginosum*, ◆/◇ = *Betula nana*, ▼/▽ = *Oxycoccus microcarpus*, Tree: ●/○ = *Picea mariana*, ■/□ = *Betula neoalaskana*, ▲/△ = *Populus tremuloides*, Moss: ●/○ = *Polytrichum commune*, ■/□ = *Ceratodon purpureus*. Note that the scales of y-axis are different between life forms and that the similarities of *P. mariana* and tree on unburned surface are overlapped.