Title	The systemic administration of an anti-miRNA oligonucleotide encapsulated pH-sensitive liposome results in reduced level of hepatic microRNA-122 in mice
Author(s)	Hatakeyama, Hiroto; Murata, Manami; Sato, Yusuke; Takahashi, Mayumi; Minakawa, Noriaki; Matsuda, Akira; Harashima, Hideyoshi
Citation	Journal of controlled release, 173, 43-50 https://doi.org/10.1016/j.jconrel.2013.10.023
Issue Date	2014-01-10
Doc URL	http://hdl.handle.net/2115/57318
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	WoS_64270_Sato_SI.pdf (Supplementary information)

[Instructions for use](#)

Supplementary information

Figure S1 Characterization of the AMO122-YSK05-MEND. (a) Gel electrophoretic analysis of the encapsulation of AMO122 in YSK05-MEND. Free AMO122 and AMO122-YSK05-MEND were treated with Triton X-100 to disrupt the lipid envelope structure. Samples were subjected to agarose gel electrophoresis. A 20 bp DNA ladder (Takara) was applied in lane 1. (b) Determination of pKa by TNS. YSK05-MENDs were diluted to 30 μ M total lipid in buffers containing 6 μ M TNS at various pH levels. The highest fluorescence was assigned as 1.0 and the relative fluorescent intensities are represented. Data represent the mean value $\pm$ SD (n=3). Curve fitting (gray line) was calculated using Sigmaplot 12.

Figure S2 The effect of AMO10b encapsulated in the YSK05-MEND (AMO10b-YSK05-MEND) on the expression of miR-122 and target genes.

AMO10b for miR-10b (5'-C_sC_sA_sA_sA_sU_sU_sC_sG_sG_sU_sU_sC_sU_sA_sC_sA_sG_sG_sG_sU_sA-3') was purchased from Hokkaido System Science (Sapporo, Japan) (2'-OMe-modified nucleotides; subscript 's' represents a phosphorothioate linkage). The diameter, pid, zeta-potential, and recovery and encapsulation efficiency of AMO10b formulated YSK-5-MEND (AMO10b-YSK05-MEND) were 67 ± 5 nm, 0.181 ± 0.04 , 0.1 ± 2.4 mV, $92 \pm 2\%$ and $95 \pm 7\%$, respectively. The characteristics of AMO10b-YSK05-MEND were comparable to that encapsulating AMO122. Either AMO122-YSK05-MEND or AMO10b-YSK05-MEND was successively administered via the tail vein of ICR mice at a dose of 1 mg/kg at day 0, 2 and 4. At day 6, liver was collected and subjected to real-time PCR for the determination of the expression level of miR-122 and target genes as described in the materials and methods section. The administration of AMO10b-YSK05-MEND showed no change in the expression of miR-122 and target genes, while AMO122-YSK05-MEND reduced the expression of miR-122 and enhanced the target *Aldoa*, *Bckdk* and *Ndr3*. These results indicate that miR-122 inhibition and the subsequent enhancement of target genes occurred in an AMO122 sequence

specific manner. Data are represented as the mean $\pm$ SD (n=3). **P<0.01.
*P<0.05 v.s. PBS.

Figure S3 Expression level of LDL receptor on Hepa1c1c7 cells.

Expression level of LDL receptor was determined by flow cytometric analysis. Hepa1c1c7 cells (1.0×10^5 cells/well) were seeded in a 6-well plate, and then incubated for 24 hr. The cells were washed with PBS twice, and detached by the treatment with 0.05% trypsin-PBS. The detached cells were incubated with anti-murine LDL receptor antibody (1:100 dilution, Abcam, Tokyo, Japan) for 30 min at 4°C, followed by the incubation with a FITC labeled anti-armerian hamster IgG antibody (1:100 dilution, Santa Cruz Biotechnology, Santa Cruz, CA, USA) for 30 min at 4°C. Ten thousand cells per sample were analyzed using a FACSCalibur flow cytometer (BD Biosciences). The results were analyzed using Cell-Quest (BD Biosciences).