



Title	Gauge origin of discrete flavor symmetries in heterotic orbifolds
Author(s)	Beye, Florian; Kobayashi, Tatsuo; Kuwakino, Shogo
Citation	Physics Letters B, 736, 433-437 https://doi.org/10.1016/j.physletb.2014.07.058
Issue Date	2014-09-07
Doc URL	http://hdl.handle.net/2115/57334
Rights	(C) 2014 The Authors. Published by Elsevier B.V.
Rights(URL)	http://creativecommons.org/licenses/by/3.0/
Type	article
File Information	PLB_736_433-.pdf



[Instructions for use](#)



Gauge origin of discrete flavor symmetries in heterotic orbifolds



Florian Beyea^a, Tatsuo Kobayashi^b, Shogo Kuwakino^{c,*}

^a Department of Physics, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8602, Japan

^b Department of Physics, Hokkaido University, Sapporo 060-0810, Japan

^c Department of Physics, Chung-Yuan Christian University, 200 Chung-Pei Rd., Chung-Li 320, Taiwan

ARTICLE INFO

Article history:

Received 1 July 2014

Received in revised form 28 July 2014

Accepted 29 July 2014

Available online 4 August 2014

Editor: M. Cvetič

ABSTRACT

We show that non-Abelian discrete symmetries in orbifold string models have a gauge origin. This can be understood when looking at the vicinity of a symmetry enhanced point in moduli space. At such an enhanced point, orbifold fixed points are characterized by an enhanced gauge symmetry. This gauge symmetry can be broken to a discrete subgroup by a nontrivial vacuum expectation value of the Kähler modulus T . Using this mechanism it is shown that the $\Delta(54)$ non-Abelian discrete symmetry group originates from a $SU(3)$ gauge symmetry, whereas the D_4 symmetry group is obtained from a $SU(2)$ gauge symmetry.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP³.

1. Introduction

It is important to understand the flavor structure of the standard model of particle physics. Quark and lepton masses are hierarchical. Two of the mixing angles in the lepton sector are large, while the mixing angles in the quark sector are suppressed, except for the Cabibbo angle. Non-Abelian discrete flavor symmetries may be useful to understand this flavor structure. Indeed, many works have considered field-theoretical model building with various non-Abelian discrete flavor symmetries (see [1–3] for reviews).

Understanding the origin of non-Abelian flavor symmetries is an important issue we have to address. It is known that several phenomenologically interesting non-Abelian discrete symmetries can be derived from string models.¹ In intersecting and magnetized D-brane models, the non-Abelian discrete symmetries D_4 , $\Delta(27)$ and $\Delta(54)$ can be realized [5–8]. Also, their gauge origins have been studied [6]. In heterotic orbifold compactifications [9–17] (also see a review [18]), non-Abelian discrete symmetries appear due to geometrical properties of orbifold fixed points and certain properties of closed string interactions [19]. First, there are permutation symmetries of orbifold fixed points. Then, there are string selection rules which determine interactions between orbifold sectors. The combination of these two kinds of discrete sym-

metries leads to a non-Abelian discrete symmetry. In particular, it is known that the D_4 group emerges from the one-dimensional orbifold S^1/Z_2 , and that the $\Delta(54)$ group is obtained from the two-dimensional orbifold T^2/Z_3 . The phenomenological applications of the string-derived non-Abelian discrete symmetries are analyzed e.g. in [20].

In this paper we point out that these non-Abelian discrete flavor symmetries originate from a gauge symmetry. To see this, we consider a heterotic orbifold model compactified on some six-dimensional orbifold. The gauge symmetry G_{gauge} of this orbifold model is, if we do not turn on any Wilson lines, a subgroup of $E_8 \times E_8$ which survives the orbifold projection. In addition, from the argument in [19], we can derive a non-Abelian discrete symmetry G_{discrete} . Then, the effective action of this model can be derived from $G_{\text{gauge}} \times G_{\text{discrete}}$ symmetry invariance.² However, this situation slightly changes if we set the model to be at a symmetry enhanced point in moduli space. At that special point, the gauge symmetry of the model is enlarged to $G_{\text{gauge}} \times G_{\text{enhanced}}$, where G_{enhanced} is a gauge symmetry group. The maximal rank of the enhanced gauge symmetry G_{enhanced} is six, because we compactify six internal dimensions. At this specific point in moduli space, orbifold fixed points are characterized by gauge charges of G_{enhanced} , and the spectrum is extended by additional massless fields charged under G_{enhanced} . Furthermore, the Kähler moduli fields T in the untwisted sector obtain G_{enhanced} -charges and a non-zero vacuum expectation value (VEV) of T corresponds to a movement away

* Corresponding author.

E-mail addresses: fbeye@eken.phys.nagoya-u.ac.jp (F. Beyea), kobayashi@particle.sci.hokudai.ac.jp (T. Kobayashi), kuwakino@cycu.edu.tw (S. Kuwakino).

¹ In [4], field theoretical models where non-Abelian discrete groups are embedded into non-Abelian gauge groups are considered.

² Here we do not consider the R -charge invariance since this is not relevant to our discussion.

from the enhanced point. This argument suggests the possibility that the non-Abelian discrete symmetry G_{discrete} is enlarged to a continuous gauge symmetry G_{enhanced} at the symmetry enhanced point. In other words, it suggests a gauge origin of the non-Abelian discrete symmetry. Moreover, the group G_{enhanced} originates from a larger non-Abelian gauge symmetry that exists before the orbifolding. We will show this explicitly in the following.

2. Gauge origin of non-Abelian discrete symmetry

In this section we demonstrate the gauge origin of non-Abelian discrete symmetries in heterotic orbifold models. We concentrate on the phenomenologically interesting non-Abelian discrete symmetries D_4 and $\Delta(54)$ which are known to arise from orbifold models.

2.1. D_4 non-Abelian discrete symmetry

First, we study a possible gauge origin of the D_4 non-Abelian discrete symmetry. This symmetry is associated with the one-dimensional S^1/Z_2 orbifold. Here, we consider the heterotic string on a S^1/Z_2 orbifold, but it is straightforward to extend our argument to T^2/Z_2 or $T^6/(Z_2 \times Z_2)$. The coordinate corresponding to the one dimension of S^1 is denoted by X . It suffices to discuss only the left-movers in order to develop our argument. Let us start with the discussion on S^1 without the Z_2 orbifold. There is always a $U(1)$ symmetry associated with the current $H = i\partial X$. At a specific point in the moduli space, i.e. at a certain radius of S^1 , two other massless vector bosons appear and the gauge symmetry is enhanced from $U(1)$ to $SU(2)$. Their currents are written as

$$E_{\pm} = e^{\pm i\alpha X}, \tag{1}$$

where $\alpha = \sqrt{2}$ is a simple root of the $SU(2)$ group. These currents, H and E_{\pm} , satisfy the $su(2)$ Kac–Moody algebra.

Now, let us study the Z_2 orbifolding $X \rightarrow -X$. The current $H = i\partial X$ is not invariant under this reflection and the corresponding $U(1)$ symmetry is broken. However, the linear combination $E_+ + E_-$ is Z_2 -invariant and the corresponding $U(1)$ symmetry remains on S^1/Z_2 . Thus, the $SU(2)$ group breaks down to $U(1)$ by orbifolding. Note that the rank is not reduced by this kind of orbifolding. It is convenient to use the following basis,

$$H' = i\partial X' = \frac{1}{\sqrt{2}}(E_+ + E_-), \tag{2}$$

$$E'_{\pm} = e^{\pm i\alpha X'} = \frac{1}{\sqrt{2}}H \mp \frac{1}{2}(E_+ - E_-). \tag{3}$$

The introduction of the boson field X' is justified because H' and E'_{\pm} satisfy the same operator product expansions (OPEs) as the original currents H and E_{\pm} . The invariant current H' corresponds to the $U(1)$ gauge boson. The E'_{\pm} transform as

$$E'_{\pm} \rightarrow -E'_{\pm} \tag{4}$$

under the Z_2 reflection and correspond to untwisted matter fields U_1 and U_2 with $U(1)$ charges $\pm\alpha$. In addition, there are other untwisted matter fields U which have vanishing $U(1)$ charge, but are charged under an unbroken subgroup of $E_8 \times E_8$.

From (4), it turns out that the Z_2 reflection is represented by a shift action in the X' coordinate,

$$X' \rightarrow X' + 2\pi \frac{w}{2}, \tag{5}$$

where $w = 1/\sqrt{2}$ is the fundamental weight of $SU(2)$. That is, the Z_2 -twisted orbifold on X is equivalent to a Z_2 -shifted orbifold on

Table 1

Field contents of $U(1) \times Z_2$ model from Z_2 orbifold. $U(1)$ charges are shown. Charges under the Z_4 unbroken subgroup of the $U(1)$ group are also shown.

Sector	Field	$U(1)$ charge	Z_4 charge
U	U	0	0
U	U_1	α	0
U	U_2	$-\alpha$	0
T	M_1	$\frac{\alpha}{4}$	$\frac{1}{4}$
T	M_2	$-\frac{\alpha}{4}$	$-\frac{1}{4}$

X' with the shift vector $s = w/2$ (see e.g., [21]). In the twist representation, there are two fixed points on the Z_2 orbifold, to each of which corresponds a twisted state. Note that the one-dimensional bosonic string X with the Z_2 -twisted boundary condition has a contribution of $h = 1/16$ to the conformal dimension. In the shift representation, the two twisted states can be understood as follows. Before the shifting, X' also represents a coordinate on S^1 at the enhanced point, so the left-mover momenta p_L lie on the momentum lattice

$$\Gamma_{SU(2)} \cup (\Gamma_{SU(2)} + w), \tag{6}$$

where $\Gamma_{SU(2)}$ is the $SU(2)$ root lattice, $\Gamma_{SU(2)} \equiv n\alpha$ with integer n . Then, the left-mover momenta in the Z_2 -shifted sector lie on the original momentum lattice shifted by the shift vector $s = w/2$, i.e.

$$\left(\Gamma_{SU(2)} + \frac{w}{2}\right) \cup \left(\Gamma_{SU(2)} + \frac{3w}{2}\right). \tag{7}$$

Thus, the shifted vacuum is degenerate and the ground states have momenta $p_L = \pm\alpha/4$. These states correspond to charged matter fields M_1 and M_2 . Note that $p_L^2/2 = 1/16$, which is exactly the same as the conformal dimension $h = 1/16$ of the twisted vacuum in the twist representation. Indeed, the twisted states can be related to the shifted states by a change of basis [21]. Notice that the twisted states have no definite $U(1)$ charge, but the shifted states do. Table 1 shows corresponding matter fields and their $U(1)$ charges.

From Table 1, we find that there is an additional Z_2 symmetry of the matter contents at the lowest mass level (in a complete model, these can correspond to massless states): Transforming the $U(1)$ -charges q as

$$q \rightarrow -q, \tag{8}$$

while at the same time permuting the fields as $U_1 \leftrightarrow U_2$ and $M_1 \leftrightarrow M_2$ maps the spectrum onto itself. The action on the U_i and M_i fields is described by the 2×2 matrix

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \tag{9}$$

This Z_2 symmetry does not commute with the $U(1)$ gauge symmetry and it turns out that one obtains a symmetry of semi-direct product structure, $U(1) \times Z_2$.

In the twist representation, this model contains the Kähler modulus field T , which corresponds to the current H and is charged under the $U(1)$ group. In the shift representation, the field T is described by the fields U_i as

$$T = \frac{1}{\sqrt{2}}(U_1 + U_2). \tag{10}$$

Now we consider the situation where our orbifold moves away from the enhanced point by taking a specific VEV of the Kähler modulus field T which corresponds to the VEV direction

$$\langle U_1 \rangle = \langle U_2 \rangle. \tag{11}$$

Note that this VEV relation maintains the Z_2 discrete symmetry (9). Moreover, since the fields U_1 and U_2 are charged under the $U(1)$ gauge symmetry and due to the presence of the M_i fields, the VEV breaks $U(1)$ down to a discrete subgroup Z_4 . The Z_4 charge is $1/4$ for M_1 and $-1/4$ for M_2 as listed in Table 1. Written as a 2×2 matrix, the Z_4 action is described by

$$\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}. \tag{12}$$

The matrices (9) and (12) are nothing but the generators of $D_4 \simeq Z_4 \times Z_2$. After the VEV, the field U transforms as the trivial singlet $\mathbf{1}$, and (M_1, M_2) forms a $\mathbf{2}$ representation under the D_4 group. This reproduces the known result for a general radius of S_1 [19]. The pattern of symmetry breaking we have shown here is summarized as follows:

$$SU(2) \xrightarrow{\text{orbifolding}} U(1) \times Z_2 \xrightarrow{(T)} D_4. \tag{13}$$

The other VEV directions of U_1 and U_2 break $U(1) \times Z_2$ to Z_4 . However, while the VEV direction defined by Eq. (11) is D-flat, the other cases do not correspond to D-flat directions and the resulting symmetries have no geometrical interpretation.

2.2. $\Delta(54)$ non-Abelian discrete symmetry

Next, we consider the two-dimensional T^2/Z_3 orbifold case which is associated with the $\Delta(54)$ non-Abelian discrete symmetry. Here, we study the heterotic string on a T^2/Z_3 orbifold. However, our argument straightforwardly extends to orbifolds such as T^6/Z_3 . The coordinates on T^2 are denoted by X^1 and X^2 . We start with the discussion of the two-dimensional torus, T^2 , without orbifolding. There is always a $U(1)^2$ symmetry corresponding to the two currents, $H_1 = i\partial X^1$ and $H_2 = i\partial X^2$. At a certain point in the moduli space of T^2 , there appear additional six massless gauge bosons. Then, the gauge symmetry is enhanced from $U(1)^2$ to $SU(3)$. The corresponding Kac–Moody currents are

$$E_{\pm 1,0}, \quad E_{0,\pm 1}, \quad E_{\pm 1,\pm 1}, \tag{14}$$

with

$$E_{n_1,n_2} = e^{i \sum_{i=1,2} (n_1 \alpha_1^i + n_2 \alpha_2^i) X^i}, \tag{15}$$

where α_1 and α_2 denote simple roots of $SU(3)$, i.e. $\alpha_1 = (\sqrt{2}, 0)$ and $\alpha_2 = (-\sqrt{2}/2, \sqrt{6}/2)$. These currents, H_i and E_{n_1,n_2} , satisfy the $su(3)$ Kac–Moody algebra.

Now, let us study the Z_3 orbifolding,

$$Z \rightarrow \omega^{-1}Z, \tag{16}$$

where $Z = X^1 + iX^2$ and $\omega = e^{2\pi i/3}$. The currents H_i and their linear combinations are not Z_3 -invariant and the corresponding gauge symmetries are broken. On the other hand, two independent linear combinations of E_{n_1,n_2} are Z_3 -invariant and correspond to a $U(1)^2$ symmetry that remains on the T^2/Z_3 orbifold. Thus, the $SU(3)$ gauge group is broken to $U(1)^2$ by the orbifolding. It is convenient to use the following basis,

$$H'_1 = \frac{i}{\sqrt{2}}(E_1^1 - E_1^2), \tag{17}$$

$$H'_2 = -\frac{1}{\sqrt{2}}(E_1^1 + E_1^2), \tag{18}$$

$$E'_{1,0} = \frac{1}{\sqrt{3}}(iH_{\omega^{-1}} + E_{\omega^{-1}}^1 + E_{\omega^{-1}}^2), \tag{19}$$

$$E'_{0,1} = \frac{1}{\sqrt{3}}(iH_{\omega^{-1}} + \omega E_{\omega^{-1}}^1 + \omega^{-1} E_{\omega^{-1}}^2), \tag{20}$$

$$E'_{-1,-1} = \frac{1}{\sqrt{3}}(iH_{\omega^{-1}} + \omega^{-1} E_{\omega^{-1}}^1 + \omega E_{\omega^{-1}}^2), \tag{21}$$

$$E'_{-1,0} = \frac{1}{\sqrt{3}}(-iH_{\omega} + E_{\omega}^1 + E_{\omega}^2), \tag{22}$$

$$E'_{0,-1} = \frac{1}{\sqrt{3}}(-iH_{\omega} + \omega E_{\omega}^1 + \omega^{-1} E_{\omega}^2), \tag{23}$$

$$E'_{1,1} = \frac{1}{\sqrt{3}}(-iH_{\omega} + \omega^{-1} E_{\omega}^1 + \omega E_{\omega}^2), \tag{24}$$

where

$$H_{\omega^{-1}} = \frac{1}{\sqrt{2}}(H_1 + iH_2), \tag{25}$$

$$H_{\omega} = \frac{1}{\sqrt{2}}(H_1 - iH_2), \tag{26}$$

$$E_{\omega^{-k}}^1 = \frac{1}{\sqrt{3}}(E_{1,0} + \omega^k E_{0,1} + \omega^{-k} E_{-1,-1}), \tag{27}$$

$$E_{\omega^{-k}}^2 = \frac{1}{\sqrt{3}}(E_{-1,0} + \omega^k E_{0,-1} + \omega^{-k} E_{1,1}). \tag{28}$$

The E'_{n_1,n_2} correspond to states with charges $(n_1 \alpha_1^1 + n_2 \alpha_2^1, n_1 \alpha_1^2 + n_2 \alpha_2^2)$ under the unbroken $U(1)^2$. They transform under the Z_3 twist action as follows:

$$\begin{aligned} E'_{-1,0} &\rightarrow \omega E'_{-1,0}, & E'_{0,-1} &\rightarrow \omega E'_{0,-1}, \\ E'_{1,1} &\rightarrow \omega E'_{1,1}, & E'_{1,0} &\rightarrow \omega^{-1} E'_{1,0}, \\ E'_{0,1} &\rightarrow \omega^{-1} E'_{0,1}, & E'_{-1,-1} &\rightarrow \omega^{-1} E'_{-1,-1}. \end{aligned} \tag{29}$$

Thus, the first three E'_{n_1,n_2} correspond to untwisted matter fields with charges $-\alpha_1$, $-\alpha_2$ and $\alpha_1 + \alpha_2$ under the unbroken $U(1)^2$. We denote them as U_1, U_2 and U_3 , respectively. The other three are their CPT conjugates. In addition, there are other untwisted matter fields U which have vanishing $U(1)^2$ charges, but are charged under an unbroken subgroup of $E_8 \times E_8$.

Now, since the primed currents fulfill the same OPEs as their unprimed counterparts, it is justified to introduce bosons X'^i , so that

$$\begin{aligned} H'^i &= i\partial X'^i \\ E'_{n_1,n_2} &= e^{i \sum_{i=1,2} (n_1 \alpha_1^i + n_2 \alpha_2^i) X'^i}. \end{aligned} \tag{30}$$

The Z_3 twist action on X^i can then be realized as a shift action on X'^i as

$$X'^i \rightarrow X'^i + 2\pi \frac{\alpha_1^i}{3}. \tag{31}$$

In the twist representation there are three fixed points on the T^2/Z_3 orbifold, to each of which corresponds a twisted state. The two-dimensional bosonic string with the Z_3 boundary condition has a contribution of $h = 1/9$ to the conformal dimension. As in the previous one-dimensional case, the twisted states can be described in the shift representation as follows. The left-moving momentum modes p_L of the torus-compactified $SU(3)$ model lie on the momentum lattice

$$\Gamma_{SU(3)} \cup (\Gamma_{SU(3)} + w_1) \cup (\Gamma_{SU(3)} - w_1), \tag{32}$$

where $\Gamma_{SU(3)}$ denotes the $SU(3)$ root lattice which is spanned by the simple roots of $SU(3)$, $\Gamma_{SU(3)} \equiv n_1 \alpha_1 + n_2 \alpha_2$, and $w_1 = (\sqrt{2}/2, \sqrt{6}/6)$ is the fundamental weight corresponding to α_1 . Then, the momenta p_L in the k -shifted sector lie on the momentum lattice shifted by the Z_3 shift vector $s = \alpha_1/3$,

Table 2
Field contents of $U(1)^2 \times S_3$ model from Z_3 orbifold. $U(1)^2$ charges are shown. Charges under the Z_3^2 unbroken subgroup of the $U(1)^2$ group are also shown.

Sector	Field	$U(1)^2$ charge	Z_3^2 charge
U	U	(0, 0)	(0, 0)
U	U_1	$-\alpha_1$	(0, 0)
U	U_2	$-\alpha_2$	(0, 0)
U	U_3	$\alpha_1 + \alpha_2$	(0, 0)
T	M_1	$\frac{\alpha_1}{3}$	$(\frac{1}{3}, \frac{1}{3})$
T	M_2	$\frac{\alpha_2}{3}$	$(-\frac{1}{3}, 0)$
T	M_3	$-\frac{\alpha_1 + \alpha_2}{3}$	$(0, -\frac{1}{3})$

$$\left(\Gamma_{SU(3)} + k \frac{\alpha_1}{3} \right) \cup \left(\Gamma_{SU(3)} + w_1 + k \frac{\alpha_1}{3} \right) \cup \left(\Gamma_{SU(3)} - w_1 + k \frac{\alpha_1}{3} \right). \tag{33}$$

For $k = 1$, there are three ground states with $p_L \in \{\alpha_1/3, \alpha_2/3, -(\alpha_1 + \alpha_2)/3\}$. They correspond to (would-be-massless) matter fields which we denote by M_1, M_2 and M_3 , respectively. These matter fields are shown in Table 2. The states for $k = -1$ correspond to CPT-conjugates. As expected, the shifted ground states have conformal dimension $h = p_L^2/2 = 1/9$, which coincides with the twisted ground states. Indeed, the shifted states are related to the twisted states by a change of basis [21]. The shifted states have definite $U(1)^2$ charges.

From Table 2, it turns out that the matter contents at the lowest mass level possess a S_3 permutation symmetry (in a complete model, these can correspond to massless states). Let S_3 be generated by a and b , with $a^3 = b^2 = (ab)^2 = 1$. Then, for a point (q_1, q_2) on the two-dimensional $U(1)^2$ charge plane, a and b shall act as

$$a: \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} \rightarrow \begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix}, \tag{34}$$

$$b: \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix}. \tag{35}$$

The action of a is equivalent to the replacement $\alpha_1 \rightarrow \alpha_2 \rightarrow -(\alpha_1 + \alpha_2) \rightarrow \alpha_1$. Then, the spectrum is left invariant if at the same time we transform the fields $F_i = (U_i, M_i)$ as $F_1 \rightarrow F_2 \rightarrow F_3 \rightarrow F_1$. The action of a on the F_i is described by the 3×3 matrix

$$\begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \tag{36}$$

The action of b corresponds to $\alpha_1 \leftrightarrow \alpha_1$ and $\alpha_2 \leftrightarrow -(\alpha_1 + \alpha_2)$, so simultaneously transforming $F_1 \leftrightarrow F_1$ and $F_2 \leftrightarrow F_3$ results in a symmetry of the spectrum. This transformation corresponds to the matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \tag{37}$$

The S_3 symmetry just shown does not commute with $U(1)^2$. Rather, S_3 and $U(1)^2$ combine to semi-direct product $U(1)^2 \rtimes S_3$.

Next we shall consider the situation where our orbifold moves away from the enhanced point by taking a certain VEV of the Kähler modulus field T , which corresponds to H_ω . The Kähler modulus can be described by the U_i fields as

$$T = \frac{1}{\sqrt{3}}(U_1 + U_2 + U_3). \tag{38}$$

The deformation is realized by the following VEV direction,

$$\langle U_1 \rangle = \langle U_2 \rangle = \langle U_3 \rangle. \tag{39}$$

Note that this VEV relation preserves the S_3 discrete symmetry generated by (36) and (37). However, the $U(1)^2$ gauge symmetry breaks down to a discrete Z_3^2 subgroup due to the presence of the M_i fields. The two Z_3 charges (z_1, z_2) are determined by $U(1)^2$ charges (u_1, u_2) as $z_1 = q_1/\sqrt{2} - q_2/\sqrt{6}, z_2 = q_1/\sqrt{2} + q_2/\sqrt{6}$. The Z_3^2 charges are listed in Table 2. The Z_3 actions are described by

$$\begin{pmatrix} \omega & 0 & 0 \\ 0 & \omega^{-1} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{40}$$

$$\begin{pmatrix} \omega & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega^{-1} \end{pmatrix}. \tag{41}$$

The matrices (36), (37), (40) and (41) are nothing but the generators of $\Delta(54) \simeq (Z_3 \times Z_3) \rtimes S_3$ in the $\mathbf{3}_{1(1)}$ representation [22]. Thus, the fields (M_1, M_2, M_3) transform as the $\mathbf{3}_{1(1)}$ under $\Delta(54)$, and the field U is the $\Delta(54)$ trivial singlet $\mathbf{1}$. This reproduces the known properties of ordinary Z_3 orbifold models at a general point in moduli space [19]. Summarizing, the origin of the $\Delta(54)$ discrete symmetry in orbifold models can be explained as follows:

$$SU(3) \xrightarrow[\text{orbifolding}]{} U(1)^2 \times S_3 \xrightarrow[\langle T \rangle]{} \Delta(54). \tag{42}$$

There are other VEV directions that one might consider. For $\langle U_1 \rangle \neq \langle U_2 \rangle = \langle U_3 \rangle = 0$ the $U(1)^2 \times S_3$ symmetry is broken to $(U(1) \times Z_2) \times Z_6$. In the case where $\langle U_1 \rangle \neq \langle U_2 \rangle = \langle U_3 \rangle \neq 0$ one obtains $Z_3 \times S_3$. Finally, when all VEVs are different, i.e. $\langle U_1 \rangle \neq \langle U_2 \rangle \neq \langle U_3 \rangle \neq \langle U_1 \rangle$ the symmetry is broken to $Z_3 \times Z_3$. However, while the VEV direction defined by (39) is D-flat, the other directions are not D-flat and do not allow for a geometrical interpretation.

3. Conclusion

We showed that non-Abelian discrete symmetries in heterotic orbifold models originate from a non-Abelian continuous gauge symmetry. The non-Abelian continuous gauge symmetry arises from torus-compactified extra dimensions at a special enhanced point in moduli space. In the two-dimensional orbifold case, by acting with Z_3 on the torus-compactified $SU(3)$ model, the non-Abelian gauge group $SU(3)$ is broken to a $U(1)^2$ subgroup. We observed that the matter contents of the orbifold model possess a S_3 symmetry which is understood to act on the two-dimensional $U(1)^2$ charge plane. The resulting orbifold model then has a symmetry of semi-direct product structure, $U(1)^2 \rtimes S_3$. In the untwisted sector, the orbifold model contains a Kähler modulus field which is charged under the unbroken Abelian gauge group. By assigning a VEV to the charged Kähler modulus field, the orbifold moves away from the enhanced point and the $U(1)^2$ gauge symmetry breaks to a discrete Z_3^2 subgroup. Thus, effectively the non-Abelian discrete symmetry $\Delta(54) \simeq (Z_3 \times Z_3) \rtimes S_3$ is realized. The other VEV directions of the untwisted scalar fields break the symmetry to $(U(1) \times Z_2) \times Z_6, Z_3 \times S_3$ or $Z_3 \times Z_3$. In the one-dimensional Z_2 orbifold case, we showed that the non-Abelian gauge symmetry $SU(2)$ is the origin of the discrete symmetry $D_4 \simeq Z_4 \rtimes Z_2$. The other VEV directions of the untwisted scalar fields break the symmetry to Z_4 .

The resulting non-Abelian discrete flavor symmetries are exactly those that have been obtained from heterotic string theory on symmetric orbifolds at a general point in moduli space [19]. In [19], the geometrical symmetries of orbifolds were used to derive these discrete flavor symmetries. However, in this paper, we have not used these geometrical symmetries on the surface, although obviously the gauge symmetries and geometrical symmetries are

tightly related with each other. At any rate, our results also indicate a procedure to derive non-Abelian discrete symmetries for models where there is no clear geometrical picture to begin with, such as in asymmetric orbifold models [23–26] or Gepner models [27].

We give a comment on anomalies. Anomalies of non-Abelian discrete symmetries are an important issue to consider (see e.g. [28]). We start with a non-Abelian (continuous) gauge symmetry and break it by orbifolding and by moduli VEVs to a non-Abelian discrete symmetry. The original non-Abelian (continuous) gauge symmetry is anomaly-free and if it were broken by the Higgs mechanism, the remaining symmetry would also be anomaly-free. That is because only pairs vector-like under the unbroken symmetry gain mass terms. But this does not hold true for orbifold breaking, as it is possible to project out chiral matter fields. Thus, in our approach the anomalies of the resulting non-Abelian discrete symmetries are a priori nontrivial. However, in our mechanism we obtain semi-direct product structures such as $U(1)^2 \times S_3$. Since the corresponding $U(1)^2$ is broken by the Higgs mechanism, the remnant Z_3^2 symmetry is expected to be anomaly-free if the original $U(1)^2$ is anomaly-free (the semi-direct product structure automatically ensures cancellation of $U(1)$ -gravity-gravity anomalies, but other anomalies have to be checked). Thus, the only discrete anomalies that remain to be considered are those involving S_3 .

We also comment on applications of our mechanism to phenomenological model building. In our construction the non-Abelian gauge group is broken by the orbifold action. This situation could be realized in the framework of field-theoretical higher-dimensional gauge theory with orbifold boundary conditions. Furthermore, our mechanism indicates that $U(1)^m \times S_n$ or $U(1)^m \times Z_n$ gauge theory can be regarded as a UV completion of non-Abelian discrete symmetries.³ Thus, it may be possible to embed other phenomenologically interesting non-Abelian discrete symmetries into such a gauge theory and investigate their phenomenological properties.

Acknowledgements

F.B. was supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT), Japan (No. 23104011). T.K. was supported in part by the Grant-in-Aid for Scientific Research No. 25400252 from the Ministry of Education, Culture, Sports, Science, and Technology of Japan. S.K. was supported by the National Science Council Taiwan under grant NSC102-2811-M-033-008.

References

- [1] G. Altarelli, F. Feruglio, *Rev. Mod. Phys.* **82** (2010) 2701, arXiv:1002.0211 [hep-ph].
- [2] H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada, M. Tanimoto, *Prog. Theor. Phys. Suppl.* **183** (2010) 1, arXiv:1003.3552 [hep-th]; H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada, M. Tanimoto, *Lect. Notes Phys.* **858** (2012) 1; H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada, M. Tanimoto, *Fortsch. Phys.* **61** (2013) 441.
- [3] S.F. King, C. Luhn, *Rep. Prog. Phys.* **76** (2013) 056201, arXiv:1301.1340 [hep-ph].
- [4] A. Adulpravitchai, A. Blum, M. Lindner, *J. High Energy Phys.* **0909** (2009) 018, arXiv:0907.2332 [hep-ph]; C. Luhn, *J. High Energy Phys.* **1103** (2011) 108, arXiv:1101.2417 [hep-ph]; A. Merle, R. Zwicky, *J. High Energy Phys.* **1202** (2012) 128, arXiv:1110.4891 [hep-ph].
- [5] H. Abe, K.-S. Choi, T. Kobayashi, H. Ohki, *Nucl. Phys. B* **820** (2009) 317, arXiv:0904.2631 [hep-ph]; H. Abe, K.-S. Choi, T. Kobayashi, H. Ohki, *Phys. Rev. D* **80** (2009) 126006, arXiv:0907.5274 [hep-th]; H. Abe, K.-S. Choi, T. Kobayashi, H. Ohki, *Phys. Rev. D* **81** (2010) 126003, arXiv:1001.1788 [hep-th].
- [6] M. Berasaluce-Gonzalez, P.G. Camara, F. Marchesano, D. Regalado, A.M. Uranga, *J. High Energy Phys.* **1209** (2012) 059, arXiv:1206.2383 [hep-th]; F. Marchesano, D. Regalado, L. Vazquez-Mercado, *J. High Energy Phys.* **1309** (2013) 028, arXiv:1306.1284 [hep-th].
- [7] Y. Hamada, T. Kobayashi, S. Uemura, *J. High Energy Phys.* **1405** (2014) 116, arXiv:1402.2052 [hep-th].
- [8] H. Abe, T. Kobayashi, H. Ohki, K. Sumita, Y. Tatsuta, arXiv:1404.0137 [hep-th].
- [9] L.J. Dixon, J.A. Harvey, C. Vafa, E. Witten, *Nucl. Phys. B* **261** (1985) 678; L.J. Dixon, J.A. Harvey, C. Vafa, E. Witten, *Nucl. Phys. B* **274** (1986) 285.
- [10] L.E. Ibañez, H.P. Nilles, F. Quevedo, *Phys. Lett. B* **187** (1987) 25; L.E. Ibañez, J.E. Kim, H.P. Nilles, F. Quevedo, *Phys. Lett. B* **191** (1987) 282.
- [11] Y. Katsuki, Y. Kawamura, T. Kobayashi, N. Ohtsubo, Y. Ono, K. Tanioka, *Nucl. Phys. B* **341** (1990) 611.
- [12] T. Kobayashi, S. Raby, R.J. Zhang, *Phys. Lett. B* **593** (2004) 262, arXiv:hep-ph/0403065; T. Kobayashi, S. Raby, R.J. Zhang, *Nucl. Phys. B* **704** (2005) 3, arXiv:hep-ph/0409098.
- [13] W. Buchmüller, K. Hamaguchi, O. Lebedev, M. Ratz, *Phys. Rev. Lett.* **96** (2006) 121602, arXiv:hep-ph/0511035; W. Buchmüller, K. Hamaguchi, O. Lebedev, M. Ratz, *Nucl. Phys. B* **785** (2007) 149, arXiv:hep-th/0606187.
- [14] J.E. Kim, B. Kyae, *Nucl. Phys. B* **770** (2007) 47, arXiv:hep-th/0608086.
- [15] O. Lebedev, H.P. Nilles, S. Raby, S. Ramos-Sanchez, M. Ratz, P.K.S. Vaudrevange, A. Wingerter, *Phys. Lett. B* **645** (2007) 88, arXiv:hep-th/0611095; O. Lebedev, H.P. Nilles, S. Raby, S. Ramos-Sanchez, M. Ratz, P.K.S. Vaudrevange, A. Wingerter, *Phys. Rev. D* **77** (2008) 046013, arXiv:0708.2691 [hep-th].
- [16] M. Blaszczyk, S. Nibbelink Groot, M. Ratz, F. Ruehle, M. Trapletti, P.K.S. Vaudrevange, *Phys. Lett. B* **683** (2010) 340, arXiv:0911.4905 [hep-th].
- [17] S. Groot Nibbelink, O. Loukas, *J. High Energy Phys.* **1312** (2013) 044, arXiv:1308.5145 [hep-th].
- [18] H.P. Nilles, S. Ramos-Sanchez, M. Ratz, P.K.S. Vaudrevange, *Eur. Phys. J. C* **59** (2009) 249, arXiv:0806.3905 [hep-th].
- [19] T. Kobayashi, H.P. Nilles, F. Ploger, S. Raby, M. Ratz, *Nucl. Phys. B* **768** (2007) 135, arXiv:hep-ph/0611020.
- [20] P. Ko, T. Kobayashi, J.-h. Park, S. Raby, *Phys. Rev. D* **76** (2007) 035005, arXiv:0704.2807 [hep-ph]; P. Ko, T. Kobayashi, J.-h. Park, S. Raby, *Phys. Rev. D* **76** (2007) 059901 (Erratum).
- [21] R. Dijkgraaf, E.P. Verlinde, H.L. Verlinde, *Commun. Math. Phys.* **115** (1988) 649; K.-i. Kobayashi, M. Sakamoto, *Z. Phys. C* **41** (1988) 55; E.J. Chun, J.E. Kim, *Phys. Lett. B* **238** (1990) 265; E.J. Chun, J. Lauer, H.P. Nilles, *Int. J. Mod. Phys. A* **7** (1992) 2175; T. Kobayashi, N. Ohtsubo, K. Tanioka, *Int. J. Mod. Phys. A* **8** (1993) 3553; T. Kobayashi, *Prog. Theor. Phys. Suppl.* **110** (1992) 277; H. Kawabe, T. Kobayashi, N. Ohtsubo, *Prog. Theor. Phys.* **88** (1992) 431.
- [22] J.A. Escobar, C. Luhn, *J. Math. Phys.* **50** (2009) 013524, arXiv:0809.0639 [hep-th].
- [23] K.S. Narain, M.H. Sarmadi, C. Vafa, *Nucl. Phys. B* **288** (1987) 551.
- [24] L.E. Ibañez, J. Mas, H.-P. Nilles, F. Quevedo, *Nucl. Phys. B* **301** (1988) 157.
- [25] M. Ito, S. Kuwakino, N. Maekawa, S. Moriyama, K. Takahashi, K. Takei, S. Teraguchi, T. Yamashita, *Phys. Rev. D* **83** (2011) 091703, arXiv:1012.1690 [hep-ph]; M. Ito, S. Kuwakino, N. Maekawa, S. Moriyama, K. Takahashi, K. Takei, S. Teraguchi, T. Yamashita, *J. High Energy Phys.* **1112** (2011) 100, arXiv:1104.0765 [hep-th].
- [26] F. Beye, T. Kobayashi, S. Kuwakino, *Nucl. Phys. B* **875** (2013) 599, arXiv:1304.5621 [hep-th]; F. Beye, T. Kobayashi, S. Kuwakino, *J. High Energy Phys.* **1401** (2014) 013, arXiv:1311.4687 [hep-th].
- [27] D. Gepner, *Nucl. Phys. B* **296** (1988) 757.
- [28] T. Araki, T. Kobayashi, J. Kubo, S. Ramos-Sanchez, M. Ratz, P.K.S. Vaudrevange, *Nucl. Phys. B* **805** (2008) 124, arXiv:0805.0207 [hep-th].
- [29] H. Abe, K.-S. Choi, T. Kobayashi, H. Ohki, M. Sakai, *Int. J. Mod. Phys. A* **26** (2011) 4067, arXiv:1009.5284 [hep-th].

³ See also [29].