



Title	Production of slow-released nitrogen fertilizer from urine
Author(s)	Ito, Ryusei; Takahashi, Eri; Funamizu, Naoyuki
Citation	Environmental technology, 34(20), 2809-2815 https://doi.org/10.1080/09593330.2013.790069
Issue Date	2013-10-01
Doc URL	http://hdl.handle.net/2115/57595
Rights	This is an Accepted Manuscript of an article published by Taylor & Francis in Environmental Technology on 2014, available online: http://www.tandfonline.com/10.1080/09593330.2013.790069
Type	article (author version)
File Information	production of slow release_rev3.pdf



[Instructions for use](#)

PRODUCTION OF SLOW-RELEASED NITROGEN FERTILIZER FROM URINE

Ryusei Ito

Water, Decontamination, Ecosystem and Health Laboratory, International Institute for Water and Environmental Engineering (2iE), 01, BP 594, Ouagadougou, 01, Burkina Faso,

Laboratory on Engineering for Sustainable Sanitation, Faculty of Engineering, Hokkaido University, Kita 13 Nishi 8, Kita-ku, Sapporo, Hokkaido, 060-8628, Japan,

e-mail: ryuusei@eng.hokudai.ac.jp

Eri Takahashi

Laboratory on Engineering for Sustainable Sanitation, Graduate School of Engineering, Hokkaido University, Kita 13 Nishi 8, Kita-ku, Sapporo, Hokkaido, 060-8628, Japan

Naoyuki Funamizu

Laboratory on Engineering for Sustainable Sanitation, Graduate School of Engineering, Hokkaido University, Kita 13 Nishi 8, Kita-ku, Sapporo, Hokkaido, 060-8628, Japan

PRODUCTION OF SLOW-RELEASED NITROGEN FERTILIZER FROM URINE


Human excreta, especially urine is rich in nitrogen to be able to utilize into the agricultural purposes, while the slow-release fertilizer allows effective utilization of nutrients in agricultural production. The direct formation of slow-release fertilizer —methylene urea— from urine was being proposed in this study. The experiments were tried to prove formation of methylene urea from human urine, and to investigate the effect of the pH and salt concentration on the reaction rate. The synthetic urine and real urine were used for urea source of the reaction. As a result, the precipitates were prepared from synthetic urine, while the small molecule fractions generated then they growth to precipitate. The NMR, IR, element analyses showed the precipitation in synthetic urine must be same compound from urea solution which is methylene urea. The reaction rate was high at low pH value. The reaction rate in buffer solution was lower than synthetic urine at same pH, because some salts may work as catalyst. The urea concentration reduction rate in real urine showed same trend with synthetic urine at same pH, while the precipitation was quite similar to methylene urea.

Keywords: methylene urea, nitrogen recovery, human urine, pH, salt concentration

Subject classification codes: include these here if the journal requires them

Introduction

Nitrogen fertilizers, such as ammonia, urine nitrate salts are the great elements for agriculture. They are able to increase the productivity of fruits or grains on the poor lands. Currently, 100M ton-N of nitrogen fertilizers are produced and consumed in the world par year [1] for agricultural purposes while their market is still developing to satisfy the growing food demands especially in developing countries. Here, the production of ammonia, which is main resource of other nitrogen fertilizers, requires 34.5GJ/ton-N of energy [2] by steam reforming method. This fact shows the energy



Both reactions are electrophilic addition reaction which catalyzed by proton of acids resulting in high reaction rate at low pH value. In usual industry, heating process is applied for accelerating the reaction rates and increasing the degree of the polymerization, however the reaction advances under room temperature. The solubility and biodegradation rate of the methylene urea are controlled by the degree of polymerization.

Focusing on human excreta, 850g of nitrogen is evacuated from a person per year [16], while human being on the earth excretes about 6M ton of nitrogen per year. The human urine contains 86 % of nitrogen in the excreta [17]. Most of the nitrogen exists as urea formation which easily decomposes into ammonium formation by biological activity and hydrolysis reaction resulting from contamination of faces [18]. So, recovery of nitrogen from urine has potential to reduce energy consumption for producing nitrogen fertilizer and the production of controlled release fertilizer has larger impact to energy problem by increasing nitrogen utilization efficiency.

We are proposing the production of controlled release fertilizer by a direct formation process of methylene urea from human urine. The real urine contains many inorganic salt and unknown organic matters, which are concerned to causes sub-reaction of undesired compounds. Considering actual reaction system, controlling pH is one of the easiest processes and recovery ratio should be evaluated for find effective reaction conditions. The aims of this paper are to prove formation of methylene urea from synthetic urine and real human urine, and to investigate the effect of the pH and salt concentration on the reaction rate and recovery of urea form nitrogen. The structure of the production was analyzed by C13-NMR, IR spectrum and element analysis.

Methods

Reaction in synthetic solution

The composition of the synthetic urine is summarized in Table 1 [19]. Agents of JIS special grade provided by Wako Pure Chemical Industries, Ltd. were utilized for preparation of synthetic urine. The pH values of the solutions were controlled by addition of 1 mol/L sulfuric acid (JIS special grade, Wako Pure Chemical Industries, Ltd.) with monitoring pH meter before starting the reactions. 600 ml of synthetic urine was taken in beakers, then was kept at 20 °C in a water bath. 30 ml of 30% formaldehyde solution (JIS special grade, Wako Pure Chemical Industries, Ltd.) was added to the beakers to start the reactions, while the solutions were mixed by magnetic stirrer during the reaction. After the reaction duration, 1 mol/L sodium hydroxide solution (JIS special grade, Wako Pure Chemical Industries, Ltd.) was added to put its pH value at 7 for stopping the reaction. The liquid and precipitates were filtrated by 0.45µm membrane filter then weight of the residue on the filter was measured by

electric balance. The filtrates were dialyzed with UF membrane tubes which had cut off of 100kDa, 10kDa and 1kDa (Spectra/Por Biotech Cellulose Ester (CE) Dialysis Membranes). The 10ml of liquid sample was put into the tubes then was sealed. The tubes were put into the ion exchanged water in the refrigerator for 2 days and the water was changed every 6–12 hours. The nitrogen concentrations in the tubes were measured by Kjeldahl method. The precipitates on the filter were well washed with ion exchanged water. The molecule structure of the precipitation was analyzed by C13-NMR (ECX-400, JEOL), solid IR spectrum and elemental analyzer (CHN corderMT-6, YANACO). For the reference, the 50g/L of urea solution was prepared and add 1 mol/L sulfuric acid to set the pH at 2. 200 ml of urea solution and 10 ml of 30% formaldehyde solution were mixed by magnetic stirrer at 20 °C for 240 hours. The precipitates in the solution were recovered by filtration with 0.45 µm membrane filter, well washed by ion exchanged water, and then their molecular structure was analyzed by same methods of the precipitates from the synthetic urine solutions.

Effect of pH and salt concentration on the reaction rate

The synthetic urine solutions with pH 1, 2, 3, 4 and 5 and without pH control were prepared by following the protocol above. 200 ml of synthetic urine solutions were taken in beakers kept at 20 °C in a water bath. 10 ml of 30% formaldehyde solution was added to the beakers and mixed by magnetic stirrers during the reaction. Some part of liquid was sampled periodically. The sample solutions were filtrated with 0.45µm membrane filter. The urea concentrations in the filtrates were measured by the diacetyl monoxime test [20]. The molecular distribution of the synthetic solution with pH 5 follows same procedure mentioned above.

Application to real urine

The real urine was taken from 4 volunteers then stored at 2 °C refrigerator, and was applied to experiment in 1 day. The reaction follows the protocol of the experiment for different pH values. 200 ml of urine was taken in a beaker, while the pH is adjusted at 2 with addition of the 1 mol/L sulfuric acid. The urea concentration in the liquid phase and molecular structure of the precipitates were analyzed.

Results and discussion

Production of methylene urea from synthetic urine

The methylene urea formation from urea solution and formaldehyde is well known to be used for industrial production. Urine is rich in urea with high concentration of salts and organic matters, which probably affect the reaction.

Figure 1 shows the time course of the molecular distribution of derivatives of urea and other compounds that contains nitrogen element at pH 2. 1600 mmol of urea and 100 mmol of creatinine and ammonium chloride were found in the synthetic urine at initial. During the experiments in this paper, the concentrations of creatinine and ammonium chloride were assumed not to change, because the concentration was enough lower than urea concentration. The urea concentration decreased with elapse of the time to form the components of larger molecular size and precipitates, but total amount of nitrogen element did not change. The components with the range of 10 k - 100 kDa were the largest at 1 hour then decreased. These phenomena show the polymerization process of urea derivatives. Therefore, it is gathered that the reaction follows the process of methylene urea production. Then, the precipitates from synthetic urine was analyzed by the means of C13-NMR, IR spectrometry and element analysis. Figure 2 shows the NMR chart of the precipitates. There are two peaks which represent the methylene group (-CH₂- 50 ppm) and the ketone group (-C=O- 160 ppm). This chart is similar to

the precipitates from urea solution as illustrated in Figure 3. As mentioned above, the reaction of formaldehyde in urea solution at acidic condition prepares precipitates of methylene urea. So, this compound from synthetic urine should be near to methylene urea. The IR spectrum of the precipitates of synthetic urine is shown in Figure 4. Three groups of peaks, such as N-H bound (3500 cm^{-1}), C=O bound (1800 cm^{-1}) and C-N bound ($>1500\text{ cm}^{-1}$) are found in it. These peak groups were also found in the IR spectrum of methylene urea from urea solution as illustrated in Figure 5. From these data, we can assume that the production is the derivatives of methylene urea described by equation (1). The element analysis of the precipitates from synthetic urine and urea solution respectively showed the weight ratio of each element as C: 28.52%, H: 5.57%, N: 35.33% and C: 29.78%, H: 5.59%, N: 36.82%. The rest should be Oxygen or other elements. These values are almost equal, so that it is seems that the same substances are produced from synthetic urine and urea solution.

Effect of pH and salt concentration on the reaction rate of methylene urea

Figure 6 shows the time course of the urea concentration in the liquid phase with different pH values. The urea concentration decreased with elapse of time at all pH values, while the rate was high at low pH. This is because, the electrophilic addition reaction which is catalyzed by proton occurs on the reaction of urea and formaldehyde resulting in high reaction rate at high proton concentration. The production and reaction rates of the component with the range of 10 k – 100 kDa at pH 5 are lower than them at pH2 as illustrated in Figure 7. This also shows the effect of the proton concentration. The urea concentration of pH 5 condition at the steady state was higher than pH 2. Some sub-reaction to stop the main reaction might occur, but about this point, more analysis is required.

To investigate the effect of salt concentration, the time courses of the urea concentration in the urea buffer solution at pH 2 and pH 5 are presented in Figure 8. In urea solution, the reaction rate was fast at low pH value, and lower than in the synthetic urine at same pH values. The buffer solution has low salt concentration, so that the salts may have catalytic activity to accelerate the reaction.

Application to real urine

Figure 9 represents the reaction rate in the real urine which pH was adjusted at 2. There was slight difference between the reaction rate and urea concentration in liquid phase in the real and in synthetic urine, while the precipitates also obtained. The precipitation from real urine was analyzed by the means of C13-NMR, IR spectrum and element analysis to check its quality as illustrated in Figures 10 and 11. The NMR chart has 2 big peaks and 3 small peaks around 60 ppm. The big peaks seems to be same function groups (-CH₂- 50 ppm and -CO- 160 ppm) in Figure 2 (precipitate from synthetic urine) and 3 (precipitates from urea). The small peaks may suggest existence of unknown functional groups or unknown derivative of -CH₂- group. In the IR spectrum of precipitates from real urine, three bound groups as N-H bound, C=O bound and C-N bound were also found and this IR analysis did not show the difference between the real urine case and the synthetic urine, urea cases (Figures 4 and 5). The element analysis showed the result of the contents, C: 29.61%, H: 5.85% and N: 30.35% and this means that the nitrogen content was lower than other precipitates. Even the lower nitrogen content of precipitate from real urine, the nitrogen content of the precipitates is sufficient level for nitrogen fertilizer [21]. From the above results, it might be concluded that the main constituents of the precipitation from real urine was methylene urea and the precipitates may contain trace unknown function groups which is deficiency of

nitrogen. It requires to more analysis for other elements or functional groups in the precipitation.

Conclusions

The slow-release fertilizer of methylene urea was tried to prepare from urine solutions. As a result, the precipitates were prepared from synthetic urine, while the small molecule fractions generated then they growth to precipitate. The NMR, IR analyses showed the precipitation in synthetic urine must be same compound from urea solution which is methylene urea. Element analysis showed the weight ratios of the compounds from synthetic and buffer solution, which were respectively C: 28.52%, H: 5.57%, N: 35.33% and C: 29.78%, H: 5.59%, N: 36.82%. The reaction rate was high at low pH value. This is because the reaction was electrophilic addition reaction as described by Eq. (1) which catalyzed by proton of acids. The reaction rate in buffer solution was lower than synthetic urine at same pH, because some salts may work as catalyst. The urea concentration reduction rate in real urine was same with synthetic urine at same pH. The NMR, IR analysis showed the precipitation should be quite similar compound to methylene urea. The weight composition of the precipitates was C: 29.61%, H: 5.85% and N: 30.35% by the element analysis.

References

- [1] FAO, *FAOSTAT*, (2009). Available at <http://faostat.fao.org/>.
- [2] G. Kongshaug, *Energy Consumption and Greenhouse Gas Emissions in Fertilizer Production*, in *IFA Technical Conference*, (1998).
- [3] M. E. Trenkel, *Use Efficiency Controlled-Release and Stabilized Fertilizers in Agriculture*. (1997).

- [4] Toray Research Center, *Jyoho Gijutu to Yoto Tenkai*. (1997), pp. 1 – 8.
- [5] R. W. Baker and H. K. Lonsdale, *Controlled Release of Biologically Active Agents*. Plenum, New York, (1973), pp. 15 – 71.
- [6] C. Berkland, M. King, A. Cox, K. Kim, and D. W. Pack, *Precise control of PLG microsphere size provides enhanced control of drug release rate*, *Journal of Controlled Release* 82 (2002), pp. 137 – 147.
- [7] R. Ito, B. Golman, and K. Shinohara, *Controlled Release of Core Particle Coated with Soluble Particles in Impermeable Layer*, *Journal of Chemical Engineering of Japan* 35 (2002), pp. 40–45.
- [8] R. Ito, B. Golman, and K. Shinohara, *Controlled release with coating layer of permeable particles*, *Journal of Controlled Release* 92 (2003), pp. 361–368.
- [9] R. Ito, B. Golman, and K. Shinohara, *Effect of the Porous Structure of a Coating Layer on the Release Rate of Core Particle*, *Journal of Chemical Engineering of Japan* 36 (2003), pp. 676–682.
- [10] R. Ito, B. Golman, and K. Shinohara, *Formation of a sigmoidal release pattern of core particles coated with layers of soluble and permeable particles*, *Advanced Powder Technology* 15 (2004), pp. 377–390.
- [11] R. Ito, B. Golman, and K. Shinohara, *Multiple-Layer Coating of Solids with Impermeable Film Dispersing Permeable Particles for Controlled Release*, *Journal of Chemical Engineering of Japan* 37 (2004), pp. 496–502.
- [12] R. Ito and K. Shinohara, *Design of multi-layer coated particles with sigmoidal release pattern*, *Chemical Engineering Science* 60 (2005), pp. 5415–5424.
- [13] U. Shavit and A. S. M. Reiss, *Wetting mechanisms of gelbased controlled-release fertilizers*, *Journal of Controlled Release* 88 (2003), pp. 71 – 83.
- [14] N. Shimono, T. Takatori, M. Ueda, M. Mori, Y. Higashi, and Y. Nakamura, *Chitosan dispersed system for colon-specific drug delivery*, *International Journal of Pharmaceutics* 245 (2002), pp. 45 – 54.
- [15] S. Shoji, *Meister: Controlled Release Fertilizer: Properties and Utilization*. Konno Printing, Sendai, (1999), pp. 16 – 20.
- [16] M. Almeida, D. Butler, and E. Friedler, *At-source domestic wastewater quality*, *Urban Water* 1 (1999), pp. 49–55.
- [17] A. F. Malisie, M. Prihandrijanti, and R. Otterpohl, *The potential of nutrient reuse from a source-separated domestic wastewater system in Indonesia--case study: ecological sanitation pilot plant in Surabaya.*, *Water science and technology : a journal of the International Association on Water Pollution Research* 56 (2007), pp. 141–148.

- [18] S. Hotta and N. Funamizu, *Evolution of ammonification potential in storage process of urine with fecal contamination.*, *Bioresource technology* 99 (2008), pp. 13–17.
- [19] J. A. Wilsenach, C. A. H. Schuurbiers, and M. C. M. van Loosdrecht, *Phosphate and potassium recovery from source separated urine through struvite precipitation.*, *Water research* 41 (2007), pp. 458–66.
- [20] WHO, *Standard Operating Procedures for CLINICAL CHEMISTRY – Diacetyl Monoxime Method.* Available at http://www.searo.who.int/en/Section10/Section17/Section53/Section481_1754.htm.
- [21] Ministry of Agriculture Forestry and Fisheries, *Fertilizers Regulation Act.* (1950).




Figure 1 Molecular distribution of urea derivatives at pH 2




Figure 2 NMR spectrum of precipitates from synthetic urine




Figure 3 NMR spectrum of precipitates from urea solution




Figure 4 IR spectrum of precipitates from synthetic urine




Figure 5 IR spectrum of precipitates from urea solution




Figure 6 Urea concentration of normal synthetic urine




Figure 7 Molecular distribution of urea derivatives at pH 5




Figure 8 time courses of the urea concentration in the urea buffer solution




Figure 9 Urea concentration of real urine




Figure 10 NMR spectrum of precipitates from synthetic urine




Figure 11 IR spectrum of precipitates from real urine

Table 1 components of synthetic urine

Chemicals	Concentration (mmol/L)
$\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$	3.2
NaCl	78.7
Na_2SO_4	16.2
$\text{Na}_3(\text{C}_6\text{H}_5\text{O}_7) \cdot 2\text{H}_2\text{O}$	2.6
KCl	21.5
$\text{C}_4\text{H}_7\text{N}_3\text{O}$	9.7
$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	4.4
KH_2PO_4	30.9
NH_4Cl	18.7
$(\text{NH}_2)_2\text{CO}$	417
$\text{Na}_2(\text{COO})_2$	0.15
$\text{C}_6\text{H}_8\text{O}_6$	0.57