Title	Vertical distribution of major sulfate-reducing bacteria in a shallow eutrophic meromictic lake
Author(s)	Kubo, Kyoko; Kojima, Hisaya; Fukui, Manabu
Citation	Systematic and Applied Microbiology, 37(7), 510-519 https://doi.org/10.1016/j.syapm.2014.05.008
Issue Date	2014-10
Doc URL	http://hdl.handle.net/2115/57661
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	manuscript.pdf

[Instructions for use](#)

**Vertical distribution of major sulfate-reducing bacteria in a eutrophic
shallow meromictic lake**

Kyoko Kubo^{*}, Hisaya Kojima^{*} and Manabu Fukui

The Institute of Low Temperature Science, Hokkaido University, Kita-19, Nishi-8, Kita-ku,
Sapporo 060-0819, Japan

^{*} Corresponding authors at: The Institute of Low Temperature Science, Hokkaido University,
Kita-19, Nishi-8, Kita-ku, Sapporo 060-0819, Japan.

E-mail addresses: kkubo@pop.lowtem.hokudai.ac.jp (K. Kubo),
kojimah@pop.lowtem.hokudai.ac.jp (H. Kojima)

Running title: SRB in a shallow meromictic lake

Abstract

The vertical distribution of sulfate-reducing bacteria was investigated in a shallow, eutrophic, meromictic lake, Lake Harutori, which is located in a residential area of Kushiro, Japan. A steep chemocline, which is characterized by gradients of oxygen, sulfide, and salinity, was found at a depth of 3.5–4.0 m. The sulfide concentration at the bottom of the lake was high (up to a concentration of 10.7 mM). Clone libraries were constructed using the *aprA* gene, which encodes adenosine-5'-phosphosulfate reductase subunit A to monitor sulfate-reducing bacteria. In the *aprA* clone libraries, the most abundant sequences were those from the *Desulfosarcina-Desulfococcus* (DSS) group. A primer set for a DSS group-specific 16S rRNA gene was used to construct another clone library, analysis of which revealed that the uncultured group of sulfate-reducing bacteria, SEEP SRB-1, accounted for nearly half of the obtained sequences. Quantification of the major bacterial groups by catalyzed reporter deposition-fluorescence *in situ* hybridization demonstrated that the DSS group accounted for 3.2–4.8% of the total bacterial community below the chemocline. Our results suggest that below the chemocline of Lake Harutori, DSS group is one of the major groups of sulfate-reducing bacteria and that these presumably metabolically versatile bacteria might play an important role in sulfur cycling in the lake.

Keywords: *Meromictic lake, sulfate-reducing bacteria, aprA, sulfide*

Introduction

The global expansion of anoxic aquatic environments has a massive influence on biological activities. For instance, in the California Current marine ecosystem, serious damage to the fish and benthic invertebrate communities is reported to have occurred due to anoxic water, and there is concern about possible further damage to the highly productive coastal environment [7]. The formation of anoxic water masses is thought to be connected to human-induced eutrophication and global warming [1], and it is speculated that the occurrence of hypoxia will increase in the coming decades [21]. Sulfide accumulation in anoxic water columns is an additional concern. In anoxic environments, sulfides are normally produced via microbial sulfate reduction. Especially in the marine environment, the sufficient supply of sulfate from seawater and of organic compounds support the growth of sulfate-reducing bacteria. Most of the sulfide is usually converted into colloidal sulfur or sulfate by bacteria, via either anaerobic or aerobic oxidation before it reaches the surface layer of the water column [21]. Therefore, the analysis of microbial sulfur cycling in oxic-anoxic interfaces is attracting increasing attention.

Meromictic lakes can provide a suitable environment to investigate the potential importance of sulfate-reducing bacteria (SRB) in anoxic water columns. In meromictic lakes, the surface layer of the lake, called the mixolimnion circulates but does not intermix with the deeper layer, called monimolimnion. Therefore, such lakes stay stratified throughout the year. At the interface between the mixolimnion and the monimolimnion, a steep chemical gradient called a chemocline is formed. A clear shift of the microbial community structures below and above the chemocline has been shown in many meromictic environments [17, 38]. Lake Harutori is located in a residential area of Kushiro, Hokkaido, Japan. This meromictic lake is consisting of water layers based on salinity. The bottom layer of Lake Harutori is known to

contain a very high amount of sulfide. The lake was reported to contain up to 670 mg L⁻¹ sulfide at one point [44], so that the toxic effects on indigenous animals and plants were a matter of concern. In addition, most of the wastewater from the adjoining residential area was directly discharged into the lake; therefore, the accumulation of organic compounds was an environmental problem. An analysis of photosynthetic pigments revealed anaerobic anoxygenic phototrophic bacteria (green sulfur bacteria) in the Lake Harutori chemocline [35]. Aside from that, not much is known about the microbial community in the lake.

In this study, we aimed to identify and quantify the main sulfate-reducing bacteria beneath the Lake Harutori chemocline. To find the key players in sulfur cycling in Lake Harutori, we investigated the phylogenetic diversity of sulfate-reducing and sulfur-oxidizing bacteria by sequence analysis of the gene encoding adenosine 5'-phosphosulfate reductase (*aprA*). Furthermore, the vertical distribution of the predominant group in *Deltaproteobacteria* was revealed by using catalyzed reporter deposition-fluorescence *in situ* hybridization (CARD-FISH). The study shows that *Desulfosarcina-Desulfococcus* (DSS) group is one of the major groups of sulfate-reducing bacteria in the monimolimnion of Lake Harutori.

Materials and methods

Study sites and sampling

Lake Harutori is a shallow meromictic lake located in Kushiro, Hokkaido, Japan (Fig. 1). The lake (surface area 0.36 km²) is a coastal lagoon in which the low salinity water is overlaying water of marine origin, resulting in a permanent stratification of the water body [2]. The lake is completely covered with ice in winter, usually from December to the end of March [2]. The center of the lake (N42°58'20.6", E144°24'6.6"), which has a maximum depth of 5.75 m, was

selected as the sampling point. Sampling took place in May 2012 after the ice had melted. Water samples were collected using a horizontal Van Dorn water sampler (3L, Rigosha, Tokyo, Japan) along the depths. Vertical profiles of temperature, specific conductivity and pH were measured *in situ*, with a multiparameter sensor (YSI 600XLM; YSI Inc., Yellow Springs, OH, USA). Photosynthetically active radiation (400–700 nm) was measured using a quantum meter (Model QMSS-S; Apogee instruments Inc., Logan, UT, USA). The dissolved oxygen profile was measured *in situ* with an optical dissolved oxygen meter (ProODO; YSI Inc., Yellow Springs, OH, USA).

Chemical measurements

Sulfide concentrations were measured colorimetrically with the methylene blue formation method [3]. To measure dissolved organic carbon (DOC), sulfate, and chloride, water samples were filtered through cellulose acetate syringe filter units with a 0.2 µm pore size (DISMIC-13CP; Toyo Roshi Kaisha, Tokyo). Concentrations of DOC were measured with a total organic carbon analyzer (TOC-V; Shimadzu, Kyoto, Japan). Sulfate and chloride concentrations were determined using an ion chromatograph (ICS-1500, column: IonPacAS12A, Dionex, Sunnyvale, CA, USA) with appropriate dilutions.

DNA extraction

The water samples (approximately 200 mL for each sample) were filtered through Sterivex-GV 0.22 µm pore-sized filter cartridges (Millipore, Billerica, MA, USA) immediately after the collection. The filter cartridges were frozen on dry ice and brought back into the laboratory without thawing, and stored at -20°C until DNA extraction. The filters were transferred into a 2 mL screw-capped tube with 0.5 g of sterile glass beads, 0.6 mL TE buffer

(10 mM Tris-HCl, 1 mM EDTA, pH 8), 30 μ L 20% SDS and 0.6 mL phenol-chloroform-isoamyl alcohol (25:24:1; v/v/v). The tube was vigorously shaken with a beads-beater (FastPrep24; MP Biomedicals, Santa Ana, CA, USA) twice at 4.0 m s⁻¹ for 30 sec. DNA in the water phase was purified using the CTAB method followed by isopropanol precipitation [43]. Extracted DNA was stored at -20°C until use.

aprA clone libraries and sequence analysis

For amplification of *aprA* genes, the primers AprA-1-FW and AprA-5-RV [25] were used. PCR conditions were as follows: an initial denaturation step at 94°C for 2 min, followed by 34 cycles of denaturation at 94°C for 30 sec, annealing at 55°C for 30 sec, and elongation at 72°C for 45 sec, with a final extension step at 72°C for 10 min. PCR products were inserted into the pCR2.1-TOPO vector (TOPO TA cloning kit; Invitrogen, Carlsbad, CA, USA) and cloned into competent *Escherichia coli* TOP10 cells according to the manufacturer's instructions. Clones that had inserts of the predicted size were screened by the direct PCR amplification with vector primers (M13F and M13R), and the PCR products were purified with isopropanol precipitation and used for sequencing reaction. The sequencing reactions were done with the ABI BigDye chemistry and analyzed with an ABI 3130 Genetic Analyzer (Applied Biosystems, Tokyo, Japan). The *aprA* sequences were translated and the deduced amino acid sequences were aligned using ClustalW implemented in MEGA5.05 software [36]. A pairwise distance matrix was calculated based on the Poisson model. Based on the distance matrix, sequences were classified into operational taxonomic units (OTUs) using mothur software [32], with 97% sequence identity as a threshold. The AprA tree was constructed from deduced amino acid sequences using the neighbor-joining method with bootstrap analysis with 1,000 replications.

DSS-specific 16S rRNA gene clone library

The primers DCC305 and DCC1165 [4] were used to amplify DSS-specific 16S rRNA genes. The PCR conditions were as follows: an initial denaturation step at 95°C for 5 min, followed by 30 cycles of denaturation at 94°C for 1 min, annealing at 65°C for 1 min, and elongation at 72°C for 1 min, with a final extension step at 72°C for 7 min. The PCR products were cloned and sequenced as described above. The sequences were aligned using ClustalW implemented in MEGA5.05 software and classified using mothur software into OTUs based on 98% sequence identity. Final phylogenetic analyses of the 16S rRNA genes were performed with the ARB software package [23] using the ARB reference database SILVA SSU release 108. All sequences were automatically aligned with the SINA web aligner (<http://www.arb-silva.de>) [28] and the alignments were subsequently optimized manually. The phylogenetic tree was constructed by neighbor-joining and maximum-likelihood analysis with different sets of filters. For construction of the reference tree, only nearly full-length sequences (>1300 bp) were considered. Partial sequences obtained in this study were inserted into the tree by parsimony criteria without allowing changes in the overall tree topology.

Nucleotide sequence accession numbers

Sequence data reported in this study are available in the DDBJ, GenBank and EMBL databases under the following accession numbers: 16S rRNA genes (AB894629-AB894655), and *aprA* genes (AB894656-AB894821).

Probe design

An oligonucleotide probe SEEP1d-468 was designed using the probe design tool in the ARB software package [23]. The probe specifically targets the 16S rRNA sequences of SEEP SRB-1d group obtained in Lake Harutori and several environmental clones (Fig. 4). The probe was tested for coverage (target group hits) and specificity (outgroup hits) in silico with the ARB probe match tool [23]. Probe specificity was checked on 618,442 prokaryotic sequences of the SILVA SSU Ref dataset Release 108 [29]. Specific CARD-FISH conditions were determined by hybridizing the probe to the lake water sample at formamide concentrations of 40%, 50%, 60%, and 70%, as explained in the following section. A deltaproteobacterial strain PL12 [8] was used as a control having a single mismatch.

Catalyzed reporter deposition fluorescence in situ hybridization (CARD-FISH)

Water samples were fixed with paraformaldehyde solution (final concentration 0.9% [v/v]) at 4°C. Afterwards, 200 µL aliquots were filtered onto polycarbonate membrane filters (Cyclopore Track Etched Membrane; pore size, 0.22 µm; 25-mm diameter; Whatman, NJ, USA) within 40 hours and stored at -20°C until further processing. Inactivation of endogenous peroxidases was done by incubating the filters in 0.01 M HCl with 0.3% H₂O₂ for 10 min at room temperature. Cells were permeabilized by incubating the filters in 15 µg mL⁻¹ proteinase K (dissolved in 0.1 M Tris-HCl, 0.05 M EDTA [pH 8.0]) for 3 min at 37°C [37]. CARD-FISH and subsequent staining with 4',6-diamidino-2-phenylindole (DAPI) followed a previously published protocol [27] with the following modification: hybridization was performed for 2 h, and tyramide signal amplification was performed for 15 min, both at 46°C. The washing temperature was 48°C. For the probes SEEP1d-1420 and SEEP1d-468, hybridization and washing temperatures were 35°C, and tyramide signal amplification was performed for 30 min at room temperature. Oligonucleotide probes labeled with horseradish

peroxidase were purchased from Biomers (Ulm, Germany). Probe sequences, probe concentrations and formamide concentrations required for specific hybridization are given in Table S1. Hybridized samples were examined with an epifluorescence microscope (Axioplan2; Zeiss, Germany). The given CARD-FISH counts are the means calculated from 40–70 randomly chosen microscopic fields, corresponding to 2,000 DAPI-stained cells.

Results

Physico-chemical properties of Lake Harutori

In Lake Harutori, a steep chemocline was found between 3.5 and 4.0 m below the surface. Physicochemical parameters are summarized in Figure 2. The thermocline almost coincided with the chemocline. The temperature of the lake bottom was 9°C. Photosynthetically active radiation decreased with depth in the mixolimnion and, in the chemocline, declined to 0% of the surface intensity. Oxygen and sulfide formed opposite concentration gradients. The sulfide concentration increased below the chemocline, reaching a concentration of 10.7 mM at the bottom of the lake. The chloride concentration in the middle of the monimolimnion was about half of the concentration of that in seawater, meaning that the water was brackish. The sulfate concentration was much lower than that of seawater and the maximal concentration was 1.4–2.1 mM in the chemocline. The concentration of DOC increased below the chemocline.

AprA diversity in around the chemocline

Clone libraries for *aprA* were generated from water taken from three different depths (3.0, 3.5, and 4.0 m) and contained 62, 42, and 62 clones, respectively. Based on a 97% identity cut-off of the deduced amino acid sequences, 51 distinct OTUs were identified (Table 1). Representative sequences of each OTU were selected to be shown in the phylogenetic tree in

Figure 3. The phylogenetic analysis of the deduced AprA amino acid sequences revealed a clear difference in the dominant sequences in each library. In the 3.0 m library, from water taken from above the chemocline, OTU A2 was the most dominant OTU (Table 1). The closest cultured species to OTU A2 was *Desulfatitalea tepidiphila* (94–97% sequence identity) [9]. In the upper part of the chemocline at 3.5 m, OTU A1, A3, and A7 were the dominant OTUs (Table 1). Sequences from OTU A3 were closely related to *Desulfocapsa thiozymogenes* (98–100%), which is known to disproportionate inorganic sulfur compounds [10]. Sequences of OTU A7 were related to *Desulfobacterium vacuolatum* (96–97%), which can oxidize a wide variety of organic acids completely [20]. OTU A1 was also the most abundant OTU in the lower part of the chemocline (4.0 m). The closest, but still distantly related, cultured organism was *Desulfofaba gelida* (88–92%), which is a psychrophilic incomplete oxidizer [16]. All the above-mentioned species and OTUs belong to the family *Desulfobacteraceae*. A few sequences related to spore-forming sulfate reducers in the genus *Desulfotomaculum* were found, however, they were all distantly related to cultured species (Table 1). Sequences related to those of known sulfur oxidizers were found at a depth of 3.0–3.5 m, which is above and in the upper part of the chemocline. Of these sulfur-oxidizing bacteria, OTU A11 was related to “*Candidatus Pelagibacter ubique*” (85% identity; EAS84493; Fig. S3), which is one of the most abundant and common species in the pelagic zone of the ocean [30]. Some OTUs were related to *Sulfuricella denitrificans* [18] (Fig. S3), and OTU A5 was related to *Thiobacillus denitrificans* [12] (Fig. S3). OTUs A13 and A20 formed a distinct clade with other environmental clones; however, their function is uncertain (Fig. 3a).

Abundance of SEEP SRB-1d in DSS-specific 16S rRNA gene clone library

The most abundant sequences in the *aprA* clone libraries were those related to the DSS group (Fig. 3b; OTU A1). Therefore, a 16S rRNA gene clone library using a DSS-specific primer set was constructed from a water sample taken at 4.5 m. This clone library contained a total of 27 sequences of ca. 820 bp in size. The sequences are grouped into 10 different OTUs based on 98% identity (Fig. 4). All but one of the sequences (OTU7) are in the DSS group. Almost half of the sequences (13 out of 27 sequences) were grouped into OTU1, which is distantly related to *Desulfofaba fastidiosa* (94.2–94.7% identity). An environmental clone obtained in the Guaymas Basin has high sequence identity (98.5–99.6%) with the sequences in OTU1. OTU1 sequences were grouped into the SEEP SRB-1d cluster in the phylogenetic tree shown in Figure 4, and had no mismatch with the probe DSS658 (Fig. S1) used for the CARD-FISH experiments.

Design and evaluation of new probe SEEP1d-468

An oligonucleotide probe SEEP1d-468 was designed to specifically detect OTU1 in SEEP SRB-1d group, which comprised nearly half of the sequences obtained from the above mentioned DSS group-specific 16S rRNA gene clone library (Fig. 4). The new probe SEEP1d-468 has coverage of nearly half of the SEEP SRB-1d group and two outgroup hits in OTU4 (Fig. 4). When the probe tested with 20% formamide concentration with hybridization temperature at 46°C, only spotty signals were observed. Bright signals were observed at 40% formamide concentration with the hybridization and washing temperature at 35°C. No signal was observed from the reference strain PL12, which has a single mismatch to probe SEEP1d-468, under the same hybridization condition. A previously published probe SEEP1d-1420 [33] showed bright signals in the lake water sample with the same hybridization condition as the probe SEEP1d-468. Since the sequence information about the probe-binding site of

SEEP1d-1420 was not available from the DSS specific clone library, the newly designed probe SEEP1d-468 was mainly used for the further analyses.

Abundance of DSS bacteria in the monimolimnion

Total cell numbers (DAPI) ranged from 1.3×10^7 to 1.5×10^7 cells mL⁻¹ in the mixolimnion, and decreased somewhat in a depth of 3.0 to 3.5 m (Fig. 5). Total cell numbers increased below 3.5 m, reaching up to 3.5×10^7 cells mL⁻¹ in the bottom part of the water column (4.8 m depth). Total bacterial cell numbers (probe EUB I-III) also increased below the chemocline, reaching up to 1.9×10^7 cells mL⁻¹ at 4.8 m. Deltaproteobacteria comprised 7.2–22.1% of total bacteria below 3.0 m (Fig. 5). Among deltaproteobacteria, cells from the DSS group were predominant (probe DSS658). At 4.5 m, in the middle of the monimolimnion, the ratio of DSS was 67% of the deltaproteobacteria (Fig. 6). The cell numbers of SEEP SRB-1d group (probe SEEP1d-468) ranged from 2.3×10^5 – 6.3×10^5 cells mL⁻¹ below the chemocline, and comprised 43.5–79.0% of DSS group (Fig. 6). Another probe SEEP1d-1420 also showed the similar cell numbers, which ranged from 2.6×10^5 – 5.7×10^5 cells mL⁻¹ below the chemocline. The typical morphology of the hybridized cells with probes DSS658 and SEEP1d-468 were coccoid (2–3 µm), as shown in Figure S1. Cells were dispersed or formed small aggregates.

Discussion

Unique characteristics of Lake Harutori

The monimolimnion of Lake Harutori contained a very high concentration of sulfide—up to 10.7 mM. Below the chemocline, the seawater seems to be diluted because the chloride concentration is nearly half of the seawater. If the water were diluted homogeneously, the

sulfate concentration should remain at half of that in seawater, around 14 mM. However, only 0.4–2 mM of sulfate was detected *in situ* (Fig. 2). The sum of the sulfate and sulfide concentrations comes close to 13 mM, which possibly means that most of the sulfate in the monimolimnion had been converted into sulfide. Active sulfate reduction by bacteria would explain the presence of such a high sulfide concentration in the monimolimnion.

Diversity of sulfate-reducing and sulfur-oxidizing bacteria in Lake Harutori

Analysis of the *aprA* clone libraries revealed a high diversity of sulfate-reducing and sulfur-oxidizing bacteria. The dominant groups of those bacteria differ around the three different depths of the chemocline. It appears that the availability of electron acceptors and salinity affected the composition of the bacterial community in the lake. Sulfur-oxidizing bacteria were found only where oxygen was available. Among those, some sequences obtained from 3.0 m were related to *Sulfuricella denitrificans* [18] and *Thiobacillus denitrificans* [12], which have been isolated from freshwater. Both of them are facultatively anaerobic chemolithoautotrophs. It has been shown that not all of sulfur oxidizers have *aprA* gene [24]; therefore we cannot exclude other sulfur oxidizers, such as *Epsilonproteobacteria*, are also present.

In the deeper layer of the chemocline, the sulfate-reducing bacterial community is dominated by a single OTU, OTU A1 (Table 1). OTU A1 is in the family *Desulfobacteraceae* and is distantly related to species in the DSS group.

Abundance of the DSS group in sulfate-reducing bacteria

In the high diversity of sulfate-reducing and sulfur-oxidizing bacteria observed in the *aprA* clone library, the most dominant sulfate-reducing bacterial OTU was distantly related to

species in the DSS group (OTU A1, Table 1). Nearly half of the sequences obtained from the DSS-specific 16S rRNA gene clone library were related to an uncultured group of *Deltaproteobacteria* (OTU1, Fig. 4), previously called as SEEP SRB-1d [33]. Both OTUs 1 and A1 are members of the *Desulfobacteraceae*, and the relative abundance of the sequences was very high in the libraries from the anoxic monimolimnion. The OTUs might correspond to each other, but require additional evidence to demonstrate genomic linkage.

Members of the DSS group are known as nutritionally versatile SRB [42]. Most of them oxidize organic compounds completely to CO₂, and several species can grow autotrophically with H₂ and sulfate. The DSS group of bacteria is abundant in the upper layer of sediments in many aquatic environments, such as permanently cold marine sediment [31], sediments above gas hydrate [15], temperate tidal flat sediment [26], and a meromictic Lake Cadagno sediment [34]. However, in anoxic water columns, only a few reports about the abundance of the DSS group are available. For example in a permanently anoxic water column in the Black Sea, SRB detected with the SRB385 probe are abundant below 100 m, where sulfide appears, and comprise up to 2–8% of total DAPI counts [22]. The abundance of DSS almost coincides with the cell counts obtained with the SRB385 probe [22, 40]; therefore, DSS is the most dominant group of SRB in the anoxic layer of the Black Sea water column. Some sequences from the DSS group are found in an anoxic brine pool of the Gulf of Mexico [11], below the chemocline of the stratified lagoon in Clipperton atoll [6], and below the chemocline of the meromictic Lake Suigetsu [19]. In contrast, very few sequences from the DSS group have been detected in seasonal anoxic water masses such as those in Saanich Inlet [41], and the Arabian Sea [5], which indicates that the DSS group might prefer permanently anoxic environments in which reduced sulfur compounds can accumulate.

Many SEEP SRB-1d cells were observed in the anoxic layer of the water column (Fig. 6). It has been suggested that SEEP SRB-1 subgroups interact with ANME-2 [15, 33] or use hydrocarbons including short-chain alkanes [13,14]. However, in Lake Harutori, we could not detect ANME-2 cells by CARD-FISH with the specific probe ANME-2 538 [39] (data not shown). In addition, the cells detected with the probe SEEP1d-468 and DSS658 were often dispersed as single cells or formed chain-like small aggregates by themselves. Hence, the specific association of the SEEP SRB-1d with ANME-2 cells is unlikely in this lake. There is a possibility that SEEP SRB-1 have a preference for the availability of hydrocarbons. We detected methane (approximately 1.5 mM) in the monimolimnion of Lake Harutori (data not shown), but no data is available for other hydrocarbons. Further study will be required to confirm this hypothesis.

The DSS group in the chemocline of Lake Harutori can probably use various organic compounds including persistent substances and might prefer a permanently anoxic environment, and are therefore able to be one of the major SRB in the bottom of Lake Harutori. Further cultivation and incubation studies will be required to estimate the contribution of these SRB to sulfide production and carbon remineralization in anoxic water columns, which may expand in the near future.

Acknowledgments

We greatly acknowledge R. Amann and K. Knittel for their helpful discussions. We are grateful to T. Hariu and the Kushiro City Museum for their help. We thank R. Tokizawa for her technical assistance. We acknowledge M. Watanabe, K. Kogure and Y. Koizumi for their help in the sampling. This work was supported by JSPS KAKENHI Grant Number 22370005 to M. Fukui, 24870001 to K. Kubo and the F3 program of the Hokkaido University.

- [1] Chan, F., Barth, J.A., Lubchenco, J., Kirincich, A., Weeks, H., Peterson, W., Menge, B.A. (2008) Emergence of anoxia in the California Current large marine ecosystem. *Science*. 319, 920.
- [2] Chikita, K., Fukuyama, R., Sakamoto, H., Nakamichi, K. (1997) Dynamic behaviors of 'dead water' in a coastal lagoon, Lake Harutori, Kushiro, Hokkaido: Field measurements for the ice-covered period. *Geophys. Bull. Hokkaido Univ.* 60, 13-28. [In Japanese with English abstract]
- [3] Cline, J.D. (1969) Spectrophotometric determination of hydrogen sulfide in natural waters. *Limnol. Oceanogr.* 14, 454-458.
- [4] Dar, S.A., Kuenen, J.G., Muyzer, G. (2005) Nested PCR-denaturing gradient gel electrophoresis approach to determine the diversity of sulfate-reducing bacteria in complex microbial communities. *Appl. Environ. Microbiol.* 71, 2325-2330.
- [5] Fuchs, B.M., Woebken, D., Zubkov, M.V., Burkill, P., Amann, R. (2005) Molecular identification of picoplankton populations in contrasting waters of the Arabian Sea. *Aquat. Microb. Ecol.* 39, 145-157.
- [6] Galand, P.E., Bourrain, M., De Maistre, E., Catala, P., Desdevises, Y., Elifantz, H., Kirchman, D.L., Lebaron, P. (2012) Phylogenetic and functional diversity of Bacteria and Archaea in a unique stratified lagoon, the Clipperton atoll (N Pacific). *FEMS Microbiol. Ecol.* 79, 203-217.
- [7] Grantham, B.A., Chan, F., Nielsen, K.J., Fox, D.S., Barth, J.A., Huyer, A., Lubchenco, J., Menge, B.A. (2004) Upwelling-driven nearshore hypoxia signals ecosystem and oceanographic changes in the northeast Pacific. *Nature*. 429, 749-754.
- [8] Higashioka, Y., Kojima, H., Nakagawa, T., Sato, S., Fukui, M. (2009) A novel *n*-alkane-degrading bacterium as a minor member of *p*-xylene-degrading sulfate-reducing consortium. *Biodegradation*, 20, 383-390.
- [9] Higashioka, Y., Kojima, H., Watanabe, M., Fukui, M. (2013) *Desulfatitalea tepidiphila* gen. nov., sp. nov., a sulfate-reducing bacterium isolated from tidal flat sediment. *Int. J. Syst. Evol. Microbiol.* 63, 761-765.
- [10] Janssen, P.H., Schuhmann, A., Bak, F., Liesack, W. (1996) Disproportionation of inorganic sulfur compounds by the sulfate-reducing bacterium *Desulfocapsa thiozymogenes* gen. nov., sp. nov. *Arch. Microbiol.* 166, 184-192.
- [11] Joye, S.B., Samarkin, V.A., Orcutt, B.N., MacDonald, I.R., Hinrichs, K.-U., Elvert, M., Teske, A.P., Lloyd, K.G., Lever, M.A., Montoya, J.P., Meile, C.D. (2009) Metabolic variability in seafloor brines revealed by carbon and sulphur dynamics. *Nat. Geosci.* 2, 349-354.
- [12] Kelly, D.P., Wood, A.P. (2000) Confirmation of *Thiobacillus denitrificans* as a species of the genus *Thiobacillus*, in the β -subclass of the *Proteobacteria*, with strain NCIMB 9548 as the type strain. *Int. J. Syst. Evol. Microbiol.* 50, 547-550.
- [13] Kleindienst, S., Ramette, A., Amann, R., Knittel, K. (2012) Distribution and *in situ* abundance of sulfate-reducing bacteria in diverse marine hydrocarbon seep sediments. *Environ. Microbiol.* 14, 2689-2710.
- [14] Kniemeyer, O., Musat, F., Sievert, S.M., Knittel, K., Wilkes, H., Blumenberg, M., Michaelis, W., Classen, A., Bolm, C., Joye, S.B., Widdel, F. (2007) Anaerobic oxidation of short-chain hydrocarbons by marine sulphate-reducing bacteria. *Nature*. 449, 898-901.
- [15] Knittel, K., Boetius, A., Eilers, A.L.H., Lochte, K., Linke, O.P.P. (2003) Activity, distribution, and diversity of sulfate reducers and other bacteria in sediments above gas

- hydrate (Cascadia Margin, Oregon). Geomicrobiol. J. 20, 269-294.
- [16] Knoblauch, C., Sahm, K., Jørgensen, B.B. (1999) Psychrophilic sulfate-reducing bacteria isolated from permanently cold Arctic marine sediments: description of *Desulfofrigus oceanense* gen. nov., sp. nov., *Desulfofrigus fragile* sp. nov., *Desulfofaba gelida* gen. nov., sp. nov., *Desulfotalea psychrophila* gen. nov., sp. nov. and *Desulfotalea arctica* sp. nov. Int. J. Syst. Bacteriol. 49, 1631-1643.
- [17] Koizumi, Y., Kojima, H., Oguri, K., Kitazato, H., Fukui, M. (2004) Vertical and temporal shifts in microbial communities in the water column and sediment of saline meromictic Lake Kaiike (Japan), as determined by a 16S rDNA-based analysis, and related to physicochemical gradients. Environ. Microbiol. 6, 622-637.
- [18] Kojima, H., Fukui, M. (2010) *Sulfuricella denitrificans* gen. nov., sp. nov., a sulfur-oxidizing autotroph isolated from a freshwater lake. Int. J. Syst. Evol. Microbiol. 60, 2862-2866.
- [19] Kondo, R., Osawa, K., Mochizuki, L., Fujioka, Y., Butani, J. (2006) Abundance and diversity of sulphate-reducing bacterioplankton in Lake Suigetsu, a meromictic lake in Fukui, Japan. Plankton. Benthos. Res. 1, 165-177.
- [20] Kuever J., Rainey F.A., Widdel F. (2005) Genus II. *Desulfobacterium* Bak and Widdel 1988, 136^{VP} (Effective publication: Bak and Widdel 1986, 175). In: Brenner, D.J., Krieg, N.R., Staley, J.T., Garrity, G.M. (Eds.), Bergey's Manual of Systematic Bacteriology. Second edition, vol. 2 (The *Proteobacteria*), part C (The *Alpha*-, *Beta*-, *Delta*-, and *Epsilonproteobacteria*), Springer, New York, pp. 965-967.
- [21] Lavik, G., Stührmann, T., Brüchert, V., Van der Plas, A., Mohrholz, V., Lam, P., Mußmann, M., Fuchs, B.M., Amann, R., Lass, U., Kuypers, M.M.M. (2009) Detoxification of sulphidic African shelf waters by blooming chemolithotrophs. Nature. 457, 581-584.
- [22] Lin, X., Wakeham, S.G., Putnam, I.F., Astor, Y.M., Scranton, M.I., Chistoserdov, A.Y., Taylor, G.T. (2006) Comparison of vertical distributions of prokaryotic assemblages in the anoxic Cariaco Basin and Black Sea by use of fluorescence in situ hybridization. Appl. Environ. Microbiol. 72, 2679-2690.
- [23] Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, Buchner, A., Lai, T., Steppi, S., Jobb, G., Förster, W., Brettske, I., Gerber, S., Ginhart, A.W., Gross, O., Grumann, S., Hermann, S., Jost, R., König, A., Liss, T., Lüssmann, R., May, M., Nonhoff, B., Reichel, B., Strehlow, R., Stamatakis, A., Stuckmann, N., Vilbig, A., Lenke, M., Ludwig, T., Bode, A., Schleifer, K.-H. (2004) ARB: a software environment for sequence data. Nucleic Acids Res. 32, 1363-1371.
- [24] Meyer, B., Kuever, J. (2007) Molecular analysis of the distribution and phylogeny of dissimilatory adenosine-5'-phosphosulfate reductase-encoding genes (*aprBA*) among sulfur-oxidizing prokaryotes. Microbiology. 153, 3478-98.
- [25] Meyer, B., Kuever, J. (2007) Molecular analysis of the diversity of sulfate-reducing and sulfur-oxidizing prokaryotes in the environment, using *aprA* as functional marker gene. Appl. Environ. Microbiol. 73, 7664-7679.
- [26] Mußmann, M., Ishii, K., Rabus, R., Amann, R. (2005) Diversity and vertical distribution of cultured and uncultured *Deltaproteobacteria* in an intertidal mud flat of the Wadden Sea. Environ. Microbiol. 7, 405-418.
- [27] Pernthaler, A., Pernthaler, J., Amann, R. (2002) Fluorescence *in situ* hybridization and catalyzed reporter deposition for the identification of marine bacteria. Appl. Environ. Microbiol. 68, 3094-3101.
- [28] Pruesse, E., Peplies, J., Glöckner, F.O. (2012) SINA: accurate high-throughput multiple

- sequence alignment of ribosomal RNA genes. *Bioinformatics*, 28, 1823-1829.
- [29] Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O. (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucl. Acids Res.* 41, D590-D596.
- [30] Rappé, M.S., Connon, S.A., Vergin, K.L., Giovannoni, S.J. (2002) Cultivation of the ubiquitous SAR11 marine bacterioplankton clade. *Nature*. 418, 630-633.
- [31] Ravensschlag, K., Sahm, K., Knoblauch, C., Jørgensen, B.B., Amann, R. (2000) Community structure, cellular rRNA content, and activity of sulfate-reducing bacteria in marine arctic sediments. *Appl. Environ. Microbiol.* 66, 3592-3602.
- [32] Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Van Horn, D.J., Weber, C.F. (2009) Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537-7541.
- [33] Schreiber, L., Holler, T., Knittel, K., Meyerdierks, A., Amann, R. (2010) Identification of the dominant sulfate-reducing bacterial partner of anaerobic methanotrophs of the ANME-2 clade. *Environ. Microbiol.* 12, 2327-2340.
- [34] Schubert, C.J., Vazquez, F., Lösekann-Behrens, T., Knittel, K., Tonolla, M., Boetius, A. (2011) Evidence for anaerobic oxidation of methane in sediments of a freshwater system (Lago di Cadagno). *FEMS Microbiol. Ecol.* 76, 26-38.
- [35] Takahashi, M., Ichimura, S. (1968) Vertical distribution and organic matter production of photosynthetic sulfur bacteria in Japanese lakes. *Limnol. Oceanogr.* 13, 644-655.
- [36] Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S. (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731-2739.
- [37] Teira, E., Reinthaler, T., Pernthaler, A., Pernthaler, J., Herndl, G.J. (2004) Combining catalyzed reporter deposition-fluorescence in situ hybridization and microautoradiography to detect substrate utilization by bacteria and archaea in the deep ocean. *Appl. Environ. Microbiol.* 70, 4411-4414.
- [38] Teske, A., Wawer, C., Muyzer, G., Ramsing, N.B. (1996) Distribution of sulfate-reducing bacteria in a stratified fjord (Mariager Fjord, Denmark) as evaluated by most-probable-number counts and denaturing gradient gel electrophoresis of PCR-amplified ribosomal DNA fragments. *Appl. Environ. Microbiol.* 62, 1405-1415.
- [39] Treude, T., Knittel, K., Blumenberg, M., Seifert, R., Boetius, A. (2005) Subsurface microbial methanotrophic mats in the Black Sea. *Appl. Environ. Microbiol.* 71, 6375-6378.
- [40] Wakeham, S.G., Amann, R., Freeman, K.H., Hopmans, E.C., Jørgensen, B.B., Putnam, I.F., Schouten, S., Damsté, J.S.S., Talbot, H.M., Woebken, D. (2007) Microbial ecology of the stratified water column of the Black Sea as revealed by a comprehensive biomarker study. *Org. Geochem.* 38, 2070-2097.
- [41] Walsh, D.A., Zaikova, E., Howes, C.G., Song, Y.C., Wright, J.J., Tringe, S.G., Tortell, P.D., Hallam, S.J. (2009) Metagenome of a versatile chemolithoautotroph from expanding oceanic dead zones. *Science*. 326, 578-582.
- [42] Widdel, F., Bak, F. (1992) Gram-negative mesophilic sulfate-reducing bacteria. In: Balows A., Trüper H. G., Dworkin M., Harder W., Schleifer K.-H. (Eds.), *The prokaryotes*, Volume 1, Second Edition, Springer-Verlag, New York, pp. 3352-3378.
- [43] Wilson, K. (1987) Preparation of genomic DNA from bacteria. In: Ausubel, F., Brent, R., Kingston, R., Moore, D., Seidman, J., Smith, J., Struhl, K. (Eds.), *Current Protocols in*

492 Molecular Biology, Greene Publ. Associates and Wiley-Interscience, New York. pp.
493 2.4.1–2.4.5.
494 [44] Yoshimura, S. (1937) Koshougaku. Enlarged Edition, Seisan-gijyutsu center, Tokyo, pp.
495 202. [In Japanese]
496

Figure captions

- Fig. 1 Map of Lake Harutori in Kushiro, Hokkaido, Japan.
- Fig. 2 Depth profiles of temperature, pH, specific conductivity, concentrations of chloride, dissolved oxygen, sulfate, sulfide, and dissolved organic carbon in the water column of Lake Harutori in May 2012.
- Fig. 3 Phylogenetic tree based on deduced AprA amino acid sequences retrieved from Lake Harutori water column. The tree was constructed based on a distance matrix (119 amino acid positions) by using the neighbor-joining method. Bootstrap value estimation was based on 1000 replicates (only values above 50% are shown). Numbers of sequences obtained in each clone library were indicated in parentheses; 3.0 m, 3.5 m, and 4.0 m in that order. a) Overview tree. b) Magnified tree of *Desulfobacteraceae* and *Desulfobulbaceae*. The bar represents 5% estimated sequence divergence.
- Fig. 4 Phylogenetic tree showing the affiliation of Lake Harutori 16S rRNA sequences to selected reference sequences of *Desulfobacteraceae* of *Deltaproteobacteria*. Representative sequences from each OTU are in boldface type. The name of the representative clone and the number of the grouped sequences are in parentheses. Probe specificity is indicated by the vertical lines. The bar represents 10% estimated sequence divergence.
- Fig. 5 Abundance of microbial cells in Lake Harutori water column determined by CARD-FISH. DAPI (closed circle), EUBI-III (closed triangle), Delta495a (closed square), and DSS658 (open circle).

519 Fig. 6 Abundance of *Deltaproteobacteria* (probe Delta495a), DSS group (probe DSS658)
520 and SEEP SRB-1d cells (probes SEEP1d-1420 and SEEP1d-468) in the anoxic
521 layer of Lake Harutori, determined by CARD-FISH.

Fig. 1
Kubo et al.

Fig. 2
Kubo et al.

Fig. 3a

Kubo et al.

Fig. 4

Kubo et al.

Fig. 5
Kubo et al.

Fig. 6
Kubo et al.

Table 1. Phylogenetic affiliation of classified OTUs of aprA clone sequences and the number of clones in each clone libraries.

Putative function	Phylogenetic affiliation	OTU	Closest cultured strain ^a	Identity (%)	Number of clones			
					3.0 m	3.5 m	4.0 m	
Sulfate reduction	<i>Desulfobacteraceae</i>	OTU A1	<i>Desulfofaba gelida</i> PSv29 (AAL57385)	88-92	0	8	41	
		OTU A2	<i>Desulfatitalea tepidiphila</i> S28OL2 (BAM15569)	94-97	10	0	0	
		OTU A6	<i>Desulfatitalea tepidiphila</i> S28OL2 (BAM15569)	92-93	0	0	8	
		OTU A7	<i>Desulfobacterium vacuolatum</i> lbrM (AAL57391)	96-97	0	7	0	
		OTU A8	<i>Desulfofaba gelida</i> PSv29 (AAL57385)	88-89	5	0	0	
		OTU A9	<i>Desulfofaba gelida</i> PSv29 (AAL57385)	94-96	1	4	0	
		OTU A14	<i>Desulfobacterium indolicum</i> InO4 (AAL57390)	96	0	0	3	
		OTU A15	<i>Desulfobacterium indolicum</i> InO4 (AAL57390)	96	0	1	1	
		OTU A24	<i>Desulfobacterium autotrophicum</i> HRM2 (AAL57375)	97	0	1	0	
		OTU A25	<i>Desulfobacterium autotrophicum</i> HRM2 (AAL57375)	95	0	1	0	
		OTU A26	<i>Desulfofaba gelida</i> PSv29 (AAL57385)	92	0	1	0	
		OTU A27	<i>Desulfatitalea tepidiphila</i> S28bF (BAM15567)	97	0	1	0	
		OTU A28	<i>Desulfobacterium indolicum</i> InO4 (AAL57390)	94	0	1	0	
		OTU A32	<i>Desulfobacterium indolicum</i> InO4 (AAL57390)	94	0	0	1	
		OTU A33	<i>Desulfobacterium indolicum</i> InO4 (AAL57390)	93	0	0	1	
		OTU A34	<i>Desulfatitalea tepidiphila</i> S28OL1 (BAM15568)	92	0	0	1	
		OTU A35	<i>Desulfatitalea tepidiphila</i> S28OL1 (BAM15568)	92	0	0	1	
		OTU A36	<i>Desulfatitalea tepidiphila</i> S28OL1 (BAM15568)	92	0	0	1	
		OTU A47	<i>Desulfofaba gelida</i> PSv29 (AAL57385)	93	1	0	0	
		OTU A48	<i>Desulfonema magnum</i> Montpellier (AAL57389)	94	1	0	0	
		OTU A49	<i>Desulfonema magnum</i> Montpellier (AAL57389)	92	1	0	0	
		OTU A50	<i>Desulfobacterium autotrophicum</i> HRM2 (AAL57375)	94	1	0	0	
		OTU A51	<i>Desulfatitalea tepidiphila</i> S28OL2 (BAM15569)	91	1	0	0	
		<i>Desulfobulbaceae</i>	OTU A3	<i>Desulfocapsa thiozymogenes</i> Bra2 (AAL57433)	98-100	0	8	1
			OTU A31	<i>Desulforhopalus vacuolatus</i> ltk 10 (ABR92535)	99	0	0	1
		<i>Syntrophaceae</i>	OTU A16	<i>Desulfobacca acetoxidans</i> ASRB2 (ABR92549)	96	2	0	0
		Gram positive SRB and related <i>Deltaproteobacteria</i>	OTU A4	<i>Desulfobacterium anilini</i> Ani1 (AAL57425)	86-88	7	2	0
	OTU A12		<i>Desulfotomaculum kuznetsovii</i> 17 (AAL57419)	86	3	0	0	
	OTU A17		<i>Desulfotomaculum kuznetsovii</i> 17 (AAL57419)	83	1	0	0	
	OTU A18		<i>Desulfotomaculum geothermicum</i> BSD (AAL57382)	83	1	0	0	
	OTU A19		<i>Desulfotomaculum kuznetsovii</i> 17 (AAL57419)	79	1	0	0	
	OTU A29		<i>Desulfotomaculum thermoacetoxidans</i> CAMZ (ABR92588)	78	0	1	0	
	OTU A37		<i>Desulfobacterium anilini</i> Ani1 (AAL57425)	88	0	0	1	
			OTU A38	<i>Desulfotomaculum kuznetsovii</i> 17 (AAL57419)	80	0	0	1
Sulfur oxidation	Lineage 1 ^b	OTU A11	<i>Thiocapsa rosea</i> 6611 (ABV80104)	79	4	0	0	
		OTU A21	<i>Thiocapsa rosea</i> 6611 (ABV80104)	92	1	0	0	
		OTU A22	<i>Thiocapsa rosea</i> 6611 (ABV80104)	90	1	0	0	
		OTU A23	<i>Lamprocystis purpurea</i> ThSch 12 (ABV80005)	89	0	1	0	
	Lineage 2 ^b	OTU A5	<i>Thiobacillus denitrificans</i> GP3 (ABV80033)	98-100	4	4	0	
		OTU A10	<i>Sulfuricella denitrificans</i> skB26 (BAI66427)	97-98	4	0	0	
		OTU A39	<i>Sulfuricella denitrificans</i> skB26 (BAI66427)	97	1	0	0	
		OTU A40	<i>Sulfuricella denitrificans</i> skB26 (BAI66427)	97	1	0	0	
		OTU A41	<i>Sulfuricella denitrificans</i> skB26 (BAI66427)	91	1	0	0	
		OTU A42	<i>Sulfuricella denitrificans</i> skB26 (BAI66427)	92	1	0	0	
		OTU A43	<i>Sulfuricella denitrificans</i> skB26 (BAI66427)	83	1	0	0	
		OTU A44	<i>Sulfuricella denitrificans</i> skB26 (BAI66427)	85	1	0	0	
		OTU A45	<i>Sulfuricella denitrificans</i> skB26 (BAI66427)	87	1	0	0	
		OTU A46	<i>Thiothrix</i> sp. CT3 (ABV80023)	100	1	0	0	
Uncertain	Unclassified ^c	OTU A13	N.A.	N.A.	3	0	0	
		OTU A20	N.A.	N.A.	1	0	0	
		OTU A30	N.A.	N.A.	0	1	0	
			Total number of clones	62	41	62		

^a Closest cultured strains were determined according to BLASTP search result. GenBank accession number of AprA protein is in parentheses.^b Classification of the lineages of sulfur-oxidizing bacteria is according to Meyer and Kuever [24].^c Sequences have less than 70% sequence identity to any known sequences were grouped as Unclassified.

N.A., Not available