



Title	Comparative Genomic View of The Inositol-1,4,5-Trisphosphate Receptor in Plants
Author(s)	Mikami, Koji
Citation	Journal of Plant Biochemistry & Physiology, 2(3), 132 https://doi.org/10.4172/2329-9029.1000132
Issue Date	2014-9-15
Doc URL	http://hdl.handle.net/2115/57862
Type	article
File Information	comparative-genomic-view-of-the-inositoltrisphosphate-receptor.pdf



[Instructions for use](#)

Comparative Genomic View of The Inositol-1,4,5-Trisphosphate Receptor in Plants

Koji Mikami*

Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate 041 - 8611, Japan

Abstract

Terrestrial plants lack inositol-1,4,5-trisphosphate (IP₃) receptor regulating transient Ca²⁺ increase to activate cellular Ca²⁺-dependent physiological events. To understand an evolutionary route of the loss of the IP₃ receptor gene, conservation of the IP₃ receptor gene in algae was examined *in silico* based on the accumulating information of genomes and expression sequence tags. Results clearly demonstrated that the lack of the gene was observed in Rhodophyta, Chlorophyta except for Volvocales and Streptophyta. It was therefore hypothesized that the plant IP₃ receptor gene was eliminated from the genome at multiple occasions; after divergence of Chlorophyta and Rhodophyta and of Chlorophyta and Charophyta.

Keywords: Alga; Ca²⁺; Comparative genomics; Gene; Inositol-1,4,5-trisphosphate receptor

Abbreviations: DAG: Diacylglycerol; IP₃: Inositol-1,4,5-Trisphosphate; IP₆: Inositol-1,2,3,4,5,6-Hexakisphosphate, PI-PLC: Phosphoinositide-Specific Phospholipase C, PKC: Protein Kinase C.

Inositol-1,4,5-trisphosphate [Ins(1,4,5)P₃, IP₃] is a second messenger involved in transient release of Ca²⁺ from the ER that activates cytosolic Ca²⁺ signalling cascades in response to extracellular and intracellular stimuli [1,2]. Phosphatidylinositol-4,5-bisphosphate is cleaved by phosphoinositide-specific phospholipase C (PI-PLC) into the second messengers diacylglycerol (DAG) and IP₃ [3,4]. These second messengers then activate protein kinase C (PKC) and the ER-localised IP₃ receptor, respectively, in animal cells [1,2]. However, although the PI-PLC signaling cascade is present in plants [5-7], genes encoding PKC and the IP₃ receptor have not been found in terrestrial plant genomes, suggesting differences in second messenger systems between animals and plants. To date, the genomes of a variety of unicellular and multicellular algae have been sequenced [8-23] as shown in (Table 1). In addition, large-scale EST information for the red seaweeds *Porphyra umbilicalis* and *Porphyra purpurea* has been accumulated [24-26]. Such rich gene information enables us to identify the genes encoding IP₃ receptor gene homologues in algae to hypothesize the evolutionary route of the loss of the IP₃ gene in plant lineages.

The origin of the IP₃ receptor-dependent transient Ca²⁺ release system predates the divergence of animals and fungi [27,28]. Indeed, homologues of genes encoding the IP₃ receptor have been identified in protozoa such as the choanoflagellate *Monosiga brevicollis* [29], the myxomycete *Dictyostelium discoideum* [30], the ciliate *Paramecium tetraurelia* [31], and the parasite *Trypanosoma brucei* [32]. Thus, it is plausible that an ancient eukaryotic cell containing an IP₃ receptor gene was the target of endosymbiosis with an ancient cyanobacterium to produce plant cells, after which the IP₃ gene was lost from plant lineages. At present, IP₃ receptor homologues have been found in green algae, such as *Chlamydomonas reinhardtii* [10] and *Volvox carteri* [33,34], and in heterokont algae including *Aureococcus anophagefferrens* [21] and *Ectocarpus siliculosus* [22], but have not been identified in red algae or streptophytes (land plants and charophytic algae) (Figure 1). These findings have led to proposals that the IP₃ receptor gene homologue was lost on multiple occasions during plant evolution. Because an ancestor of both green and red photosynthetic algal cells appeared after the primary endosymbiosis of a cyanobacterium into an ancient non-photosynthetic eukaryotic cell [35], the IP₃ receptor homologue

was probably lost from lineages of red algae and green algae except for Volvocales (Figure 1). In fact, the genomes of unicellular *Aureococcus anophagefferrens* and multicellular *Ectocarpus siliculosus* carry an IP₃ receptor gene homologue (Figure 1). Because both photosynthetic algae arose from secondary endosymbiosis of a red algal cell into an ancient non-photosynthetic eukaryotic cell [35], it appears that red algae subsequently lost the IP₃ receptor gene homologue during their evolution, although some of *Heterokontophyta* that evolved by secondary symbiosis retain an ancient progenitor of the IP₃ receptor gene to this date. Moreover, in the green plant lineage, streptophytes have an impaired IP₃ receptor that is structurally similar to that in animals, Volvocales of chlorophytes, and brown seaweed (Figure 1). Thus, the loss of the IP₃ receptor may also occurred after the divergence of chlorophytes and streptophytes. Accordingly, there have been multiple occasions upon which the IP₃ receptor was lost from plant lineages. In contrast to the above conclusions drawn from genomic sequence information, there is evidence of IP₃-dependent Ca²⁺ release in terrestrial plants [36-42], which suggests the presence of a Ca²⁺ channel functionally resembling the IP₃ receptor in streptophytes. However, IP₃-dependent Ca²⁺ release has been reported only in green algae among plants [43,44]. Because the major intracellular store of Ca²⁺ in plant cells is the vacuole [45,46], IP₃ receptor activity is thought to be localised to vacuolar membranes in green algae and streptophytes. Such is the case in the fungus *Neurospora crassa*, in which IP₃-mediated Ca²⁺ release occurs from vacuoles [47], as it also does in protozoan ciliates and trypanosomes, in which the IP₃ receptor has been visualized on vacuolar membranes [27,28]. Thus, the green plant lineage has maintained an ancient system for transient release of Ca²⁺ from vacuoles, which is distinct from ER-mediated Ca²⁺ release in animal cells that do not possess vacuoles.

*Corresponding author: Koji Mikami, Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate 041 - 8611, Japan, Tel: +81-138-40-8899; Fax: +81-138-40-8899 E-mail: komikami@fish.hokudai.ac.jp

Received July 17, 2014; Accepted September 08, 2014; Published September 15, 2014

Citation: Mikami K (2014) Comparative Genomic View of The Inositol-1,4,5-Trisphosphate Receptor in Plants. J Plant Biochem Physiol 2: 132. doi:10.4172/2329-9029.1000132

Copyright: © 2014 Mikami K. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Phylum	Class	Order	Family	Species	Ref
Chlorophyta	Prasinophyceae	Mamiellales	Mamiellaceae	<i>Ostreococcus tauri</i>	[8]
				<i>Ostreococcus lucimarinus</i>	[9]
	Chlorophyceae	Volvocales	Chlamydomonadaceae	<i>Chlamydomonas reinhardtii</i>	[10]
			Volvocaceae	<i>Volvox carteri</i>	[11]
		Chlorococcales	Coccomyxaceae	<i>Coccomyxa subellipsoidea</i>	[12]
Rhodophyta	Cyanidiophyceae	Cyanidiales	Cyanidiaceae	<i>Cyanidioschyzon merolae</i>	[13]
				<i>Galdieria sulphuraria</i>	[14]
	Porphyridiophyceae	Porphyridiales	Porphyridiaceae	<i>Porphyridium purpureum</i>	[15]
	Rhodophyceae	Bangiales	Bangiaceae	<i>Pyropia yezoensis</i>	[16]
	Florideophyceae	Gigartinales	Gigartiniaceae	<i>Chondrus crispus</i>	[17]
Glaucophyta	Glaucophyceae	Glaucocystales	Glaucocystaceae	<i>Cyanophora paradoxa</i>	[18]
Heterokontophyta	Coccinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Thalassiosira pseudonana</i>	[19]
	Bacillariophyceae	Naviculales	Phaeodactylaceae	<i>Phaeodactylum tricorutum</i>	[20]
	Pelagophyceae	Pelagomonadales	Pelagomonadaceae	<i>Aureococcus anophagefferens</i>	[21]
	Phaeophytia	Ectocarpales	Ectocarpaceae	<i>Ectocarpus siliculosus</i>	[22]
Charophyta	Klebsormiophyceae	Klebsormidiales	Klebsormiaceae	<i>Klebsormidium flaccidum</i>	[23]

Table 1: List of algal species whose genome sequences have been analyzed.




Figure 1: Conservation of the IP₃ receptor genes in algae and terrestrial plants. The presence of the gene is indicated by "+" with highlighting order names with yellow.

IP₃-mediated Ca²⁺ release has been observed at the ER membrane that predominates in the perinuclear and apex regions of cells of the brown seaweed *Fucus serratus* [48]. Because an IP₃ receptor homologue was found in the *Ectocarpus siliculosus* genome [22], it is possible that the location of the IP₃-sensitive Ca²⁺ store shifted from the vacuole to the ER in brown seaweeds, where it is currently found in animal cells.

Thus, the brown seaweeds might possess a PI-PLC signaling system more similar to that in animals.

Although IP₃-mediated Ca²⁺ release has not yet been shown in red algae, an inhibitor of the IP₃-receptor, 2-APB, prevented establishment of cell polarity for the migration and germination of monospores in the red seaweed *Pyropia yezoensis* [49], which suggests the presence of an IP₃ receptor-mediated Ca²⁺ release system in red seaweeds. However, an IP₃-receptor homologue has not yet been identified in the *Pyropia yezoensis* genome. As there is currently no evidence indicating the presence of IP₃ in *Pyropia yezoensis*, biochemical determinations of this inositol derivative will be necessary to elucidate Ca²⁺ release upon PI-PLC action in red algae.

In plant cells, DAG is usually phosphorylated by DAG kinase [50,51] to produce phosphatidic acid, and IP₃ is phosphorylated by inositol phosphate kinases, IPK1 and IPK2 [52,53] to produce inositol-1,3,4,5,6-pentakisphosphate and inositol-1,2,3,4,5,6-hexakisphosphate [Ins(1,2,3,4,5,6)P₆; phytate; IP₆], a high-abundance molecule that is considered important for phosphorus storage in plant cells. To date, PA and IP₆ are thought to act as major second messengers in plant cells [7,54], although the function of IP₃ as a second messenger in plants has not been ruled out [42,47]. For instance, Munnik and Vermeer [54] have proposed that IP₆, which is rapidly converted from IP₃, is a major second messenger involved in abscisic acid-dependent inhibition of stomatal opening. They have also proposed a parallel between the IP₃ and IP₆ signalling systems because these two molecules are both produced by the action of PI-PLC. Although neither an IP₃ nor an IP₆ receptor have yet been identified in terrestrial plants, it is possible that an IP₃ receptor or an IP₆ receptor of unknown structure is present in streptophytes. Taken together, comparative genomic information clearly demonstrates the loss of the IP₃ receptor gene in red algae, green algae except for Volvocales and streptophytes during plant evolution. However, IP₃-dependent transient Ca²⁺ release from intracellular stores has been shown in these organisms by physiological experiments, although whether plants lacking the IP₃ receptor might both possess a common system for such transient Ca²⁺ release is uncertain. Therefore, the identification and characterization of genes encoding putative IP₃ or IP₆ receptors of unknown structure is of the highest priority for elucidating and comparing the regulation of the PI-PLC signalling cascade between IP₃ receptor-carrying and -lacking algae.

References

1. Finch EA, Augustine GJ (1998) Local calcium signalling by inositol-1,4,5-trisphosphate in Purkinje cell dendrites. *Nature* 396: 753-756.
2. Taylor CW, Tovey SC, Rossi AM, Lopez Sanjurjo CI, Prole DL, et al. (2014) Structural organization of signalling to and from IP₃ receptors. *Biochem Soc Trans* 42: 63-70.
3. Rebecchi MJ, Pentyala SN (2000) Structure, function, and control of phosphoinositide-specific phospholipase C. *Physiol Rev* 80: 1291-1335.
4. Suh PG, Park JI, Manzoli L, Cocco L, Peak JC, et al. (2008) Multiple roles of phosphoinositide-specific phospholipase C isozymes. *BMB Rep* 41: 415-434.
5. Xue HW, Chen X, Mei Y (2009) Function and regulation of phospholipid signalling in plants. *Biochem J* 421: 145-156.
6. Janda M, Planchais S, Djafi N, Martinec J, Burketova L, et al. (2013) Phosphoglycerolipids are master players in plant hormone signal transduction. *Plant Cell Rep* 32: 839-851.
7. Pokotylo I, Kolesnikov Y, Kravets V, Zachowski A, Ruelland E (2014) Plant phosphoinositide-dependent phospholipases C: variations around a canonical theme. *Biochimie* 96: 144-157.
8. Derelle E, Ferraz C, Rombauts S, Rouzé P, Worden AZ, et al. (2006) Genome analysis of the smallest free-living eukaryote *Ostreococcus tauri* unveils many unique features. *Proc Natl Acad Sci U S A* 103: 11647-11652.
9. Palenik B, Grimwood J, Aerts A, Rouzé P, Salamov A, et al. (2007) The tiny eukaryote *Ostreococcus* provides genomic insights into the paradox of plankton speciation. *Proc Natl Acad Sci U S A* 104: 7705-7710.
10. Merchant SS, Prochnik SE, Vallon O, Harris EH, Karpowicz SJ, et al. (2007) The *Chlamydomonas* genome reveals the evolution of key animal and plant functions. *Science* 318: 245-250.
11. Prochnik SE, Umen J, Nedelcu AM, Hallmann A, Miller SM, et al. (2010) Genomic analysis of organismal complexity in the multicellular green alga *Volvox carterii*. *Science* 329: 223-226.
12. Blanc G, Agarkova I, Grimwood J, Kuo A, Brueggeman A, et al. (2012) The genome of the polar eukaryotic microalga *Coccomyxa subellipsoidea* reveals traits of cold adaptation. *Genome Biol* 13:R39.
13. Matsuzaki M, Misumi O, Shin-I T, Maruyama S, Takahara M, et al. (2004) Genome sequence of the ultrasmall unicellular red alga *Cyanidioschyzon merolae* 10D. *Nature* 428: 653-657.
14. Barbier G, Oesterhelt C, Larson MD, Halgren RG, Wilkerson C, et al. (2005) Comparative genomics of two closely related unicellular thermo-acidophilic red algae, *Galdieria sulphuraria* and *Cyanidioschyzon merolae*, reveals the molecular basis of the metabolic flexibility of *Galdieria sulphuraria* and significant differences in carbohydrate metabolism of both algae. *Plant Physiol* 137: 460-474.
15. Bhattacharya D, Price DC, Chan CX, Qiu H, Rose N, et al. (2013) Genome of the red alga *Porphyridium purpureum*. *Nat Commun* 4: 1941.
16. Nakamura Y, Sasaki N, Kobayashi M, Ojima N, Yasuike M, et al. (2013) The first symbiont-free genome sequence of marine red alga, *Susabi-nori* (*Pyropia yezoensis*). *PLoS One* 8: e57122.
17. Collén J, Porcel B, Carré W, Ball SG, Chaparro C, et al. (2013) Genome structure and metabolic features in the red seaweed *Chondrus crispus* shed light on evolution of the Archaeplastida. *Proc Natl Acad Sci U S A* 110: 5247-5252.
18. Price DC, Chan CX, Yoon HS, Yang EC, Qiu H, et al. (2012) *Cyanophora paradoxa* genome elucidates origin of photosynthesis in algae and plants. *Science* 335: 843-847.
19. Armbrust EV, Berges JA, Bowler C, Green BR, Martinez D, et al. (2004) The genome of the diatom *Thalassiosira pseudonana*: ecology, evolution, and metabolism. *Science* 306: 79-86.
20. Bowler C, Allen AE, Badger JH, Grimwood J, Jabbari K, et al. (2008) The *Phaeodactylum* genome reveals the evolutionary history of diatom genomes. *Nature* 456: 239-244.
21. Gobler CJ, Berry DL, Dyhrman ST, Wilhelm SW, Salamov A, et al. (2011) Niche of harmful alga *Aureococcus anophagefferens* revealed through ecogenomics. *Proc Natl Acad Sci U S A* 108: 4352-4357.
22. Cock JM, Sterck L, Rouzé P, Scornet D, Allen AE, et al. (2010) The *Ectocarpus* genome and the independent evolution of multicellularity in brown algae. *Nature* 465: 617-621.
23. Hori K, Maruyama F, Fujisawa T, Togashi T, Yamamoto N, et al. (2014) *Klebsormidium flaccidum* genome reveals primary factors for plant terrestrial adaptation. *Nat Commun* 5:3978.
24. Chan CX, Blouin NA, Zhuang Y, Zäuner S, Prochnik SE, et al. (2012) *Porphyra* (Bangiphyceae) transcriptomes provide insights into red algal development and metabolism. *J Phycol* 48: 1328-1342.
25. Chan CX, Zäuner S, Wheeler G, Grossman AR, Prochnik SE, et al. (2012) Analysis of *Porphyra* membrane transporters demonstrates gene transfer among photosynthetic eukaryotes and numerous sodium-coupled transport systems. *Plant Physiol* 158: 2001-2012.
26. Stiller JW, Perry J, Rymarquis LA, Accerbi M, Green PJ, et al. (2012) Major developmental regulators and their expression in two closely related species of *Porphyra* (Rhodophyta). *J Phycol* 48: 883-896.
27. Docampo R, Moreno SN, Plattner H (2013) Intracellular calcium channels in protozoa. *Eur J Pharmacol* 15: 4-18.
28. Plattner H, Verkhatsky A (2013) Ca²⁺ signalling early in evolution--all but primitive. *J Cell Sci* 126: 2141-2150.
29. Cai X (2008) Unicellular Ca²⁺ signaling 'toolkit' at the origin of metazoa. *Mol Biol Evol* 25: 1357-1361.
30. Traynor D, Milne JL, Insall RH, Kay RR (2000) Ca²⁺ signalling is not required for chemotaxis in *Dictyostelium*. *EMBO J* 19: 4846-4854.
31. Ladenburger EM, Korn I, Kasielke N, Wassmer T, Plattner H (2006) An Ins(1,4,5)P₃ receptor in *Paramecium* is associated with the osmoregulatory system. *J Cell Sci* 119: 3705-3717.
32. Huang G, Bartlett PJ, Thomas AP, Moreno SN, Docampo R (2013) Acidocalcisomes of *Trypanosoma brucei* have an inositol 1,4,5-trisphosphate receptor that is required for growth and infectivity. *Proc Natl Acad Sci U S A* 110: 1887-1892.
33. Wheeler GL, Brownlee C (2008) Ca²⁺ signalling in plants and green algae--changing channels. *Trends Plant Sci* 13: 506-514.
34. Verret F, Wheeler G, Taylor AR, Farnham G, Brownlee C (2010) Calcium channels in photosynthetic eukaryotes: implications for evolution of calcium-based signalling. *New Phytol* 187: 23-43.
35. Keeling PJ (2010) The endosymbiotic origin, diversification and fate of plastids. *Philos Trans R Soc Lond B Biol Sci* 365: 729-748.
36. Drøbak BK, Ferguson IB (1985) Release of Ca²⁺ from plant hypocotyl microsomes by inositol-1,4,5-trisphosphate. *Biochem Biophys Res Commun* 130: 1241-1246.
37. Poovaiah BW, Reddy AS (1987) Calcium messenger system in plants. *CRC Crit Rev Plant Sci* 6: 47-103.
38. Ranjeva R, Carrasco A, Boudet AM (1998) Inositol trisphosphate stimulates the release of calcium from intact vacuoles isolated from *Acer* cells. *FEBS Lett* 230: 137-141.
39. Gilroy S, Read ND, Trewavas AJ (1990) Elevation of cytoplasmic calcium by caged calcium or caged inositol triphosphate initiates stomatal closure. *Nature* 346: 769-771.
40. Blatt MR, Thiel G, Trentham DR (1990) Reversible inactivation of K⁺ channels of *Vicia* stomatal guard cells following the photolysis of caged inositol 1,4,5-trisphosphate. *Nature* 346: 766-769.
41. Muir SR, Sanders D (1997) Inositol 1,4,5-trisphosphate-sensitive Ca²⁺ release across nonvacuolar membranes in cauliflower. *Plant Physiol* 114: 1511-1521.
42. Monteiro D, Liu Q, Lisboa S, Scherer GE, Quader H, et al. (2005) Phosphoinositides and phosphatidic acid regulate pollen tube growth and reorientation through modulation of [Ca²⁺]_c and membrane secretion. *J Exp Bot* 56: 1665-1674.
43. Thiel G, MacRobbie EA, Hanke DE (1990) Raising the intracellular level of inositol 1,4,5-trisphosphate changes plasma membrane ion transport in characean algae. *EMBO J* 9: 1737-1741.
44. Förster B (1990) Injected inositol 1,4,5-trisphosphate activates Ca²⁺-sensitive K⁺ channels in the plasmalemma of *Eremosphaera viridis*. *FEBS Lett* 269: 197-201.

45. Sanders D, Brownlee C, Harper JF (1999) Communicating with calcium. *Plant Cell* 11: 691-706.
46. Krinke O, Novotná Z, Valentová O, Martinec J (2007) Inositol trisphosphate receptor in higher plants: is it real? *J Exp Bot* 58: 361-376.
47. Cornelius G, Gebauer G, Techel D (1989) Inositol trisphosphate induces calcium release from *Neurospora crassa* vacuoles. *Biochem Biophys Res Commun* 162: 852-856.
48. Goddard H, Manison NF, Tomos D, Brownlee C (2000) Elemental propagation of calcium signals in response-specific patterns determined by environmental stimulus strength. *Proc Natl Acad Sci U S A* 97: 1932-1937.
49. Li L, Saga N, Mikami K (2009) Ca²⁺ influx and phosphoinositide signalling are essential for the establishment and maintenance of cell polarity in monospores from the red alga *Porphyra yezoensis*. *J Exp Bot* 60: 3477-3489.
50. Arisz SA, Testerink C, Munnik T (2009) Plant PA signaling via diacylglycerol kinase. *Biochim Biophys Acta* 1791: 869-875.
51. Dong W, Lv H, Xia G, Wang M (2012) Does diacylglycerol serve as a signaling molecule in plants? *Plant Signal Behav* 7: 472-475.
52. Stevenson-Paulik J, Odom AR, York JD (2002) Molecular and biochemical characterization of two plant inositol polyphosphate 6-/3-/5-kinases. *J Biol Chem* 277: 42711-42718.
53. Stevenson-Paulik J, Bastidas RJ, Chiou ST, Frye RA, York JD (2005) Generation of phytate-free seeds in *Arabidopsis* through disruption of inositol polyphosphate kinases. *Proc Natl Acad Sci U S A* 102: 12612-12617.
54. Munnik T, Vermeer JE (2010) Osmotic stress-induced phosphoinositide and inositol phosphate signalling in plants. *Plant Cell Environ* 33: 655-669.

Citation: Mikami K (2014) Comparative Genomic View of The Inositol-1,4,5-Trisphosphate Receptor in Plants. *J Plant Biochem Physiol* 2: 132.
doi:[10.4172/2329-9029.1000132](https://doi.org/10.4172/2329-9029.1000132)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>

