Title	Pathology of frontotemporal dementia with limb girdle muscular dystrophy caused by a DNAJB6 mutation
Author(s)	Yabe, Ichiro; Tanino, Mishie; Yaguchi, Hiroaki; Takiyama, Akihiro; Cai, Huaying; Kanno, Hiromi; Takahashi, Ikuko; Hayashi, Yukiko K.; Watanabe, Masashi; Takahashi, Hidehisa; Hatakeyama, Shigetsugu; Tanaka, Shinya; Sasaki, Hidenao
Citation	Clinical neurology and neurosurgery, 127, 10-12 https://doi.org/10.1016/j.clineuro.2014.09.013
Issue Date	2014-12
Doc URL	http://hdl.handle.net/2115/57945
Type	article (author version)
File Information	Case_report_revise_ver5.pdf

[Instructions for use](#)

Case Report

Pathology of frontotemporal dementia with limb girdle muscular dystrophy caused by a *DNAJB6* mutation

Ichiro Yabe^{a,*#}, Mishie Tanino^{b#}, Hiroaki Yaguchi^{a,c#}, Akihiro Takiyama^b, Huaying Cai^a,
Hiromi Kanno^b, Ikuko Takahashi^a, Yukiko K. Hayashi^d, Masashi Watanabe^c, Hidehisa
Takahashi^c, Shigetsugu Hatakeyama^c, Shinya Tanaka^b, Hidenao Sasaki^a

^a Department of Neurology, Hokkaido University Graduate School of Medicine,
Sapporo, Japan

^b Department of Cancer Pathology, Hokkaido University Graduate School of
Medicine, Sapporo, Japan

^c Department of Biochemistry, Hokkaido University Graduate School of Medicine,
Sapporo, Japan

^d Department of Neuromuscular Research, NCNP, Kodaira, Japan

These authors equally contributed to this work.

* Correspondence to Ichiro Yabe

e-mail; yabe@med.hokudai.ac.jp

Running head: FTD with DNAJB6 myopathy

Word counts of paper; 1,040 words

Number of tables; 0

Number of figures; 2

Number of References; 7

Supplemental figures; 2

Funding/Support: This work was supported in part by a Grant-in-Aid for the Research Committee of CNS Degenerative Diseases of the Research on Measures for Intractable Diseases from the Ministry of Health, Welfare and Labor, Japan, and by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture, Japan.

Conflict of interest: None

Key words; frontotemporal dementia, DNAJB6, brain pathology, LC-3, autophagy

Introduction

Frontotemporal lobar degeneration (FTLD) is a heterogeneous group of disorders characterized by disturbances of behavior and personality and different types of language impairment with or without concomitant features of motor neuron disease or parkinsonism [1]. FTLD is classified into three categories: frontotemporal dementia

(FTD), progressive non-fluent aphasia (PNFA), and semantic dementia (SD) [2].

Moreover, FTD is classified into two categories: pick type and motor neuron disease (MND) type. FTLD is also classified into four neuropathological categories: FTLD-tau pathology type, FTLD-TDP43 pathology type, FTLD-FUS pathology type, and FTLD-other type [2]. Clinical FTD-Pick type is associated with FTLD-tau, TDP, or FUS, and some FTLD is caused by various genetic mutations.

Limb girdle muscular dystrophy (LGMD) is a progressive myopathy with necrosis and regenerative changes in skeletal muscle. LGMD Type 1D is caused by mutations in the *DNAJB6* (DnaJ homolog subfamily B member 6) gene [3]. Muscle pathology shows aggregates of TDP-43 and DNAJB6, together with rimmed vacuoles.

Here, we describe the phenotype and pathology of a case with frontotemporal

dementia (FTD) and LGMD caused by a *DNAJB6* mutation.

Case report

A 61 year old man complained of progressive lower limb muscle weakness, similar to his 3 siblings. Muscle CT demonstrated fatty degeneration, and the patient was diagnosed as LGMD. Genetic analysis indicated a heterozygous missense mutation of c.279C>G (p. Phe93Leu) in *DNAJB6* (Family D-1 in the ref.3) [4]. At 75 years old, he became stubborn and angry toward his family, and was suggested to have a behavioral variant of FTD [2]. These symptoms gradually progressed, and brain magnetic resonance images showed frontal atrophy with mild temporal atrophy (Fig. 1A,B). However, hippocampal atrophy and cerebellar atrophy were not observed (Fig. 1C,D). He developed pneumonia frequently, and died of septic shock at 76 years old. No mutations were observed in *VCP*, *PRGN* and *MAPT* genes and no abnormal expansion of *C9ORF72* was observed, all of which are common causes of hereditary FTD [1].

Total brain weight was 1,338 g, and mild neuronal loss was observed in the frontal and temporal cortex. Axonal structure was preserved (Suppl. Fig.1. A), and no

neurofibrillary tangles, senile plaques, Pick bodies, or Lewy bodies were observed in

the brainstem, basal ganglia, frontotemporal cortex or white matter (Suppl. Fig.1. B-D).

No TDP-43, Tau or FUS reactivity was observed in the central nervous system (CNS)

(Suppl. Fig.1. E-H). However, there was a severe reduction of DNAJB6

immunoreactivity in the frontal cortex (Fig. 1E) compared to samples from normal

control (Fig. 1F), an ALS-dementia patient without underlying proteinopathy (Fig. 1G),

a TDP-43 positive ALS patient (Fig. 1H), and a TDP-43 negative ALS patient (Fig. 1I).

DNAJB6 staining in the thalamus was preserved (Fig. 1J), and was similar to that in

normal control (Fig. 1K). While intranuclear inclusion bodies were not detected,

ubiquitin-positive aggregates were observed in the frontotemporal cortex (Fig. 2A),

some of which appeared co-localized with p62 (Fig. 2B). The expression of LC3 in this

patient (Fig.2C) was reduced compared to that of normal control (Fig. 2D).

Discussion

DNAJB6 is highly enriched in the CNS and inhibits polyglutamine aggregation of

huntingtin. In addition, DNAJB6 is a component of Lewy bodies [5]. The availability of

DNAJ proteins, co-chaperones to the Hsp70 machine, may be a rate-limiting factor in

handling diseased proteins within the cell. DNAJ proteins can affect the aggregation of disease-causing proteins [6]. These reports indicate that DNAJB6 is important in the etiology of CNS neurodegenerative diseases. However, there are no reports of FTD associated with mutations in *DNAJB6*. Anti-DNAJB6 antibody recognized both wild type and mutant DNAJB6 proteins (Suppl. Fig. 2) in all samples. The tissue samples were obtained at autopsy performed at similar time intervals after death, and the samples were stained simultaneously. Therefore, the severe reduction of DNAJB6 in this patient is considered to be significant. Moreover, ubiquitin-positive aggregates were detected in the cerebral cortex, consistent with FTD. However, neither TDP-43, Tau nor FUS reactivity were observed in the CNS. Therefore, this patient was diagnosed as “FTLD-other”, or FTD with ubiquitin-positive aggregates and without tau, TDP, and FUS pathology [2]. Previous studies suggest that DNAJB6 facilitates ubiquitin-dependent degradation substrates. Therefore, this *DNAJB6* mutation may be closely related to neurodegeneration. Previously, we reported that DNAJB6 accumulated in the cytoplasm where it co-localized with members of the autophagy complex [4]. In this study, we showed a severe reduction of DNAJB6 staining, and a

mild reduction of LC3 staining of the cerebral cortex in our patient. These results suggest that this patient showed autophagy dysfunction in the brain, as well as in skeletal muscle. Although p62 immunoreactive inclusions have been observed in FTLD cases, with an underlying *C9ORF72* mutation [1], we did not find any mutations in *VCP*, *PRGN* and *MAPT* genes and no abnormal expansion of *C9ORF72*. In addition, the accumulations of ubiquitin were co-localized with p62, which is a common component of neuronal protein inclusions, a major characteristic of neurodegenerative diseases [7] and a key molecule managing autophagic clearance of polyubiquitinated proteins. In this study, the reduction of LC3 and p62 staining suggests a loss of autophagy function by the *DNAJB6* mutation. Although there were mild reductions of DNAJB6 staining and aggregation of p62 in the cerebellum, aggregation of ubiquitin and reduction of LC3 staining were not observed. This patient showed no ataxia or cerebellar atrophy. These results suggest that the cerebellar functional loss may be milder than that of the frontotemporal lobe.

Some genes are known to be associated with multisystem proteinopathy. Mutations in *VCP* cause vacuolar myopathy and FTD, and influence autophagy, an important

mechanism of dementias including FTD. Moreover, the muscle pathology of LGMD1D

has similarity with that of vacuolar myopathy. Loss of VCP activity leads to the

accumulation of ubiquitinated proteins and impaired autophagy. These studies and our

findings suggest that dysfunction of ubiquitination and autophagy may be a common

mechanism between *VCP* and *DNAJB6* mutations.

Conclusions

We report a pathological study of a frontotemporal dementia patient with LGMD caused

by a *DNAJB6* mutation. A missense mutation in *DNAJB6* may cause not only LGMD,

but also FTD through the dysfunction of autophagy. Further study is necessary to

determine whether LGMD and FTD are caused by the same pathomechanism.

Functional analysis of DNAJB6 at the molecular level should provide therapeutic

benefits for LGMD and dementia.

References

1. Sieben A, Van Langenhove T, Engelborghs S, Martin JJ, Boon P, Cras P, et al. The genetics and neuropathology of frontotemporal lobar degeneration. *Acta Neuropathol* 2012; 124: 353-372.
2. Josephs KA, Hodges JR, Snowden JS, Mackenzie IR, Neumann M, Mann DM, et al. Neuropathological background of phenotypical variability in frontotemporal dementia. *Acta Neuropathol* 2011; 122: 137-153.
3. Harms MB, Sommerville RB, Allred P, Bell S, Ma D, Cooper P, et al. Exome sequencing reveals DNAJB6 mutations in dominantly-inherited myopathy. *Ann Neurol* 2012; 71: 407-416.
4. Sato T, Hayashi YK, Oya Y, Kondo T, Sugie K, Kaneda D, et al. DNAJB6 myopathy in an Asian cohort and cytoplasmic/nuclear inclusions. *Neuromuscul Disord* 2013; 23: 269-276.
5. Durrenberger PF, Filiou MD, Moran LB, Michael GJ, Novoselov S, Cheetham ME, et al. DnaJB6 is present in the core of Lewy bodies and is highly up-regulated in parkinsonian astrocytes. *J Neurosci Res* 2009; 87: 238-245.

6. Kakkar V, Prins LC, Kampinga HH. DNAJ proteins and protein aggregation diseases. *Curr Top Med Chem* 2012; 12: 2479-2490.
7. Kragh CL, Ubhi K, Wyss-Coray T, Masliah E. Autophagy in dementias. *Brain Pathol* 2012; 22: 99-109.

Figure Legends

Figure 1. Brain MRI and immunoreactivity for DNAJB6

A-D: T2-weighted magnetic resonance images of the brain, showing bilateral frontotemporal atrophy, but not cerebellar atrophy.

E-K: Immunoreactivity for DNAJB6. Reduced DNAJB6 immunoreactivity in the left frontal lobe of this patient (E) is shown, compared with normal control (F), an ALS-dementia patient (G), a TDP-43 positive ALS patient (H) and a TDP-43 negative ALS patient (I). DNAJB6 immunoreactivity in the thalamus is not reduced (J) compared to normal control (K). Scale bar, 100 μ m.

Figure 2. Immunoreactivity for p62 and LC3

Serial sections of grey matter in the left frontal lobe of this patient (A and B). A: Immunoreactivity for ubiquitin is observed in some aggregations. B: Some p62-immunoreactive aggregations are observed. Aggregations indicated by the arrow and arrow head seem to be co-localized. C and D: LC3 immunoreactivity in the gray matter in the left frontal lobe. The expression of LC3 immunoreactivity of this patient

(C) is reduced compared to that of normal control (D). Scale bar, 100 μ m

A)

B)

C)

D)

E)

F)

G)

H)

I)

J)

K)

Yabe et al. Figure 2