Title	Calcium concentration in leaf litter alters the community composition of soil invertebrates in warm-temperate forests
Author(s)	Ohta, Tamihisa; Niwa, Shigeru; Agetsuma, Naoki; Hiura, Tsutom
Citation	Pedobiologia, 57(4-6), 257-262 https://doi.org/10.1016/j.pedobi.2014.07.003
Issue Date	2014-11
Doc URL	http://hdl.handle.net/2115/58011
Type	article (author version)
File Information	Pedobiologia_rev3.pdf

[Instructions for use](#)

1 Title: Calcium concentration in leaf litter alters the community composition of soil

2 invertebrates in warm-temperate forests.

3
4 Authors: Tamihisa Ohta¹, Shigeru Niwa², Naoki Agetsuma³ and Tsutom Hiura¹

5
6 *Affiliations:*

7 ¹Tomakomai Research Station, Field Science Center for Northern Biosphere, Hokkaido

8 University, Takaoka, Tomakomai, Hokkaido 053-0035, Japan

9 ²Network Center of Forest and Grassland Survey in Monitoring Sites 1000 Project,

10 Japan Wildlife Research Center, Takaoka, Tomakomai, Hokkaido 053-0035, Japan

11 ³Wakayama Experimental Forest, Field Science Center for Northern Biosphere,

12 Hokkaido University, Hirai, Kozagawa-cho Higashimuro-gun, Wakayama, 649-4563,

13 Japan

14
15 Running title: Calcium in leaf litter and soil organisms

16

17 *Corresponding Author: Email: tammyohta@gmail.com Phone: +81-0144-33-2171,

18 Fax: +81-0144-33-2173

19 Email:

20 S. Niwa: sniwa@fsc.hokudai.ac.jp

21 N. Agetsuma: agetsuma@fsc.hokudai.ac.jp

22 T. Hiura: hiura@fsc.hokudai.ac.jp

23

24

25

26

27

28

29

30

31

32

33 ABSTRACT

34

35 Many studies have shown the effects of aboveground plant species on soil organisms
36 due to differences in litter quality. However, the calcium concentration in soil has
37 received less attention as a controlling factor of soil invertebrate communities, despite
38 its being an essential element for many animals, especially crustaceans. Litter of
39 Japanese cedar (*Cryptomeria japonica*) plantations, which account for 19% of the
40 forested area in Japan, has a higher calcium concentration compared to other taxa such
41 as broad-leaved trees. We predicted that *C. japonica* plantations affect soil invertebrates
42 by altering calcium availability. We compared soil properties including exchangeable
43 calcium concentration and soil invertebrate communities between *C. japonica*
44 plantations and natural broad-leaved forests. Exchangeable calcium was significantly
45 higher in soil from cedar plantations than in that from broad-leaved forests. The
46 invertebrate community composition differed between the two forest types and was best
47 explained by the exchangeable calcium concentration. In particular, two major taxa of
48 soil crustaceans (Talitridae and *Ligidium japonicum*) were found only in cedar

49 plantations. Our results suggest that calcium concentrations in soil are altered in *C.*

50 *japonica* plantations and that this affects soil invertebrate communities.

51

52 *Keywords:* Evergreen broad-leaved trees, *Cryptomeria japonica*, forest management,

53 crustaceans, *Ligidium japonicum*, Talitridae

54

55

56

57

58

59

60

61

62

63

64

65 **Introduction**

66

67 Soil organisms can be affected by differences in aboveground vegetation (Bardgett
68 and Wardle 2010), often driven by the quality of the litter types (Swift et al. 1979; Berg
69 and McClaugherty 2003). Differences in litter quality among plant species can influence
70 the chemical properties of soil and act as determinants of the community structure of
71 soil invertebrates (Widden and Hsu 1987; Wardle et al. 2006). Because calcium is a
72 major structural component of the proteins forming animals, ambient calcium
73 concentration related to the animal densities in calcium poor environment (Alstad et al.
74 1999; Hessen et al. 2000; Ohta et al. 2014). Similarly, soil calcium concentrations can
75 have an important influence on soil invertebrate communities (Springett and Syers
76 1984). For example, the abundance of some soil invertebrates increase with the
77 available soil calcium concentration (Hotopp 2002; Reich et al. 2005; Skeldon et al.
78 2007). Global pattern of soil calcium concentration is governed by geological change,
79 acid deposition and annual precipitation (Potter and Conkling 2012; Binkley and Fisher
80 2013). For example, low rainfall areas (central North America) tend to have grater soil

81 calcium than humid Eastern United States (Potter and Conkling 2012). Calcium
82 availability and cycling in regional scale are governed by numerous factors including
83 forest vegetation dynamics, atmospheric deposition, soil mineral weathering, and losses
84 due to leaching (Likens et al. 1998; McLaughlin and Wimmer 1999; Dijkstra and Smits
85 2002). However, much research over the last half century has focused on the leaching of
86 calcium due to acid deposition (Likens et al. 1996; Driscoll et al. 2001), with much less
87 emphasis on other factors, such as changes in forest vegetation, even though the calcium
88 concentration in leaf litter varies greatly among tree species.

89 Litter of members of the Cupressaceae has a higher concentration of calcium
90 compared with other plant families (Kiilsgaard et al. 1987; Ohta et al. 2014). In this
91 study, we focused on Japanese cedar (*Cryptomeria japonica*, Cupressaceae) because its
92 leaf litter contains ~3% calcium (Xue and Luo 2002; Baba et al. 2004), more than three
93 times the amount in many other taxa, such as fir (*Abies* spp.) and many broad-leaved
94 trees (Kiilsgaard et al. 1987; Reich et al. 2005; Ohta et al. 2014). Japanese cedar
95 plantations cover 12% of the total land area and 19% of the forested area in Japan
96 (Forestry Agency 2011). Because soil organic matter in forests is derived mainly from

plant litter, the chemical properties of litter affect soil chemical properties (Reich et al. 2005). Indeed, the soil in Japanese cedar plantations has a calcium content that is three to four times higher than that in evergreen broad-leaved forests in some parts of Japan (Tsutsumi 1987; Ohta et al. 2014). Ohta et al. (2014) showed that the calcium concentration in soil and streams, and the density and survival of dominant aquatic crustaceans, were significantly higher in *C. japonica* plantations compared with evergreen broad-leaved forests. However, Ohta et al. (2014) did not assess the effects of forest vegetation on soil animal community through alteration of calcium availability. Soil crustaceans that are frequently dominant decomposers in soil systems (O'hanlon and Bolger 1999), and contain large amount of calcium in their body (Greenaway 1985). Terrestrial crustaceans mainly take calcium from their food (e.g. leaf litter) and soil water. Therefore, we anticipate that the calcium concentration in litter affects the community structure of soil organisms in calcium-poor environment.

Addition of inorganic calcium often increases soil pH (Likens et al. 1996; Driscoll et al. 2001; Warby et al. 2009), and therefore, higher calcium concentrations in soil due to differences in forest vegetation are also likely to increase soil pH (Reich et al. 2005).

113 Alteration of soil pH also causes changes in the abundance of soil invertebrates (Hågvar
114 1990; Myrold 1990; Kaneko and Kofuji 2000). Therefore, plantations of *C. japonica*
115 may affect the community structure of soil invertebrates via increased soil pH.

116 We examined the effect of Japanese cedar (*C. japonica*) plantations on the
117 community structure of soil invertebrates, particularly the density of crustaceans. We
118 conducted field surveys in six plots that differed in surrounding forest vegetation. We
119 predicted (1) that the calcium concentration and soil pH would be higher in *C. japonica*
120 plantations compared to evergreen broad-leaved forests, and (2) that crustacean density
121 would be higher in *C. japonica* plantations than in evergreen broad-leaved forests.

122

123 **Methods**

124

125 *Study area*

126

127 We conducted field surveys in the Wakayama Experimental Forest of Hokkaido
128 University (33°40'N, 135°40'E; 428 ha; annual mean temperature: 15.2°C) on the

129 Southern Kii Peninsula of Japan. The geological structure in this region consists of
130 sandstone and mudstone formed during the middle Tertiary (Tateishi 1976). Because of
131 the highly acidic soil and high annual rainfall (~4000 mm), the area is extremely poor in
132 calcium (Kihira et al. 2005). The forest soils are extremely thin, nearly exposing the
133 bedrock. Japanese cedar was planted in much of the area beginning in the 1960s, and
134 remnant natural evergreen broad-leaved forests are patchy.

135 We established a sampling plot (50 × 50 m) in each of six different catchments of
136 the Wakayama Experimental Forest. The plots were located on relatively flat forest
137 floors and separated by 0.2–1.5 km. Three of the six catchments were mostly covered
138 by evergreen broad-leaved forests ‘evergreen’, and the other three were covered by
139 Japanese cedar plantations ‘cedar’. Forests in the ‘evergreen’ plots were dominated by
140 *Quercus acuta*, *Quercus myrsinifolia*, *Quercus sessilifolia*, *Neolitsea aciculata*, *Eurya*
141 *japonica*, and *Machilus thunbergii* (Ohta et al. 2014). The *C. japonica* trees in the
142 ‘cedar’ plots were planted 30–82 years prior to this study. Calcium concentration in the
143 litter of *C. japonica* (3.4%) is about three times higher than in the evergreen
144 broad-leaved species (0.8–1.5%) at this study site (Ohta et al. 2014). Carbon, nitrogen,

phosphorus, and magnesium concentrations do not differ significantly among the species, whereas potassium is about three times lower in *C. japonica* compared to the broad-leaved species (Ohta et al. 2014).

Sampling

On 24 July 2012, we collected five samples at each plot from the litter and soil layers using core samplers (soil layer: 50 mm in diameter and 50 mm in height, litter layer: 113 mm in diameter and 40 mm in height) to measure soil chemical properties and mass of the litter layer. To determine the soil crustacean density, we established five sampling quadrats separated by over 10 m in each plot. We collected crustaceans within the sampling quadrats (25 × 25-cm) to a depth of 3 cm (including litter and surface soil layers) on 17 May and 21 September 2013. Soil crustaceans were separated from soil by hand-sorting and placed in 99% ethanol. To examine the community structure of ground-dwelling macroinvertebrates, we established two subplots (20 × 20 m) in each plot. We collected ground-dwelling invertebrates using five pitfall traps (8 cm in

diameter and 6 cm in depth) per subplot. We placed the pitfall traps in each subplot ~2 m apart on 17 May, 19 July, 19 October, and 18 November 2013, and collected them 3 days later. We counted and identified all invertebrates found in the traps at least to the ordinal level following Aoki (1999) and Ueno et al. (1985).

Sample processing

To measure soil nitrate and exchangeable calcium, we shook (160 rev min^{-1}) a 0.5 g (air-dried mass) subsample of each soil sample in 100 ml of 1 M KCl solution for 1 h, filtered the sample through filter paper (No. 5C; Advantec, Tokyo, Japan), and then stored the suspension at -30°C until analysis. We analyzed the soil extracts for calcium and nitrate concentration per unit air-dried mass using an inductively coupled plasma (ICP) atomic emission spectrometer (ICPE-9000; Shimadzu, Kyoto, Japan) and the absorptiometric method (Sakata 2000). We placed a 5-g (air-dried mass) subsample of each soil sample in 25 ml 1 M KCL and measured the pH using a pH meter (TOA-DKK, HM-30V; TOA Electronics, Tokyo, Japan). We dried soil subsamples in a drying oven

177 at 60°C for 24 h and then analyzed 50-mg dried soil samples for carbon and nitrogen
178 concentrations per dry mass using a CN analyzer (Sumigraph NC-900; Sumika
179 Chemical Analysis Service, Osaka, Japan). We dried a 1-g fresh subsample at 60°C for
180 48 h to calculate the soil water content as the difference in mass before and after
181 desiccation.

182

183 *Statistical analysis*

184

185 The soil properties (mass of the litter layer, water content, pH, C:N ratio,
186 exchangeable calcium, total carbon, total nitrogen, and nitrate concentration) were fit to
187 linear mixed models with forest vegetation type as a fixed factor and plot identity as a
188 random factor. The statistical significance of the effect of the fixed factor in each model
189 was evaluated by a likelihood ratio test ($\alpha = 0.05$).

190 We performed canonical correspondence analysis (CCA) to explore the
191 relationships between the soil invertebrate composition and soil properties. The
192 invertebrate data from the five pitfall traps on all four sampling dates were pooled for

each subplot. Before conducting the CCA ordination, the abundance data for each taxon were standardized to unit variance, and the most important explanatory variables from all soil properties were determined by forward stepwise selection based on Akaike's information criteria and Monte Carlo permutation tests. All statistical analyses were conducted with R version 2.9.2 software (R Development Core Team 2011).

Results

Soil exchangeable calcium concentration (likelihood ratio test: $\chi^2 = 9.13$, d.f. = 1, $P = 0.002$) and C:N ratio (likelihood ratio test: $\chi^2 = 8.90$, d.f. = 1, $P = 0.003$) were significantly higher, and nitrate concentration (likelihood ratio test: $\chi^2 = 8.53$, d.f. = 1, $P = 0.003$) was significantly lower in 'cedar' than 'evergreen'. In particular, soil in 'cedar' plots had ~2.5 times more exchangeable calcium than the 'evergreen' plots (Table 1, Fig. 1). The other soil properties including the mass of litter layer that was selected explanatory variable in community composition of soil invertebrates did not differ significantly between the forest types although soil pH was marginally higher in 'cedar'

209 than ‘evergreen’ (likelihood ratio test: $\chi^2 = 3.05$, d.f. = 1, $P = 0.08$). We found two taxa
210 of crustaceans, Talitridae (Amphipoda) and *Ligidium japonicum* (Isopoda: Ligiidae) in
211 ‘cedar’ plots by hand-sorting, but we found no crustaceans in ‘evergreen’ plots (Fig. 2).
212 Abundances of Talitridae and *L. japonicum* did not differ significantly among ‘cedar’
213 plots between the two sampling months (generalized linear models assuming Poisson
214 distribution and likelihood ratio tests: $P > 0.05$).

215 Crustaceans, spiders, ants, beetles, lepidopteran larvae, and millipedes accounted for
216 23, 19, 9, 19, 10, and 9%, respectively, of the total invertebrates captured by pitfall traps
217 (Table 2). Crustaceans (mainly Talitridae and *L. japonicum*) dominated the
218 ground-dwelling invertebrate communities in ‘cedar’ plots, while they were remarkably
219 scarce, and millipedes (Paradoxosomatidae) and beetles were relatively abundant in
220 ‘evergreen’ plots. Talitridae and *L. japonicum* were only collected in ‘cedar’ plots using
221 pitfall traps, and a few individuals of *Venezillo* sp. (Isopoda: Armadillidae) were
222 collected not only in ‘cedar’ but also ‘evergreen’ plots (Table 2). Taxonomic
223 compositions of ground invertebrates differed distinctively between ‘cedar’ and
224 ‘evergreen’ plots (Table 2, Fig. 3). From all the soil properties, exchangeable calcium

concentration and mass of the litter layer explained the most variation among the 12 subplots in invertebrate community composition as determined through the forward selection process of CCA. The first and second axes explained 17.17 and 14.04%, respectively, of the variation in community composition (Monte Carlo permutation test: $P < 0.05$). The CCA ordination showed that community composition in the ‘cedar’ plots was distinctively different from that in the ‘evergreen’ plots along the first CCA axis, which corresponded to the gradient of exchangeable calcium concentration, and the ‘cedar’ plots had much lower variation than the ‘evergreen’ plots (Fig. 3). Compositional difference within ‘evergreen’ was distinctively indicated by second axis, which weakly correlated with mass of litter layer (Fig. 3). Talitridae and *L. japonicum* had large negative values on the first axis, and their high abundance characterized the ‘cedar’ community.

Discussion

Our results show that forest vegetation might affect the community structure of soil

241 invertebrates by altering calcium availability. Our field survey showed that soil
242 exchangeable calcium was ~2.5 times higher and soil pH was not significantly higher in
243 *C. japonica* plantations compared with that in evergreen broad-leaved forests (these
244 partly supported prediction 1). The major taxa of soil crustaceans (Talitridae and *L.*
245 *japonicum*) were found only in *C. japonica* plantations, whereas only a few individuals
246 of a minor crustacean species (*Venezillo* sp.) occurred in broad-leaved forests
247 (supporting prediction 2). The community structure of soil invertebrates varied with
248 forest vegetation types, and calcium in the soil layer was the most important
249 environmental variable explaining the variation in community composition (supporting
250 prediction 2). Furthermore, both intra- and inter-plot variation in the soil invertebrate
251 community structure in *C. japonica* plantations were lower than in natural broad-leaved
252 forests, suggesting that the homogeneous environment created by the monoculture
253 plantation caused a large decrease in micro- and local-scale β -diversity of soil
254 invertebrates. This might be caused by uniform increase in the abundance of Talitridae
255 and *L. japonicum* in *C. japonica* plantations and difference in the mass of litter layer
256 within ‘evergreen’ plots (Table 1, Fig. 3).

257 The difference in calcium concentration in leaf litter produces significant
258 differences in soil calcium (Morrison 1985; Kloeppel and Abrams 1995). In fact, Ohta
259 et al. (2014) showed that the calcium concentration in *C. japonica* litter was about three
260 times higher than that of dominant evergreen broad-leaved trees in Wakayama
261 Experimental Forest. We found that total calcium in the litter layer and
262 water-extractable calcium in the soil at ‘cedar’ sites were three to four times higher than
263 at ‘evergreen’ sites. These results support our finding that calcium supplied by *C.*
264 *japonica* litter increased the calcium concentration in the soil of our study plots.

265 Field manipulations at the Hubbard Brook Experimental Forest in the northeastern
266 United States indicated that adding CaSiO_3 to a catchment area increased calcium
267 concentrations in soil (Juice et al. 2006; Minocha et al. 2010; Nezat et al. 2010) and
268 altered the community structure of terrestrial snails, which have high demand for
269 calcium (Skeldon et al. 2007). Hotopp (2002) showed that the abundance of sugar
270 maple, a calciphilic species (Likens and Bormann 1970), was positively correlated with
271 terrestrial snail density. As in snails, crustaceans must ingest a lot of calcium
272 (Greenaway 1985). Because terrestrial crustaceans lose 20% of their body calcium

273 through exuviae (Ziegler et al. 2007), they need adequate calcium to calcify their
274 exoskeleton rapidly after exuviation. Terrestrial crustaceans in soil get calcium from
275 soil water and litter (Greenaway 1985; Glötzner and Ziegler 2000). Therefore, the
276 ‘cedar’ plots, which have high calcium levels in their litter and soil layers, are well
277 suited for the survival of crustaceans (Figs. 1, 2). Indeed, both natural and artificial *C.*
278 *japonica* forests in central Japan have high densities of crustaceans (Ikeda et al. 2005).

279 Calcium addition to acidic soil increases soil pH (Groffman et al. 2006; Groffman
280 and Fink 2011). However, our result showed the soil pH was not significantly higher in
281 *C. japonica* plantations that increase the calcium concentration of the soils (Table 1).

282 Although previous studies showed soil pH may also affect the structure of invertebrate
283 communities by altering forest vegetation (e.g. Kaneko and Kofuji 2000), forward
284 selection of the CCA analysis selected exchangeable soil calcium concentration and not
285 pH in our result. Calcium concentration may have a greater impact on the community
286 structure of soil invertebrates in our study site.

287 The litter of other members of Cupressaceae, such as *Chamaecyparis* and
288 *Sequoiadendron*, also have high calcium content comparable to that of *C. japonica*

289 (Kiilsgaard et al. 1987; D'Amore et al. 2009). This might mean change in soil system
290 through difference in calcium concentration in leaf litter is caused in other forest types.
291 Furthermore, Reich et al. (2005) showed that exchangeable calcium in soils and the
292 density of earthworms were significantly higher in forests dominated by *Acer*
293 *pseudoplatanus* and *Tilia cordata* compared to forests dominated by *Larix* and *Pinus*,
294 even though calcium concentrations were ~40% lower in the litter of *A. pseudoplatanus*
295 and *T. cordata* than in *C. japonica*. This implies that the change in soil systems is
296 caused by the alteration of forest vegetation with low calcium content relative to *C.*
297 *japonica*.

298 Soil crustaceans, such as Talitridae and *L. japonicum*, are powerful litter
299 decomposers (Zimmer 2002). Their existence influences litter decomposition rates
300 (O'Hanlon and Bolger 1999), and their abundance can lead to increased turnover rates
301 of organic matter. Furthermore, soil crustaceans affect the dynamics of organic matter
302 by incorporating organic material from the forest floor into deeper soil horizons
303 (Mattson 2012) and may also enhance microbial biomass (Escher et al. 2000). Reich et
304 al. (2005) showed that plantations of tree species with high calcium concentrations in

305 their litter cause an increase in exchangeable soil calcium and earthworm densities.
306 Therefore, especially in calcium-poor environments, transformation of forest vegetation
307 might change the densities of key decomposers such as crustaceans and earthworms,
308 thereby affecting the decomposition rates of soil organic matter and nutrient dynamics.

309

310 Acknowledgments

311 We thank staff members of Wakayama Experimental Forest, Hokkaido University, for
312 their support during the study. This study was partly supported by Grant-in-Aid from
313 JSPS (12J07244 to TO and 2566011103, 25281053 to TH) and from the Ministry of
314 Environment (S-9-3 to TH).

315

316

317

318

319

320

321 References

- 322 Alstad, N.E.W., Skardal, L., Hessen, D.O., 1999. The effect of calcium concentration on
323 the calcification of *Daphnia magna*. Limnol. Oceanogr. 44, 2011-2017.
- 324 Aoki, J., 1990. Soil Animals of Japan: Manual with Keys and Illustrations (in Japanese).
325 Tokai University Press, Tokyo.
- 326 Baba, M., Kato, M., Sugiura, T., Kobayashi, H., 2004. Calcium accumulation alleviates
327 soil acidification in Japanese cedar (*Cryptomeria japonica*) stands. Soil Sci.
328 Plant Nutr. 50, 403-411.
- 329 Bardgett, R.D., Wardle, D.A., 2010. Aboveground-Belowground Linkages. Biotic
330 Interactions, Ecosystem Processes and Global Change. Oxford university
331 press, Oxford.
- 332 Berg, B., McClaugherty, C., 2003. Plant litter: Decomposition, Humus Formation and
333 Carbon Sequestration. Springer, Berlin.
- 334 Binkley, D. Fisher R.F., 2013. Ecology and Management of Forest Soils 4th ed.
335 Wiley-Blackwell, New York.
- 336 D'Amore, D.V., Hennon, P.E., Schaberg, P.G., Hawley, G.J., 2009. Adaptation to

337 exploit nitrate in surface soils predisposes yellow-cedar to climate-induced
338 decline while enhancing the survival of western redcedar: A new hypothesis.
339 Forest Ecol. Manag. 258, 2261-2268.

340 Dijkstra, F.A., Smits, M.M., 2002. Tree species effects on calcium cycling: The role of
341 calcium uptake in deep soils. Ecosystems 5, 385-398.

342 Driscoll, C.T., Lawrence, G.B., Bulger, A.J., Butler, T.J., Cronan, C.S., Eagar, C.,
343 Lambert, K.F., Likens, G.E., Stoddard, J.L., Weathers, K.C., 2001. Acidic
344 deposition in the northeastern United States: Sources and inputs, ecosystem
345 effects, and management strategies. Bioscience 51, 180-198.

346 Escher, N., Käch, B., Nentwig, W., 2000. Decomposition of transgenic *Bacillus*
347 *thuringiensis* maize by microorganisms and woodlice *Porcellio scaber*
348 (Crustacea: Isopoda). Basic Appl. Ecol. 1, 161-169.

349 Forestry Agency, 2011. 2011 Forestry Census. Forestry Agency, Tokyo.

350 Glötzner, J., Ziegler, A., 2000. Morphometric analysis of the calcium-transporting
351 sternal epithelial cells of the terrestrial isopods *Ligia oceanica*, *Ligidium*
352 *hypnorum*, and *Porcellio scaber* during molt. Arthropod Struct. Dev. 29,

353 241-257.

354 Greenaway, P., 1985. Calcium balance and molting in the crustacea. Biol. Rev. 60,

355 425-454.

356 Groffman, P.M., Fisk, M.C., 2011. Calcium constrains plant control over forest

357 ecosystem nitrogen cycling. Ecology 92, 2035-2042.

358 Groffman, P.M., Fisk, M.C., Driscoll, C.T., Likens, G.E., Fahey, T.J., Eagar, C., Pardo,

359 L.H., 2006. Calcium additions and microbial nitrogen cycle processes in a

360 northern hardwood forest. *Ecosystems* 9, 1289-1305.

361 Hågvar, S., Abrahamsen, G., 1990. Microarthropoda and Enchytraeidae (Oligochaeta)

362 in naturally lead-contaminated soil - a gradient study. Environ. Entomol. 19,

363 1263-1277.

364 Hessen, D.O., Alstad, N.E.W., Skardal, L., 2000. Ca limitation in *Daphnia magna*. J.

365 Plankton Res. 22, 553-568.

366 Hotopp, K.P., 2002. Land snails and soil calcium in central Appalachian Mountain

367 Forest. Southeast Nat. 1, 27-44.

368 Ikeda, H., Homma, K., Kubota, K., 2005. Biotic and abiotic factors affecting the
369 structures of ground invertebrate communities in Japanese cedar dominant
370 forests. *Eur. J. For. Res.* 8, 1-13.

371 Juice, S.M., Fahey, T.J., Siccama, T.G., Driscoll, C.T., Denny, E.G., Eagar, C., Cleavitt,
372 N.L., Minocha, R., Richardson, A.D., 2006. Response of sugar maple to
373 calcium addition to Northern Hardwood Forest. *Ecology* 87, 1267-1280.

374 Kaneko, N., Kofuji, R., 2000. Effects of soil pH gradient caused by stemflow
375 acidification on soil microarthropod community structure in a Japanese red
376 cedar plantation: An evaluation of ecological risk on decomposition. *J. For.*
377 *Res.* 5, 157-162.

378 Kihira, T., Yoshida, S., Hironishi, M., Miwa, H., Okamoto, K., Kondo, T., 2005.
379 Changes in the incidence of amyotrophic lateral sclerosis in Wakayama,
380 Japan. *Amyotroph. Lateral. Sc.* 6, 155-163.

381 Kiilsgaard, C.W., Greene, S.E., Stafford, S.G., 1987. Nutrient concentrations in litterfall
382 from some western conifers with special reference to calcium. *Plant Soil* 102,
383 223-227.

384 Kloeppel, B.D., Abrams, M.D., 1995. Ecophysiological attributes of the native *Acer*
 385 *saccharum* and the exotic *Acer platanoides* in urban oak forests in
 386 Pennsylvania, USA. *Tree Physiol.* 15, 739-746.

387 Likens, G.E., Bormann, F.H., 1970. Chemical analyses of plant tissue from the Hubbard
 388 Brook Ecosystem in New Hampshire. *Yale Univ. Sch. For. Bull.* 79, 1–25.

389 Likens, G.E., Driscoll, C.T., Buso, D.C., 1996. Long-term effects of acid rain: Response
 390 and recovery of a forest ecosystem. *Science* 272, 244-246.

391 Likens, G.E., Driscoll, C.T., Buso, D.C., Siccama, T.G., Johnson, C.E., Lovett, G.M.,
 392 Fahey, T.J., Reiners, W.A., Ryan, D.F., Martin, C.W., Bailey, S.W., 1998.
 393 The biogeochemistry of calcium at Hubbard Brook. *Biogeochemistry* 41,
 394 89-173.

395 Mattson, W.J., 2012. The Role of Arthropods in Forest Ecosystems (Proceedings in Life
 396 Sciences). Springer, Berlin.

397 McLaughlin, S.B., Wimmer, R., 1999. Tansley Review No. 104 - Calcium physiology
 398 and terrestrial ecosystem processes. *New Phytol.* 142, 373-417.

399 Minocha, R., Long, S., Thangavel, P., Minocha, S.C., Eagar, C., Driscoll, C.T., 2010.

400 Elevation dependent sensitivity of northern hardwoods to Ca addition at
 401 Hubbard Brook Experimental Forest, NH, USA. *Forest Ecol. Manag.* 260,
 402 2115-2124.

403 Morrison, I.K., 1985. Effect of crown position on foliar concentrations of 11 elements
 404 in *Acer Saccharum* and *Betula Alleghaniensis* trees on a till soil. *Can. J.*
 405 *Forest Res.* 15, 179-183.

406 Myrold, D.D., 1990. Effect of Acidic Deposition on Soil Organisms. In *Mechanisms of*
 407 *Forest Response to Acidic Deposition*. Lucir A.A. and Haines S.G. (eds.),
 408 Springer, New York, pp. 163-187.

409 Nezat, C.A., Blum, J.D., Driscoll, C.T., 2010. Patterns of Ca/Sr and Sr-87/Sr-86
 410 variation before and after a whole watershed CaSiO₃ addition at the Hubbard
 411 Brook Experimental Forest, USA. *Geochim. Cosmochim. Ac* 74, 3129-3142.

412 O'hanlon, R.P., Bolger, T., 1999. The importance of *Arcitalitrus dorrieni* (Hunt)
 413 (Crustacea: Amphipoda: Talitridae) in coniferous litter breakdown. *Appl. Soil*
 414 *Ecol.* 11, 29-33.

415 Ohta, T., Niwa, S., Hiura, T., 2014. Calcium concentration in leaf litter affects the

416 abundance and survival of crustaceans in streams draining warm-temperate
417 forests. *Freshw. Biol.* 59, 748-760.

418 Potter, K.M., Conkling, B. L., 2012. Forest health monitoring: 2008 national technical
419 report. Gen. Tech. Rep. SRS-158. Asheville, NC: U.S. Department of
420 Agriculture Forest Service, Southern Research Station. 179 p.

421 R Development Core Team, 2011. R: A Language and Environment for Statistical
422 Computing. R Foundation for Statistical Computing, Vienna, Austria.

423 Reich, P.B., Oleksyn, J., Modrzyński, J., Mrozinski, P., Hobbie, S.E., Eissenstat, D.M.,
424 Chorover, J., Chadwick, O.A., Hale, C.M., Tjoelker, M.G., 2005. Linking
425 litter calcium, earthworms and soil properties: a common garden test with 14
426 tree species. *Ecol. Lett.* 8, 811-818.

427 Sakata, T., 2000. The assay of nitrate nitrogen in the soil extract using the UV
428 absorbance (in Japanese). *Jpn. J. For. Environment.* 42, 53-55.

429 Skeldon, M.A., Vadeboncoeur, M.A., Hamburg, S.P., Slum, J.D., 2007. Terrestrial
430 gastropod responses to ecosystem-level calcium manipulation a northern
431 hardwood forest. *Can J. Zool.* 85, 994-1007.

432 Springett, J.A., Syers, J.K., 1984. Effect of pH and calcium content of soil on
433 earthworm cast production in the laboratory. *Soil Biol. Biochem.* 16,
434 185-189.

435 Swift, M.J., Heal, O.W., Anderson, J.M., 1979. *Decomposition in Terrestrial*
436 *Ecosystems*. Backwell, Oxford.

437 Tateishi, M., 1976. The Muro Group in the southwestern part of the Muro Belt of the
438 Shimanto Terrain (in Japanese). *J. Geol. Soc. Jpn.* 82, 395–407.

439 Tsutsumi, T., 1987. *Material Cycling in Forest* (in Japanese). University of Tokyo Press,
440 Tokyo.

441 Ueno, S., Kurosawa, Y., Sato, M., 1985. *The Coleoptera of Japan in color, vol. II* (in
442 Japanese). Hoikusha Press, Osaka.

443 Warby, R.A.F., Johnson, C.E., Driscoll, C.T., 2009. Continuing acidification of organic
444 soils across the north-eastern USA: 1984–2001. *Soil Sci. Soc. Am. J.* 73,
445 274–284.

446 Wardle, D.A., Yeates, G.W., Barker, G.M., Bonner, K.I., 2006. The influence of plant
447 litter diversity on decomposer abundance and diversity. *Soil. Biol. Biochem.*

448 38, 1052-1062.

449 Widden, P., Hsu, D., 1987. Competition between Trichoderma Species - Effects of

450 Temperature and Litter Type. Soil Biol. Biochem. 19, 89-93.

451 Xue, L., Luo, S., 2002. Seasonal changes in the nutrient concentrations of leaves and

452 leaf litter in a young Cryptomeria japonica stand. Scand. J. Forest Res. 17,

453 495-500.

454 Ziegler, A., Hagedorn, M., Ahearn, G.A., Carefoot, T.H., 2007. Calcium translocations

455 during the moulting cycle of the semiterrestrial isopod *Ligia hawaiiensis*

456 (Oniscidea, Crustacea). J. Comp. Physiol. B 177, 99-108.

457 Zimmer, M., 2002. Nutrition in terrestrial isopods (Isopoda: Oniscidea): an

458 evolutionary-ecological approach. Biological Reviews 77, 455-493.

459

460

461

462

463

Table 1 Soil properties (mean $\pm$ SE) at each plot. Significant differences between cedar plantations (cedar) and evergreen broad-leaved forests (evergreen) are denoted in the last column: ** $P < 0.01$, n.s. $P > 0.05$ (likelihood ratio tests).

Plot name	cedar1	cedar2	cedar3	evergreen1	evergreen2	evergreen3	Significant differences between 'cedar' and 'evergreen' plots
Mass of litter layer (g)	10.42 $\pm$ 0.527	11.04 $\pm$ 0.561	10.28 $\pm$ 1.560	11.70 $\pm$ 1.240	7.47 $\pm$ 0.569	7.80 $\pm$ 0.845	n.s.
Water content (g/g)	0.62 $\pm$ 0.025	0.63 $\pm$ 0.064	0.57 $\pm$ 0.057	0.53 $\pm$ 0.045	0.64 $\pm$ 0.023	0.60 $\pm$ 0.036	n.s.
pH	4.34 $\pm$ 0.166	4.81 $\pm$ 0.143	5.20 $\pm$ 0.217	4.26 $\pm$ 0.182	4.55 $\pm$ 0.226	3.58 $\pm$ 0.085	n.s. ($P = 0.08$)
Exchangeable calcium (mg/g)	1.81 $\pm$ 0.256	1.83 $\pm$ 0.349	2.21 $\pm$ 0.971	0.63 $\pm$ 0.183	1.06 $\pm$ 0.413	0.43 $\pm$ 0.112	**
Total carbon (mg/g)	154.03 $\pm$ 23.188	117.63 $\pm$ 33.999	174.65 $\pm$ 58.206	139.58 $\pm$ 28.232	257.24 $\pm$ 47.917	201.00 $\pm$ 30.483	n.s.
Total nitrogen (mg/g)	10.33 $\pm$ 1.823	6.76 $\pm$ 1.770	9.90 $\pm$ 3.079	9.90 $\pm$ 1.908	19.0 $\pm$ 3.641	14.15 $\pm$ 1.880	n.s.
Nitrate (mg/g)	0.39 $\pm$ 0.021	0.52 $\pm$ 0.059	0.58 $\pm$ 0.047	0.73 $\pm$ 0.099	1.01 $\pm$ 0.113	0.82 $\pm$ 0.081	**
C:N ratio	15.89 $\pm$ 1.990	17.00 $\pm$ 0.781	17.36 $\pm$ 1.140	14.27 $\pm$ 0.775	13.52 $\pm$ 0.638	14.13 $\pm$ 0.473	**

485 Table 2 Abundance (5 traps⁻¹ 12 days⁻¹; mean ± SE) of ground-dwelling
 486 macroinvertebrates sampled by pitfall traps. ‘cedar’ and ‘evergreen’ mean cedar
 487 plantation and evergreen broad-leaved forest.

	cedar	evergreen
Gastropoda	0.67 ± 0.67	0.00 ± 0.00
Oligochaeta		
Haplotaxida	0.33 ± 0.33	0.00 ± 0.00
Diplopoda		
Polydesmida		
Paradoxosomatidae	0.00 ± 0.00	3.33 ± 1.76
Xystodesmidae		
<i>Xystodesmus</i> sp.	0.33 ± 0.33	0.00 ± 0.00
Julida		
Julidae		
<i>Anaulaciulus</i> sp.	0.00 ± 0.00	0.33 ± 0.33
Arachnida		
Araneae	4.00 ± 0.58	4.00 ± 1.00
Crustacea		
Amphipoda		
Talitridae	4.33 ± 0.88	0.00 ± 0.00
Isopoda		
Ligiidae		
<i>Ligidium japonicum</i>	4.67 ± 0.88	0.00 ± 0.00
Armadillidae		
<i>Venezillo</i> sp.	0.67 ± 0.33	0.33 ± 0.33
Insecta		
Archaeognatha		
Machilidae	0.33 ± 0.33	0.33 ± 0.33
Orthoptera		
Rhaphidophoridae	0.00 ± 0.00	0.67 ± 0.33
Lepidoptera (larva)	2.33 ± 0.88	2.00 ± 0.00
Hymenoptera		
Formicidae		
<i>Pachycondyla chinensis</i>	0.67 ± 0.67	0.00 ± 0.00
<i>Aphaenogaster famelica</i>	0.33 ± 0.33	1.00 ± 1.00
<i>Paratrechina flavipes</i>	0.33 ± 0.33	0.67 ± 0.67
<i>Formica hayashi</i>	1.00 ± 0.58	0.33 ± 0.33
<i>Camponotus obscuripes</i>	0.00 ± 0.00	0.67 ± 0.67
Coleoptera (larva)	0.00 ± 0.00	0.67 ± 0.67
Coleoptera (adult)		
Geotrupidae		
<i>Phelotrupes laevistriatus</i>	0.33 ± 0.33	0.33 ± 0.33
<i>Phelotrupes auratus</i>	0.33 ± 0.33	0.00 ± 0.00
Scarabaeidae		
<i>Panelus parvulus</i>	0.00 ± 0.00	0.33 ± 0.33
<i>Onthophagus nitidus</i>	0.33 ± 0.33	0.67 ± 0.67
Staphylinidae		
<i>Bolitobius</i> sp.	0.33 ± 0.33	0.33 ± 0.33
Staphylininae	0.00 ± 0.00	0.33 ± 0.33
Carabidae		
<i>Carabus iwawakianus</i>	0.00 ± 0.00	0.67 ± 0.67
<i>Chlaenius costiger</i>	0.67 ± 0.67	0.00 ± 0.00
<i>Stomis prognathus</i>	0.00 ± 0.00	0.33 ± 0.33
<i>Pterostichus (Rhagadus)</i> sp.	0.33 ± 0.33	1.33 ± 0.88
<i>Rupa japonica</i>	1.00 ± 0.58	0.33 ± 0.33
<i>Synuchus picicolor</i>	0.00 ± 0.00	0.33 ± 0.33

489 Fig. 1 Concentration (mean $\pm$ SE) of exchangeable calcium in soil in each plot. White
490 and black bars indicate evergreen broad-leaved forests (evergreen) and cedar plantations
491 (cedar), respectively. Significant differences between vegetation types are denoted by
492 different letters (likelihood ratio tests, $P < 0.05$).

493

494 Fig. 2 Abundance (mean $\pm$ SE) of Talitridae (a) and *Ligidium japonicum* (b) in each plot
495 by the hand-sorting method. White and black bars indicate evergreen broad-leaved
496 forests (evergreen) and cedar plantations (cedar), respectively.

497

498 Fig. 3 Canonical correspondence analysis (CCA) ordination of soil invertebrate
499 community composition in 12 subplots by the pitfall trap method. Explanatory variables
500 selected by forward selection are shown as arrows: Ca, exchangeable calcium
501 concentration and Litter, mass of the litter layer. White and black symbols indicate
502 subplot scores (mean $\pm$ SE) of evergreen broad-leaved forests (evergreen) and cedar
503 plantations (cedar), respectively. Invertebrate taxa are abbreviated by alphabets: G,
504 Gastropoda; H, Haplotaxida; P, Paradoxosomatidae; Xs, *Xystodesmus* sp.; As,
505 *Anaulaciulus* sp.; A, Araneae; T, Talitridae; Lj, *Ligidium japonicum*; Vs *Venezillo* sp.;
506 L, Lepidoptera (larva); Pc, *Pachycondyla chinensis*; Af, *Aphaenogaster famelica*; Pf,
507 *Paratrechina flavipes*; Fh, *Formica hayashi*; Co, *Camponotus obscuripes*; C,
508 Coleoptera (larva); Pp, *Panelus parvulus*; On, *Onthophagus nitidus*; Pl, *Phelotrupes*
509 *laevistriatus*; Pa, *Phelotrupes auratus*; Bs, *Bolitobius* sp.; St, Staphylininae; Ci,
510 *Carabus iwawakianus*; Cc, *Chlaenius costiger*; Sp, *Stomis prognathous*; Ps,

511 *Pterostichus (Rhagadus) sp.*; Rj, *Rupa japonica*; Syp, *Synuchus picicolor*; M,

512 Machilidae; R, Rhaphidophoridae. Taxa indicated by boldface are crustaceans.

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

534

535

536

537

538

539

540

541

542