Title	Genetic divergence in populations of <i>Lutzomyia ayacuchensis</i> , a vector of Andean-type cutaneous leishmaniasis, in Ecuador and Peru
Author(s)	Kato, Hirotomo; Caceres, Abraham G.; Gomez, Eduardo A.; Mimori, Tatsuyuki; Uezato, Hiroshi; Hashiguchi, Yoshihisa
Citation	Acta Tropica, 141(Part A), 79-87 https://doi.org/10.1016/j.actatropica.2014.10.004
Issue Date	2015-01
Doc URL	http://hdl.handle.net/2115/58016
Type	article (author version)
File Information	Acta tropica v.141p.79.pdf

[Instructions for use](#)

Genetic divergence in populations of *Lutzomyia ayacuchensis*, a vector of Andean-type cutaneous leishmaniasis, in Ecuador and Peru

Hiroto Kato^{a*}, Abraham G. Cáceres^{b,c}, Eduardo A. Gomez^d, Tatsuyuki Mimori^e, Hiroshi Uezato^f, Yoshihisa Hashiguchi^{g,h,i}

^aLaboratory of Parasitology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Japan

^bDepartamento Académico de Microbiología Médica, Facultad de Medicina Humana, Universidad Nacional Mayor de San Marcos, Peru

^cLaboratorio de Entomología, Instituto Nacional de Salud, Peru

^dDepartamento de Medicina Tropical, Facultad de Medicina, Universidad Católica de Guayaquil, Ecuador

^eDepartment of Microbiology, Faculty of Life Sciences, Graduate School of Health Sciences, Kumamoto University, Japan

^fDepartment of Dermatology, Faculty of Medicine, University of the Ryukyus, Japan

^gDepartment of Parasitology, Kochi Medical School, Kochi University, Japan

^hPrometeo, Secretaria Nacional de Educación Superior, Ciencia, Tecnología e Innovación (SENESCYT), Ecuador

ⁱCentro de Biomedicina, Universidad Central del Ecuador, Ecuador

*Corresponding author at: Laboratory of Parasitology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, North 18 West 9, Kita-ku, Sapporo, Hokkaido, 060-0818, Japan.

Phone & Fax: +81-11-706-5196.

E-mail address: hkato@vetmed.hokudai.ac.jp (H. Kato).

Abstract

Haplotype and gene network analyses were performed on mitochondrial cytochrome oxidase I and cytochrome *b* gene sequences of *Lutzomyia* (*Lu.*) *ayacuchensis* populations from Andean areas of Ecuador and southern Peru where the sand fly species transmit *Leishmania* (*Leishmania*) *mexicana* and *Leishmania* (*Viannia*) *peruviana*, respectively, and populations from the northern Peruvian Andes, for which transmission of *Leishmania* by *Lu. ayacuchensis* has not been reported. The haplotype analyses showed higher intrapopulation genetic divergence in northern Peruvian Andes populations and less divergence in the southern Peru and Ecuador populations, suggesting that a population bottleneck occurred in the latter populations, but not in former ones. Importantly, both haplotype and phylogenetic analyses showed that populations from Ecuador consisted of clearly distinct clusters from southern Peru, and the two populations were separated from those of northern Peru.

Keywords: Phlebotomine sand fly; *Lutzomyia ayacuchensis*; cytochrome oxidase I; cytochrome *b*

1. Introduction

Phlebotomine sand flies are insects of the family Psychodidae in the order Diptera. Approximately 800 sand fly species have been recorded; of these, fewer than 10% have been confirmed as vector species of leishmaniasis. These are in two currently defined genera, *Phlebotomus* in the Old World and the genus *Lutzomyia* in the New World (Munstermann, 2004; Bates, 2007; Kato *et al.*, 2010; Alvar *et al.*, 2012). Only a restricted number of species support the development of specific *Leishmania* species and consequently transmit the parasites (Kato *et al.*, 2010; Ready, 2013). Therefore, surveillance of circulating sand flies is important for predicting the risk and expansion of the disease in endemic and surrounding areas. Sand flies are generally identified by morphologic characteristics; mainly internal structures such as the spermatheca, cibarium and pharynx in females, and terminal genitalia in males (Young and Duncan, 1994). Genetic information on sand flies is accumulating, and several genetic markers have been used to examine the systematics, relationships and evolution among sand fly species (Kato *et al.*, 2010). The molecular taxonomy of sand flies mostly supports the traditional morphological classification, and can be applied to the surveillance of circulating species as well as identification of the species responsible for the transmission of *Leishmania* parasites in given endemic areas (Aransay *et al.*, 1999; Beati *et al.*, 2004; Kato *et al.*, 2005, 2007, 2008; Terayama *et al.*, 2008; Kuwahara *et al.*, 2009; Fujita *et al.*, 2012).

In addition to the species differences, intraspecific population divergence caused by multiple environmental factors such as climate, distance, altitude, and geographic barriers is suggested to influence vector competence (Lanzaro *et al.*, 1993; Hamarsheh *et al.*, 2009; Ready 2013). Since the maternally inherited mitochondrial genes reflect the

evolutionary history more accurately because of their clonal inheritance, lack of recombination and higher mutation rate compared with nuclear DNA (Avice, 1994; Rokas *et al.*, 2003), these genes have been used to estimate the population structure of arthropod vectors, and their geographical variation among populations has been reported in sand flies (Esseghir *et al.*, 1997; Ishikawa *et al.*, 1999; Hodgkinson *et al.*, 2003; Hamarsheh *et al.*, 2007; Belen *et al.*, 2011; Florin *et al.*, 2011; Rocha *et al.*, 2011; Cohnstaedt *et al.*, 2012; Yamamoto *et al.*, 2013; Pech-May *et al.*, 2013).

Lutzomyia (Lu.) ayacuchensis is a unique sand fly species distributing mainly in the Andean highlands of Ecuador and Peru (Takaoka *et al.*, 1990; Caceres *et al.*, 2004; Kato *et al.*, 2005, 2008; Gomez *et al.*, 2014a,b). The species is a proven vector of *Leishmania (Leishmania) mexicana* in the Ecuadorian Andes (Takaoka *et al.*, 1990; Hashiguchi *et al.*, 1991; Kato *et al.*, 2005, 2008; Gomez *et al.*, 2014a,b), whereas the same species transmits *Leishmania (Viannia) peruviana* in Andean areas of southern Peru (Caceres *et al.*, 2004). The sand fly populations from Ecuador and Peru were indistinguishable by morphological observation and genomic analysis of the 18S rRNA genes and rRNA internal transcribed spacer (ITS) sequences (Kuwahara *et al.*, 2009). *Lutzomyia ayacuchensis* is also distributed in the northern Peruvian Andes where cutaneous leishmaniasis is endemic; however, no transmission of *Leishmania* parasites by sand fly species has been reported in these areas. In the present study, mitochondrial cytochrome oxidase I (COI) and cytochrome *b* (*cyt b*) genes were compared in *Lu. ayacuchensis* populations from Andean areas of Ecuador and northern and southern Peru to assess genetic divergence among populations with different vector competence.

2. Materials and methods

2.1. Sand fly collection

Sand flies were collected in 4 Andean areas of Ecuador; Huigra (1,200 m above sea level), Chanchan (1,500 m a.s.l.), and Alausi (2,300 m a.s.l.), the Province of Chimborazo; and Paute (2,750 m a.s.l.), Province of Azuay (Kato et al., 2008; Kuwahara et al., 2009); and 4 Andean areas of Peru; Higosniyocc, Province of Lucanas, and Saquihuacca (2,250 m a.s.l.), Province of Parinacochas, Department of Ayacucho; Zapote (340 m a.s.l.), Province of Lambayeque, Department of Lambayeque; Viza (1,750 m) and El Paraiso (1,400 m a.s.l.), Province of Cutervo, Department of Cajamarca; and La Perla (1,930 m a.s.l.), Province of Huancabamba, Department of Piura (Kato et al., 2008; Kuwahara et al., 2009; Fujita et al., 2012) (Fig.1). All flies were captured between 18:30 and 21:00 by protected human bait, between 18:00 and 22:00 with Shannon traps, and between 19:00 and the next morning at 6:00 by CDC light traps. The sand flies were identified based on the morphology of their spermathecae, measurements of wing veins, the ratio of the palpus length to antenna and the thorax color (Young and Duncan, 1994). These morphologically identified specimens were fixed in absolute ethanol and stored at room temperature for further molecular analysis.

2.2. DNA extraction

Ethanol-fixed sand flies were individually lysed in 50 µl of DNA extraction buffer [150 mM NaCl, 10 mM Tris-HCl (pH 8.0), 10 mM EDTA and 0.1 % sodium dodecyl sulfate (SDS)] with 100 µg/ml of proteinase K. The samples were incubated at 37°C overnight, heated at 95°C for 5 min, and then 0.5 µl portions were directly used as the

templates for PCR amplification.

PCR amplification and sequence analysis of Lutzomyia ayacuchensis cytochrome oxidase I and cytochrome b genes

The *Lu. ayacuchensis* COI gene fragment was amplified with universal COI primers (LCO1490: GGTCAACAAATCATAAAGATATTGG and HCO2198: TAAACTTCAGGGTGACCAAAAAATCA) (Folmer et al., 1994), and the *cyt b* gene fragment was amplified with primers prepared based on the *Lu. ayacuchensis cyt b* gene sequences (Lay cyt S: TGTCGAGATGTAACTATGG and Lay cyt R: TGCTATTTAAGCTTATTAAC) (Yamamoto et al., 2013). PCR amplification was carried out in a volume of 15 µl with the primers (0.4 µM each), Ampdirect Plus (Shimadzu Biotech, Tsukuba, Japan), and high fidelity DNA polymerase (KOD-Plus-ver.2; TOYOBO, Tokyo, Japan). After an initial denaturation at 95°C for 5 min, amplification was performed with 35 cycles of denaturation (95°C, 1 min), annealing (55°C, 1 min) and polymerization (72°C, 1 min), followed by a final extension at 72°C for 10 min. The PCR products were purified using a FastGene Gel/PCR Extraction kit (NIPPON Genetics, Tokyo, Japan) to remove excessive primers, and the sequences were directly determined with a forward primer by the dideoxy chain termination method using a BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA).

2.3. Data analysis

The sequences were aligned with CLUSTAL W software (Thompson et al., 1994) and examined using the MEGA program (Molecular Evolutionary Genetics Analysis)

version 5.2 (Tamura et al., 2011). The pairwise genetic distances between groups were analyzed by MEGA 5.2 using the Kimura two-parameter (Tamura et al., 2011). Phylogenetic analyses were performed by the Maximum Likelihood (ML) method with the distance algorithms available in the MEGA package (Tamura et al., 2011). The number of segregating sites, number of haplotypes, haplotype diversity, average number of differences, and nucleotide diversity for each population were calculated using DnaSP 5.0 (Rozas et al., 2003), and genetic diversity between populations was determined using MEGA 5.2 (Tamura et al., 2011). A haplotype network was constructed using the median-joining methods as implemented in the program NETWORK 4.6.1.2 (<http://www.fluxus-engineering.com/sharenet.htm>) (Bandelt et al., 1999).

3. Results

3.1. Haplotype analysis of the *Lutzomyia ayacuchensis* COI gene

COI gene sequences of 215 *Lu. ayacuchensis* collected from 4 Ecuadorian Andes sites (Huigra, Chanchan, Alausi, and Paute) and 4 Peruvian Andes sites (Ayacucho, Lambayeque, Cajamarca, and Piura) were determined by direct sequencing, and the sequences of 628bp-fragments were aligned. The sequence analysis identified the presence of 103 haplotypes with 92 segregating sites (Fig. 2). No more than two different nucleotides were identified at each segregating site except for three (nucleotide positions at 310, 349 and 607), and neither nonsense mutations nor insertions/deletions were observed (Fig. 2). The number of segregating sites (S), number of haplotypes (H), haplotype diversity (Hd), average number of differences (K), and nucleotide diversity (Pi) were calculated for each population (Table 1). Haplotype analysis showed that 48 flies from Huigra, 23 from Chanchan, and 21 from Alausi, Province of Chimborazo, belonged to 14, 7, and 7 haplotypes, respectively (Fig. 2, Table 1). On the other hand, 14 flies from Paute, Province of Azuay, consisted of 6 haplotypes (Fig. 2, Table 1). Interestingly, dominant haplotypes, Hap1 and Hap2, were noted in all these areas of Ecuador where *Lu. ayacuchensis* transmits *L. (L.) mexicana* (Fig. 2 and 3). In Ayacucho of Peru, where *Lu. ayacuchensis* is a proven vector of *L. (V.) peruviana*, 45 sand flies composed 28 haplotypes, of which a haplotype, Hap27, was relatively dominant, showed distinct haplotypes from those of endemic areas in Ecuador (Fig. 2 and 3). In other Peruvian areas where *Leishmania* species transmitted by *Lu. ayacuchensis* have not been reported, 16 sand flies from Lambayeque, 16 flies from Cajamarca, and 32 flies from Piura belonged to 16, 12, and 24 haplotypes, respectively, each including less than 4 individuals in the absence of an apparently dominant haplotype (Fig. 2 and 3).

All these haplotypes of *Lu. ayacuchensis* were distinct from those of the *L. (L.) mexicana*-prevalent areas of Ecuador (Huigra, Chanchan, Alausi, and Paute) and *L. (V.) peruviana*-prevalent areas of Ayacucho (Fig. 2 and 3). Statistical analyses supported a higher genetic diversity in populations of Lambayeque ($K=12.758$, $P_i=0.020$), Cajamarca ($K=3.308$, $P_i=0.005$) and Piura ($K=11.921$, $P_i=0.019$), when compared to populations of Ecuador ($K=0.909-1.202$, $P_i=0.001-0.002$) and Ayacucho of Peru ($K=1.538$, $P_i=0.002$) (Table 1). Intrapopulation genetic divergence was lower in the areas of Ecuador (Huigra, Chanchan, Alausi, and Paute: 0.1-0.2%) and Ayacucho of Peru (0.4%), when compared to the other Peruvian areas, Lambayeque, Cajamarca, and Piura (0.5-2.1%) (Table 2). The genetic divergence between populations was lower between Ecuadorian populations (Huigra, Chanchan, Alausi, and Paute) (0.2-0.3%), and the highest between Ayacucho and Lambayeque (5.3%) (Table 3). The genetic divergence between *Lu. ayacuchensis* populations and *Lu. hartmanni* from Huigra, both of which belong to the subgenus *Helcocyrtomyia*, was 19.0-20.4%. Phylogenetic analysis showed that *Lu. ayacuchensis* from the Ecuadorian Andes (Huigra, Chanchan, Alausi, and Paute) and from Ayacucho of Peru where the species transmits *L. (L.) mexicana* and *L. (V.) peruviana*, respectively, consisted of clearly distinct clusters, which were separated from those of Lambayeque, Cajamarca, and Piura of Peru, where transmission of *Leishmania* by this species has not been reported (Fig. 4).

3.2. Haplotype analysis of the *Lutzomyia ayacuchensis* cyt b gene

The *cyt b* gene sequences of 216 *Lu. ayacuchensis* collected from 4 Ecuadorian sites and 4 Peruvian Andes sites were determined by direct sequencing, and the sequences of 596bp-fragments were aligned. The sequence analysis identified the presence of 97

haplotypes with 136 segregating sites, and neither nonsense mutations nor insertions/deletions were observed (Fig. 5). Haplotype analysis showed that 48 flies from Huigra, 23 from Chanchan, and 24 from Alausi, Province of Chimborazo, belonged to 13, 4, and 7 haplotypes, respectively, whereas 13 flies from Paute, Province of Azuay, consisted of 3 haplotypes (Fig. 5, Table 4). A dominant haplotype, Hap1, was noted in all these areas of Ecuador (Fig. 5). In Ayacucho in Peru, 46 sand flies had 20 haplotypes, of which a haplotype Hap23 was dominant, and showed distinct haplotypes from those of endemic areas in Ecuador (Fig. 5 and 6). In other Peruvian areas (Lambayeque, Cajamarca, and Piura), 16, 12, and 34 sand flies belonged to 15, 9, and 33 haplotypes, respectively, each including less than 3 individuals in the absence of an apparently dominant haplotype (Fig. 5 and 6). All these haplotypes were distinct from those of Ecuador (Huigra, Chanchan, Alausi, and Paute) and Ayacucho of Peru where *Lu. ayacuchensis* transmits *Leishmania* species (Fig. 5 and 6). Statistical analyses supported a higher genetic diversity in populations of Lambayeque ($K=9.650$, $P_i=0.016$), Cajamarca ($K=5.379$, $P_i=0.009$) and Piura ($K=13.865$, $P_i=0.023$), when compared to populations of Ecuador ($K=0.498-1.051$, $P_i=0.001-0.002$) and Ayacucho of Peru ($K=1.569$, $P_i=0.003$) (Table 4). Intrapopulation genetic divergence was lower in the areas of Ecuador (Huigra, Chanchan, Alausi, and Paute: 0.1%) and Ayacucho of Peru (0.2%), when compared to other areas, Lambayeque, Cajamarca, and Piura (0.7-1.9%) (Table 2). The genetic divergence between populations was lower between Ecuadorian populations (Huigra, Chanchan, Alausi, and Paute) (0.1%), and the highest between Paute and Piura (4.5%) (Table 3). Similar to the results obtained from phylogenetic analysis of the COI gene, *Lu. ayacuchensis* from the Ecuadorian Andes (Huigra, Chanchan, Alausi, and Paute) and from Ayacucho of Peru made up distinct clusters,

which were isolated from those of Lambayeque, Cajamarca, and Piura of Peru (Fig. 7).

4. Discussion

It has been suggested that genetic divergence caused by genetic drift and/or selection may affect the vectorial capacity of sand flies, as well as other arthropods (Lanzaro *et al.*, 1993). In the present study, mitochondrial COI and *cyt b* genes were comparatively analyzed in *Lu. ayacuchensis* populations from Andean areas of Ecuador (Huigra, Chanchan, Alausi, and Paute) and southern Peru (Ayacucho), where sand fly species transmit *L. (L.) mexicana* and *L. (V.) peruviana*, respectively, and those from the northern Peruvian Andes (Lambayeque, Cajamarca, and Piura), in which transmission of *Leishmania* by *Lu. ayacuchensis* has not been reported. The results showed apparent genetic divergence among populations with different vector competence.

Lutzomyia ayacuchensis is a rare species with regard to vector competence; this species is a proven vector of *L. (L.) mexicana* in the Ecuadorian Andes whereas it transmits another subgenus species, *L. (V.) peruviana* in southern Peru (Takaoka *et al.*, 1990; Hashiguchi *et al.*, 1991; Caceres *et al.*, 2004; Kato *et al.*, 2005, 2008; Gomez *et al.*, 2014a,b). Initially, we suspected that sand flies from Ecuador and Peru may be closely related, but not identical species; however, they were indistinguishable after comparative morphologic classification and genomic analysis (Caceres, unpublished; Kuwahara *et al.*, 2009). These observations, along with a restricted range of sand fly activity through their development stages (Alexander, 1987; Alexander and Young, 1992; Morrison *et al.*, 1993) encouraged the analysis of the genetic divergence among populations. Haplotype analyses targeting the COI and *cyt b* genes revealed higher intrapopulation genetic divergence in Lambayeque, Cajamarca and Piura of Peru, when compared to populations of Ecuador and Ayacucho of Peru. Particularly, 4 Ecuadorian populations (Huigra, Chanchan, Alausi, and Paute) showed extremely low genetic

divergence despite their geographical isolation (e.g. more than 60 km between Alausi and Paute). In general, higher genetic diversity within a geographical location is found in populations that have not undergone drastic size reductions over evolutionary time, whereas the effect of genetic drift is reduced when the population size is expanding. Therefore, the data suggest that a population bottleneck may have occurred in Ayacucho in Peru and especially, in Ecuador (Huigra, Chanchan, Alausi, and Paute), but not in Lambayeque, Cajamarca and Piura. Since sand flies are typically active at night, they may need to adapt to the severe climate in the Andean highlands where active time is limited due to a temperature drop after dark. Less genetic divergence may have some relation to the higher ratio of *L. (L.) mexicana* infection in Ecuador populations.

In the phylogenetic and haplotype analyses of Peruvian specimens, the Ayacucho population was clearly separated from the populations of Lambayeque, Cajamarca and Piura. This may simply reflect the geographic distance between Ayacucho and other Peruvian areas studied; however, it may not be so because the genetic divergence in Ecuadorian populations was comparable in Peruvian populations regardless of the geographic distance. The sample collection sites of Ayacucho and Lambayeque are located on the western Andean slopes at altitudes of 2,250 m a.s.l. and 340 m a.s.l., respectively, and those of Piura and Cajamarca are located on eastern Andean slopes at altitudes of 1,930 m a.s.l. and 1,400-1,750 m a.s.l., respectively. The climate is different between the slopes; it is very dry at the western Andean slope sites regardless of the altitude and becomes cold after dark at highland areas of Ayacucho, whereas eastern Andean slope areas have a subtropical climate and are humid. In a recent study, *Lu. verrucarum* from each valley of Peru was reported to have unique genotypes with genetic divergence of 0.2-1.0% that was not shared with sand flies from other valleys or

from more distant regions (genetic divergence of 1.6-3.1%) (Cohnstaedt *et al.*, 2012). The genetic divergence rates of the *Lu. ayacuchensis* *cyt b* gene between western slope sites, Ayacucho and Lambayeque, and between eastern slope areas, Cajamarca and Piura, were 3.8% and 2.2%, respectively, which were comparable between western and eastern slope areas (1.6-4.4%). The results indicated that slope-specific variation was not observed in *Lu. ayacuchensis* of Peru, which differed from *Lu. verrucarum*. This may partly reflect higher intrapopulation genetic variation in *Lu. ayacuchensis* of Peru, especially in Lambayeque and Piura (1.3-1.9%).

The present study clearly demonstrated apparent genetic divergence among populations of *Lu. ayacuchensis* with different vector competence. Although laboratory studies do not always reflect the natural conditions, experimental infection of *Lu. ayacuchensis* from Ecuador populations by *L. (V.) peruviana* or those from Ayacucho population by *L. (L.) mexicana* may help to understand the vectorial capacity of these sand flies.

Funding

This study was financially supported by the Ministry of Education, Culture and Sports, Science and Technology (MEXT) of Japan (Grant Nos. 23256002 and 25257501), and the Prometeo Project of the Secretaria Nacional de Educacion Superior, Ciencia, Tecnologia e Innovacion (SENESCYT), Ecuador.

Conflict of interest

The authors have no conflicts of interest to declare.

Acknowledgements

We are indebted to Flavio-Valeriano Zambrano C. (Servicio Nacional de Erradicacion de la Malaria, Guayaquil, Ecuador), Kazue Hashiguchi (Centro de Biomedicina, Universidad Central del Ecuador, Quito, Ecuador), and Roberto Sud A. (Ministerio de Salud Publica y Asistencia Social, Guayaquil, Ecuador) for their technical assistance during the field phase of the present study. We would like to thank the following person for their invaluable support in the collection of *Lu. ayacuchensis* and other sand fly species at different localities in Peru: Arrunátegui-Jimenez, J. M. (Gerencia Regional de Salud Lambayeque), Holgín-Mauricci, C. (Dirección Regional de Salud de Piura), Luna-Caipo, V. D. (Dirección Regional de Salud de Cajamarca), Lañas-Rosas, M. F. (Dirección Regional de Salud de Piura) and Giovana De La Cruz-Vásquez (Instituto Nacional de Salud, Lima).

References

- Alexander, J.B., 1987. Dispersal of *phlebotomine* sand flies (Diptera: Psychodidae) in a Colombian coffee plantation. J. Med. Entomol. 24, 552-558.
- Alexander, B., Young, D.G., 1992. Dispersal of phlebotomine sand flies (Diptera: Psychodidae) in a Colombian focus of *Leishmania (Viannia) braziliensis*. Mem. Inst. Oswaldo Cruz. 87, 397-403.
- Alvar, J., Vélez, I. D., Bern, C., Herrero, M., Desjeux, P., Cano, J., Jannin, J., den Boer, M.; WHO Leishmaniasis Control Team., 2012. Leishmaniasis worldwide and global estimates of its incidence. PLoS One. 7, e35671.
- Aransay, A.M., Scoulica, E., Chaniotis, B., Tselentis, Y., 1999. Typing of sand flies from Greece and Cyprus by DNA polymorphism of 18S rRNA gene. Insect Mol. Biol. 8, 179-184.
- Awise, J.C., Bowen, B.W., 1994. Investigating sea turtle migration using DNA markers. Curr. Opin. Genet. Dev. 4, 882-886.
- Bandelt, H.J., Forster, P., Röhl, A., 1999. Median-joining networks for inferring intraspecific phylogenies. Mol. Biol. Evol. 16, 37-48.
- Bates, P.A., 2007. Transmission of *Leishmania* metacyclic promastigotes by phlebotomine sand flies. Int. J. Parasitol. 37, 1097-1106.
- Beati, L., Caceres, A.G., Lee, J.A., Munstermann, L.E., 2004. Systematic relationships among *Lutzomyia* sand flies (Diptera: Psychodidae) of Peru and Colombia based on the analysis of 12S and 28S ribosomal DNA sequences. Int. J. Parasitol. 34, 225-234.

- Belen, A., Kucukyildirim, S., Alten, B., 2011. Genetic structures of sand fly (Diptera: Psychodidae) populations in a leishmaniasis endemic region of Turkey. *J. Vector Ecol.* 1, 32-48.
- Caceres, A.G., Villaseca, P., Dujardin, J.C., Bañuls, A.L., Inga, R., Lopez, M., Arana, M., Le Ray, D., Arevalo, J., 2004. Epidemiology of Andean cutaneous leishmaniasis: incrimination of *Lutzomyia ayacuchensis* (Diptera: psychodidae) as a vector of *Leishmania* in geographically isolated, upland valleys of Peru. *Am. J. Trop. Med. Hyg.* 70, 607-612.
- Cohnstaedt, L.W., Caceres, A.G., Beati, L., Munstermann, L.E., 2012. The population structure of *Lutzomyia verrucarum* (Diptera: Psycodidae), a *Bartonella bacilliformis* and *Leishmania peruviana* vector in Peru. *J. Med. Entomol.* 49, 77-84.
- Esseghir, S., Ready, P.D., Killick-Kendrick, R., Ben-Ismail, R., 1997. Mitochondrial haplotypes and phylogeography of *Phlebotomus* vectors of *Leishmania major*. *Insect Mol. Biol.* 6, 211-225.
- Florin, D.A., Davies, S.J., Olsen, C., Lawyer, P., Lipnick, R., Schultz, G., Rowton, E., Wilkerson, R., Keep, L., 2011. Morphometric and molecular analyses of the sand fly species *Lutzomyia shannoni* (Diptera: Psychodidae: Phlebotominae) collected from seven different geographical areas in the southeastern United States. *J. Med. Entomol.* 46, 154-166.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* 3, 294-299.
- Fujita, M., Kato, H., Caceres, A.G., Gomez, E.A., Velez, L., Mimori, T., Zhang, F., Iwata, H., Korenaga, M., Sakurai, T., Katakura, K., Hashiguchi, Y., 2012.

- Genotyping of sand fly species in Peruvian Andes where leishmaniasis is endemic. *Acta Trop.* 121, 93-98.
- Gomez, E.A., Kato, H., Mimori, T., Hashiguchi, Y., 2014a. Distribution of *Lutzomyia ayacuchensis*, the vector of Andean-type cutaneous leishmaniasis, at different altitudes on the Andean slope of Ecuador. *Acta Trop.* 137, 118-122.
- Gomez, E.A., Kato, H., Hashiguchi, Y., 2014b. Man-biting sand fly species and natural infection with the *Leishmania* promastigote in leishmaniasis-endemic areas of Ecuador. *Acta Trop.* in press
- Hamarsheh, O., Presber, W., Abdeen, Z., Sawalha, S., Al-Lahem, A., Schonian, G., 2007. Genetic structure of Mediterranean populations of the sandfly *Phlebotomus papatasi* by mitochondrial cytochrome *b* haplotype analysis. *Med. Vet. Entomol.* 21, 270-277.
- Hamarsheh, O., Presber, W., Al-Jawabreh, A., Abdeen, Z., Amro, A., Schonian, G., 2009. Molecular markers for *Phlebotomus papatasi* (Diptera: Psychodidae) and their usefulness for population genetic analysis. *Trans. R. Soc. Trop. Med. Hyg.* 103, 1085-1086.
- Hashiguchi, Y., Gomez, E.A., de Coronel, V.V., Mimori, T., Kawabata, M., Furuya, M., Nonaka, S., Takaoka, H., Alexander, J.B., Quizhpe, A.M., Grimaldi, G. Jr., Kreutzer, R.D., Tesh, R.B., 1991. Andean leishmaniasis in Ecuador caused by infection with *Leishmania mexicana* and *L. major*-like parasites. *Am. J. Trop. Med. Hyg.* 44, 205-217.
- Hodgkinson, V.H., Birungi, J., Quintana, M., Dietze, R., Munstermann, L.E., 2003. Mitochondrial cytochrome *b* variation in populations of the visceral leishmaniasis vector *Lutzomyia longipalpis* across eastern Brazil. *Am. J. Trop. Med. Hyg.* 69,

386-392.

- Ishikawa, E.A., Ready, P.D., de Souza, A.A., Day, J.C., Rangel, E.F., Davies, C.R., Shaw, J.J., 1999. A mitochondrial DNA phylogeny indicates close relationships between populations of *Lutzomyia whitmani* (Diptera: Psychodidae, Phlebotominae) from the rain-forest regions of Amazonia and northeast Brazil. Mem. Inst. Oswaldo Cruz. 94, 339-345
- Kato, H., Uezato, H., Katakura, K., Calvopiña, M., Marco, J.D., Barroso, P.A., Gomez, E.A., Mimori, T., Korenaga, M., Iwata, H., Nonaka, S., Hashiguchi, Y., 2005. Detection and identification of *Leishmania* species within naturally infected sand flies in the Andean areas of Ecuador by a polymerase chain reaction. Am. J. Trop. Med. Hyg. 72, 87-93.
- Kato, H., Cáceres, A.G., Gomez, E.A., Mimori, T., Uezato, H., Marco, J.D., Barroso, P.A., Iwata, H., Hashiguchi, Y., 2008. Molecular mass screening to incriminate sand fly vectors of Andean-type cutaneous leishmaniasis in Ecuador and Peru. Am. J. Trop. Med. Hyg. 79, 719-721.
- Kato, H., Gomez, E.A., Cáceres, A.G., Uezato, H., Mimori, T., Hashiguchi, Y., 2010. Molecular epidemiology for vector research on leishmaniasis. Int. J. Environ. Res. Public Health. 7, 814-826.
- Kato, H., Uezato, H., Gomez, E.A., Terayama, Y., Calvopiña, M., Iwata, H., Hashiguchi, Y., 2007. Establishment of a mass screening method of sand fly vectors for *Leishmania* infection by molecular biological methods. Am. J. Trop. Med. Hyg. 77, 324-329.
- Kuwahara, K., Kato, H., Gomez, E.A., Uezato, H., Mimori, T., Yamamoto, Y., Calvopina, M., Caceres, A.G., Iwata, H., Hashiguchi, Y., 2009. Genetic diversity of

- ribosomal RNA internal transcribed spacer sequences in *Lutzomyia* species from areas endemic for New World cutaneous leishmaniasis. *Acta Trop.* 112, 131-136.
- Lanzaro, G.C., Ostrovska, K., Herrero, M.V., Lawyer, P.G., Warburg, A., 1993. *Lutzomyia longipalpis* is a species complex: genetic divergence and interspecific hybrid sterility among three populations. *Am. J. Trop. Med. Hyg.* 48, 839-847.
- Morrison, A.C., Ferro, C., Morales, A., Tesh, R.B., Wilson, M.L., 1993. Dispersal of the sand fly *Lutzomyia longipalpis* (Diptera: Psychodidae) at an endemic focus of visceral leishmaniasis in Colombia. *J. Med. Entomol.* 30, 427-435.
- Munstermann, L.E., 2004. Phlebotomine sand flies, the Psychodidae. In: Marquardt, W.C., Black, W.C., Freier, J.E., Hagedorn, H.H., Hemingway, J., Higgs, S., James, A.A., Kondratieff, B., Moore, C.G. (Eds.), *Biology of Disease Vectors*, 2nd ed. Elsevier, San Diego, pp.141-151.
- Pech-May, A., Marina, C.F., Vázquez-Domínguez, E., Berzunza-Cruz, M., Rebollar-Téllez, E.A., Narváez-Zapata, J.A., Moo-Llanes, D., Ibáñez-Bernal, S., Ramsey, J.M., Becker, I., 2013. Genetic structure and divergence in populations of *Lutzomyia cruciata*, a phlebotomine sand fly (Diptera: Psychodidae) vector of *Leishmania mexicana* in southeastern Mexico. *Infect. Genet. Evol.* 16, 254-262.
- Ready, P.D., 2013. Biology of phlebotomine sand flies as vectors of disease agents. *Annu. Rev. Entomol.* 58, 227-250.
- Rocha, L.de S., Falqueto, A., Dos Santos, C.B., Grimaldi, G.J., Cupolillo, E., 2011. Possible implication of the genetic composition of the *Lutzomyia longipalpis* (Diptera: Psychodidae) populations in the epidemiology of the visceral leishmaniasis. *J. Med. Entomol.* 48, 1016-1022.

- Rokas, A., Melika, G., Abe, Y., Nieves-Aldrey, J.L., Cook, J.M., Stone, G.N., 2003. Lifecycle closure, lineage sorting, and hybridization revealed in a phylogenetic analysis of European oak gallwasps (Hymenoptera: Cynipidae: Cynipini) using mitochondrial sequence data. *Mol. Phylogenet. Evol.* 26, 36-45.
- Rozas, J., Sanchez-DelBarrio, J.C., Messeguer, X., Rozas, R., 2003. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics.* 19, 2496-2497.
- Takaoka, H., Gomez, E.A., Alexander, J.B., Hashiguchi, Y., 1990. Natural infections with *Leishmania* promastigotes in *Lutzomyia ayacuchensis* (Diptera: Psychodidae) in an Andean focus of Ecuador. *J. Med. Entomol.* 27, 701-702.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28: 2731-2739.
- Terayama, Y., Kato, H., Gomez, E.A., Uezato, H., Calvopiña, M., Iwata, H., Hashiguchi, Y., 2008. Molecular typing of sand fly species (Diptera, Psychodidae, Phlebotominae) from areas endemic for leishmaniasis in Ecuador by PCR-RFLP of 18S ribosomal RNA gene. *J. Vet. Med. Sci.* 70, 907-913.
- Thompson, J.D., Higgins, D.G., Gibson, T.J., 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22, 4673-4680.
- Yamamoto, K., Cáceres, A.G., Gomez, E.A., Mimori, T., Iwata, H., Korenaga, M., Sakurai, T., Katakura, K., Hashiguchi, Y., Kato, H., 2013. Genetic diversity of the mitochondrial cytochrome *b* gene in *Lutzomyia* spp., with special reference to

Lutzomyia peruensis, a main vector of *Leishmania (Viannia) peruviana* in the Peruvian Andes. Acta Trop. 126, 156-163.

Young, D.G., Duncan, M.A., 1994. Guide to the Identification and Geographic Distribution of *Lutzomyia* Sand Flies in Mexico, the West Indies, Central and South America (Diptera: Psychodidae), Memoirs of the American Entomological Institute, vol. 54, Associated Publishers—American Entomological Institute, Gainesville, FL.

Figure Legends

Fig.1. (A) Map of Ecuador and Peru showing the geographic locations where *Lutzomyia ayacuchensis* were collected. HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute; PI, Piura; CA, Cajamarca; LA, Lambayeque; AY, Ayacucho.

Fig.2. Variable nucleotides found in the alignment of the *Lutzomyia ayacuchensis* cytochrome oxidase I gene. The cytochrome oxidase I (COI) gene sequence of a 628bp-fragment was analyzed in 215 *Lutzomyia (Lu.) ayacuchensis* collected from 4 Ecuadorian (Huigra, Chanchan, Alausi, and Paute) and 4 Peruvian Andes regions (Ayacucho, Lambayeque, Cajamarca, and Piura). Dots denote identical sequences and numbers show their corresponding positions from the *Lu. ayacuchensis* COI gene fragment obtained in this study. HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute; PI, Piura; CA, Cajamarca; LA, Lambayeque; AY, Ayacucho.

Fig.3. Haplotype network of the cytochrome oxidase I sequences of *Lutzomyia ayacuchensis* collected from 4 Ecuadorian (Huigra, Chanchan, Alausi, and Paute) and 4 Peruvian Andes regions (Ayacucho, Lambayeque, Cajamarca, and Piura). Each haplotype is represented by a circle sized in proportion to the frequency of the haplotypes. Each crossbar represents one nucleotide substitution. Small black circles indicate one nucleotide substitution between haplotype. HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute; PI, Piura; CA, Cajamarca; LA, Lambayeque; AY, Ayacucho.

Fig.4. Phylogenetic tree of the cytochrome oxidase I sequences among *Lutzomyia ayacuchensis* populations. The scale bar represents 0.005% divergence. HU, Huigra;

CH, Chanchan; AL, Alausi; PA, Paute; PI, Piura; CA, Cajamarca; LA, Lambayeque; AY, Ayacucho.

Fig.5. Variable nucleotides found in the alignment of the *Lutzomyia ayacuchensis* cytochrome *b* gene. The cytochrome *b* (*cyt b*) gene sequence of a 596bp-fragment was analyzed in 216 *Lutzomyia (Lu.) ayacuchensis* collected from 4 Ecuadorian (Huigra, Chanchan, Alausi, and Paute) and 4 Peruvian Andes regions (Ayacucho, Lambayeque, Cajamarca, and Piura). Dots denote identical sequences and numbers show their corresponding positions from the *Lu. ayacuchensis cyt b* gene fragment obtained in this study. HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute; PI, Piura; CA, Cajamarca; LA, Lambayeque; AY, Ayacucho.

Fig.6. Haplotype network of the cytochrome *b* sequences of *Lutzomyia ayacuchensis* collected from 4 Ecuadorian (Huigra, Chanchan, Alausi, and Paute) and 4 Peruvian Andes regions (Ayacucho, Lambayeque, Cajamarca, and Piura). Each haplotype is represented by a circle sized in proportion to the frequency of the haplotypes. Each crossbar represents one nucleotide substitution. Small black circles indicate one nucleotide substitution between haplotype. HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute; PI, Piura; CA, Cajamarca; LA, Lambayeque; AY, Ayacucho.

Fig.7. Phylogenetic tree of the cytochrome *b* sequences among *Lutzomyia ayacuchensis* populations. The scale bar represents 0.005% divergence. HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute; PI, Piura; CA, Cajamarca; LA, Lambayeque; AY, Ayacucho.

Table 1.
Summary statistics for the COI gene from *Lutzomyia ayacuchensis* populations

population ^a	N	S	H	Hd	K	Pi
HU	48	11	14	0.778	1.202	0.002
CH	23	7	7	0.522	0.909	0.001
AL	21	7	7	0.767	1.200	0.002
PA	14	6	6	0.791	1.538	0.002
AY	45	28	30	0.956	2.644	0.004
LA	16	42	16	1.000	12.758	0.020
CA	16	12	12	0.958	3.308	0.005
PI	32	56	24	0.974	11.921	0.019
Total	215	92	103	0.933	18.511	0.029

^apopulation: HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute; AY, Ayacucho;
LA, Lambayeque; CA, Cajamarca; PI, Piura

N, number of sequences; S, number of segregating sites; H, number of haplotypes;
Hd, haplotype diversity; K, average number of differences; Pi, nucleotide diversity
The number of segregating sites, number of haplotypes, haplotype diversity,
average number of differences, and nucleotide diversity for each population were
calculated using DnaSP 5.0 (Rozas et al., 2003).

Table 2.

Genetic divergence of *Lutzomyia ayacuchensis* within populations

	HU ^a	CH	AL	PA	AY	LA	CA	PI
COI	0.002	0.001	0.002	0.002	0.004	0.021	0.005	0.019
<i>cyt b</i>	0.001	0.001	0.001	0.001	0.002	0.013	0.007	0.019

^apopulation: HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute;

AY, Ayacucho; LA, Lambayeque; CA, Cajamarca; PI, Piura

Genetic divergence was determined using MEGA 5.2 (Tamura et al., 2011).

Table 3.

Genetic divergence of *Lutzomyia ayacuchensis* between populations

		cyt b							
		HU ^a	CH	AL	PA	AY	LA	CA	PI
COI	HU		0.001	0.001	0.001	0.044	0.040	0.037	0.044
	CH	0.002		0.001	0.001	0.044	0.040	0.037	0.044
	AL	0.002	0.002		0.001	0.044	0.040	0.037	0.044
	PA	0.002	0.002	0.003		0.044	0.040	0.036	0.045
	AY	0.035	0.036	0.035	0.036		0.038	0.037	0.044
	LA	0.049	0.049	0.049	0.049	0.053		0.019	0.016
	CA	0.050	0.051	0.050	0.050	0.048	0.024		0.022
	PI	0.049	0.049	0.049	0.049	0.052	0.020	0.024	

^apopulation: HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute;

AY, Ayacucho; LA, Lambayeque; CA, Cajamarca; PI, Piura

Genetic divergence was determined using MEGA 5.2 (Tamura et al., 2011).

Table 4.

Summary statistics for the *cyt b* gene from the *Lutzomyia ayacuchensis* populations

population ^a	N	S	H	Hd	K	Pi
HU	48	14	13	0.501	0.741	0.001
CH	23	4	4	0.383	0.498	0.001
AL	24	6	7	0.558	0.652	0.001
PA	13	3	3	0.590	1.051	0.002
AY	46	17	20	0.824	1.569	0.003
LA	16	34	15	0.992	9.650	0.016
CA	12	20	9	0.939	5.379	0.009
PI	34	86	33	0.998	13.865	0.023
Total	216	136	97	0.868	20.300	0.034

^apopulation: HU, Huigra; CH, Chanchan; AL, Alausi; PA, Paute; AY, Ayacucho; LA, Lambayeque; CA, Cajamarca; PI, Piura

N, number of sequences; S, number of segregating sites; H, number of haplotypes; Hd, haplotype diversity; K, average number of differences; Pi, nucleotide diversity
The number of segregating sites, number of haplotypes, haplotype diversity, average number of differences, and nucleotide diversity for each population were calculated using DnaSP 5.0 (Rozas et al., 2003).

Fig. 1

Fig. 3

Fig. 4

Fig. 6

Fig. 7