Title	Projected coral bleaching in response to future sea surface temperature rises and the uncertainties among climate models
Author(s)	Yara, Yumiko; Fujii, Masahiko; Yamano, Hiroya; Yamanaka, Yasuhiro
Citation	Hydrobiologia, 733(1), 19-29 https://doi.org/10.1007/s10750-014-1838-0
Issue Date	2014-07
Doc URL	http://hdl.handle.net/2115/58019
Rights	The final publication is available at link.springer.com . © Springer International Publishing Switzerland 2014, Published online: 5 March 2014
Type	article (author version)
File Information	Yara_Hydrobiologia_final.pdf

[Instructions for use](#)

Projected coral bleaching in response to future sea surface temperature rises and the uncertainties among climate models

Yumiko Yara^{1,2}, Masahiko Fujii^{2,*}, Hiroya Yamano¹ and Yasuhiro Yamanaka²

¹Center for Environmental Biology and Ecosystem Studies, National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan

²Faculty of Environmental Earth Science, Hokkaido University, N10W5, Kita-ku, Sapporo, Hokkaido 060-0810, Japan

* Corresponding author: M. Fujii, E-mail: mfujii@ees.hokudai.ac.jp, Tel./Fax.: +81 11 706 2359

Abstract

We quantitatively evaluated the effects of rising sea surface temperature (SST) on coral bleaching and the uncertainties resulting from differences in global warming projections. To do so, we used monthly SSTs in the 21st century obtained from 23 climate models under the A1B scenario (from the Special Report on Emissions Scenarios) and SST-based indices for coral bleaching. All of the projections indicated that severe bleaching or death of corals will be common and severe in wide areas of the tropical and subtropical oceans by the middle of

this century. However, decadal oscillation could modify the exact timing by around ± 10 years. Such projections are important for conserving marine biodiversity and designing future strategies to avoid tropical and subtropical coral extinction. To obtain more reliable projections and reduce uncertainties, climate models should be improved by using higher spatiotemporal resolutions and more realistic biological indices should be embedded into existing models.

Keywords: Biodiversity, Coral bleaching, Future projection, Global warming, Rise in water temperature, Uncertainty

Introduction

Corals play a fundamental role in primary production and habitat formation for numerous other species in tropical and subtropical oceans. Thus, the degradation of coral habitats can cause fundamental modifications to coastal ecosystems. Although small-scale coral bleaching has been reported for at least 75 years (Yonge & Nichols, 1931), since the early 1980s, mass coral bleaching that might be connected to global climate change has increased rapidly in frequency, intensity, and geographical extent across tropical and subtropical oceans (e.g., Glynn, 1984, 1988, 1991, 1993; Brown, 1997; Hough-Guldberg, 1999, 2011, Nakano, 2004; Hough-Guldberg et al., 2007; Nojima & Okamoto, 2008). The largest such bleaching event, which occurred in 1998, is estimated to have killed 16% of the world's corals, primarily in the western Pacific and Indian Ocean (Wilkinson, 2004; IPCC, 2007b).

Coral bleaching is a general response to stress (Hough-Guldberg, 2011). Corals bleach in response to a range of conditions including sudden changes in light, temperature, and salinity, the presence of toxins, and microbial infections. The causal relationship between sea temperature and mass coral bleaching has been proven empirically (Hough-Guldberg and Smith, 1989; Glynn & D'Croz, 1990) and *in situ* (Brown, 1997; Hough-Guldberg, 1999). Using this causal relationship, mass bleaching events can be predicted with greater than 95% accuracy from satellite measurements of sea surface temperature (SST) anomalies relative to the maximum summer temperatures (e.g., Goreau & Hayes, 1994; Toscano et al., 2000).

Global warming and associated increases in seawater temperatures necessitate urgent precise projections of future coral bleaching and death, not only to conserve marine biodiversity but also to plan for the adaptation of human societies to these changes. Therefore, long-term future projections of the effects of global warming on corals derived from climate models, as well as short-term predictions from satellite SST measurements, are sought as guidelines to design our adaptive measures to climate change and global warming.

Using climate model outputs and simplified indices to express coral bleaching in response to future rises in water temperature, several modeling studies have attempted to project the future probability of coral bleaching and death (e.g., Done et al., 2003; Donner et al., 2005, 2009; Guinotte et al., 2003; Hoegh-Guldberg, 1999, 2005, 2011; Meissner et al., 2012; Sheppard et al., 2003; Tevena et al., 2012; Wooldridge et al., 2005; Yara et al., 2009, 2011, 2012; Frieler et al., 2012). These projected results are all qualitatively identical in that both the frequency and extent of the severe bleaching or death of corals are expected to intensify. Particularly, intermittent high water temperatures, which result in the severe bleaching or death of present-day corals, will appear perpetually in the latter half of the 21st century (e.g., Yara et al., 2009).

However, water temperature is projected differently by climate models with different spatial resolutions, which may generate uncertainties in the results, as discussed by Yara et al. (2009). Moreover, if the climate scenarios and indices used in projections are all different, it is

difficult to directly examine the uncertainties underlying such projections. In substance, through the comparison of 23 different climate model outputs, Yara et al. (2011) demonstrated that there exists a high uncertainty in the projected poleward range expansion of coral habitats in response to rising water temperatures. Therefore, we may need to pay special attention to evaluating coral bleaching projections based on climate model water temperature outputs.

In this study, using procedures similar to those of Yara et al. (2011), we quantitatively examine the potential effects of SST increases on coral bleaching, as well as the uncertainties resulting from differences in the SST warming trends identified among models and locations. The following section describes the experimental design of a SST-based index for coral bleaching and the SST datasets of the multiple climate models used in this study. The third section includes the results and discussion of projections of coral bleaching and their uncertainties. The last section draws conclusions based on the results and discussion.

Materials and methods

Simplified index for coral bleaching

The algorithm used with satellite data predicts that coral bleaching starts when a threshold of 1°C above a region's mean SST during the warmest month is exceeded for more than 4 weeks (e.g., Goreau & Hayes, 1994; Toscano et al., 2000; Hoegh-Guldberg, 2011; Meissner et al., 2012). Several previous studies use Degree Heating Weeks (DHW (°C week)), a product

of exposure intensity ($^{\circ}\text{C}$ above threshold) and duration (in weeks), developed by the National Oceanic and Atmospheric Administration (NOAA) Coral Reef Watch Program (Liu et al., 2003) to predict coral bleaching events (NOAA Hotspot Program; Hoegh-Guldberg, 1999; Strong et al., 2000). In this metric, coral bleaching is predicted to occur when $\text{DHW} > 4$, a condition that indicates that the period over which the threshold temperature is exceeded by 1°C lasts for more than 4 weeks. Coral bleaching becomes progressively worse at higher temperatures or for longer periods over which the threshold temperature is exceeded. Severe coral bleaching, which may lead to the extinction of corals, is predicted to occur by this metric when $\text{DHW} > 8$, that is, the period over which the threshold temperature is exceeded by 1°C (2°C) lasts for more than 8 (4) weeks.

However, most climate model outputs are available monthly rather than weekly, and the DHW cannot be applied to these outputs. Alternatively, the Degree Heating Month (DHM ($^{\circ}\text{C}$ month)) metric, derived from the DHW, has been used in modeling studies that only have access to monthly SST outputs (e.g., Donner et al., 2005; Yara et al., 2009; Tevena et al., 2012). By this metric, coral bleaching is predicted to occur when $\text{DHM} > 1$, i.e., the threshold temperature is exceeded by 1°C for more than 1 month. Similarly, severe coral bleaching is predicted to occur when $\text{DHM} > 2$, i.e., the threshold temperature is exceeded by 1°C (2°C) for more than 2 (1) months. The DHM value has proved to be a reasonable proxy for DHW value (Donner et al., 2005). In this study, DHM was used as a simplified index for predicting

coral bleaching, basically following the procedure introduced by Yara et al. (2009).

Datasets of modeled water temperatures

We used SST outputs provided by multiple climate model projections from the World Climate Research Programme's (WCRP's) phase 3 of the Coupled Model Intercomparison Project (CMIP3; Meehl et al., 2007), which was performed for the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC AR4; IPCC, 2007a). As noted by Yara et al. (2011), when evaluating projections based on the SST warming trends obtained from the CMIP3 models, it is important to consider the uncertainties in the SST trends.

Monthly mean SSTs from 23 CMIP3 model projections (Table 1; Yara et al., 2011) were used and combined with the DHM metric for coral bleaching projections. The climate models have different ocean models with different spatial resolutions. For example, the horizontal resolutions range from 0.2° to 5°. We employed the “20th century climate in coupled models” (20C3M) simulations from 1980 to 1999 as predicted by the models using the global warming projections under the Special Report on Emissions Scenarios (SRES) A1B scenario, which assumes a future world of rapid economic growth with a balanced emphasis on all energy sources (IPCC, 2007a).

Values obtained by each model may depart from the real values (or the expected values in future projections), which are referred to as the model's biases. Such biases need to be

corrected for the period of discussion, which is 2000 through 2099 in this study. We corrected biases in the monthly mean SST in each of the CMIP3 models as follows (Yara et al., 2009, 2011): First, we calculated monthly mean SST anomalies during 2000–2099 (i.e., 1,200 months) under the SRES A1B scenario projection ($SST_{SRESA1B}$) relative to the monthly mean climatology (the 20-year mean SST from 1980 to 1999) of the 20C3M simulation ($\overline{SST_{20C3M}}$). Second, the SST anomaly for each month during 2000–2099 was added to the observed monthly mean climatology (18-year mean SST from 1982 to 1999) of the NOAA Optical Interpolation Sea Surface Temperature (OISST; Reynolds et al. 2007), interpolated to a horizontal grid point in each of the CMIP3 models ($\overline{OISST}$). The modeled SST discussed in this study after the bias correction process described above is expressed as:

$$SST(x, y, t, n) = \underbrace{\overline{OISST}(x, y, t, n)}_{Climatology} + \underbrace{\{SST_{SRESA1B}(x, y, t, n) - \overline{SST_{20C3M}}(x, y, t, n)\}}_{Anomaly} \quad (1)$$

where x and y are the number of longitudinal and latitudinal grids in each model, respectively; t is the number of months, from the starting point of future simulations (January 2000) for SST and $SST_{SRESA1B}$, and the corresponding months from January to December for the monthly mean climatology ($\overline{SST_{20C3M}}$ and $\overline{OISST}$); n is the model number in Table 1.

We calculated the bias-corrected monthly mean SST from 23 CMIP3 models (Table 1) during 2000–2099. The CMIP3 multi-model SST outputs combined with the DHM were also compared to one another. This comparison was performed for four tropical/subtropical coral reefs for which coral bleaching has been monitored or projected in previous studies (e.g.,

148 Hoegh-Guldberg, 1999, 2011; Yara et al., 2009), the Sekisei Lagoon in the Ryukyu Islands,
 149 Japan (124.0°E, 24.3°N); Phuket, Thailand (98.4°E, 7.9°N); the US Virgin Islands in the
 150 Caribbean Sea (64.8°W, 18.3°N); and Heron Island, Australia, on the Great Barrier Reef
 151 (151.9°E, 23.4°S).

152 Although projected results are similar using any model outputs, different regional patterns
 153 and magnitudes in the SST warming trends of various model outputs lead to uncertainties in
 154 the projected results. This is because different models project different responses to the same
 155 external forcing as a result of their treatments of physical processes, numerical schemes, and
 156 other factors (e.g., Yara et al., 2011; Brown et al., 2012).

157 To evaluate quantitatively the uncertainties arising from these factors, using the same
 158 procedure as that of Yara et al. (2011), we divided the temporal fluctuation in modeled SST in
 159 the warmest months in the 21st century into four components: the climatology ($\overline{SST_c}$)
 160 obtained by averaging model results from 1980 to 1999, the global warming trend (ΔSST_{gw}),
 161 the decadal oscillation (ΔSST_d), and the interannual fluctuation (ΔSST_i). Then, the SST was
 162 expressed as the sum of the four temporal components as follows:

$$163 \quad SST(x, y, t, n) = \underbrace{\overline{SST_c}(x, y, t, n)}_{\text{Climatology}} + \underbrace{\Delta SST_{gw}(x, y, t, n)}_{\text{Global warming trend}} + \underbrace{\Delta SST_d(x, y, t, n)}_{\text{Decadal oscillation}} + \underbrace{\Delta SST_i(x, y, t, n)}_{\text{Interannual fluctuation}}, \quad (2)$$

164 where x and y are the number of longitudinal and latitudinal grids in each model, respectively;
 165 t is the number of months, from the starting point of future simulations (January 2000) for
 166 SST, ΔSST_{gw} , ΔSST_d , and ΔSST_i , and the corresponding month from January to December
 167 for the monthly mean climatology ($\overline{SST_c}$); n is the model number in Table 1.

168

169 **Results**

170 The projected frequency of the severe bleaching or death of corals in the four sites from
171 the 2000s to 2090s was obtained using SST outputs from all 23 CMIP3 climate models (Table
172 1) and DHM (Fig. 1). The frequency of the severe bleaching or death of corals is projected to
173 be as low as zero in the 2000s and 2010s, but is projected to rise thereafter under the SRES
174 A1B scenario.

175 In Equation (2), ΔSST_{gw} was calculated from the linear trend in monthly mean SST from
176 2000 to 2099. ΔSST_d was the decadal oscillation component generated by the
177 ocean-atmosphere climate system, such as the Pacific Decadal Oscillation, and was defined
178 by a 5-year running mean component of $(SST - \overline{SST}_c - \Delta SST_{gw})$. The remainder $(SST -$
179 $\overline{SST}_c - \Delta SST_{gw} - \Delta SST_d)$ was regarded as the interannual fluctuation component (ΔSST_i),
180 but is not discussed here because we considered the projected effects of global warming and
181 the uncertainties with time scales longer than 10 years in this study. To evaluate the range of
182 uncertainty derived from the decadal oscillation, we calculated the standard deviation of
183 ΔSST_d and compared the values of the +2 and -2 standard deviation cases (+2SD and -2SD
184 cases, respectively) to the standard case, which is defined as $\overline{SST}_c + \Delta SST_{gw}$. The difference
185 between the +2SD and -2SD cases indicates the possible range in the timing caused by the
186 decadal oscillation, that is, the uncertainty due to decadal variations in the timing of the

continuous severe bleaching or death of corals.

Uncertainties in the projected effects of SST warming on the severe bleaching or death of corals in the four sites were assessed for the warmest months in the CMIP3 multi-model projections (Fig. 2). The simulated results show that the severe bleaching or death of corals tends to start to occur continuously a decade earlier or later in the +2SD and -2SD cases, respectively, compared to the standard case. The time at which the probability of the severe bleaching or death of corals will exceed 50% (i.e., predicted by more than half of the total climate models) for the +2SD and -2SD cases is in the 2070s and 2090s in Sekisei Lagoon, in the 2070s and 2090s in Phuket, in the 2050s and 2090s in the US Virgin Islands, and in the 2080s and later than the 2090s in Heron Island, respectively.

Discussion

Projected coral bleaching and its uncertainties

Most of the climate models predict that extremely high SSTs will appear every year by the end of the 21st century. This means that the severe bleaching or death of corals will be a common and crucial issue over wide areas of tropical and subtropical oceans by the middle of this century, and we will need to take action to mitigate and adapt to global warming to avoid tropical and subtropical coral extinctions. Considering uncertainties in projected results, the difference in the timing of bleaching occurrence between the two cases is 20-40 years,

206 although the timing is different among models, being mostly 10-20 years earlier and later,
207 respectively, than the timing predicted by the standard case at each site. This means that the
208 timing could be modified by around ± 10 years by the decadal oscillation, which is the
209 uncertainty relevant to global warming.

210

211 Limitations of this study

212 There are, however, a number of limitations that lead to uncertainties in our model
213 results. Some result from the insufficient spatiotemporal resolution of climate models,
214 whereas others stem from biological indices that are too simplified when combined with
215 climate models.

216 The temporal and spatial resolutions of the model outputs are monthly and ~ 100 km,
217 respectively (Table 1). A higher temporal resolution ($\sim$ weekly) would project bleaching more
218 accurately, as shown by favorable predictions based on DHW (e.g., Liu et al., 2003). A higher
219 spatial resolution is required to reproduce physical processes in coral habitats in shallow
220 coastal areas. Moreover, ocean current patterns are an important factor when considering the
221 existence of corals in coastal areas because ocean currents transport coral eggs and larvae. To
222 reproduce the future distribution of corals under rising water temperature and changing ocean
223 currents with fewer uncertainties, climate models with higher spatial and temporal resolutions
224 are required. Preferable models for such aims include climate models from phase 5 and later

of the Coupled Model Intercomparison Project and those that embed the Regional Ocean Modeling System (ROMS), which provide results with high horizontal resolutions (e.g., Gruber et al., 2012).

Although all of these climate models use a set of primitive equations to reproduce physical processes in the ocean, different models tend to have strengths in different areas and at different spatial scales (Brown et al., 2012). Some models reproduce the Kuroshio Current well, whereas others reproduce El Niño-Southern Oscillation (ENSO) events well (e.g. Guilyardi et al., 2009; Brown et al., 2012; Ganachaud et al., 2013). For example, as oceanic conditions are controlled strongly by the Kuroshio Current in the Sekisei Lagoon, coral bleaching in this oceanic domain is expected to be reproduced well by the MIROC3.2_hires (Table 1) which has relative strength in reproducing the Kuroshio Current amongst the climate models. Yet, each model performance will continue to improve in future as climate models are updated. For example, both frequency and duration of ENSO events were reported to be reproduced better in the updated versions of MIROC (Watanabe et al., 2010; Sakamoto et al., 2012).

So far, we have not considered any potential for the thermal adaptation and acclimatization of corals to warming events. However, several previous studies have suggested that coral acclimatization and adaptation to extreme warming events will increase thermal tolerance (e.g., Brown et al., 2002; Castillo & Helmuth, 2005; Teneva et al., 2012;

244 Guest et al., 2012; Howells et al., 2012; Oliver & Palumbi, 2011; Hoegh-Guldberg et al.,
245 2007; Csaszar et al., 2010; Maynard et al., 2008; Brown & Cossins, 2011; Tsuchiya & Fujita,
246 2009; Frieler et al., 2012), especially in regions subject to more variable temperature regimes.
247 Such biological responses have the potential to alleviate much of the impact of warming on
248 corals (e.g., Pandolfi et al., 2011). Recently, future projections of coral bleaching in response
249 to global warming have included new, simplified indices of the adaptation of corals to future
250 global warming (Frieler et al., 2012). We will need to further develop new indices of the
251 adaptability of corals to coral bleaching events and embed these into climate models to further
252 reduce the uncertainties and increase the reproducibility of simulated results.

253 In addition to these emerging insights, we should also pay attention to understanding
254 the responses of coral communities to multiple environmental stressors (e.g., Manzello, 2010).
255 Although there is no doubt that high water temperatures are a major cause of coral bleaching,
256 other factors are also considered to be multiple stressors that can cause coral bleaching
257 simultaneously (e.g., Langdon & Atkinson, 2005; Anthony et al., 2008 and 2011; Meissner et
258 al., 2012; Yara et al., 2012). For example, experiments have shown that exposure to low pH
259 and the saturation state of the mineral carbonate aragonite caused by ocean acidification,
260 another global phenomenon, makes corals more prone to bleaching (Anthony et al., 2011).
261 Other regional and local factors such as destructive fishing, overfishing, siltation, pollution,
262 crown-of-thorns starfish predation, and diseases presumably co-affect future coral bleaching

events along with global warming. Improving such biological knowledge, along with improving physical projections as described above, would contribute significantly to projecting the future status of corals.

Acknowledgements

The authors appreciate the helpful comments of two anonymous reviewers. We thank the modeling groups, the Program for Climate Model Diagnosis and Intercomparison (PCMDI), and the WCRP's Working Group on Coupled Modelling (WGCM) for their roles in making the WCRP phase 3 of the Coupled Model Intercomparison Project (CMIP3) multi-model dataset available. Support for this dataset was provided by the Office of Science of the US Department of Energy. This study was supported by the Global Environment Research Fund (S-5 and S-9) of the Ministry of the Environment, Program for Risk Information on Climate Change of Ministry of Education, Culture, Sports, Science and Technology, Japan, and Hokkaido University's Sustainable Low Carbon Society Project.

References

- Anthony, K. R. N., D. I. Kline, G. Diaz-Pulido, S. Dove & O. Hoegh-Guldberg, 2008. Ocean acidification causes bleaching and productivity loss in coral reef builders. *Proceedings of the National Academy of Sciences of the United States of America* 105: 17442–17446.
- Anthony, K. R. N., J. A. Maynard, G. Diaz-Pulido, P. J. Mumby, P. A. Marshall, L. Cao & O. Hoegh-Guldberg, 2011. Ocean acidification and warming will lower coral reef resilience. *Global Change Biology* 17: 1798-1808.
- Bleck, R., 2002: An oceanic general circulation model framed in hybrid isopycnic-Cartesian coordinates. *Ocean Modelling* 37, 55-88.
- Brown, B. E, 1997. Coral bleaching: causes and consequences. *Coral Reefs*, 16s: S129-138.
- Brown, B. E. & A. R. Cossins, 2011. The potential for temperature acclimatisation of reef corals in the face of climate change. In Dubinsky, Z. & N. Stambler (eds), *Coral reefs: an ecosystem in transition*. Springer, Berlin, 421-433.
- Brown, B. E., R. P. Dunne, M. S. Goodson & A. E. Douglas, 2002. Experience shapes the susceptibility of a reef coral to bleaching. *Coral Reefs* 21: 119-126.
- Brown, J. N., A. S. Gupta, J. R. Brown, L. C. Muir, J. S. Risbey, P. Whetton, X Zhang, A. Ganachaud, B. Murphy & S. E. Wijffels, 2012. Implications of CMIP model biases and uncertainties for climate projections in the western tropical Pacific. *Climatic Change* 119: 147-161.
- Castillo, K. D. & B. S. T. Helmuth, 2005. Influence of thermal history on the response of *Montastrea annularis* to short-term temperature exposure. *Marine Biology* 148: 261-270.
- Csaszar, N. B. M., P. J. Ralph, R. Frankham, R. Berkelmans & M. J. H. van Oppen , 2010. Estimating the potential for adaptation of corals to climate warming. *Plos One* 5: e9751.

Diansky, N. A., A.V. Bagno & V. B. Zalesny, 2002. Sigma model of global ocean circulation and its sensitivity to variations in wind stress. *Izvestiya, Atmospheric and Oceanic Physics* (English Translation) 38(5): 477-494.

Diansky, N.A. & E. M. Volodin, 2002. Simulation of present-day climate with a coupled atmosphere-ocean general circulation model. *Izvestiya, Atmospheric and Oceanic Physics* (English Translation) 38(6): 732-747.

Done, T. J., P. Whetton, R. Johns, R. Berkelmans, J. Lough, W. Skirving & S. Wooldridge , 2003. Global climate change and coral bleaching on the Great Barrier Reef. In: *Development of Natural Resources and Mines*, Queensland, Australia.

Donner, S. D., 2009: Coping with commitment: Projected thermal stress on coral reefs under different future scenarios. *PLoS One* 4(6): e5712.

Donner, S. D., W. J. Skirving, C. M. Little, M. Oppenheimer & O. Hoegh-Guldberg, 2005. Global assessment of coral bleaching and required rates of adaptation under climate change. *Global Change Biology* 11: 2251-2265.

Flato, G.M. & G.J. Boer, 2001. Warming asymmetry in climate change simulations. *Geophysical Research Letter* 28: 195-198.

Frieler, K., M. Meinshausen, A. Golly, M. Mengel, K. Lebek, S. D. Donner & O. Hoegh-Guldberg, 2012. Limiting global warming to 2°C is unlikely to save most coral reefs. *Nature Climate Change* 3: 165–170.

Ganachaud, A., A. Sen Gupta, J. N. Brown, K. Evans, C. Maes, L. C. Muir & F. S. Graham, 2013. Projected changes in the tropical Pacific Ocean of importance to tuna fisheries. *Climatic Change* 119: 163-179.

Glynn, P. W., 1984. Widespread coral mortality and the 1982/83 El Ninjo warming event. *Environmental Conservation* 11: 133-146.

Glynn, P. W., 1988. Coral bleaching and mortality in the tropical eastern Pacific during the 1982-1983 El Nino warming event. In: Ogden J., & R. Wicklund (eds), Mass bleaching of coral reefs in the Carribean: a research strategy. NOAA's Undersea Research Program. St Croix, US Virgin Islands, Research Reports 88(2): 42-45.

Glynn, P. W., 1991. Coral reef bleaching in the 1980s and possible connections with global warming. *Trends in Ecology and Evolution* 6: 175-179.

Glynn, P. W., 1993. Coral reef bleaching: ecological perspectives. *Coral Reefs* 12: 1-17.

Glynn, P. W. & L. D'Croz, 1990. Experimental for high temperature stress as the cause of El Nino warming event. *Science* 253: 69-71.

Gnanadesikan, A., K. W. Dixon, S. M. Griffies, V. Balaji, M. Barreiro, J. A. Beesley, W. F. Cooke, T. L. Delworth, R. Gerdes, M. J. Harrison, I. M. Held, W. J. Hurlin, H.-C. Lee, Z. Liang, G. Nong, R. C. Pacanowski, A. Rosati, J. Russell, B. L. Samuels, Q. Song, M. J. Spelman, R. J. Stouffer, C. O. Sweeney, G. Vecchi, M. Winton, A. T. Wittenberg, F. Zeng, R. Zhang & J. P. Dunne, 2006. GFDL's CM2 global coupled climate models. Part II: The baseline ocean simulation. *Journal of Climate* 19: 675-697.

Gordon, C., C. Cooper, C.A. Senior, H. Banks, J.M. Gregory, T.C. Johns, J.F.B. Mitchell & R.A. Wood, 2000. The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments. *Climate Dynamics* 16: 147-168.

Gordon, H. B., L. D. Rotstayn, J. L. McGregor, M. R. Dix, E. A. Kowalczyk, S. P. O'Farrell, L. J. Waterman, A. C. Hirst, S. G. Wilson, M. A. Collier, I. G. Watterson & T. I. Elliott, 2002. The CSIRO Mk3 Climate System Model. CSIRO Atmospheric Research Technical Paper 60: 134pp.

Goreau, T. J. & R. L. Hayes, 1994. Coral bleaching and ocean hot-spots. *Ambio* 23: 176-180.

Gruber, N., C. Hauri, Z. Lachkar, D. Loher, T. L. Frölicher, and G.-K. Plattner, 2012. Rapid progression of ocean acidification in the California Current System, *Science* 337: 220-223.

379

380 Guest, J. R., A. H. Baird, J. A. Maynard, E. Muttaqin, A. J. Edwards, S. J. Campbell, K.
381 Yewdall, Y. A. Affendi & L. M. Chou ,2012. Contrasting patterns of coral bleaching
382 susceptibility in 2010 suggest an adaptive response to thermal stress. PLoS One 7(3): e33353.

383

384 Guilyardi E., A. Wittenberg, A. Fedorov, M. Collins, C. Wang, A. Capotondi, G.J. van
385 Oldenborgh & T. Stockdale, 2009. Understanding El Niño in Ocean-Atmosphere General
386 Circulation Model: progress and challenges. Bulletin of the American Meteorological Society
387 90: 325-340.

388

389 Guinotte, J. M., R. W. Buddemeier & J. A. Kleypas, 2003. Future coral reef habitat
390 marginality: temporal and spatial effects of climate change in the Pacific basin. Coral Reefs
391 22: 551–558.

392

393 Haak, H., J. Jungclaus, U. Mikolajewicz & M. Latif, 2003. Formation and propagation of great
394 salinity anomalies, Geophysical Research Letters 30(9) 1473: doi:10.1029/2003GL17065.

395

396 Hoegh-Guldberg, O., 1999. Climate change, coral bleaching and the future of the world's
397 coral reefs. Marine Freshwater Research 50: 839-866.

398

399 Hoegh-Guldberg, O., 2005: Low coral cover in a high-CO₂ world. Journal of
400 Geophysical Research 110: C09S06, doi:10.1029/2004JC002528.

401

402 Hoegh-Guldberg, O., 2011: Coral reef ecosystems and anthropogenic climate change.
403 Regional Environmental Change 11 (Supplement 1): S215-S227,
404 doi:10.1007/s10113-010-0189-2.

405

406 Hoegh-Guldberg, O., P. J. Mumby, A. J. Hooten, R. S. Steneck, P. Greenfield, E. Gomez, C. D.
407 Harvell, P. F. Sale, A. J. Edwards, K. Caldeira, N. Knowlton, C. M. Eakin, R. Iglesias-Prieto,
408 N. Muthiga, R. H. Bradbury, A. Dubi & M. E. Hatzioios, 2007. Coral reefs under rapid
409 climate change and ocean acidification. Science 318: 1737-1742.

410

411 Hoegh-Guldberg, O. & G. J. Smith, 1989. Light, salinity, and temperature and the population

density, metabolism and export of zooxanthellae from *Stylophora pistillata* and *Seriatopora hystrix*. *Journal of Experimental Marine Biology and Ecology* 129: 279-303.

Howells, E. J., V. H. Beltran, N. W. Larsen, L. K. Bay, B. L. Willis & M. J. H. van Oppen, 2012. Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nature Climate Change* 2: 116-120, doi: 10.1038/NCLIMATE1330.

IPCC, 2007a. The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor & H. L. Miller (eds.), Cambridge University Press, Cambridge, UK and New York, NY, USA, 996pp.

IPCC, 2007b. Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Parry, M. L., O. F. Canziani, J. P. Palutikof, P. J. van der Linden & C. E. Hanson (eds.), Cambridge University Press, Cambridge, UK and New York, NY, USA, 976pp.

Johns, T. C., R. E. Carnell, J. F. Crossley, J. M. Gregory, J. F. B. Mitchell, C. A. Senior, S. F. B. Tett & R. A. Wood, 1997. The second Hadley Centre coupled ocean-atmosphere GCM: model description, spinup and validation. *Climate Dynamics* 13: 103-134.

Johns, T. C., C. F. Durman, H. T. Banks, M. J. Roberts, A. J. McLaren, J. K. Ridley, C. A. Senior, K. D. Williams, A. Jones, G. J. Rickard, S. Cusack, W. J. Ingram, M. Crucifix, D. M. H. Sexton, M. M. Joshi, B-W. Dong, H. Spencer, R. S. R. Hill, J. M. Gregory, A.B. Keen, A. K. Pardaens, J. A. Lowe, A. Bodas-Salcedo, S. Stark & Y. Searl, 2006. The new Hadley Centre climate model (HadGEM1): Evaluation of coupled simulations. *Journal of Climate* 19(7): 1327-1353.

K-1 model developers, 2004. K-1 coupled model (MIROC) description, K-1 technical report, 1, H. Hasumi and S. Emori (eds.), Center for Climate System Research, University of Tokyo: 34pp.

Kim, S.-J., G.M. Flato & G.J. Boer, 2003. A coupled climate model simulation of the Last Glacial Maximum, Part 2: approach to equilibrium. *Climate Dynamics* 20: 635-661.

Kim, S.-J., G.M. Flato, G.J. Boer & N.A. McFarlane, 2002: A coupled climate model simulation of the Last Glacial Maximum, Part 1: transient multi-decadal response. *Climate Dynamics* 19: 515-537.

- Langdon, C. & M. J. Atkinson, 2005. Effect of elevated pCO₂ on photosynthesis and calcification of corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment. *Journal of Geophysical Research* 110: C09S07, doi:10.1029/2004JC002576.
- Legutke, S. & E. Maier-Reimer, 1999. Climatology of the HOPE-G Global Ocean General Circulation Model. Technical report, No. 21, German Climate Computing Centre (DKRZ), Hamburg: 90 pp.
- Liu, G., A. E. Strong & W. Skirving, 2003. Remote sensing of sea surface temperatures during 2002 Barrier Reef coral bleaching. *EOS, Transactions, American Geophysical Union* 84:, 137.
- Madec G., P. Delecluse, M. Imbard & C. Lévy, 1998. OPA version 8.1 Ocean General Circulation Model reference manual. Notes du Pôle de Modélisation, Institut Pierre-Simon Laplace (IPSL), France, N°11: 91 pp.
- Manzello, D. P., 2010. Coral growth with thermal stress and ocean acidification: lessons from the eastern tropical Pacific. *Coral Reefs* 29: 749-758, DOI 10.1007/s00338-010-0623-4, .
- Marsland, S. J., H. Haak, J. H. Jungclaus, M. Latif, & F. Röske, 2003. The Max Planck Institute global ocean/sea-ice model with orthogonal curvilinear coordinates. *Ocean Modelling* 5: 91-127.
- Maynard, J. A, K. R. N. Anthony, P. A. Marshall & I. Masiri, 2008. Major bleaching events can lead to increased thermal tolerance in corals. *Marine Biology* 155: 173–182.
- Meehl, G. A., T. F. Stocker, W. D. Collins, P. Friedlingstein, A. T. Gaye, J. M. Gregory, A. Kitoh, R. Knutti, J. M. Murphy, A. Noda, S. C. B. Raper, I. G. Watterson, A. J. Weaver & Z.-C. Zhao, 2007. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, chap. Global Climate Projections, Cambridge University Press, Cambridge, UK and New York, NY, USA, 747–846.
- Meissner, K. J., T. Lippmann & A. Sen Gupta, 2012. Large-scale stress factors affecting coral

reefs: open ocean sea surface temperature and surface seawater aragonite saturation over the next 400 years. *Coral Reefs*, doi:10.1007/s00338-011-0866-8.

Nakano, Y., 2004. Global environmental change and coral bleaching, In: *The Japanese Coral Reef Society and Ministry of the Environment (eds), Coral Reefs of Japan*, Ministry of Environment, Tokyo, 42-48.

Nojima, S. & M. Okamoto, 2008. Enlargement of habitats of scleractinian corals to north and coral bleaching events. *Nippon Suisan Gakkaishi* 74(5): 884-888.

Oliver, T. A. & S. R. Palumbi, 2011. Do fluctuating temperature environments elevate coral thermal tolerance? *Coral Reefs* 30: 429–440.

Pandolfi, J. M., S. R. Connolly, D. J. Marshall, A. L. Cohen, 2011. Projecting coral reef futures under global warming and ocean acidification. *Science* 333: 418–422.

Reynolds, R. W., T. M. Smith, C. Liu, D. B. Chelton, K. S. Casey & M. G. Schlax, 2007. Daily high-resolution-blended analyses for sea surface temperature. *Journal of Climate* 20: 5473-5496.

Russell, G. L., J. R. Miller & D. Rind, 1995. A coupled atmosphere-ocean model for transient climate change studies. *Atmosphere-Ocean* 33 (4): 683-730.

Russell, G. L., J. R. Miller, D. Rind, R. A. Ruedy, G. A. Schmidt & S. Sheth, 2000. Comparison of model and observed regional temperature changes during the past 40 years. *Journal of Geophysical Research* 105: 14891-14898.

Sakamoto, T. T., Y. Komuro, T. Nishimura, M. Ishii, H. Tatebe, H. Shiogama, A. Hasegawa, T. Toyoda, M. Mori, T. Suzuki, Y. Imada, T. Nozawa, K. Takata, T. Mochizuki, K. Ogochi, S. Emori, H. Hasumi, and M. Kimoto, 2012. MIROC4h –A new high-resolution atmosphere-ocean coupled general circulation model. *Journal of the Meteorological Society of Japan* 90(3): 325-359.

518 Salas-Mélia, D., F. Chauvin, M. Déqué, H. Douville, J.F. Guérémy, P. Marquet, S. Planton,
519 J.F. Royer & S. Tyteca, 2005. Description and validation of the CNRM-CM3 global coupled
520 model. CNRM Technical Report 103: 36pp.

521

522 Sheppard, C. R. C., 2003. Predicted recurrences of mass coral mortality in the Indian Ocean.
523 *Nature* 425: 294-297.

524

525 Smith, R. D., S. Kortas & B. Meltz, 1995. Curvilinear coordinates for global ocean models,
526 LA-UR-95-1146, Los Alamos National Laboratory, Los Alamos, New Mexico: 38pp.

527

528 Strong, A. E., E. J. Kearns & K. K. Gjovig, 2000. Sea surface temperature signals from
529 satellites: an update. *Geophysical Research Letters* 27(11): 1667-1670.

530

531 Teneva, L., M. Karnauskas, C. A. Logan, L. Bianucci, J. C. Currie & J. A. Kleypas, 2012.
532 Predicting coral bleaching hotspots: the role of regional variability in thermal stress and
533 potential adaptation rates. *Coral Reefs* 31: 1-12, doi:10.1007/s00338-011-0812-9.

534

535 Toscano, M. A., G. Liu, I. C. Guch, K. S. Casey & A. E. Strong, 2000. Improved prediction of
536 coral bleaching using high-resolution HotSpot anomaly mapping. *Proceedings of the 9th*
537 *International Coral Reef Symposium*: 1143-1147.

538

539 Tsuchiya, M. & Y. Fujita, 2009. Anguish of coral reefs. Tokai University Press 203pp.

540

541 Washington, W. M., J. W. Weatherly, G. A. Meehl, A. J. Semtner Jr., T. W. Bettge, A. P.
542 Craig, W. G. Strand Jr., J. Arblaster, V. B. Wayland, R. James, and Y. Zhang, 2000. Parallel
543 climate model (PCM) control and transient simulations, *Climate Dynamics* 16: 755-774.

544

545 Watanabe, M., T. Suzuki, R. O'ishi, Y. Komuro, S. Watanabe, S. Emori, T. Takemura, M.
546 Chikira, T. Ogura, M. Sekiguchi, K. Takata, D. Yamazaki, T. Yokohata, T. Nozawa, H.
547 Hasumi, H. Tatebe, and M. Kimoto, 2010. Improved climate simulation by MIROC5: Mean
548 states, variability, and climate sensitivity. *Journal of Climate* 23: 6312-6335.

549

550 Wilkinson, C. (ed.), 2004. Status of coral reefs of the world: 2004. Australian Institute of
551 Marine Science, Townsville, Queensland, Australia: 301pp.

552

553 Wooldridge, S., T. Done, R. Berkelman, R. Jones & P. Marshall, 2005. Precursors for
554 resilience in coral communities in a warming climate: a belief network approach. *Marine*
555 *Ecology Progress Series* 295: 157-169.

556

557 Yara, Y., M. Fujii, Y. Yamanaka, N. Okada, H. Yamano, K. Oshima, 2009. Projected effects of
558 global warming on coral reefs in seas close to Japan: *Journal of the Japanese Coral Reef*
559 *Society* 11: 131-140 (in Japanese with English abstract).

560

561 Yara, Y., K. Oshima, M. Fujii, H. Yamano, Y. Yamanaka & N. Okada, 2011. Projection and
562 uncertainty of the poleward range expansion of coral habitats in response to sea surface
563 temperature warming: A multiple climate model study. *Galaxea, Journal of Coral Reef Studies*
564 13: 11–20.

565

566 Yara, Y., M. Vogt, M. Fujii, H. Yamano, C. Hauri, M. Steinacher, N. Gruber, and Y.
567 Yamanaka, 2012. Ocean acidification limits temperature-induced poleward expansion of coral
568 habitats around Japan. *Biogeosciences* 9: 4955-4968.

569

570 Yonge, C. M. & A. G. Nichols, 1931. Studies on the physiology of corals. IV. The structure,
571 distribution and physiology of the zooxanthellae. *Scientific Reports of the Great Barrier Reef*
572 *Expedition, 1928-29(1)*, 135-176.

573

574 Yongqiang, Y., Y. Rucong, Z. Xuehong & L. Hailong, 2002. A flexible global coupled
575 climate model. *Advances in Atmospheric Sciences* 19: 169-190.

576

577 Yongqiang, Y., Z. Xuehong & G. Yufu, 2004. Global coupled ocean-atmosphere general
578 circulation models in LASG/IAP. *Advances in Atmospheric Sciences* 21:444-455.

579

580 Yukimoto, S., A. Noda, A. Kitoh, M. Hosaka, H. Yoshimura, T. Uchiyama, K. Shibata, O.
581 Arakawa & S. Kusunoki, 2006a. Present-Day Climate and Climate Sensitivity in the
582 Meteorological Research Institute Coupled GCM Version 2.3 (MRI-CGCM2.3). *Journal of*
583 *the Meteorological Society of Japan* 84: 333-363.

584

585 Yukimoto, S., A. Noda, T. Uchiyama, S. Kusunoki & A. Kitoh, 2001. Climate changes of the
586 twentieth through twenty-first centuries simulated by the MRI-CGCM2.3. *Papers in*
587 *Meteorology and Geophys* 51: 47-88.

588

589 Yukimoto, S., A. Noda, T. Uchiyama, S. Kusunoki & A. Kitoh, 2006b. Climate Changes of
590 the Twentieth through Twenty-first Centuries Simulated by the MRI-CGCM2.3. Papers in
591 Meteorology and Geophysics 56: 9-24.

592

593 Table 1. List of climate models of which monthly mean SSTs are used in this study and the
594 spatial resolution, ocean model and references

Model (Country)	Spatial resolution (longitude×latitude×the number of vertical layer)	Ocean model	References
1. BCCR-BCM2.0 (Norway)	1.5°×0.5-1.5°×35	NBRSC-MICOM1.0	(1)
2. CGCM3.1_T47 (Canada)	1.85°×1.85°×29	CCCMA(OGCM3.1)	(2),(3),(4)
3. CGCM3.1_T63 (Canada)	1.4°×0.94°×29	CCCMA(OGCM3.1)	(2),(3),(4)
4. CNRM-CM3 (France)	2°×0.5-2°×31	OPA8.1	(5),(6)
5. CSIRO-MK3.0 (Australia)	1.88°×0.84°×31	MOM2.2	(7)
6. CSIRO-MK3.5 (Australia)	1.88°×0.84°×31	MOM2.2	(7)
7. GFDL-CM2.0 (USA)	1.0°×0.33-1.0°×50	OM3	(8)
8. GFDL-CM2.1 (USA)	1.0°×0.33-1.0°×50	OM3.1	(8)
9. GISS-AOM (USA)	4°×3°×16	AOM 4×3	(9)
10. GISS-EH (USA)	2°×2°×16	HYCOM	(10)
11. GISS-ER (USA)	5°×4°×13	Russell Ocean	(9),(11)
12. FGOALS-g1.0 (China)	1.0°×1.0°×33	LICOM1.0	(12),(13)
13. INGV-SXG (Italy)	2°×1-2°×31	OPA 8.2	(5)
14. INM-CM3.0 (Russia)	2.5°×2°×33	INM-CM3.0	(14),(15)
15. IPSL-CM4 (France)	2°×1-2°×31	OPA	(5)
16. MIROC3.2_hires (Japan)	0.28°×0.19°×47	COCO3.3	(16)
17. MIROC3.2_medres (Japan)	1.4°×0.5-1.4°×43	COCO3.3	(16)
18. ECHO-G (Germany/Korea)	2.8°×0.5-2.8°×20	HOPE-G	(17)
19. ECHAM5-MPI-OM (Germany)	1.5°×1.5°×40	MPI-OM	(18),(19)
20. MRI-CGCM2.3.2 (Japan)	2.5°×0.5-2.0°×23	MRI-CGCM2.3.2a	(20),(21),(22)
21. NCAR-PCM1 (USA)	1-1.13°×0.27-1°×32	POP1.0	(23),(24)
22. UKMO-HadCM3 (UK)	1.25°×1.25°×20	HadCM3	(25),(26)
23. UKMO-HadGEM1 (UK)	1.0°×0.3-1.0°×40	HadGEM1	(27)

595 References noted here are: (1) www.bcm.uib.no; (2) Flato & Boer (2001); (3) Kim et al.
596 (2002); (4) Kim et al. (2003); (5) Madec et al. (1998); (6) Salas-Mélia et al. (2005); (7)
597 Gordon et al. (2002); (8) Gnanadesikan et al. (2006); (9) Russell et al. (1995); (10) Bleck
598 (2002); (11) Russell et al. (2000); (12) Yongqiang et al. (2002); (13) Yongqiang et al. (2004);
599 (14) Diansky et al. (2002); (15) Diansky & Volodin (2002); (16) K-1 model developers
600 (2004); (17) Legutke & Maier-Reimer (1999); (18) Haak et al. (2003); (19) Marsland et al.
601 (2003); (20) Yukimoto et al. (2001); (21) Yukimoto et al. (2006a); (22) Yukimoto et al.
602 (2006b); (23) Smith et al. (1995); (24) Washington et al. (2000); (25) Johns et al. (1997); (26)

603 Gordon et al. (2000); (27) Johns et al. (2006).

604

Sekisei Lagoon, Japan (124.0°E, 24.3°N)

Virgin Islands, USA (64.8°W, 18.3°N)

605

Phuket, Thailand (98.4°E, 7.9°N)

Heron Island, Australia (151.9°E, 23.4°S)

606

607 Fig. 1. Projected frequency and probability (%) of high SST that would potentially induce
 608 severe coral bleaching or death in Sekisei Lagoon in the Ryukyu Islands, Japan (124.0°E,
 609 24.3°N), Phuket, Thailand (98.4°E, 7.9°N), the US Virgin Islands in the Caribbean Sea
 610 (64.8°W, 18.3°N), and Heron Island, Australia, on the Great Barrier Reef (151.9°E, 23.4°S)
 611 for the 2000s to the 2090s, obtained using the projected monthly-mean SSTs of multiple
 612 climate models and a simplified evaluation metric of Degree Heating Month (DHM). For
 613 example, a frequency of 1 or 0.5 indicates that such high SSTs appear every year or five times
 614 a decade, respectively. The probability of occurrence of the severe bleaching or death of
 615 corals for each frequency in each decade is evaluated by how many climate models predicted
 616 the occurrence for each frequency in each decade. For example, the predicted probability of
 617 the continuous severe bleaching or death of corals in the 2090s is 70% because 16 of 23
 618 climate models predict this with a frequency of 1 for that decade.

619

621

622 Fig. 2. Cumulative probability (%) distribution of the timing of the continuous severe
 623 bleaching or death of corals in Sekisei Lagoon, Phuket, Virgin Islands, and Heron Island for
 624 the -2SD case (in gray bars), standard case (in white bars), and +2SD case (in black bars),
 625 respectively, from the 2000s through the 2090s, as projected by the climate models. A
 626 probability of 50%, for example, indicates that half of the total climate models project the
 627 timing of the continuous severe bleaching or death of corals.