Title	DNA barcoding for identification of sand fly species (Diptera: Psychodidae) from leishmaniasis-endemic areas of Peru
Author(s)	Nzelu, Chukwunonso O.; Cáceres, Abraham G.; Arrunátegui-Jiménez, Martín J.; Lañas-Rosas, Máximo F.; Yañez-Trujillano, Henry H.; Luna-Caiipo, Deysi V.; Holguín-Mauricci, Carlos E.; Katakura, Ken; Hashiguchi, Yoshihisa; Kato, Hirotomo
Citation	Acta Tropica, 145, 45-51 https://doi.org/10.1016/j.actatropica.2015.02.003
Issue Date	2015-03
Doc URL	http://hdl.handle.net/2115/58020
Type	article (author version)
File Information	ActaTropica v.145p.45.pdf

[Instructions for use](#)

1 DNA barcoding for identification of sand fly species (Diptera: Psychodidae) from
2 leishmaniasis-endemic areas of Peru

3 Chukwunonso O. Nzelu^a, Abraham G. Cáceres^{b,c}, Martín J. Arrunátegui-Jiménez^d, Máximo F.
4 Lañas-Rosas^e, Henry H. Yañez-Trujillano^f, Deysi V. Luna-Caipo^g, Carlos E. Holguín-Mauricci^h,
5 Ken Katakura^a, Yoshihisa Hashiguchi^{i,j,k} and Hiroto Kato^a

6

7 ^aLaboratory of Parasitology, Department of Disease Control, Graduate School of Veterinary
8 Medicine, Hokkaido University, Sapporo, Japan

9 ^bDepartamento Académico de Microbiología Médica, Facultad de Medicina Humana, Universidad
10 Nacional Mayor de San Marcos, Lima, Perú

11 ^cLaboratorio de Entomología, Instituto Nacional de Salud, Lima, Perú

12 ^dDirección Ejecutiva de Salud Ambiental, Gerencia Regional de Salud Lambayeque, Perú Área de
13 Entomología Laboratorio Referencial, Dirección Regional de Salud Piura, Perú

14 ^eÁrea de Entomología, Laboratorio Referencial, Dirección Regional de Salud Piura, Perú

15 ^fDirección Ejecutiva de Salud Ambiental, Dirección Regional de Salud Cusco, Perú

16 ^gDirección Ejecutiva de Salud Ambiental, Dirección Sub Regional de Salud Cutervo, Dirección
17 Regional de Salud Cajamarca, Perú

18 ^hLaboratorio Referencial, Dirección Regional de Salud Piura, Perú

19 ⁱDepartment of Parasitology, Kochi Medical School, Kochi University, Kochi, Japan

20 ^jCentro de Biomedicina, Universidad Central del Ecuador, Quito, Ecuador

21 ^kPrometeo, Secretaría Nacional de Educacion Superior, Ciencia, Tecnologia e Innovacion
22 (SENESCYT), Ecuador

23

24 *Corresponding author: Hirotomo Kato, Laboratory of Parasitology, Department of Disease
25 Control, Graduate School of Veterinary Medicine, Hokkaido University, North 18 West 9, Kita-ku,
26 Sapporo, Hokkaido, 060-0818, Japan.

27 Phone & Fax: +81-11-706-5196.

28 E-mail address: hkato@vetmed.hokudai.ac.jp

29

Abstract

Phlebotomine sand flies are the only proven vectors of leishmaniasis, a group of human and animal diseases. Accurate knowledge of sand fly species identification is essential in understanding the epidemiology of leishmaniasis and vector control in endemic areas. Classical identification of sand fly species based on morphological characteristics often remains difficult and requires taxonomic expertise. Here, we generated DNA barcodes of the cytochrome c oxidase subunit 1 (COI) gene using 159 adult specimens morphologically identified to be 19 species of sand flies, belonging to 6 subgenera/species groups circulating in Peru, including the vector species. Neighbor-joining (NJ) analysis based on Kimura 2-Parameter genetic distances formed non-overlapping clusters for all species. The levels of intraspecific genetic divergence ranged from 0 to 5.96%, whereas interspecific genetic divergence among different species ranged from 8.39 to 19.08%. The generated COI barcodes could discriminate between all the sand fly taxa. Besides its success in separating known species, we found that DNA barcoding is useful in revealing population differentiation and cryptic diversity, and thus promises to be a valuable tool for epidemiological studies of leishmaniasis.

Keywords: DNA barcode, sand flies, leishmaniasis, vector, Peru

1. Introduction

Phlebotomine sand flies (Diptera: Psychodidae, Phlebotominae) are of significant public health importance in many parts of the world as vectors of the causative agents of leishmaniasis, bartonellosis and sand fly fever (Maroli et al., 2013). Leishmaniasis remains one of the world's most neglected diseases; endemic in 98 countries, putting 350 million people at risk, with some two million new cases every year (WHO, 2010; Alvar et al., 2012). The disease is caused by several species of cell-infecting parasites of the genus *Leishmania*, and presents three major distinct clinical manifestations: cutaneous, mucocutaneous and visceral forms (Bates, 2007). Sand flies are the only proven vectors of leishmaniasis and approximately 800 species have been recorded in five major genera; *Phlebotomus* and *Sergentomyia* in the Old World, and *Lutzomyia*, *Brumptomyia* and *Warileya* in the New World (Munstermann, 2004). However, only species belonging to the genera *Phlebotomus* and *Lutzomyia* are the putative vectors of *Leishmania* (Munstermann, 2004; Bates, 2007; Kato et al., 2010). Therefore, correct sand fly species identification is very important to design strategies for surveillance and control of leishmaniasis in given endemic areas.

In Peru, three sand fly species, *Lutzomyia* (Lu.) *peruensis*, *Lu. ayacuchensis* and *Lu. verrucarum* have been incriminated as the vectors of *Leishmania* (*Viannia*) *peruviana*, the aetiological agent of Andean-type cutaneous leishmaniasis in the endemic areas (Davies et al., 1993; Perez et al., 1991, 1994, 2007; Caceres et al., 2004; Kato et al., 2008, 2011). Although these sand fly species role in disease transmission has been well studied, detailed molecular taxonomic knowledge of each species is far from complete. Presently, sand flies are typically identified based on morphological features; mainly by internal structures such as the cibarium, pharynx, spermatheca of females and terminal genitalia of males (Young and Duncan, 1994).

However, this morphological classification is laborious, time-consuming and often complicated by phenotypic plasticity and cryptic species complexes, as well as demanding considerable skill and taxonomic expertise. To overcome these technical limitations, in the last three decades molecular approaches have been increasingly employed to explore the taxonomy, population structure and phylogeny in insect vectors including sand flies (Mukhopadhyay et al., 2000; Uribe Soto et al., 2001; Depaquit et al., 2002; Beati et al., 2004; Khalid et al., 2010; Fujita et al., 2012; Yamamoto et al., 2013; Nzelu et al., 2014; Gomez et al., 2014; Kato et al., 2015).

Recently, DNA barcoding has been widely shown to be an effective molecular tool for species identification of many insects (Hebert et al., 2003a, b). Among the mitochondrial genes, cytochrome c oxidase I (COI) is reported to be the most conserved gene and hence has distinct advantages for taxonomic studies (Knowlton and Weigt, 1998). Studies have demonstrated the use of DNA barcoding as a valid molecular tool for identifying dipterans; mosquitoes (Cywinska et al., 2006), black flies (Pramual et al., 2011) and sand flies (Azpurua et al., 2010; Kumar et al., 2012; Contreras Gutiérrez et al., 2014; Nzelu et al., 2014), and revealing sand fly cryptic species (Cohnstaedt et al., 2011; Scarpassa et al., 2012; Kumar et al., 2012). Construction of DNA barcodes for each species of sand fly would provide an efficient tool for their identification. The aim of the present study was to explore the utility of the DNA barcode approach for identification of sand fly species in Peru, where leishmaniasis is endemic, by analyzing morphologically identified adult specimens.

2. Materials and methods

2.1 Sand fly collection

Sand flies were caught using CDC light traps and Shannon traps peridomiciliary, intradomiciliary and extradomiciliary at 12 Departments; Piura, Lambayeque, Cajamarca, La Libertad, San Martin, Huánuco, Lima, Huancavelica, Ayacucho, Apurimac, Cusco and Puno in Peru between 2001 and 2013 (Fig. 1, Table 1). All the flies were captured throughout the night between 18:00 p.m. and 6:00 a.m. with CDC light traps and between 18:00 p.m. and 22:00 p.m. with Shannon traps. The sand flies were morphologically identified mainly based on measurements of wing veins, the ratio of palpus length to length of antenna and the color of the thorax (Young and Duncan, 1994), and then fixed in absolute ethanol and stored at room temperature.

2.2 DNA extraction

Individual ethanol-fixed sand flies were lysed in 50 µl of DNA extraction buffer [150 mM NaCl, 10 mM Tris-HCl (pH8.0), 10 mM EDTA and 0.1% sodium dodecyl sulfate (SDS)] with 100 µg/ml proteinase K. The samples were incubated at 37 °C overnight. Afterwards, 25 µl of distilled water was added, heated at 95 °C for 5 min, and then, 0.5 µl portions were directly used as templates for PCR amplification.

2.3 Polymerase chain reaction and sequence analysis

The mitochondrial COI was amplified using the primers LCO 1490 (5'-GGTCAACAAATCATAAAGATATTGG3') and HCO 2198

113 (5'TAAACTTCAGGGTGACCAAAAATCA-3') (Folmer et al., 1994). The reaction was carried
114 out in a volume of 15 µl using a pair of primers (0.4 µM each), Ampdirect Plus (Shimadzu
115 Biotech, Tsukuba, Japan), and BIOTaq HS DNA polymerase (Bioline, London, UK). After an
116 initial denaturation at 95 °C for 5 min, amplification was performed with 35 cycles consisting of
117 denaturation at 95 °C for 1 min, annealing at 49 °C for 30 sec, extension at 72 °C for 2 min,
118 followed by a final extension at 72 °C for 10 min. PCR products were purified using a FastGene
119 Gel/PCR Extraction kit (NIPPON Genetics) and sequenced with a forward primer by the dideoxy
120 chain termination method using a BigDye Terminator v3.1 Cycle Sequencing Kit (Applied
121 Biosystems, Foster City, CA). To confirm the authenticity of the COI sequences obtained, some
122 individuals from each sand fly species (in total 40 specimens) were subjected to PCR using a high
123 fidelity DNA polymerase (KOD-Plus-ver.2; TOYOBO, Tokyo, Japan), and the sequences were
124 determined as described above.

125

126 2.4 Data analysis

127 The COI sequences were aligned by CLUSTAL W software (Thompson et al., 1997)
128 incorporated into MEGA (Molecular Evolutionary Genetics Analysis) version 5.2 (Tamura et al.,
129 2011). The nucleotide compositions and sequence divergences within and between species were
130 calculated using the distance model Kimura 2-Parameter (K2P) (Kimura, 1980). This model
131 provides the best estimate of divergence when genetic distances are low, as in recently diverged
132 taxa (Nei and Kumar, 2000). The 'best match' method in the SpeciesIdentifier module of
133 TaxonDNA software v1.7.8 (Meier et al., 2006) was used to test the frequency of successful
134 identification. A Neighbor-joining (NJ) tree of K2P distances was created to provide a graphic
135 representation of the clustering pattern among different species (Saitou and Nei, 1987). Branch

136 support for NJ was calculated using the bootstrapping method with 1,000 replicates in MEGA
137 5.2 (Tamura et al., 2011). All sequences and other specimen information are available in the
138 dataset project 'DBSFP' (Process IDs: DBSFP001-15 to DBSFP159-15) on the Barcode of Life
139 Data System (BOLD) at <http://www.boldsystems.org>. Sequence data are also available in the
140 DDBJ/EMBL/GenBank databases (<http://www.ncbi.nlm.nih.gov/genbank/>) under the accession
141 numbers AB984357–AB984520 (Table 1).

142

3. Results

3.1 Sequence analysis

A total of 159 COI sequences from 19 morphologically identified species of phlebotomine sand flies belonging to 6 subgenera/species groups were obtained. Each species was analyzed using 1 to 17 specimens (Table 1, 2). The COI sequences obtained varied from 650 to 672 bp by direct sequencing. No deletions, insertions or stop codons were found indicating that all sequences constituted functional mitochondrial products (Funk and Omland, 2003). As these COI sequences contained no indels, alignments were straight forward. The COI sequences from 19 species of sand flies had high adenosine and thymine (AT)-rich content bias (65.9%), similar to that observed for other insects (Simon et al., 1994), with an average nucleotide composition of A=28.2%, T=37.7% , C=18.0%, and G=16.2%.

Intraspecific genetic distances based on Kimura 2-Parameter values for the sand fly species ranged from 0 to 5.96% (Table 2). Maximum intraspecific genetic diversity was observed in 7 of the 19 species, especially within the three vector species of *Leishmania (Viannia) peruviana*; *Lutzomyia (Lu.) peruensis* (5.96%), *Lu. verrucarum* (3.48%) and *Lu. ayacuchensis* (2.76%) (Table 2). The high level of intraspecific divergence was related to geographical distribution in some species. The mean intraspecific K2P distance ranged from 0.05% (*Lu. caballeroi*) to 3.36% (*Lu. peruensis*) (Table 2). Interspecific K2P genetic distances for the sand fly species included in the study ranged from 8.39 to 19.08%. The overall nucleotide diversity recorded among the 19 species under the Subfamily Phlebotominae included in the study was found to be 13.9%. Low levels of minimum interspecific divergence typically occurred in closely related species, such as *Lu. tejadai* and *Lu. quillabamba* (8.39%), *Lu. bicornuta* and *Lu. battistinii* (8.88%), and *Lu. castanea* and *Lu. caballeroi* (10.15%). Collectively, the distances between species belonging to

the subgenus *Lutzomyia* (*Lu. bicornuta* and *Lu. battistinii*), and the *Verrucarum* group (*Lu. verrucarum* and *Lu. nuneztovari*) were 8.88 and 12.27%, respectively, whereas the distances between species belonging to the subgenus *Pifanomyia* (*Lu. maranonensis*, *Lu. robusta* and *Lu. nevesi*) ranged from 11.46 to 15.43%. On the other hand, the diversity within the subgenus *Helcocyrtomyia* (*Lu. munaypata*, *Lu. quillabamba*, *Lu. pescei*, *Lu. noguchii*, *Lu. castanea*, *Lu. caballeroi*, *Lu. tejadai*, *Lu. ayacuchensis* and *Lu. peruensis*) ranged from 8.39 to 16.5%. Using SpeciesIdentifier, the overall percentage of correct species identification was 100% as recognized by traditional taxonomy based on morphological features. Generally, the COI divergence within species did not exceed the genetic distance between species. As a result, there was a distinction between intraspecific and interspecific genetic divergences, which is critical for DNA barcoding success in taxa discrimination (Fig. 2).

3.2 Neighbor-joining tree

The Neighbor-joining tree showed shallow intraspecific and deep interspecific divergences (Fig. 3). Specimens of the same species were grouped closely together, regardless of the collection site. However, in exceptional cases, obvious geographic differences in sequences within the same species were found in three species, namely, *Lu. peruensis*, *Lu. verrucarum* and *Lu. maranonensis*. Each of these species was monophyletic, but the individuals were divided into two clades corresponding to geographic origin (Fig. 3), namely, *Lu. peruensis*: Cusco (southeast) clade, and Lima (central) and La Libertad (northwest) clade; *Lu. verrucarum*: Lima (central) clade, and Piura (northwest) clade; *Lu. maranonensis*: Querocotillo (Cajamarca-northwest) clade, and San José de Lourdes (Cajamarca-northwest) clade. Additionally, the *Lu. robusta* cluster was

188 monophyletic, but the individuals were divided into two clades: Cajamarca I (northwest) clade,
189 and Cajamarca II (northwest) clade (Fig. 3).

190

4. Discussion

Adequate knowledge of the ecological and medical importance of sand flies relies on correct identification of these minute and fragile insects. The gold standard for any robust taxonomic system is its ability to deliver reliable and accurate species identifications (Hebert et al., 2003a). In the present study, we were able to recover and align the targeted COI fragment from all sand fly species examined. The strongly biased AT-richness noted within the DNA barcode sequences was consistent with results from other insects including different sand fly genera (Cywinska et al., 2006; Pramual et al., 2011; Kumar et al., 2012; Contreras Gutiérrez et al., 2014). The range of intraspecific variation was between 0 and 5.96%, while the interspecific variability for species differentiation ranged between 8.39 and 19.08%. The range of intraspecific genetic divergence in this study, 0 to 5.96%, was very similar to values reported for New World sand flies in Colombia (0 to 6%; Contreras Gutiérrez et al., 2014), whereas the 5.96% maximum intraspecific divergence was higher than that earlier reported for Old World sand flies in India (2%; Kumar et al., 2012). High intraspecific sequence divergence was found among the three vector species of *Leishmania* (*Viannia*) *peruviana* in Peru (Davies et al., 1993; Perez et al., 1991, 1994, 2007; Caceres et al., 2004; Kato et al., 2008, 2011); *Lu. (Helcocyrtomyia) peruensis* (5.96% maximum intraspecific divergence), *Lu. (Verrucarum) verrucarum* (3.48%), and *Lu. (Helcocyrtomyia) ayacuchensis* (2.76%) (Table 2). Overall, there were no overlaps between the intra- and inter-specific divergence levels, despite the high values of intraspecific divergences observed in some species, supporting the utility of mtDNA COI in discriminating species (Hebert et al., 2003b).

Inclusion of DNA sequence data in phylogenetic analysis of individuals allows for quick determination of monophyletic groups and recognition of hidden or potential species. In addition to delineation of species, DNA barcoding can reveal cryptic diversity (Hebert et al., 2004;

Pramual et al., 2011; Kumar et al., 2012). The Neighbor-joining (NJ) tree analysis distinctively clustered the COI sequences obtained from individual *Lutzomyia* sand flies as previously determined by morphological features; specimens of single species formed barcode clusters with tight cohesion that were clearly distinct from those of allied species. However, in this study four taxa, *Lu. peruensis*, *Lu. verrucarum*, *Lu. maranonensis* and *Lu. robusta* were found to be complexes of at least 2 genetically distinct groups. The NJ tree revealed two deep-divergence reciprocally monophyletic clades in *Lu. peruensis*, the main vector of *L. (V.) peruviana* (Perez et al., 1991, 2007; Kato et al., 2008, 2011) that clustered by geographic collection area (Fig. 3), and indicated that the *Lu. peruensis* group from Cusco (southeast) was genetically distinct from those from Lima (central) and La Libertad (northwest). A similar scenario was previously observed in *Lu. peruensis* based on the analysis of *cyt b* gene, among specimens from two regions, Ancash and Lima (central), and La Libertad (northwest) (Yamamoto et al., 2013). Additionally, we also observed a high genetic distance between the Lima (central) and Piura (northwest) groups of *Lu. verrucarum*. Therefore, these observed genetic distances among *Lu. peruensis* and *Lu. verrucarum* groups indicated that the groups in each taxa were highly isolated, with little or no genetic exchange between the groups.

Phlebotomine sand flies are known to have limited sustained flight and tend to remain associated with their emergence sites, typically not dispersing more than a half kilometer (Alexander, 1987; Killick-Kendrick, 1990; Morrison et al., 1993). Such putatively low flight capacity and dispersal range of sand flies coupled with the heterogeneous terrain of the Andes may cause the isolation of these populations throughout their distributions, thereby producing genetically distinct but morphologically similar individuals. Consequently, genetic differentiation observed within some species, may be ascribable to local adaptation or genetic

drift. Moreover, the observation of genetic differentiation among the vector species (*Lu. peruensis*, *Lu. verrucarum*, and *Lu. ayacuchensis*) may have serious implications for epidemiological studies and vector control strategies. Indeed it has been shown that insecticide susceptibility and vector capacity can vary within evolutionary lineages, even within the same species (Lanzaro et al., 1993; O' Loughlin et al., 2008; Hassan et al., 2012). Collectively, the fact that some species showed higher conspecific divergence does not compromise the use of COI sequences for their identification; rather, it allows delineation of the regional lineages that constitute them (Hebert et al., 2003b). Nonetheless, further investigations with more specimens from various localities will be required to provide additional information on the levels of divergence within these vector species and other prevalent sand fly species in Peru. Such studies may also contribute to more effective leishmaniasis control in endemic areas.

In conclusion, this study provided the COI barcodes for several Peruvian sand flies and showed their effectiveness in discriminating species recognized through prior conventional taxonomic work. Aside from its success in discriminating known species, DNA barcoding was found to be useful in revealing population differentiation. A better understanding of the implication of the genetic differentiation among the vector species will require further molecular (genetic information from multiple loci), ecological and morphological studies.

255 **Acknowledgements**

256 This study was financially supported by the Ministry of Education, Culture, Sports, Science and
257 Technology (MEXT) of Japan (Grant nos. 23256002 and 25257501), the Program for Leading
258 Graduate Schools “Fostering Global Leaders in Veterinary Science for Contributing to One Health”
259 (FO1), MEXT, Japan. CON is a recipient of the Japanese Government Ministry of Education,
260 Culture, Science and Technology (MEXT) PhD fellowship.

261

References

- Alexander, J.B., 1897. Dispersal of phlebotomine sand flies (Diptera: Psychodidae) in a Colombian coffee plantation. *J. Med. Entomol.* 24, 552-558.
- Alvar, J., Vélez, I.D., Bern, C., Herrero, M., Desjeux, P., Cano, J., Jannin, J., den Boer, M., WHO Leishmaniasis Control Team, 2012. Leishmaniasis worldwide and global estimates of its incidence. *PLoS one* 7, e35671.
- Azpurua, J., De La Cruz, D., Valderama, A., Windsor, D., 2010. *Lutzomyia* sand fly diversity and rates of infection by *Wolbachia* and an exotic *Leishmania* species on Barro Colorado Island, Panama. *PLoS Negl. Trop. Dis.* 4, e627.
- Bates, P.A., 2007. Transmission of *Leishmania* metacyclic promastigotes by phlebotomine sand flies. *Int. J. Parasitol.* 37, 1097-1106.
- Beati, L., Caceres, A.G., Lee, J.A., Munstermann, L.E., 2004. Systematic relationships among *Lutzomyia* sand flies (Diptera: Psychodidae) of Peru and Colombia based on the analysis of 12S and 28S ribosomal DNA sequences. *Int. J. Parasitol.* 34, 225-234.
- Caceres, A.G., Villaseca, P., Dujardin, J.C., Bañuls, A.L., Inga, R., Lopez, M., Arana, M., Le Ray, D., Arevalo, J., 2004. Epidemiology of Andean cutaneous leishmaniasis: incrimination of *Lutzomyia ayacuchensis* (Diptera: Psychodidae) as a vector of *Leishmania* in geographically isolated, upland valleys of Peru. *Am. J. Trop. Med. Hyg.* 70, 607- 612.
- Cohnsteadt, L.W., Beati, L., Caceres, A.G., Ferro, C., Munstermann, L.E., 2011. Phylogenetics of the phlebotomine sand fly group *Verrucarum* (Diptera: Psychodidae: *Lutzomyia*). *Am. J. Trop. Med. Hyg.* 84, 913-922.

283 Contreras Gutiérrez, M.A., Vivero, R.J., Vélez, I.D., Porter, C.H., Uribe, S., 2014. DNA
 284 Barcoding for the identification of sand fly species (Diptera, Psychodidae, Phlebotominae) in
 285 Colombia. PLoS one 9, e85496.

286 Cywinska, A., Hunter, F.F., Hebert, P.D.N., 2006. Identifying Canadian mosquito species through
 287 DNA barcodes. Med.Vet. Entomol. 20, 413-424.

288 Davies, C.R., Fernandez, M., Paz, L., Roncal, N., Llanos-Cuentas, A., 1993. *Lutzomyia*
 289 *verrucarum* can transmit *Leishmania peruviana*, the aetiological agent of Andean cutaneous
 290 leishmaniasis. Trans. R. Soc. Trop. Med. Hyg. 87, 603-606.

291 Depaquit, J., Ferté, H., Léger, N., Lefranc, F., Alves-Pires, C., Hanafi, H., Maroli, M., Morillas-
 292 Marquez, F., Rioux, J.A., Svobodova, M., Volf, P., 2002. ITS 2 sequences heterogeneity in
 293 *Phlebotomus sergenti* and *Phlebotomus similis* (Diptera: Psychodidae): possible consequences
 294 in their ability to transmit *Leishmania tropica*. Int. J. Parasitol. 32, 1123-1131.

295 Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., 1994. DNA primers for amplification
 296 of mitochondrial cytochrome oxidase subunit I from diverse metazoan invertebrates. Mol.
 297 Mar. Biol. Biotechnol. 3, 294-299.

298 Fujita, M., Kato, H., Caceres, A.G., Gomez, E.A., Velez, L., Mimori, T., Zhang, F., Iwata, H.,
 299 Korenaga, M., Sakurai, T., Katakura, K., Hashiguchi, Y., 2012. Genotyping of sand fly
 300 species in Peruvian Andes where leishmaniasis is endemic. Acta Trop. 121, 93-98.

301 Funk, D.J., Omland, K.E., 2003. Species-level paraphyly and polyphyly: frequency, causes, and
 302 consequences, with insights from animal mitochondrial DNA. Annu. Rev. Ecol. Syst. 34, 397-
 303 423.

304 Gomez, E.A., Kato, H., Mimori, T., Hashiguchi, Y., 2014. Distribution of *Lutzomyia ayacuchensis*,
305 the vector of Andean-type cutaneous leishmaniasis, at different altitudes on the Andean slope
306 of Ecuador. *Acta Trop.* 137, 118-122.

307 Hassan, M.M., Widaa, S.O., Osman, O.M., Numiary, M.S., Ibrahim, M.A., Abshama, H.M., 2012.
308 Insecticide resistance in the sand fly, *Phlebotomus papatasi* from Khartoum State, Sudan.
309 *Parasit. Vectors* 5, 46.

310 Hebert, P.D.N., Cywinska, A., Ball, S.L., deWaard, J.R., 2003a. Biological identifications through
311 DNA barcodes. *Proc. R. Soc. Lond. B. Biol. Sci.* 270, 313-321.

312 Hebert, P.D.N., Ratnasingham, S., deWaard, J.R., 2003b. Barcoding animal life: cytochrome c
313 oxidase subunit I divergences among closely related species. *Proc. R. Soc. Lond. B. Biol. Sci.*
314 270, S96-S99.

315 Hebert, P.D.N., Stoeckle, M.Y., Zemplak, T.S., Francis, C.M., 2004. Identification of birds through
316 DNA. *PLoS Biol.* 2, e312.

317 Kato, H., Cáceres, A.G., Gomez, E.A., Mimori, T., Uezato, H., Marco, J.D., Barroso, P.A., Gomez,
318 E.A., Iwata, H., Hashiguchi, Y., 2008. Molecular mass screening to incriminate sand fly
319 vectors of Andean-type cutaneous leishmaniasis in Ecuador and Peru. *Am. J. Trop. Med. Hyg.*
320 79, 719-721.

321 Kato, H., Gomez, E.A., Cáceres, A.G., Uezato, H., Mimori, T., Hashiguchi, Y., 2010. Molecular
322 epidemiology for vector research on leishmaniasis. *Int. J. Environ. Res. Public Health* 7, 814-
323 826.

324 Kato, H., Gomez, E.A., Cáceres, A.G., Vargas, F., Mimori, T., Yamamoto, K., Iwata, H.,
325 Korenaga, M., Velez, L., Hashiguchi, Y., 2011. Natural infection of man-biting sand flies by
326 *Leishmania* and *Trypanosoma* species in the Northern Peruvian Andes. *Vector Borne Zoonotic*

327 Dis. 11, 515-521.

328 Kato, H., Cáceres, A.G., Gomez, E.A., Mimori, T., Uezato, H., Hashiguchi, Y., 2015. Genetic
 329 divergence in populations of *Lutzomyia ayacuchensis*, a vector of Andean-type cutaneous
 330 leishmaniasis, in Ecuador and Peru. *Acta Trop.* 141, 79-87.

331 Khalid, N., El Naiem, D., Aboud, M., Al Rabba, F., Tripet, F., 2010. Morphometric and
 332 molecular differentiation of *Phlebotomus* (*Phlebotomus*) sand flies. *Med. Vet. Entomol.* 24,
 333 352-360.

334 Killick-Kendrick, R., 1990. Phlebotomine vectors of the leishmaniasis: a review. *Med. Vet.*
 335 *Entomol.* 4, 1-24.

336 Knowlton, N., Weigt, L.A., 1998. New dates and new rates for divergence across the Isthmus of
 337 Panama. *Proc. R. Soc. Lond. B. Biol. Sci.* 265, 2257-2263.

338 Kimura, M., 1980. A simple method for estimating evolutionary rate of base substitutions through
 339 comparative studies of nucleotide sequences. *J. Mol. Evol.* 16, 111-120.

340 Kumar, N.P., Srinivasan, R., Jambulingam, P., 2012. DNA barcoding for identification of sand
 341 flies (Diptera: Psychodidae) in India. *Mol. Ecol. Res.* 12, 414-420.

342 Lanzaro, G.C., Ostrovskaya, K., Herrero, M.V., Lawyer, P.G., Warburg, A., 1993. *Lutzomyia*
 343 *longipalpis* is a species complex: genetic divergence and interspecific hybrid sterility among
 344 three populations. *Am. J. Trop. Med. Hyg.* 48, 839-847.

345 Maroli, M., Feliciangeli, M.D., Bichaud, L., Charrel, R., Gradoni, L., 2013. Phlebotomine sand
 346 flies and the spreading of leishmaniasis and other diseases of public health concern. *Med. Vet.*
 347 *Entomol.* 27, 123-147.

348 Meier, R., Kwong, S., Vaidya, G., Ng, P.K.L., 2006. DNA barcoding and taxonomy in Diptera: a
 349 tale of high intraspecific variability and low identification success. *Syst. Biol.* 55, 715-728.

350 Morrison, A.C., Ferro, C., Morales, A., Tesh, R.B., Wilson, M.L., 1993. Dispersal of the sand fly
 351 *Lutzomyia longipalpis* (Diptera: Psychodidae) at an endemic focus of visceral leishmaniasis in
 352 Colombia. J. Med. Entomol. 30, 427-435.

353 Mukhopadhyay, J., Ghosh, K., Braig, H., 2000. Identification of cutaneous leishmaniasis vectors,
 354 *Phlebotomus papatasi* and *P. duboscqi* using random amplified polymorphic DNA. Acta
 355 Trop. 76, 277-283.

356 Munstermann, L.E., 2004. Phlebotomine sand flies, the Psychodidae. In: Marquardt WC, Black
 357 WC, Freier JE, Hagedorn HH, Hemingway J, et al., editor. Biology of Disease Vectors,
 358 Second ed. Elsevier, San Diego, CA, pp. 141-151.

359 Nei, M., Kumar, S., 2000. Molecular Evolution and Phylogenetics. Oxford University Press, New
 360 York, P 333.

361 Nzelu, C.O., Kato, H., Puplampu, N., Desewu, K., Odoom, S., Wilsom, M.D., Sakurai, T.,
 362 Katakura K., Boakye, D.A., 2014. First detection of *Leishmania tropica* DNA and
 363 *Trypanosoma* species in *Sergentomyia* sand flies (Diptera: Psychodidae) from an outbreak
 364 area of cutaneous leishmaniasis in Ghana. PLoS Negl. Trop. Dis. 8, e2630.

365 O'Loughlin, S.M., Okabayashi, T., Hond, M., Kitazo, Y., Kishin, H., Somboon, P., Sochantha, T.,
 366 Nambanya S., Saiki, P., Dev, V., Walton, C., 2008. Complex population history of two
 367 *Anopheles dirus* mosquito species in Southeast Asia suggests the influence of Pleistocene
 368 climate change rather than human-mediated effects. J. Evol. Biol. 21, 1555-1569.

369 Perez, J.E., Villaseca, P., Caceres, A., Lopez, M., Zolessi, A., Campos, M., Guerra, H., Llanos
 370 Cuentas, A., 1991. *Leishmania* (*Viannia*) *peruviana* isolated from the sand fly *Lutzomyia*
 371 *peruensis* (Diptera: Psychodidae) and a sentinel hamster in the Huayllacallán Valley, Ancash.
 372 Peru. Trans. R. Soc. Trop. Med. Hyg. 85, 60.

373 Perez, J.E., Ogusuku, E., Inga, R., Lopez, M., Monje, J., Pas, L., Nieto, E., Arevalo, J., Guerra,
 374 H., 1994. Natural *Leishmania* infection of *Lutzomyia* spp. in Peru. Trans. R. Soc. Trop. Med.
 375 Hyg. 88, 161-164.

376 Perez, J.E., Veland, N., Espinosa, D., Torres, K., Ogusuku, E., Llanos-Cuentas, A., Gamboa, D.,
 377 Arévalo, J., 2007. Isolation and molecular identification of *Leishmania* (*Viannia*) *peruviana*
 378 from naturally infected *Lutzomyia peruensis* (Diptera: Psychodidae) in the Peruvian Andes.
 379 Mem. Inst. Oswaldo Cruz. 102, 655-658.

380 Pramual, P., Wongpakam, K., Alder, P.H., 2011. Cryptic biodiversity and phylogenetic
 381 relationship revealed by DNA barcoding of Oriental black flies in the subgenus *Gomphostilbia*
 382 (Diptera: Simuliidae). Genome, 54, 1-9.

383 Saitou, N., Nei, M., 1987. The Neighbour-joining method: a new method for reconstructing
 384 phylogenetic trees. Mol. Biol. Evol. 4, 406-425.

385 Scarpassa, V.M., Alencar, R.B., 2012. *Lutzomyia umbratilis*, the main vector of *Leishmania*
 386 *guyanensis*, represents a novel species complex? PLoS one 7, e37341.

387 Simon, C., Frati, F., Bechenbach, A., Crespi, B., Liu, H., Flook, P., 1994. Evolution, weighting,
 388 and a phylogenetic utility of mitochondrial gene sequences and a compilation of conserved
 389 polymerase chain reaction primers. Ann. Entomol. Soc. Am. 87, 651-701.

390 Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: molecular
 391 evolutionary genetics analysis using maximum likelihood, evolutionary distance, and
 392 maximum parsimony methods. Mol. Biol. Evol. 28, 2731-2739.

393 Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., Higgins, D.G., 1997. The Clustal X
 394 windows interface: flexible strategies for multiple sequence alignment aided by quality
 395 analysis tools. Nucleic Acids Res. 25, 4876-4882.

396 Uribe Soto, S.I., Lehmann, T., Rowton, E.D., Velez, I.D., Porter, C.H., 2001. Speciation and
 397 population structure in the morphospecies *Lutzomyia longipalpis* (Lutz and Neiva) as derived
 398 from the mitochondrial ND4 gene. Mol. Phylogenet. Evol. 18, 84-93.
 399 WHO, 2010. Control of the leishmaniasis: report of a meeting of the WHO Expert Committee
 400 on the control of leishmaniasis. World Health Organ Tech Rep Ser 949, Geneva.
 401 Yamamoto, K., Cáceres, A.G., Gomez, E.A., Mimori, T., Iwata, H., Korenaga, M., Sakurai, T.,
 402 Katakura, K., Hashiguchi, Y., Kato, H., 2013. Genetic diversity of the mitochondrial
 403 cytochrome *b* gene in *Lutzomyia* spp., with special reference to *Lutzomyia peruensis*, a main
 404 vector of *Leishmania* (*Viannia*) *peruviana* in the Peruvian Andes. Acta Trop. 126, 156-163.
 405 Young, D.G., Duncan, M.A., 1994. Guide to the identification and geographic distribution of
 406 *Lutzomyia* sand flies in Mexico, the West Indies Central and South America (Diptera:
 407 Psychodidae). Memoirs of the American Entomological Institute, 54. Associated Publishers-
 408 American Entomological Institute, Gainesville, FL.
 409

Figure Legends

Fig. 1. Map of Peru showing the geographic locations where sand flies were collected along with the respective species analyzed in this study. Bat: *Lu. battistinii*; Bic: *Lu. bicornuta*; Ver: *Lu. verrucarum*; Nun: *Lu. nuneztovari*; Mar: *Lu. maranonensis*; Nev: *Lu. nevesi*; Rob: *Lu. robusta*; Tej: *Lu. tejadai*; Qui: *Lu. quillabamba*; Cab: *Lu. caballeroi*; Cas: *Lu. castanea*; Pes: *Lu. pescei*; Aya: *Lu. ayacuchensis*; Per: *Lu. peruensis*; Mun: *Lu. munaypata*; Nog: *Lu. noguchii*; Fis: *Lu. fischeri*; Gor: *Lu. gorbitzi*; Eun: *Warileya euniceae*.

Fig. 2. Distribution of intraspecific (grey) and interspecific (black) genetic distances for the cytochrome c oxidase subunit I (COI) sequences of sand fly species from Peru.

Fig. 3. Neighbor-joining tree based on Kimura 2-parameter genetic distances of mitochondrial COI sequences of *Lutzomyia* sand fly species from Peru. The bar scale represents 0.01% divergences. Bootstrap values are shown above or below branches. Specimen IDs (in parentheses).

Fig. 1

Fig. 2

Fig. 3

Fig. 3 (Continued)

Table 1

Details of sand fly specimens used for DNA barcoding and analysis in the study along with the BOLD Process IDs and GenBank accession numbers.

Sand fly species	No. of samples	Location		Trapping method	Year of capture	BOLD Process IDs	GenBank accession nos.
		District	Department				
<i>Lu. battistinii</i>	10	Abancay	Apurimac	CDC	2012	DBSFP001-15- DBSFP010-15	AB984357-AB984366
<i>Lu. bicornuta</i>	3	Luricocha	Ayacucho	CDC	2009	DBSFP011-15- DBSFP013-15	AB984367-AB984369
<i>Lu. bicornuta</i>	5	San Marcos de Rocchac	Huancavelica	CDC	2011	DBSFP014-15- DBSFP018-15	AB984370-AB984374
<i>Lu. munaypata</i>	10	Maranura	Cusco	CDC	2007	DBSFP019-15- DBSFP028-15	AB984375-AB984384
<i>Lu. quillabamba</i>	2	Calca	Cusco	CDC	2011	DBSFP029-15- DBSFP030-15	AB984385-AB984386
<i>Lu. tejadai</i>	11	Huánuco	Huánuco	CDC	2007, 2012	DBSFP031-15- DBSFP041-15	AB984387-AB984397
<i>Lu. pescei</i>	9	Luricocha	Ayacucho	CDC	2009	DBSFP042-15- DBSFP050-15	AB984398-AB984406
<i>Lu. pescei</i>	1	Chacoche	Apurimac	CDC	2012	DBSFP051-15	AB984407
<i>Lu. caballeroi</i>	10	La Cuesta	La Libertad	CDC	2007	DBSFP052-15- DBSFP061-15	AB984408-AB984417
<i>Lu. castanea</i>	5	San José de Lourdes	Cajamarca	ShT	2001	DBSFP062-15- DBSFP066-15	AB984418-AB984422
<i>Lu. castanea</i>	4	San Ignacio	Cajamarca	CDC	2007	DBSFP067-15- DBSFP070-15	AB984423-AB984426
<i>Lu. fischeri</i>	12	Ambo	Huánuco	CDC	2013	DBSFP071-15- DBSFP082-15	AB984428-AB984439
<i>Lu. gorbitzi</i>	2	Motupe	Lambayeque	CDC	2013	DBSFP083-15- DBSFP084-15	AB984440-AB984441
<i>Lu. ayacuchensis</i>	7	Motupe	Lambayeque	CDC	2013	DBSFP085-15- DBSFP091-15	AB984442-AB984448
<i>Lu. ayacuchensis</i>	1	San José de Lourdes	Cajamarca	CDC	2013	DBSFP092-15	AB984449
<i>Lu. peruensis</i>	2	Ambar	Lima	CDC	2007	DBSFP093-15- DBSFP094-15	AB984450-AB984451
<i>Lu. peruensis</i>	3	La Cuesta	La Libertad	CDC	2007	DBSFP095-15- DBSFP097-15	AB984452-AB984454
<i>Lu. peruensis</i>	1	Cusipata	Cusco	CDC	2011	DBSFP098-15	AB984455
<i>Lu. peruensis</i>	1	Urubamba	Cusco	CDC	2012	DBSFP099-15	AB984456
<i>Lu. peruensis</i>	3	Ollantaytambo	Cusco	CDC	2012	DBSFP100-15- DBSFP102-15	AB984457-AB984459
<i>Lu. verrucarum</i>	3	Huancabamba	Piura	CDC	2004	DBSFP103-15- DBSFP105-15	AB984460-AB984462
<i>Lu. verrucarum</i>	2	San Antonio de Chaclla	Lima	CDC	2013	DBSFP106-15- DBSFP107-15	AB984463-AB984464
<i>Lu. verrucarum</i>	6	Santa Eulalia	Lima	CDC	2013	DBSFP108-15- DBSFP113-15	AB984465-AB984470
<i>Lu. nuneztovari</i>	3	Maranura	Cusco	CDC	2007	DBSFP114-15- DBSFP116-15	AB984471-AB984473
<i>Lu. noguchii</i>	9	Pullo	Ayacucho	CDC	2007	DBSFP117-15- DBSFP125-15	AB984474-AB984482
<i>Lu. nevesi</i>	2	Tarapoto	San Martin	CDC	2007	DBSFP126-15- DBSFP127-15	AB984483-AB984484
<i>Lu. maranonesis</i>	2	San José de Lourdes	Cajamarca	CDC	2001	DBSFP128-15- DBSFP129-15	AB984485-AB984486
<i>Lu. maranonesis</i>	12	Querocotillo	Cajamarca	CDC	2006	DBSFP130-15- DBSFP141-15	AB984487-AB984498
<i>Lu. robusta</i>	2	San José de Lourdes	Cajamarca	ShT	2001	DBSFP142-15- DBSFP143-15	AB984501-AB984502
<i>Lu. robusta</i>	10	San José de Lourdes	Cajamarca	CDC	2001	DBSFP144-15- DBSFP153-15	AB984503-AB984512
<i>Lu. robusta</i>	5	San Juan de Cutervo	Cajamarca	CDC	2007	DBSFP154-15- DBSFP158-15	AB984513-AB984517
<i>Wa. euniceae</i>	1	Alto Inambari	Puno	ShT	2013	DBSFP159-15	AB984520

Lu.: *Lutzomyia*; *Wa.*: *Warileya*; CDC: CDC light trap; ShT: Shannon trap; BOLD: Barcode of Life Data System.

Table 2

Intraspecific Kimura 2-Parameter (K2P) pairwise distances of sand fly species from Peru.

Subgenera/Species group	Sand fly species	No. of Specimens	Minimum (%)	Mean (%)	Maximum (%)
<i>Lutzomyia</i>	<i>Lu. battistinii</i>	10	0	0.27	0.46
	<i>Lu. bicornuta</i>	8	0	0.76	1.25
<i>Verrucarum</i>	<i>Lu. verrucarum</i>	11	0	1.59	3.48
	<i>Lu. nuneztovari</i>	3	0	0.3	0.46
<i>Pifanomyia</i>	<i>Lu. maranonesis</i>	14	0	0.81	1.87
	<i>Lu. nevesi</i>	2		0.93	0.93
	<i>Lu. robusta</i>	17	0	0.37	1.55
<i>Helcocyrtomyia</i>	<i>Lu. tejadai</i>	11	0	0.16	0.46
	<i>Lu. quillabamba</i>	2		0.15	0.15
	<i>Lu. caballeroi</i>	10	0	0.05	0.15
	<i>Lu. castanea</i>	9	0	0.5	1.24
	<i>Lu. pescei</i>	10	0	0.58	1.69
	<i>Lu. ayacuchensis</i>	8	0.2	1.85	2.76
	<i>Lu. peruensis</i>	10	0	3.36	5.96
	<i>Lu. munaypata</i>	10	0	0.28	0.46
	<i>Lu. noguchii</i>	9	0.2	0.92	1.72
<i>Pintomyia</i>	<i>Lu. fischeri</i>	12	0	0.2	0.75
<i>Blancasmyia</i>	<i>Lu. gorbitzi</i>	2		0.15	0.15
—	<i>Wa. euniceae</i>	1		N/A	

N/A: not applicable.