Title	Respiration and ammonia excretion of euphausiid crustaceans: synthesis toward a global-bathymetric model
Author(s)	Ikeda, Tsutomu
Citation	Marine Biology, 160(2), 251-262 https://doi.org/10.1007/s00227-012-2150-z
Issue Date	2013-02
Doc URL	http://hdl.handle.net/2115/59727
Rights	The final publication is available at Springer via http://dx.doi.org/10.1007/s00227-012-2150-z
Type	article (author version)
File Information	HUSCUP-Euphausiids.pdf

[Instructions for use](#)

Mar. Biol. 160: 251–262 (2013)

Respiration and ammonia excretion of euphausiid crustaceans: synthesis toward a
global-bathymetric model

Tsutomu Ikeda*

16-3-1001 Toyokawa-cho, Hakodate, 040-0065 Japan

ikedatutomu@sepia.plala.or.jp

Tel: +81-138-22-5612

Running head: Global-bathymetric model of euphausiid metabolism

Keywords: ammonia excretion, euphausiids, global-bathymetric model, O:N ratio,
respiration

ABSTRACT

Respiration and ammonia excretion rates of 19–24 euphausiids from the epipelagic through bathypelagic zones of the world's oceans were compiled. Body mass (expressed in terms of dry mass, carbon or nitrogen), habitat temperature and sampling depth were designated as parameters in multiple regression analysis. Results suggested that the three parameters were highly significant, contributing 71–89 % of the variance in respiration rates and 69–81% of the variance in ammonia excretion rates. Atomic O:N ratios derived from simultaneous measurements of respiration and ammonia excretion rates ranged from 11 to 90 (median: 27), and no appreciable effects of the three parameters on O:N ratios were detected. If global-bathymetric models for the metabolism and chemical composition of copepods and chaetognaths are compared with those of euphausiids, it becomes evident that euphausiids are unique in that they maintain high metabolic rates and accumulate moderate amounts of energy reserves (lipids).

Introduction

The order Euphausiacea (euphausiids) has 85 species and is distributed from epipelagic to bathypelagic depths in the world's oceans (Mauchline 1980). Euphausiids are an important link between primary production and production of fish, squids, marine mammals and seabirds (Mauchline and Fisher 1969; Mauchline 1980). Moreover, they are a major contributor to world plankton community biomass (21%), second only to copepods (47%), in terms of their total organic carbon (Longhurst 1985). Many euphausiids undertake diel vertical migrations (DVMs) of 400-500 m (Brinton 1967), a behavior which is regarded as an important mechanism of rapid vertical transport of materials in the pelagic realm of the ocean (Longhurst et al. 1990; Hidaka et al. 2001).

Information about metabolism [respiration rates, ammonia excretion rates, and O:N (as $\text{NH}_4\text{-N}$) ratios] has proven to be useful in understanding the energy demand, major metabolic substrates and nutritional condition of marine zooplankton (cf. Ikeda et al. 2000). Although body mass and temperature have been regarded as the two major parameters for defining the metabolic characteristics of marine epipelagic animals (Ivleva 1980; Ikeda 1985), habitat depth has emerged as an additional parameter since the observation that metabolic rates decrease rapidly with depth for larger pelagic animals with image-forming eyes such as micronektonic fishes, crustaceans, and cephalopods (Childress 1995; Seibel and Drazen 2007). To date, the effect of habitat depth on respiration rates and O:N ratios of euphausiids has only been analyzed together with mysids, decapods and other crustacean taxa (Childress 1975; Quetin et al. 1980; Ikeda 1988; Torres et al. 1994); no analyses have attempted for euphausiids as an individual taxon.

To evaluate global-bathymetric patterns in the metabolism of euphausiids, I

compiled published and unpublished data of respiration, ammonia excretion and O:N ratios of euphausiids living at various depths in polar, temperate and tropical/subtropical seas, and significant parameters affecting the variance were explored. Body mass, habitat temperature, sampling depth and ambient oxygen saturation have been used as determinants of metabolic rates as in the global-bathymetric model for pelagic copepods and chaetognaths (Ikeda et al. 2007; Ikeda and Takahashi 2012). Finally, the present results were compared with those of copepods and chaetognaths to highlight any unique global-bathymetric features of euphausiids.

Materials and methods

The data

The metabolic rate of animals is defined with respect to the activity of animals as ‘standard’ or ‘basal’ metabolism (maintenance only), ‘routine’ (uncontrolled but minimum motor activity), and ‘active’ metabolism (enforced activity at a maximal level). In most studies on euphausiid metabolism, no attempts have been made to relate the metabolic data to the level of activity on the premise that the measured rates are close to routine metabolism (Ikeda et al. 2000). As a notable exception, Torres and Childress (1983) established the relationship between respiration rates and swimming speed in the migrating euphausiid *Euphausia pacifica*. To date, intensive studies of the metabolic rates have been made on some euphausiids, including *E. pacifica* in the northern North Pacific (Ross 1982 and citations therein), *E. superba* in the Southern Ocean (Clarke and Morris 1983) and *Meganyctiphanes norvegica* in the northern North Atlantic (Saborowski et al. 2002). However, available information about metabolism on the other euphausiids is scarce and widely spread in the literature. For the present

analyses, the data compiled were those which met the following criteria:

1. Data represented post-larvae (juveniles and adults) collected from the field and used for experiments without considerable time delay (< 24 h).
2. Measurements were made in the absence of food at near *in situ* temperatures in the dark [and at normal pressure (1 atm) for deep-sea euphausiids on the premise that the hydrostatic pressure affects little to the metabolic rates of deep-sea pelagic crustaceans, cf. review of Ikeda et al. (2000)]. This practice, combined with the criterion above, makes it possible to compare the data of various euphausiids for which information about feeding conditions in the field prior to the experiments is not available.
3. O:N ratios were computed from simultaneous measurements of respiration rates and ammonia excretion rates.
4. Body mass in terms of wet mass (WM), dry mass (DM), carbon (C) or nitrogen (N) units were given together with metabolic data (note: body mass specific rates without body mass data are not useful).
5. The depth of sampling of euphausiids was described or deducible.

In cases where multiple papers (including those of the present author) were available on the same species from similar regions, preference was given to mine to reduce possible between-workers bias in the data. As a result, a total of 39 data sets from 24 euphausiids for respiration rates, and 31 data sets from 19 euphausiids were selected and analyzed in the present study (Table 1). Data sets were separated into juveniles, females, gravid females, and males if the information was available. The same euphausiids but from different locations were treated as independent data sets, though mere repetition of the data on the same euphausiids was carefully avoided. The data expressed in the form of regression equations only were converted to the

respiration rates of a specimen at mid-body mass ranges. Study sites of all euphausiids are plotted on the world map (Fig. 1) to illustrate the worldwide coverage of the data sets in the present study.

Regression model

In addition to the 2 conventional independent variables (X_1 : body mass; and X_2 : habitat temperature) used in the previous global respiration model for marine epipelagic crustaceans (Ivleva 1980; Ikeda 1985), one new independent variables (X_3 : mid-sampling depth) were introduced to the present analyses. X_1 was expressed as DM, C or N since the choice of the body mass unit is known to affect the results (Ivleva 1980; Ikeda 1985). Initially, ambient oxygen saturation was introduced as an another new independent variable, but removed in the following analyses because of high correlation ($R = -0.758$) with the sampling depth. For the species on which C and/or N composition data are not reported, the missing values were substituted with those of the same species studied by other workers, or with grand mean values for all euphausiids of the present study.

Two regression models were adopted according to the mathematical form of the temperature and body mass effects. One was a theoretical model characterized by the Arrhenius relationship ($R = R_0 M^{3/4} e^{-E_a/kT}$ or $E = E_0 M^{3/4} e^{-E_a/kT}$, where R is respiration rate, E is ammonia excretion rate, M is body mass, T is absolute temperature, $3/4$ is theoretical body mass exponent, E_a is an average activation energy for the rate-limiting enzyme-catalyzed biochemical reactions of metabolism, k is Boltzmann's constant ($8.62 \times 10^{-5} \text{ eV K}^{-1}$) and R_0 or E_0 is a normalization constant (cf. Gillooly et al. 2001) and the

other was empirical (or log/linear) model characterized by the Van't Hoff rule (Q_{10})

(Ikeda 1985; Ikeda and Takahashi 2012);

Theoretical model: $\ln Y = a_0 + a_1 \ln X_1 + a_2(1000X_2^{-1}) + a_3 \ln X_3$

Empirical model: $\ln Y = a_0 + a_1 \ln X_1 + a_2 X_2 + a_3 \ln X_3$

It is noted that a_1 was 0.75 (= 3/4) for the theoretical model. As indices of temperature effects, E_a of the theoretical model and Q_{10} of empirical model could be computed as $E_a = a_2 \times 1000 \times 8.62 \times 10^{-5}$ and $Q_{10} = \exp(10 \times a_2)$, respectively. The attributes of these variables were analyzed simultaneously by using stepwise multiple regression (forward selection) (Sokal and Rohlf 1995). Independent variables were added and removed at $p = 0.05$. The calculation was conducted using SYSTAT version 10.2.

Results

Metabolic rates

Of the euphausiids considered in the present study, the smallest and largest species were *Euphausia krohnii* (0.29 mgDM) and *Thysanopoda cornuta* (1405 mgDM), respectively (Table 2). Respiration rates at *in situ* temperature ranged from 0.43 μLO_2 ind.⁻¹ h⁻¹ (*Thysanoessa macrura* from the Scotia Sea) to 109 μLO_2 ind.⁻¹ h⁻¹ (*Thysanopoda cornuta*), and ammonia excretion rates from 0.026 μgN ind.⁻¹ h⁻¹ (*Thysanoessa macrura* juveniles from the Scotia Sea) to 7.44 μgN ind.⁻¹ h⁻¹ (*Euphausia superba* gravid females off Wilkes Land, Antarctica) (Table 2).

Prior to the stepwise multiple regression analyses, preliminary analysis was made for the effects of temperature and sampling depth on the rates of respiration (R) and

ammonia excretion (E) by first plotting the rates standardized to 1 mg DM ($R_0 = R \times DM^{-0.75}$ or $E_0 = E \times DM^{-0.75}$) against temperature (1000/K or $^{\circ}\text{C}$) where the scale coefficient of body mass was assumed as 0.75 (as in the theoretical model) (Fig. 2). To facilitate analysis, the data were separated into two groups depending on the depth of euphausiids sampled (< 500 m and ≥ 500 m). It is clear that the values for species from ≥ 500 m depth distribute well below the rates < 500 m at equivalent inverse temperature or temperature. Because of this result, only the data of < 500 m were used for the analysis of temperature effects on R_0 or E_0 . The resultant slope (-4.119 for respiration rates, and -5.214 for ammonia excretion rates) of the regression lines was used to compute R_0 or E_0 at a given temperature (designated as 10°C) of the euphausiids from these sampling depths (< 500 m + ≥ 500 m), which were plotted against the mid-sampling depth (Fig. 3). The standardized rates (R_0 or E_0 at 10°C) of these euphausiids were correlated negatively with the sampling depth ($p < 0.01$).

The overall results of stepwise multiple regressions showed that the three independent variables were highly significant (F -test, $p < 0.001$). As judged by R^2 values, the empirical model was superior to the theoretical model, accounting for 85.7–89.0% vs 71.2–74.0%, respectively, of the variability in respiration, and 78.2–80.6% vs 69.0–70.6%, respectively, for the ammonia excretion rates (Table 3). As a body mass unit, N yielded the best fit, excepting the theoretical model of ammonia excretion rates for which C yielded the best fit. Overall, the choice of body mass units affected little to R^2 values.

The multiple regression analyses yielded results similar to those of the simple regression analyses (Figs. 2, 3) in which respiration rates or ammonia excretion rates were standardized by body mass and temperature (e.g., R_0 or E_0 at 10°C , respectively)

and grouped based on a single criterion (mid-sampling depth or ambient oxygen saturation).

O:N ratios

The atomic ratio of oxygen consumption rate to ammonia-nitrogen excretion rate (O:N ratio) has been used as an index of the proportion of protein in the diet of marine zooplankton (Mayzaud and Conover 1988; Ikeda et al. 2000). When only protein is metabolized, the O:N ratio is 7 (Table 10.3 in Ikeda et al. 2000). When protein and lipid or carbohydrate are catabolized in equal quantities O:N ratios are calculated as 21 or 13. Hence, O:N ratios of 7–17 (mid-point of 21 and 13) may be used as an index of protein-oriented metabolism and ratios of >17 as lipid/carbohydrate-oriented metabolism.

A total of 31 O:N ratios from 19 euphausiids ranged from 11 (*Euphausia mutica*) to 90 (*Euphausia pacifica* female from the western subarctic Pacific). The O:N data were separated into two depth groups (< 500 m and $\geq$ 500 m) and plotted against habitat temperature of the euphausiids (Fig. 4). While the number of data from the $\geq$ 500 m group were limited to only two, they fall well within the range of those of the < 500 m group (*U*-test, $p = 0.904$). The correlation between the O:N ratios (pooled data of the two depth groups) and temperature was significant ($p < 0.05$) but the contribution of temperature to the variance in the O:N ratios was only 15% (R^2). Mean and median O:N ratio were $30.0 (\pm 17.4, \text{SD})$ and 27.1, respectively.

Discussion

Body mass and temperature as traditional parameters

The effects of body mass and temperature on respiration rates of euphausiids have been studied most intensively in *Euphausia pacifica*, an intermediate-sized species that is important throughout the North Pacific including west coast of North America (Small and Hebard 1967; Paranjape 1967; Ross 1982). Earlier studies showed that the respiration rates of *E. pacifica* were not a power function of body mass but scaled directly with it (e.g. $a_3 = 1.0$ in the regression models of the present study, cf. Table 3). Later, Ross (1982), working on a broader size range of *E. pacifica* specimens raised in the laboratory (0.01–3 mgC or 2 orders of magnitude as compared with 1–10 mg DM or 1 order of magnitude for the earlier workers), demonstrated that the rates were indeed a power function of body mass ($a_3 = 0.7$ – 0.8 , as has been reported for diverse animal taxa (Zeuthen 1947). The range of DM for the 24 euphausiids considered in the present study was 4 orders of magnitude (Table 2) and resultant a_3 from the empirical models in which the attributes of body mass and ambient oxygen saturation were taken into account was 0.78–0.79 (depending on the choice of body mass units; DM, C or N) for respiration rates and 0.80–0.81 for ammonia excretion (Table 3).

The effect of temperature on metabolism has been studied in individual euphausiid species at graded temperatures within the range of their habitats. According to the definition by Clarke (1987), this is “acclimation” (adjustment of an organism to a new temperature in the laboratory), in contrast to “adaptation” (the evolutionary adjustment of an organism’s physiology to environment). The activation energy (E_a) or Q_{10} values thus obtained for acclimated euphausiids by previous workers are 0.63–0.67 eV or 2.0–3.5 (Table 4). Similar intra-specific Q_{10} values (2–3) are typical for the respiration rates measured at graded temperatures within the ranges of natural habitats of acclimated aquatic fishes and crustaceans living in arctic and tropical regions

(Scholander et al. 1953). On the other hand, inter-specific E_a or Q_{10} values (0.35 or 1.7) derived from global data sets of adapted euphausiids are less than these intra-specific values, but similar to those from global compilation of inter-specific data of adapted pelagic copepods, chaetognaths, teleost fishes and crustaceans (Table 4). Analyzing the effect of temperature on 69 teleost fishes from polar to tropical regions, Clarke and Johnston (1999) noted that a $Q_{10} = 1.83$ derived from inter-specific data over the temperature range 0–30°C was smaller than typical intra-specific acclimated Q_{10} values in the literature (median: 2.40, $N = 138$), as is the case for euphausiids of the present study. Those results suggest that evolutionary temperature adaption has produced an inter-specific relationship that has a lower thermal sensitivity than is typical of intra-specific relationships (cf. Clarke and Johnston 1999).

Habitat (sampling) depth as a new parameter

The present results of negative effects of habitat depth on respiration rates of euphausiids are consistent with those of larger pelagic animals with image-forming eyes such as micronektonic crustaceans, cephalopods and fishes (Torres et al. 1994; Childress 1995; Seibel and Drazen 2007), and zooplankton with no such the eyes including copepods (Ikeda et al. 2006, 2007) and chaetognaths (Kruse et al. 2011; Ikeda and Takahashi 2012). For the progressive decline in respiration rates in deeper-living micronekton and zooplankton, the “visual-interactions hypothesis” (Childress 1995) or “predation-mediated selection hypothesis” (Ikeda et al. 2007) have been proposed respectively. These two hypotheses are similar as both interpret the phenomena as a result of lowered selective pressure for high activity at depth because of the decrease in visual predators in the dark. However, these two hypotheses are different in that the

former applies strictly to micronekton with functional eyes, and the latter to both micronekton and zooplankton irrespective of presence/absence of functional eyes.

Torres et al. (1994) compiled the relationship between respiration rates and the depth of occurrence for pelagic crustaceans (amphipods, decapods, euphausiids, isopods, mysids and ostracods) off California, the Gulf of Mexico, off Hawaii and Antarctic waters. According to their results, the respiration rates [standardized to a body size of 1 mg wet mass by using the body mass exponent of 0.75 (equivalent to the theoretical model adopted in the present study), and at 0.5°C by assuming $Q_{10} = 2.0$], the reduction in the rates of a specimen due to the increase of its habitat depth from 1 m to 1000 m depth is in the order of 0.1–0.5 times. Similar calculations for euphausiids based on the present results (theoretical models in Table 3) showed that the reduction was 0.5–0.6 times, depending on the choice of DM, C or N body mass unit, which is close to the upper range of the values for the mixed crustacean taxa by Torres et al. (1994).

Accompanied with reduction in respiration rates, lowered swimming activity of deeper-living pelagic crustaceans has also been demonstrated by higher water, lower protein or lower N content (= reduced muscles) of their bodies (Childress and Nygaard 1973, 1974; Ikeda et al. 2006). For euphausiids only, presently available data are too small to allow precise analysis of depth-related patterns in the body chemical composition of this taxon.

Ammonia excretion and O:N ratios

While no global-bathymetric model is presently available for ammonia excretion rates of marine zooplankton including euphausiids, global ammonia excretion models with two parameters (body mass, habitat temperature) for mixed zooplankton taxa

(Ikeda 1985) and for planktonic copepods (Ikeda et al. 2001) yielded a body mass exponent and Q_{10} similar to those for respiration rates. With regard to the effect of habitat depth, lowered ammonia excretion rates of deep-sea euphausiids have already been reported on the bathypelagic *Bentheuphausia ambloyops* (Quetin et al. 1980) and mesopelagic *Euphausia tricantha* (Ikeda 1988). Parallel changes in respiration and ammonia excretion rates are anticipated in other species. In support of this hypothesis, O:N ratios (mean; 30.0 ± 17.4 , median; 27.1) and body C:N ratios (4.0 ± 0.7 , see below) were observed to be largely stable across the 24 euphausiids considered in this study (Table 2). Insignificant effects of body mass and habitat temperature on O:N ratios have already been reported on mixed zooplankton taxa (Ikeda et al. 2001) and pelagic copepods (1985).

It is noted that the O:N ratios of euphausiids listed in Table 2 are derived from experiments in which they were placed in filtered seawater, a common practice when using the sealed-chamber method (Ikeda et al. 2000). Use of filtered seawater is imperative to determine the rates of respiration and ammonia excretion accurately without any corrections for complex uptake/release of oxygen and ammonia by food organisms, but starvation of euphausiids may influence their normal metabolism. Ikeda and Dixon (1984) examined the effect of methodological problems associated with the sealed chamber method on *Euphausia superba*. According to their results, respiration and ammonia excretion rates of wild *E. superba* in which daily ration is 5% are 1.6 and 4.5, respectively, times the rates of non-feeding krill in 24 h laboratory experiments. From these results, the O:N ratios of wild *E. superba* are calculated as 0.36 ($= 1.6/4.5$) times less than those of the non-feeding *E. superba*. Generalization of the results of *E. superba* to other euphausiids requires caution since no comparable data are available for

the other euphausiids.

Euphausiids as compared with copepods and chaetognaths

The present results for euphausiids can be put into a wider perspective if compared with global-bathymetric models presented for pelagic copepods and chaetognaths (Table 5). As an important group within the marine zooplankton, euphausiids are different from copepods and chaetognaths in that they are active swimmers, with well developed compound eyes and respiratory organs (gills). For a specimen with similar body mass (1 mg N) living in the epipelagic zone (100 m depth, and 100% oxygen saturation) of temperate latitudes (20°C), predicted respiration rates from the theoretical or empirical model (17.9 or $18.3 \mu\text{O}_2 \text{ ind.}^{-1} \text{ h}^{-1}$) of euphausiids are the greatest followed by that (10.2 or $12.3 \mu\text{O}_2 \text{ ind.}^{-1} \text{ h}^{-1}$) of copepods and that (5.1 or $7.9 \mu\text{O}_2 \text{ ind.}^{-1} \text{ h}^{-1}$) of chaetognaths. The rank of high to low remains unchanged but for the specimen living in the bathypelagic zone (1000 m, 5°C and 10% oxygen saturation). Higher respiration rates of euphausiids may be related to their active swimming as compared with copepods or chaetognaths. Large standard deviations (SD) associated with the mean O:N ratios of the three taxa suggest non-normal distribution of the O:N data. Thus, the medians rather than means are thought to provide better index of the central trend. High metabolic O:N ratios (median; 27.1) of euphausiids are indicative of their carbohydrate/lipid-oriented metabolism in contrast to protein-oriented metabolism in chaetognaths (median; 12.2), which are consistent with the food habits of these taxa. In terms of body chemical composition, chaetognaths are different from euphausiids and copepods by their considerably higher water content (84–95% of WM) and low upper range values of C:N ratios (< 5.1) as compared with respective values ($< 84\%$ of WM

and 8.6–9.4) of euphausiids and copepods. According to Ventura (2006), C:N ratio of crustacean plankton protein is 3.3, and it is 8.4 for organic matter composed of equal amounts of protein and lipids (as wax esters). Carbohydrate is omitted as it comprises < 8.5% of DM. On this basis, very high C:N ratios (3.7–9.4) of copepods are due to the accumulation of lipids by species inhabiting cold thermal regimes (high latitude seas, deep-seas) where they function as an energy reserve for coping with temporal food scarcity, for reproduction or for achieving neutral buoyancy as is seen in diapausing copepods (Lee et al. 2006). Judging from the C:N ratios, presently available data suggest that the accumulation of lipids in the body of euphausiids is modest among the three taxa (Table 5). To date, the maximum lipid content reported is as high as 70% of DM or more for *Rhincalanus*, *Metridia*, *Calanus* and *Neocalanus* copepods (Lee et al. 2000), and is around 50% for *Thysanoessa* euphausiids (Lee et al. 2000; Harvey et al. 2012), all in high latitude seas. Major lipid class is wax esters for copepods while this is phospholipids and triacylglycerols for euphausiids.

In conclusion, global-bathymetric compilation of respiration and ammonia excretion rate data in 19–24 euphausiids revealed that the rates are a function of body mass, habitat temperature and habitat depth. The O:N ratios were largely independent from these parameters. The present results from inter-specific analyses of respiration data are compared with those from intra-specific analyses by previous workers, indicating similar body mass effects but dissimilar thermal sensitivities (lower E_a or Q_{10} of the former than the latter). The reduction in metabolic rates of deeper-living euphausiids was close to the upper range of those having been reported on pelagic crustaceans. Global-bathymetric comparison of the present results with those of pelagic copepods and chaetognaths revealed taxon-specific respiration rates and metabolic O:N,

354 as the possible reflect of dissimilar swimming patterns, food habits and body structures

355 (water content, C:N ratios) among these taxa.

356

357 Acknowledgments

358 I am grateful to Jose Torres for reviewing earlier drafts of this paper.

359

References

- Båmstedt U (1986) Chemical composition and energy content. In: Corner EDS, O'Hara SCM (eds), The biological chemistry of marine copepods. Clarendon Press, Oxford, pp 1–58
- Brinton E (1967) Vertical migration and avoidance capability of euphausiids in the California Current. Limnol Oceanogr 12: 451–483
- Childress JJ (1975) The respiratory rates of midwater crustaceans as a function of depth occurrence and relation to the oxygen minimum layer off Southern California. Comp Biochem Physiol A 50: 787–799
- Childress JJ (1995) Are there physiological and biochemical adaptation of metabolism in deep-sea animals? Trends Ecol Evol 10: 30–36
- Childress JJ, Nygaard M (1973) The chemical composition of midwater fishes as a function of depth of occurrence off southern California. Deep-Sea Res 20: 1093–1109
- Childress JJ, Nygaard M (1974) The chemical composition and relative buoyancy of midwater crustaceans as a function of depth off Southern California. Mar Biol 27: 225–238
- Clarke A (1987) The adaptation of aquatic animals to low temperatures. In: Grout BWW, Morris GJ (eds), The effects of low temperatures on biological systems, Edward Arnold, London, pp 315–348
- Clarke A, Johnston NM (1999) Scaling of metabolic rate with body mass and temperature in teleost fish. J Anim Ecol 68: 893–905
- Clarke A, Morris DJ (1983) Towards an energy budget for krill: The physiology and biochemistry of *Euphausia superba* Dana. Polar Biol 2: 69–86

384 Conover RJ, Corner EDS (1968) Respiration and nitrogen excretion by some marine
385 zooplankton in relation to their life cycles. J Mar Biol Ass UK, 48: 49–75

386 Cowles DL, Childress JJ, Wells ME (1991) Metabolic rates of midwater crustaceans as
387 a function of depth of occurrence off the Hawaii Islands: food availability as a
388 selective factor? Mar Biol 110: 75–83

389 Davenport J, Trueman ER (1985) Oxygen uptake and buoyancy in zooplanktonic
390 organisms from the tropical eastern Atlantic. Comp Biochem Physiol 81A:
391 857–863

392 Donnelly J, Torres JJ (1988) Oxygen consumption of midwater fishes and crustaceans
393 from the eastern Gulf of Mexico. Mar Biol 97: 483–494

394 Gillooly JF, Brown JH, West GB, Savage VM, Charnov EL (2001) Effects of size and
395 temperature on metabolic rate. Science 293: 2248–2251

396 Harvey HR, Pleuthner RL, Lessard EJ, Bernhardt MJ, Shaw CT (2012) Physical and
397 biochemical properties of the euphausiids *Thysanoessa inermis*, *Thysanoessa*
398 *raschii*, and *Thysanoessa longipes* in the eastern Bering Sea. Deep-Sea Res II
399 65–70: 173–183

400 Hirche H-J (1984) Temperature and metabolism of plankton-I. Respiration of Antarctic
401 zooplankton at different temperatures with a comparison of Antarctic and
402 Nordic krill. Comp Biochem Physiol 77A: 361–368

403 Hidaka K, Kawaguchi K, Murakami M, Takahashi M (2001) Downward transport of
404 organic carbon by diel migratory micronekton in the western equatorial Pacific:
405 its quantitative importance. Deep-Sea Res I 48: 1923–1939

406 Iguchi N, Ikeda T (1995) Growth, metabolism and growth efficiency of a euphausiid
407 crustacean *Euphausia pacifica* in the southern Japan Sea, as influenced by

408 temperature. J Plankton Res 17: 1757–1769

409 Ikeda T (1974) Nutritional ecology of marine zooplankton. Mem Fac Fish Hokkaido
 410 Univ 22: 1–97

411 Ikeda T (1985) Metabolic rates of epipelagic marine zooplankton as a function of body
 412 mass and temperature. Mar Biol 85: 1–11

413 Ikeda T (1988) Metabolism and chemical composition of crustaceans from the Antarctic
 414 mesopelagic zone. Deep-Sea Res 35: 1991–2002

415 Ikeda T (2012) Metabolism and chemical composition of zooplankton from 500 to
 416 5,000 m depth of the western subarctic Pacific Ocean. J Oceanogr

417 Ikeda T, Bruce B (1986) Metabolic activity and elemental composition of krill and
 418 other zooplankton from Prydz Bay, Antarctica, during early summer
 419 (November–December). Mar Biol.92: 545–555

420 Ikeda T, Dixon P (1984) The influence of feeding on the metabolic activity of Antarctic
 421 krill (*Euphausia superba* Dana) Polar Biol 3: 1–9

422 Ikeda T, Kirkwood R (1989) Metabolism and elemental composition of a giant
 423 chaetognath *Sagitta gazellae* from the Southern Ocean. Mar Biol 100:
 424 261–267

425 Ikeda T, Mckinnon DA (2012) Metabolism and chemical composition of zooplankton
 426 and hyperbenthos from the Great Barrier Reef waters, North Queensland,
 427 Australia. Plankton Benthos Res 7: 8–19

428 Ikeda T, Mitchell AW (1982) Oxygen uptake, ammonia excretion, and phosphate
 429 excretion of krill and other Antarctic zooplankton, in relation to their body size
 430 and chemical composition. Mar Biol.71: 283–298

431

432 Ikeda T, Skjoldal HR (1989) Metabolism and elemental composition of zooplankton
 433 from the Barents Sea during early arctic summer. *Mar Biol* 100: 173–183
 434 Ikeda T, Takahashi T. (2012) Synthesis towards a global-bathymetric model of
 435 metabolism and chemical composition of marine pelagic chaetognaths. *J Exp*
 436 *Mar Biol Ecol* 424–425: 78–88
 437 Ikeda T, Kanno Y, Ozaki K, Shinada A (2001) Metabolic rates of epipelagic marine
 438 copepods as a function of body mass and temperature. *Mar Biol* 139: 587–586
 439 Ikeda T, Sano F, Yamaguchi A (2007) Respiration in marine pelagic copepods: a
 440 global-bathymetric model. *Mar Ecol Prog Ser* 339: 215–219
 441 Ikeda T, Torres JJ, Hernández-León S, Geiger SP (2000) Metabolism. In: Harris RP,
 442 Wiebe PH, Lenz J, Skjoldal HR, Huntley M (eds), ICES zooplankton
 443 methodology manual. Academic Press, San Diego, pp 455–532
 444 Ikeda T, Yamaguchi A, Matsuishi T (2006) Chemical composition and energy content
 445 of deep-sea calanoid copepods in the western North Pacific Ocean. *Deep-Sea*
 446 *Res I* 53: 1791–1809
 447 Ivleva IV (1980) The dependence of crustacean respiration rate on body mass and
 448 habitat temperature. *Int Revue ges Hydrobiol* 65: 1–47
 449 James MR, Wilkinson VH (1988) Biomass, carbon ingestion, and ammonia excretion
 450 by zooplankton associated with an upwelling plume in western Cook Strait,
 451 New Zealand. *NZ J Mar Freshw Res* 22: 249–257
 452 Kim HS, Yamaguchi A, Ikeda T (2010) Metabolism and elemental composition of the
 453 euphausiids *Euphausia pacifica* and *Thysanoessa inspinata* during the
 454 phytoplankton bloom season in the Oyashio region, western subarctic Pacific
 455 Ocean. *Deep-Sea Res* 57: 1733–1741

456 Kruse S, Brey T, Bathmann U (2010) Role of midwater chaetognaths in Southern
 457 Ocean pelagic energy flow. *Mar Ecol Prog Ser* 416: 105–113
 458 Lee RF, Hagen W, Kattner G (2006) Lipid storage in marine zooplankton. *Mar Ecol*
 459 *Prog Ser* 307: 273–306
 460 Lindley JA, Robins DB, Williams R (1999) Dry weight carbon and nitrogen of some
 461 euphausiids from the north Atlantic Ocean and the Celtic Sea. *J Plankton Res*
 462 21: 2053–2066
 463 Longhurst AR, Bedo A, Harrison WG, Head EJH, Sameoto DD (1990) Vertical flux of
 464 respiratory carbon by oceanic diel migrant biota. *Deep-Sea Res* 37: 685–694
 465 Mauchline J (1980) The biology of mysids and euphausiids. *Adv Mar Biol* 18: 1–681
 466 Mauchline J, Fisher LR (1967) The biology of euphausiids. *Adv Mar Biol* 7: 1–454
 467 Mayzaud P (1973) Respiration and nitrogen excretion of zooplankton. II. Studies of the
 468 metabolic characteristics of starved animals. *Mar Biol* 21: 19–28
 469 Mayzaud P, Conover RJ (1988) O:N atomic ratio as a tool to describe zooplankton
 470 metabolism. *Mar Ecol Prog Ser* 45: 289–302
 471 Morris MJ, Hopkins TL (1983) Biochemical composition of crustacean zooplankton
 472 from the eastern Gulf of Mexico. *J Exp Mar Biol Ecol* 69: 1–19
 473 Nemoto T, Kamada K, Hara K (1972) Fecundity of a euphausiid crustacean,
 474 *Nematoscelis difficilis*, in the North Pacific Ocean. *Mar Biol* 14:41–47
 475 Paranjape MA (1967) Molting and respiration of euphausiids. *J Fish Res Bd Canada* 24:
 476 1229–1239
 477 Quetin LB, Ross RM, Uchio K (1980) Metabolic characteristics of midwater
 478 zooplankton: ammonia excretion, O:N ratios, and the effect of starvation. *Mar*
 479 *Biol* 59: 201–209

480 Ross RM (1982) Energetics of *Euphausia pacifica*. I. Effects of body carbon and
 481 nitrogen and temperature on measured and predicted production. Mar Biol 68:
 482 1–13

483 Saborowski R, Bröhl S, Tarling GA, Buchholz F (2002) Metabolic properties of
 484 Northern krill, *Meganyctiphanes norvegica*, from different climatic zones. I.
 485 Respiration and excretion. Mar Biol 140: 547–556

486 Sameoto DD (1976) Respiration rates, energy budgets, and molting frequencies of
 487 three species of euphausiids found in the Gulf of St. Lawrence. J Fish Res Bd
 488 Canada 33: 2568–2576

489 Scholander PF, Flagg W, Walters V, Irving L (1953) Climatic adaptation in arctic and
 490 tropical poikilotherms. Physiol Zool 26: 67–92

491 Seibel, BA, Drazen JC (2007) The rate of metabolism in marine animals:
 492 environmental constraints, ecological demands and energetic opportunities.
 493 Phil Trans.R Soc B 362: 2061–2078

494 Small LF, Hebard JF (1967) Respiration of a vertically migrating marine crustacean
 495 *Euphausia pacifica* Hansen. Limnol Oceanogr 12: 272–280

496 Sokal RR, Rohlf FJ (1995) Biometry. The principles and practice of statistics in
 497 biological research. Freeman, New York

498 Stuart V (1986) Feeding and metabolism of *Euphausia lucens* (Euphausiacea) in the
 499 southern Benguela current. Mar Ecol Prog Ser 30: 117–125

500 Teal JM, Carey FG (1967) Effects of pressure and temperature on the respiration of
 501 euphausiids. Deep-Sea Res 14: 725–733

502 Torres JJ, Childress JJ (1983) Relationship of oxygen consumption to swimming speed
 503 in *Euphausia pacifica*. 1. Effects of temperature and pressure. Mar Biol 74: 79–86

504 Torres JJ, Childress JJ (1985) Respiration and chemical composition of the
 505 bathypelagic euphausiid *Bentheuphausia amblyops*. Mar Biol 87: 267–272
 506 Torres JJ, Aarset AV, Donnelly J, Hopkins TL, Lancraft TM, Ainley DJ (1994)
 507 Metabolism of Antarctic micronektonic Crustacea as a function of depth of
 508 occurrence and season. Mar Ecol Prog Ser 113: 207–219
 509 Ventura M (2006) Linking biochemical and elemental composition in freshwater and
 510 marine crustacean zooplankton. Mar Ecol Prog Ser 327: 233–246
 511 Vidal J, Whitley (1982) Rates of metabolism of planktonic crustaceans as relation to
 512 body weight and temperature of habitat. J Plankton Res 4: 77–84
 513 Zeuthen E (1947) Body size and metabolic rate in the Animal Kingdom with special
 514 regard to the marine microfauna. C r Trav Lab. Carlsberg (Sér chim) 26: 17–161
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531

Figure captions

Fig. 1. Study sites of metabolic rates of euphausiids. The sites were separated into two groups depending on the depth where euphausiids collected (< 500 m and ≥ 500 m). The number and associated character alongside the symbol correspond the code of each euphausiid listed in Table 1.

Fig. 2. Euphausiids. Relationship between the respiration rate (top) or ammonia excretion rate (bottom) standardized to a body size of 1 mg body DM (R_0 or E_0) and temperature (T^{-1} : $1000/K$, or T : $^{\circ}C$) of the specimens from shallow (< 500 m) and deep layers (≥ 500 m). The data points represent means from the data sets in Table 2, and the regression line is derived from shallow layer species only. **: $p < 0.01$.

Fig. 3. Euphausiids. Relationship between respiration rates (top) or ammonia excretion rates (bottom) standardized to a body size of 1 mgDM (R_0 or E_0) at $10^{\circ}C$ and mid-sampling depth. The data points represent means derived from the data sets in Table 2. For symbols see Fig. 2. **: $p < 0.01$.

Fig. 4. Euphausiids. Relationships between O:N (as NH_4-N) ratios and habitat temperature (T) of euphausiids from various regions of the world's oceans. The data points represent means in Tables 2. For symbols see Fig. 2. *: $p < 0.05$.

Table 1. A list of euphausiid species of which metabolic data were used for the present analyses. J = juveniles, M = males, F = females, FG = gravid females. Blank = no data.

Code	Species	Region	Date	Reference
1a	<i>Bentheuphausia amblyops</i>	W. subarctic Pacific	May 2005	Ikeda (2012)
1b	<i>Bentheuphausia amblyops</i>	Off S. California	1976–1978	Torres & Childress (1985)
1c	<i>Bentheuphausia amblyops</i>	Off S. California	1976–1977	Quetin et al. (1980)
2a	<i>Euphausia crystallorophias</i>	Prydz Bay, Antarctica	Nov–Dec 1982	Ikeda & Bruce (1986)
2b	<i>Euphausia crystallorophias</i> (J)	Off Enderby Land, Antarctica	Nov 1985	Ikeda & Kirkwood (1989)
3	<i>Euphausia diomedea</i>	Tropical Atlantic	Nov 1971	Ikeda (1974)
4	<i>Euphausia distinguenda</i>	Tropical Pacific	Feb 1972	Ikeda (1974)
5	<i>Euphausia gibba</i>	Coral Sea	May 2010	Ikeda & McKinnon (2012)
6	<i>Euphausia gibboides</i>	Off W. Africa	Aug 1983	Davenport & Trueman (1985)
7	<i>Euphausia krohnii</i>	Topical Atlantic	Nov 1971	Ikeda (1974)
8	<i>Euphausia lucens</i>	Off W. South Africa		Stuart (1986)
9	<i>Euphausia mutica</i>	Coral Sea	May 2010	Ikeda & McKinnon (2012)
10a	<i>Euphausia pacifica</i>	E. subarctic Pacific	July 1975	Ikeda, unpublished
10b	<i>Euphausia pacifica</i> (F)	W. subarctic Pacific	Mar–Apr 2007	Kim, Yamaguchi & Ikeda (2010)
10c	<i>Euphausia pacifica</i> (M)	W. subarctic Pacific	Mar–Apr 2007	Kim, Yamaguchi & Ikeda (2010)
11a	<i>Euphausia superba</i>	Off Wilkes Land, Antarctica	Jan 1980	Ikeda & Mitchell (1982)
11b	<i>Euphausia superba</i> (J)	Off Enderby Land, Antarctica	Nov 1985	Ikeda & Kirkwood (1989)
11c	<i>Euphausia superba</i> (FG)	Off Wilkes Land, Antarctica	Jan 1980	Ikeda & Mitchell (1982)
11d	<i>Euphausia superba</i> (FG)	Off Enderby Land, Antarctica	Nov 1985	Ikeda & Kirkwood (1989)
11e	<i>Euphausia superba</i> (M)	Off Enderby Land, Antarctica	Nov 1985	Ikeda & Kirkwood (1989)
12	<i>Euphausia tenera</i>	Tropical Indian & Atlantic	Oct–Dec 1971	Ikeda (1974)
13a	<i>Euphausia triacantha</i>	Off Wilkes Land, Antarctica	Jan 1980	Ikeda & Mitchell (1982)
13b	<i>Euphausia triacantha</i>	Prydz Bay, Antarctica	Dec 1984–Feb 1985	Ikeda (1988)
14a	<i>Meganyctiphanes norvegica</i>	Gulf of Maine	Jan–Mar 1966	Conover & Corner (1968)
14b	<i>Meganyctiphanes norvegica</i>	Mediterranean	Jan 1971	Mayzaud (1973)
14c	<i>Meganyctiphanes norvegica</i>	Kattegat Bay, Denmark	Jul 1996, Mar 1997	Saborowski et al. (2002)
15	<i>Nyctiphanes australis</i>	E. Cook Strait, NZ	Mar–Apr 1983	James & Wilkinson (1988)
16	<i>Tessarabrachion oculatus</i>	W. subarctic Pacific	Jun 2003	Ikeda (2012)
17	<i>Thysanoessa inermis</i>	Barents Sea	May–June 1987	Ikeda & Skjoldal (1989)
18a	<i>Thysanoessa macrura</i> (J)	Scotia Sea	Jan–Mar 1981	Ikeda, unpublished data
18b	<i>Thysanoessa macrura</i>	Prydz Bay, Antarctica	Nov–Dec 1982	Ikeda, unpublished data
18c	<i>Thysanoessa macrura</i>	Off Enderby Land, Antarctica	Sept–Dec 1985	Ikeda, unpublished data
19	<i>Thysanoessa raschii</i>	Gulf of St. Lawrence	June 1973	Sameoto (1976)
20	<i>Thysanoessa</i> spp.	Mid-Atlantic Bight	Aug–Sep 1980	Vidal & Whitedge (1982)
21a	<i>Thysanoessa spinifera</i> (F)	W. subarctic Pacific	Mar–Apr 2007	Kim, Yamaguchi & Ikeda (2010)
21b	<i>Thysanoessa spinifera</i> (M)	W. subarctic Pacific	Mar–Apr 2007	Kim, Yamaguchi & Ikeda (2010)
22	<i>Thysanopoda cornuta</i>	Off Hawaii, USA	July 1983, 1986, 1987	Cowles et al. (1991)
23	<i>Thysanopoda monocantha</i>	E. Gulf of Mexico	Jun 1981–July 1985	Donnelly & Torres (1988)
24	<i>Thysanopoda tricuspidata</i>	Coral Sea	Nov 2009	Ikeda & McKinnon (2012)

576

577

578

579

580

581

582

583

584

585

586

Table 2. Body mass, metabolic rates (oxygen consumption and ammonia excretion), O:N ratio and elemental composition (C, N) of euphausiids from various depths and regions of the world's oceans. For species codes, see Table 1. Values are means $\pm$ SD. *Italic* values for sampling depth are those estimated by the present author. Blank = no data.

Code	Sampling depth (range)		T (°C)	N	Body mass (mg DM ind. ⁻¹)	Respiration consumption rate (μ l O ₂ ind. ⁻¹ h ⁻¹)	Ammonia excretion rate (μ g N ind. ⁻¹ h ⁻¹)	O:N ratio (atomic)	Elemental composition		
	(m)								C (% of DM)	N (% of DM)	C:N
1a	4000	(3000–5000)	1.5	1	41.7	8.73			52.1	9.0	5.8
1b	1500	(500–2500)	3.5	16	27.9	3.82 $\pm$ 0.18			39.2	6.4	6.1
1c	600	(400–800)	5.5	1	60.0	10.0	0.30	41	[39.2] ^a	[6.4] ^a	
2a	10	(0–21)	–1.6	24	30.8 $\pm$ 18.9	16.4 $\pm$ 9.4	0.68 $\pm$ 0.61	42 $\pm$ 22	45.5	11.0	4.1
2b	5	(2–8)	–1.3	42	9.10 $\pm$ 5.17	4.68 $\pm$ 2.41	0.24 $\pm$ 0.21	35 $\pm$ 18	41.9	11.7	3.6
3	2	(0–<5)	26	2	1.89 $\pm$ 0.62	5.55 $\pm$ 0.21	0.42 $\pm$ 0.23	19 $\pm$ 10	41.3	10.6	3.9
4	2	(0–<5)	27	2	3.54 $\pm$ 2.23	15.6 $\pm$ 4.5	1.19 $\pm$ 0.11	16 $\pm$ 3	39.4	10.6	3.7
5	1	surface	28	7	3.29 $\pm$ 0.69	11.5 $\pm$ 1.7	0.70 $\pm$ 0.24	23 $\pm$ 7	41.7	11.3	3.7
6	200		18	3	5.10	12.7			[41.9] ^b	[10.6] ^b	
7	2	(0–<5)	20	2	0.29 $\pm$ 0.02	1.46 $\pm$ 0.16	0.13 $\pm$ 0.01	15 $\pm$ 1	40.7	10.0	4.1
8	40	(0–80)	13	28–40	5.00	10.5	0.85	15	40.4	10.8	3.7
9	1	surface	28	10	1.82 $\pm$ 0.66	5.88 $\pm$ 1.35	0.79 $\pm$ 0.48	11 $\pm$ 4	40.9	11.4	3.6
10a	2	(0–<5)	13	6	4.97 $\pm$ 1.71	6.03 $\pm$ 1.34	0.35 $\pm$ 0.12	23 $\pm$ 7	41.4	10.9	3.8
10b	100	(0–200)	5	39	12.8 $\pm$ 5.4	9.65 $\pm$ 3.82	0.18 $\pm$ 0.16	142 $\pm$ 148 (median: 90)	35.8	9.6	3.7
10c	100	(0–200)	5	19	11.5 $\pm$ 4.8	8.59 $\pm$ 3.10	0.15 $\pm$ 0.12	137 $\pm$ 138 (median: 76)	34.8	9.2	3.8
11a	35	(0–70)	–1.1	58	86.3 $\pm$ 57.8	35.5 $\pm$ 22.1	2.79 $\pm$ 1.56	17 $\pm$ 8	44.5	10.2	4.4
11b	3		–1.7	24	40.3 $\pm$ 11.1	20.0 $\pm$ 4.9	0.83 $\pm$ 0.42	34 $\pm$ 10	40.8	11.9	3.4
11c	35	(0–70)	–1.1	19	247.1 $\pm$ 96.7	94.2 $\pm$ 30.0	7.44 $\pm$ 3.48	17 $\pm$ 4	46.9	10.0	4.7
11d	3		–1.7	23	80.5 $\pm$ 30.1	34.6 $\pm$ 13.3	1.66 $\pm$ 0.93	29 $\pm$ 13	40.3	11.9	3.4
11e	3		–1.7	25	97.7 $\pm$ 36.1	42.8 $\pm$ 18.0	1.83 $\pm$ 0.82	32 $\pm$ 12	40.4	11.7	3.5
12	2	(0–<5)	27	9	0.77 $\pm$ 0.48	3.40 $\pm$ 2.51	0.31 $\pm$ 0.18	14 $\pm$ 5	40.2	10.1	4.0
13a	100	(0–200)	–0.8	15	23.0 $\pm$ 3.03	8.35 $\pm$ 1.43	0.86 $\pm$ 0.20	12 $\pm$ 2	41.2	11.6	3.5
13b	600	(200–1000)	0	9	82.8 $\pm$ 13.0	25.2 $\pm$ 6.3	1.27 $\pm$ 0.63	29 $\pm$ 12	47.3	10.7	4.4
14a	250	(0–500)	4	3–4	74.1	37.8	1.13	42	[35.0] ^c	10.3	
14b	300	(0–600)	13	27	32.5 $\pm$ 2.7	41.3 $\pm$ 7.2	[11.04 $\pm$ 0.51] ^d	[4.7] ^d	39.4	11.5	3.4
14c	50		4	6–10	45.8	17.1	1.36	16	[37.2] ^e	[10.9] ^e	
15	2		15	21–31	2.00	4.71	0.27	22	43.5	11.6	3.8
16	750	(500–1000)	3	2	9.17 $\pm$ 0.23	4.07 $\pm$ 0.62			43.8	10.3	4.2
17	50	(0–100)	1	19	17.3 $\pm$ 16.2	7.64 $\pm$ 5.91	0.40 $\pm$ 0.44	44 $\pm$ 27	51.3	8.9	5.7
18a	2	(0–<5)	–0.4	27	0.47 $\pm$ 0.14	0.43 $\pm$ 0.13	0.026 $\pm$ 0.015	26 $\pm$ 16	42.6	11.2	3.8
18b	2	(0–<5)	–1.7	3	5.49 $\pm$ 0.36	4.46 $\pm$ 1.34	0.21 $\pm$ 0.08	27 $\pm$ 2	40.7	9.9	4.1
18c	300	(0–<1000)	–0.5	20	6.89 $\pm$ 2.32	3.21 $\pm$ 1.08	0.20 $\pm$ 0.10	22 $\pm$ 8	43.1	12.0	3.6
19	25		4	17	10.0	12.9			[41.9] ^b	[10.6] ^b	
20	15	(0–30)	20	9–13	5.00	8.59	0.33	32	[41.9] ^b	[10.6] ^b	
21a	100	(0–200)	5	34	12.2 $\pm$ 4.5	12.2 $\pm$ 4.5	0.40 $\pm$ 0.37	71 $\pm$ 82 (median: 42)	36.6	9.8	3.7
21b	100	(0–200)	5	32	7.02 $\pm$ 3.29	6.56 $\pm$ 2.04	0.24 $\pm$ 0.13	46 $\pm$ 27 (median: 35)	37.1	10.1	3.7
22	650	(100–1200)	5	2	1405	109.3 $\pm$ 28.3			[41.9] ^b	[10.6] ^b	
23	500	(0–1000)	7	5	46.1	15.0	3.3		[41.9] ^b	[10.6] ^b	
24	1	surface	26	6	12.8 $\pm$ 3.5	32.3 $\pm$ 7.5	1.51 $\pm$ 0.76	31 $\pm$ 13	42.7	12.7	3.4

^a after Torress and Childress (1985)

^b grand mean values of 19 euphausiids in this table on which C and N compositions are determined; C = 41.9% ($\pm$ 3.8) of DM, N = 10.6% ($\pm$ 1.2) of DM.

^c calculated from C:N ratio given by Mayzaud (1973)

^d assumed as an artifact (cf. Mayzaud and Conover 1988) and excluded in the present analysis

^e estimated from Conover and Corner (1968) and Mayzaud (1973)

Table 3. Stepwise (forward selection) multiple regression statistics of theoretical and empirical models of respiration rates (Y : $\mu\text{l O}_2 \text{ ind.}^{-1}\text{h}^{-1}$) or ammonia excretion rates (Y : $\mu\text{gN ind.}^{-1}\text{h}^{-1}$) of euphausiids on body mass (X_1 : mg ind.^{-1}), habitat temperature (X_2 : $1000/\text{K}$ for the former, $^{\circ}\text{C}$ for the latter) and depth sampled (X_3 : m).

Regression model	Body mass unit	N	Step No.	Regression equation: $\ln Y = a_0 + a_1 \ln X_1 + a_2 X_2 + a_3 \ln X_3$				R^2 (adjusted R^2)	
				a_0	a_1	a_2	a_3		
Oxygen consumption									
Theoretical	DM	39	1		0.75	-4.628		0.628	
			2	14.017	0.75	-3.717	-0.109	0.735 (0.720)	
	C	39	1		0.75	-4.708		0.624	
			2	14.971	0.75	-3.798	-0.109	0.727 (0.712)	
	N	39	1		0.75	-4.530		0.679	
			2	15.992	0.75	-3.818	-0.085	0.753 (0.740)	
	Empirical	DM	39	1		0.517			0.680
				2		0.690	0.051		0.837
				3	0.392	0.753	0.046	-0.107	0.877 (0.867)
C		39	1		0.510			0.672	
			2		0.684	0.052		0.830	
			3	1.060	0.745	0.046	-0.105	0.868 (0.857)	
N		39	1		0.533			0.713	
			2		0.707	0.052		0.874	
			3	2.002	0.749	0.047	-0.082	0.899 (0.890)	
Ammonia excretion									
Theoretical	DM	31	1		0.75	-5.346		0.633	
			2	13.061	0.75	-4.248	-0.141	0.710 (0.690)	
	C	31	1		0.75	-5.400		0.657	
			2	14.142	0.75	-4.371	-0.132	0.726 (0.706)	
	N	31	1		0.75	-5.293		0.668	
			2	15.240	0.75	-4.407	-0.114	0.723 (0.703)	
	Empirical	DM	31	1		0.464			0.498
				2		0.729	0.064		0.753
				3	-2.606	0.788	0.056	-0.145	0.803 (0.782)
C		31	1		0.466			0.508	
			2		0.736	0.066		0.772	
			3	-1.940	0.788	0.057	-0.136	0.817 (0.796)	
N		31	1		0.479			0.525	
			2		0.750	0.066		0.792	
			3	-0.927	0.790	0.058	-0.117	0.826 (0.806)	

Table 4. Temperature effects on respiration rates of various zooplankton taxa and euphausiids in terms of activation energy (E_a) of Arrhenius equation and Q_{10} of Van't Hoff rule. Values are based on DM as a measure of body mass, and those in parentheses denote the range of 95% CI.

Data set type/zooplankton taxa	E_a (eV)	Temp range (°C)	Q_{10}	Temp range (°C)	Reference
Global/interspecific					
Euphausiids (24 species)	0.35 (0.28–0.42)	–1.7 to 28	1.7 (1.5–1.9)	–1.7 to 28	This study
Copepods (108 species)	0.45 (0.36–0.54)	–1.7 to 28.5	1.9 (1.7–2.2)	–1.7 to 28.5	Ikeda et al (2007)
Chaetognaths (17 species)	0.38 (0.18–0.58)	–1 to 28	1.7 (1.4–2.4)	–1 to 28	Ikeda & Takahashi (2012)
Chaetognaths (14 species)	0.59	0–30			Kruse et al. (2010)
Teleost fishes (69 species)	0.43	–2 to 40	1.8	0–30	Clarke & Johnson (1999)
Crustaceans (many species**)	0.57*	0–30	2.1–2.4	0–30	Ivleva (1980)
Regional/intraspecific					
<i>Euphausia superba</i>	0.67*	0–5	2.8	0–5	Hirsche (1984)
<i>Meganyctiphanes norvegica</i>	0.63*	0–5	2.6	0–5	
<i>Euphausia pacifica</i>			2.2	1–20	Iguchi & Ikeda (1995)
<i>Euphausia pacifica</i>			2.4	5–15	Paranjape (1967)
<i>Euphausia pacifica</i>			2.8	5–15	Small & Hebard (1967)
<i>Euphausia americana</i>			2.3	15–25	Teal & Carey (1967)
<i>Euphausia hemigibba</i>			2.8	10–25	
<i>Thysanopoda monacantha</i>			2.5	10–20	
<i>Thysanopoda obtusifrons</i>			2.5	10–20	
<i>Thysanopoda tricuspidata</i>			3.5	10–20	
<i>Meganyctiphanes norvegica</i>			2.0	5–10	

*converted from μ (kcal mole^{–1}) by an equation $E_a = \mu/23.06$ (cf. Brown et al. 2004).

**some benthic species included

Table 5. Global-bathymetric comparisons of ecological and physiological characteristics of euphausiids, copepods and chaetognaths living in the world's oceans. For respiration rate, T and E denote Theoretical and Empirical models, respectively. N is the number of data and N_{sp} the number of species.

Ecological/physiological characteristics		Euphausiids	Copepods	Chaetognaths
Behavior in the water column		active swimming	suspended, up-and-down or continuous gliding	suspended
Food habit		Herbivore, omnivore, carnivore	Herbivore, omnivore, carnivore	Carnivore
Respiration rate ($\mu\text{l O}_2 \text{ ind.}^{-1} \text{ h}^{-1}$)				
A specimen of 1 mgN body mass inhabiting 100 m depth (20°C, O ₂ saturation = 1.0)	T-model	13.1 ^a	10.2 ^b	5.1 ^c
	E-model	13.0 ^a	12.3 ^b	7.9 ^c
A specimen of 1 mgN body mass inhabiting 1000 m depth (5°C, O ₂ saturation = 0.1)	T-model	5.3 ^a	2.2 ^b	1.3 ^c
	E-model	5.3 ^a	2.5 ^b	1.5 ^c
Metabolic O:N ratio (by atoms)	Range	11–142 ^a	4.8–49 ^d	6.8–36 ^e
	Mean ($\pm$ SD)	30.0 (17.4)	20.7(11.3)	15.6(8.9)
	Median	27.1	16.9	12.2
	N (N _{sp})	31(19)	37(29)	12(10)
Body water content (% of WM)	Range	62.6–82.3 ^f	78.3–83.8 ^g	83.7–94.7 ^c
	Mean ($\pm$ SD)	76.9 (3.7)		90.4 (3.0)
	N (N _{sp})	36(27)	93(93)	19(14)
Body C:N (by mass)	Range	3.4–8.6 ^f	3.7–9.4 ^g	2.6–5.1 ^c
	Mean ($\pm$ SD)	4.2 (1.1)		4.0 (0.7)
	N (N _{sp})	41(26)	94(94)	30(24)

^a calculated from the equations in Table 3 of this study
^b calculated from the equations in Ikeda et al. (2007). Note that ambient oxygen saturation is a parameter for the model of copepods, but not for those of chaetognaths and euphausiids
^c calculated from the equations in Ikeda and Takahashi (2012)
^d from Ikeda et al. (2001)
^e from literatures listed in Table 2 of Ikeda and Takahashi (2012) in which both respiration and ammonia excretion rates determined simultaneously
^f from Table 2 of this study supplemented by those of Nemoto et al. (1972), Lindley et al. (1999), Morris and Hopkins (1983) and Harvey et al. (2012)
^g from Båmstedt (1986)

Ikeda Fig. 1

609
610
611
612
613
614

Ikeda Fig. 2

615
616
617
618
619
620
621
622

Ikeda Fig. 3

638

639

Ikeda Fig. 4

640

641

642

643

644