



| | |
|------------------------|---|
| Title | Equine major histocompatibility complex class I molecules act as entry receptors that bind to equine herpesvirus-1 glycoprotein D. |
| Author(s) | Sasaki, Michihito; Hasebe, Rie; Makino, Yoshinori; Suzuki, Tadaki; Fukushi, Hideto; Okamoto, Minoru; Matsuda, Kazuya; Taniyama, Hiroyuki; Sawa, Hirofumi; Kimura, Takashi |
| Citation | Genes to Cells, 16(4), 343-357 https://doi.org/10.1111/j.1365-2443.2011.01491.x |
| Issue Date | 2011-04 |
| Doc URL | http://hdl.handle.net/2115/59847 |
| Rights(URL) | https://creativecommons.org/licenses/by-nc-sa/4.0/deed.ja |
| Type | article |
| Additional Information | There are other files related to this item in HUSCAP. Check the above URL. |
| File Information | GTC_1491_sm_FigureS1-S2.pdf |



[Instructions for use](#)

1 **FIGURE LEGENDS OF SUPPORTING INFORMATION**

2 **Figure S1.** Flow cytometric detection of MHC class I expression on E. Derm, EBMECs
3 and PBMC cells stained with anti-MHC class I antibody PT85A (red), H58A (blue),
4 B5C (orange), or isotype controls IgG2a and IgG2b (black and gray, respectively).

5

6 **Figure S2.** Cellular ATP levels of each cell line with or without the ATP depletion
7 treatment. RK13, 3T3-A68, CHO-K1 and E. Derm cells were incubated with ATP
8 depletion media (open circles) or DMEM control media (solid squares) for the indicated
9 time periods. Cellular ATP was measured with the CellTiter-Glo luminescent cell
10 viability assay. Error bars represent standard deviations.

11

Fig. S1




Fig. S2

