



Title	Heterogeneous Pt Catalysts for Direct Synthesis of Chemicals from Alcohols by Borrowing-Hydrogen and Acceptorless Dehydrogenation Reactions [an abstract of dissertation and a summary of dissertation review]
Author(s)	CHAUDHARI, CHANDAN SUBHASH
Citation	北海道大学. 博士(総合化学) 甲第12036号
Issue Date	2015-09-25
Doc URL	http://hdl.handle.net/2115/59993
Rights(URL)	http://creativecommons.org/licenses/by-nc-sa/2.1/jp/
Type	theses (doctoral - abstract and summary of review)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Chandan_Chaudhari_Subhash_abstract.pdf (論文内容の要旨)



[Instructions for use](#)

学 位 論 文 内 容 の 要 旨

博士の専攻分野の名称 博士(総合化学) チャンダン サブハッシュュ チャドハリ

学 位 論 文 題 名

Heterogeneous Pt Catalysts for Direct Synthesis of Chemicals from Alcohols by

Borrowing Hydrogen and Acceptorless Dehydrogenation Reactions

(不均一系白金触媒による水素移動・脱水素反応を利用したアルコールからの直接的化学
品合成)

The limited fossil fuel and awareness of green chemistry inspire researchers to develop new green methodology for the synthesis of chemicals with atom-efficiency and step-efficiency. The classical method for the synthesis of C-C and C-N bonds is based on activated substrate such as alkyl halide which produce large amount of salt wastes. With respect to green chemistry, the alcohols based borrowing hydrogen and acceptorless dehydrogenation methodology can be good option for the classical synthetic method of C-C and C-N bonds formation because this methodologies produce H₂O and/or H₂ as byproducts. This thesis focuses on direct C-C and C-N bonds formation using alcohols as green alkylating reagents via borrowing hydrogen and acceptorless dehydrogenation methodology. The objectives of this work is the development of efficient and reusable heterogeneous catalyst for the synthesis of chemicals under neutral conditions which include the preparation of various metals supported and Pt-loaded catalysts and characterization of catalysts using XANES, EXAFS and XPS spectra.

In chapter 2, I examined various metal loaded-Al₂O₃ and supported Pt catalysts for alkylation of 2-methylquinoline with benzyl alcohol under additive free condition. Among screened catalysts, Pt/Al₂O₃ was found to be most effective catalyst for the alkylation of 2-methylquinoline with benzyl alcohol. The catalyst was reusable and showed good to moderate yield of for the alkylation of 2-methylquinoline with various alcohols. Mechanistic study show that the reaction was driven by the borrowing hydrogen pathway which showed a sequence of dehydrogenation-condensation-hydrogenation reactions. This results demonstrate the first heterogeneous catalytic system for this reaction.

In chapter 3, I investigated various metal loaded-CeO₂ and supported Pt catalysts for alkylation of oxindole with alcohols under additive free condition. I found that Pt/CeO₂ was best catalyst for alkylation of oxindole with 1-octanol. The catalyst was reusable and showed good to

moderate yield of for the alkylation of oxindoles with various alcohols. Mechanistic study show that the reaction was driven by the borrowing hydrogen pathway which showed a sequence of dehydrogenation-condensation-hydrogenation reactions. This results demonstrate the first additive-free catalytic system for this reaction.

In chapter 4, a series of transition metal-loaded metal oxide catalysts examined for self-coupling of 2-octanol. Pt/CeO₂ showed the highest activity among various metal loaded-CeO₂ and supported Pt catalysts for self-coupling of 2-octanol. This results demonstrated that Pt/CeO₂ was the most effective catalyst for self-coupling of aliphatic alcohols. Pt/ CeO₂ was also effective for α -alkylation of methyl ketone.

In chapter 5, I demonstrated the first acceptorless method for synthesise of 2-quinazoline from 2-aminobenzylamine with primary alcohols or aldehydes under additive-free condition. CeO₂-supported Pt nanoparticle catalysts showed high activity among various metal supported and Pt loaded catalysts. I investigated the reusability of catalyst and general applicability of the present catalytic system. Mechanistic study show that the reaction was driven by the acceptorless dehydrogenation pathway which include dehydrogenation-condensation and cyclization-dehydrogenation steps.

In chapter 6, Pt/Al₂O₃ and Pt/TiO₂ were effective catalysts for the synthesis of 2-substituted benzothiazoles and benzimidazoles from 2-aminothiophenol and 1,2-phenylenediamine with alcohols or aldehydes under acceptor-free and additive-free conditions. With optimized Pt/TiO₂ and Pt/Al₂O₃ catalysts, I investigated the general applicability of present catalytic system using various alcohols for the synthesis of 2-substituted benzothiazole and 2-substituted benzimidazole respectively.

Chapter 7 is the general summary. Chapters 2-6 show the first systematic examples of heterogeneous catalysis for the synthesis of chemicals via alkylation of nucleophiles by alcohols under additive-free conditions. The borrowing hydrogen methodology include dehydrogenation-condensation-hydrogenation steps with elimination of water as byproduct. The acceptorless dehydrogenation methodology include dehydrogenation-condensation and cyclization-dehydrogenation steps with elimination of water and liberation of H₂ as byproduct.