Title	Isolation of a major genetic interaction associated with an extreme phenotype using assorted F2 populations in rice
Author(s)	Ota, Yuya; Ishiguro, Seiya; Aoyama, Eiko; Aiba, Ryosuke; Iwashiro, Reika; Tanabata, Takanari; Takamure, Itsuro; Fujino, Kaiken; Kishima, Yuji
Citation	Molecular Breeding, 33(4), 997-1003 https://doi.org/10.1007/s11032-013-9998-6
Issue Date	2014-04
Doc URL	http://hdl.handle.net/2115/60043
Rights	The final publication is available at Springer via http://dx.doi.org/10.1007/s11032-013-9998-6
Type	article (author version)
File Information	Ota et al Molucular Breeding MS.pdf

[Instructions for use](#)

Short communication

Isolation of a major genetic interaction associated with an extreme phenotype using assorted F2 populations in rice

**Yuya Ota¹, Seiya Ishiguro¹, Eiko Aoyama¹, Ryosuke Aiba¹, Reika Iwashiro¹,
Takanari Tanabata², Itsuro Takamure¹, Kaiken Fujino¹, Yuji Kishima¹**

¹ Research Faculty of Agriculture, Hokkaido University, Sapporo 060-8589, Japan

² National Institute of Agrobiological Sciences, Tsukuba, Ibaraki 305-8602, Japan

Present address of T Tanabata

RIKEN Center for Sustainable Resource Science, Tsukuba, Ibaraki 3050074, Japan

Corresponding author: Yuji Kishima. E-mail address: kishima@abs.agr.hokudai.ac.jp

Tel: +81-11-706-2439

Abstract

Detection of quantitative trait loci (QTLs) is dependent on materials used in the analysis, as different combinations of parental materials may lead to different outcomes in QTLs for the same trait. On the other hand, an extreme phenotype associated with a given trait implies the potential involvement of a particular allele in various allelic interactions. A genetic factor associated with such an extreme phenotype may frequently be identified from various genetic populations consisting of different parental combinations. In this study, we attempted to uncover the genetic factor associated with extremely early heading date in rice using various F2 populations. Heading date in rice has been characterized by at least 19 QTLs, from which 12 genes have been identified. A58, a rice strain with an extremely early heading date, is adapted to Hokkaido, the northernmost limit of rice cultivation. Six F2 populations derived from crosses of A58 with six other strains displayed a range of heading dates. Genotyping using 19 QTL markers indicated that the A58 allele of the *Ghd7* locus was present in most F2 individuals exhibiting extremely early heading dates. This analysis also demonstrated that when the wild-type *Ehd1* allele was present, the *Ghd7* allele from A58 accelerated floral induction. The results of this study demonstrate that assorted F2 populations are valuable materials for comprehensive genotyping to explore major genetic factors for extreme phenotypes, and that this methodology is broadly applicable to other unknown traits.

Keywords

Assorted F2 populations, Extreme phenotype, Heading date, *Ghd7*, *Ehd1*, Rice

Text

Analyses for quantitative trait loci (QTLs) have broadly and profoundly contributed to the discovery of a complex of genes with minor effects for a particular phenotypic trait (Borevitz and Chory 2004). An enormous number of QTLs have been identified using second filial generation (F₂) populations, recombinant inbred lines (RILs), backcross inbred lines (BILs), and chromosomal substitution lines in various crops (Yamamoto et al. 2009; Wurschum 2012). As technical development has boosted the ability to produce high-resolution genomic markers throughout the genome, QTL detection accuracy and frequency have greatly advanced (Ganal et al. 2009; Varshney et al. 2009). In rice, QTL analyses have generally employed RILs and BILs (Yamamoto et al. 2009). Unlike F₂ populations, RILs and BILs are genetically stable after establishment, allowing recurrent use under different conditions. Such populations, however, possess only the two parental alleles at each locus, limiting genetic variation to two entities. Consequently, a trait of interest examined using RILs or BILs is defined by their parental characteristics.

Ebana et al. (2011) successfully characterized known QTLs for rice heading date using multiple F₂ populations derived from different cross combinations. The different allele combinations resulted in different phenotypic variations for the corresponding trait in the F₂ populations. Their study caused us to consider whether common genetic factors can be extracted from individuals exhibiting phenotypes extremely different from the majority in various F₂ populations. Different F₂ populations are expected to show a wide range of phenotypic variations. If particular F₂ populations have distinct distributions of phenotypic variations, we might be able to explore the loci associated with such distinct phenotype using F₂ individuals and isolate major genetic factors.

To attempt isolation of major genetic factors from various F₂ populations, we focused on rice heading date, a well-characterized, important agricultural trait. Approximately two dozen QTLs (*Hd1-Hd17*, *Ehd1*, *Ehd2*, *Ehd3*, *Edh4*, and *Ghd7*) associated with heading date have been reported over roughly the last decade (Yano et al. 1997; Yamamoto et al. 2000; Lin et al. 2003; Takeuchi et al. 2003; Uga et al. 2007; Matsubara et al. 2008a; Matsubara et al. 2008b; Matsubara et al. 2011; Shibaya et al. 2011; Gao et al. 2013), and genes have been identified and characterized in 12 QTLs: *Hd1*, *Hd2*, *Hd3a*, *Hd5* (*Ghd8*), *Hd6*, *Hd16*, *Hd17* (*Hd3b*), *Ehd1*, *Ehd2*, *Ehd3*, *Ehd4*, and *Ghd7* (Yano et al. 2000; Takahashi et al. 2001; Kojima et al. 2002; Doi

et al. 2004; Matsubara et al. 2008b; Xue et al. 2008; Matsubara et al. 2011; Yan et al. 2011; Matsubara et al. 2012; Hori et al. 2013; Koo et al. 2013). Comprehensive gene identifications in a quantitative trait facilitate direct mapping of genes having unknown genetic contributions to their associated phenotypes.

An old rice strain, A58, is characterized by extremely early heading date, which thought to be required for adaptation to northernmost limits of cultivation such as Hokkaido (Baruah et al. 2011). In studies of F2 and BC1F1 populations between Hokkaido variety Hoshinoyume and Nipponbare (Fujino and Sekiguchi 2005b), alleles of both *Hd2* and *Ghd7* (= *Hd4*) were found to be prerequisite to conferral of extremely early heading date. Some other Hokkaido varieties showing extremely early heading also possess these recessive alleles (Fujino and Sekiguchi 2005a; Nonoue et al. 2008). We did not know, however, whether Hokkaido strain A58 also contained the same functional alleles possessed by the other Hokkaido varieties. In this study, we produced F1 seeds derived from six combinations, in which A58 as the seed parent was crossed with three japonica, Nipponbare, Koshihikari, and Taichung 65 (T65), and three indica strains, #108, IR36, and Kasalath. The F1 plants were cultivated in a greenhouse equipped with a short-day chamber at Hokkaido University (Sapporo, Japan) in 2012. To evaluate F2 populations, about 200 seeds from each F1 population were sown in nursery beds in the greenhouse, with 4-week-old seedlings, then transplanted into a paddy field at the university. Sowing and transplanting were performed on May 7 and June 4, 2012. Days to heading of the earliest heading panicle among individual panicles were recorded for each plant to represent the number of days required from sowing to heading.

Heading status was checked every 2 d, beginning in the middle of July, 2012. All six pollen parents showed later heading dates (114–126 d in Sapporo, Hokkaido, Japan, 43°N latitude) relative to A58 (81 d). The latest strain was IR36, which headed at 126 d. The six F2 populations derived from the cross of A58 with the six pollen parents showed different segregation patterns in terms of heading date (Supplementary Fig. S1). Based on earliest observed heading date compared with that of A58, F2 populations were classified into three groups: 1) those contained individuals earlier than A58 (F2 populations of A58/Kasalath [F2-A58/Kasalath] and F2-A58/#108); 2) the earliest individuals in those were comparable to A58 (F2-A58/Nipponbare and F2-A58/Koshihikari); and 3) those were later than A58 (F2-A58/IR36 and F2-A58/T65) (Supplementary Fig. S1).

We randomly selected 93 individuals from each F2 population (comprising about 200 individuals each), and performed genotyping using SSR markers (McCouch et al. 2002; Fujino and Sekiguchi 2005a; Matsubara et al. 2008a; Matsubara et al. 2008b) for 19 loci related to heading date (Table 1). Polymorphisms at each locus between A58 and the other strains were assessed by PCR. No polymorphisms were detected between A58 and japonica strains at 11 or 8 loci because of genetic similarities to A58. Between A58 and indica strains, polymorphisms were detected at all 19 loci (Table 1). Average heading dates were calculated for all individuals that were homozygous or heterozygous at each locus. The significance of correlations between heading date and alleles at each locus was assessed by ANOVA. The *Ghd7* locus was significantly associated with heading date in four F2 populations (F2-A58/Nipponbare, F2-A58/Kasalath, F2-A58/IR36, and F2-A58/#108) (Table 1), while a correlation with *Hd2* was detected in three (F2-A58/Nipponbare, F2-A58/T65, and F2-A58/IR36) (Table 1). Other significant correlations with heading date were found with *Hd1* (F2-A58/Nipponbare); *Hd6*, *Hd16*, and *Ehd1* (F2-A58/Kasalath); *Hd17* (F2-A58/IR36); and *Hd3* and *Hd5* (F2-A58/#108) (Table 1). Two-way ANOVA detected a strong interaction ($p < 0.01$) between *Ghd7* and *Ehd1* in F2-A58/Kasalath (Supplementary Table S1). We subsequently focused on 20 individuals from the four F2 populations (4 plants from F2-A58/Nipponbare, 2 plants from F2-A58/Koshihikari, 12 plants from F2-A58/Kasalath, and 2 plants from F2-A58/#108) that had earlier or identical heading dates to A58. These extreme individuals were genotyped at the 19 loci, and the resulting data were statistically evaluated for non-Mendelian distributions. As shown in Table 2, the distribution of *Ghd7* genotypes was clearly divergent from the expected Mendelian segregation ratio (1:2:1 for two alleles). Eighteen individuals were homozygous at *Ghd7* for the allele from A58, implying its contribution to their extremely early heading dates. In contrast, no contribution to early heading date was observed from the A58 allele at the *Ehd1* locus (Table 2), implying that the homozygous *Ehd1* alleles from A58 tended to inhibit early heading. In fact, a remarkable epistatic interaction ($P = 0.0044$) was detected between *Ghd7* (A58 allele) and *Ehd1* (Kasalath allele) in F2-A58/Kasalath as suggested by two-way ANOVA (Fig. 1). The trade-off interaction between both the wild-type alleles from *Ghd7* and *Ehd1* has actually been observed with respect to their expressions under long-day conditions (Matsubara et al. 2011). In the cited study, expression of *Ghd7* suppressed that of *Ehd1*, leading to delayed floral induction under long-day conditions. Sequence analysis has determined that A58 contains the same recessive *Ghd7* allele found in Hoshinoyume (Lu et al. 2012) (Supplementary Fig. S2),

1 and Doi et al. (2004) have reported that Kasalath contains a functional *Ehd1*. It has thus been
2 suggested that the combination of the A58 allele of *Ghd7* and the Kasalath allele of *Ehd1*
3 accelerates floral induction following basic vegetative growth (Tsuji et al. 2011). In our study,
4 this allele combination promoted a significantly shorter heading date than did A58 alone.

5
6 It is curious that when A58 was crossed with indica strains, some loci in the corresponding F2
7 populations did not segregate in a Mendelian manner irrespective to heading date, i.e., marked
8 genetic distortion occurred in *Hd8* and *Hd13* in F2-A58/IR36 and *Hd1* and *Ghd7* in
9 F2-A58/#108 (Table 1). Such genetic distortion has often been observed in hybrid progenies of
10 distantly related strains. Because of this genetic distortion, a large number of F2 individuals are
11 required to satisfy statistical testing.

12
13 The purpose of this pilot study was to demonstrate the use of assorted F2 populations, rather
14 than RILs or BILs, for rapid extraction of major genetic factors and/or interactions associated
15 with extreme phenotypes in complex QTLs. Heading date was a feasible trait with which to
16 search for functionally important genes: there are at least 19 heading date-related genes or loci
17 currently characterized, which allowed us to perform their genotyping directly. We successfully
18 detected one major genetic interaction associated with extremely early heading date from F2
19 populations derived from crosses between A58, a strain adapted to the northernmost rice
20 cultivation limit, and assorted other strains. The F2 populations examined here exhibited
21 different distributional patterns of heading date frequencies, implying that various combinations
22 of genetic factors were present. As shown in this study, individuals with extreme phenotypes in
23 assorted F2 populations are valuable materials for exploring the major genetic factor(s) involved
24 in such phenotypes. This approach has both strengths and weaknesses; the strengths are to select
25 the most effective one(s) among a broad range of the genetic interactions derived from multiple
26 alleles, and to accomplish isolation of such interaction(s) in a short period as F2 generation,
27 while the weakness is to required a number of F2 plants from the different cross combinations
28 to obtain the conclusive results. Combined with deciphered genomic information, this
29 methodology is broadly applicable to the investigation of various polygenic traits associated
30 with extreme phenotypes, such as biotic and abiotic stress tolerance and disease and insect
31 resistance.

Acknowledgments

We are especially grateful to Ms. Takako Takeuchi (Research Faculty of Agriculture, Hokkaido University) for technical assistance and Drs. Kaworu Ebana and Masahiro Yano (QTL Genomic Research Center, National Institute of Agrobiological Sciences) for technical advice regarding the field data recording system. This research was supported by funding from the Scientific Technique Research Promotion Program for Agriculture, Forestry, Fisheries and Food Industry.

References

- Baruah AR, Onishi K, Oguma Y, Ishigo-Oka N, Uwatoko N, Sano Y (2011) Effects of acclimation on chilling tolerance in Asian cultivated and wild rice. *Euphytica* 181:293-303.
- Borevitz JO, Chory J (2004) Genomics tools for QTL analysis and gene discovery. *Curr Opin Plant Biol* 7:132-136.
- Doi K, Izawa T, Fuse T, Yamanouchi U, Kubo T, Shimatani Z, Yano M, Yoshimura A (2004) Ehd1, a B-type response regulator in rice, confers short-day promotion of flowering and controls FT-like gene expression independently of Hd11. *Gene Dev* 18:926-936.
- Ebana K, Shibaya T, Wu JZ, Matsubara K, Kanamori H, Yamane H, Yamanouchi U, Mizubayashi T, Kono I, Shomura A, Ito S, Ando T, Hori K, Matsumoto T, Yano M (2011) Uncovering of major genetic factors generating naturally occurring variation in heading date among Asian rice cultivars. *Theor Appl Genet* 122:1199-1210.
- Fujino K, Sekiguchi H (2005a) Identification of QTLs conferring genetic variation for heading date among rice varieties at the northern-limit of rice cultivation. *Breeding Sci* 55:141-146.
- Fujino K, Sekiguchi H (2005b) Mapping of QTLs conferring extremely early heading in rice (*Oryza sativa* L.). *Theor Appl Genet* 111:393-398.
- Ganal MW, Altmann T, Roder MS (2009) SNP identification in crop plants. *Curr Opin Plant Biol* 12:211-217.
- Gao H, Zheng XM, Fei GL, Chen J, Jin MN, Ren YL, Wu WX, Zhou KN, Sheng PK, Zhou F, Jiang L, Wang J, Zhang X, Guo XP, Wang JL, Cheng ZJ, Wu CY, Wang HY, Wan JM (2013) Ehd4 encodes a novel and *Oryza*-genus-specific regulator of photoperiodic flowering in rice. *Plos Genet* 9.

1 Hori K, Ogiso-Tanaka E, Matsubara K, Yamanouchi U, Ebana K, Yano M (2013) Hd16, a gene
2 for casein kinase I, is involved in the control of rice flowering time by modulating the
3 day-length response. *Plant J*.

4 Kojima S, Takahashi Y, Kobayashi Y, Monna L, Sasaki T, Araki T, Yano M (2002) Hd3a, a
5 rice ortholog of the Arabidopsis FT gene, promotes transition to flowering downstream
6 of Hd1 under short-day conditions. *Plant Cell Physiol* 43:1096-1105.

7 Koo BH, Yoo SC, Park JW, Kwon CT, Lee BD, An G, Zhang Z, Li J, Li Z, Paek NC (2013)
8 Natural variation in OsPRR37 regulates heading date and contributes to rice cultivation
9 at a wide range of latitudes. *Mol Plant*.

10 Lin HX, Liang ZW, Sasaki T, Yano M (2003) Fine mapping and characterization of quantitative
11 trait loci Hd4 and Hd5 controlling heading date in rice. *Breeding Sci* 53:51-59.

12 Lu L, Yan WH, Xue WY, Shao D, Xing YZ (2012) Evolution and association analysis of Ghd7
13 in rice. *Plos One* 7.

14 Matsubara K, Kono I, Hori K, Nonoue Y, Ono N, Shomura A, Mizubayashi T, Yamamoto S,
15 Yamanouchi U, Shirasawa K, Nishio T, Yano M (2008a) Novel QTLs for photoperiodic
16 flowering revealed by using reciprocal backcross inbred lines from crosses between
17 japonica rice cultivars. *Theor Appl Genet* 117:935-945.

18 Matsubara K, Ogiso-Tanaka E, Hori K, Ebana K, Ando T, Yano M (2012) Natural variation in
19 Hd17, a homolog of Arabidopsis ELF3 that is involved in rice photoperiodic flowering.
20 *Plant Cell Physiol* 53:709-716.

21 Matsubara K, Yamanouchi U, Nonoue Y, Sugimoto K, Wang ZX, Minobe Y, Yano M (2011)
22 Ehd3, encoding a plant homeodomain finger-containing protein, is a critical promoter of
23 rice flowering. *Plant J* 66:603-612.

24 Matsubara K, Yamanouchi U, Wang ZX, Minobe Y, Izawa T, Yano M (2008b) Ehd2, a rice
25 ortholog of the maize INDETERMINATE1 gene, promotes flowering by up-regulating
26 Ehd1. *Plant Physiol* 148:1425-1435.

27 McCouch SR, Teytelman L, Xu YB, Lobos KB, Clare K, Walton M, Fu BY, Maghirang R, Li
28 ZK, Xing YZ, Zhang QF, Kono I, Yano M, Fjellstrom R, DeClerck G, Schneider D,
29 Cartinhour S, Ware D, Stein L (2002) Development and mapping of 2240 new SSR
30 markers for rice (*Oryza sativa* L.). *DNA Res* 9:199-207.

31 Nonoue Y, Fujino K, Hirayama Y, Yamanouchi U, Lin SY, Yano M (2008) Detection of
32 quantitative trait loci controlling extremely early heading in rice. *Theor Appl Genet*
33 116:715-722.

Shibaya T, Nonoue Y, Ono N, Yamanouchi U, Hori K, Yano M (2011) Genetic interactions involved in the inhibition of heading by heading date QTL, Hd2 in rice under long-day conditions. *Theor Appl Genet* 123:1133-1143.

Takahashi Y, Shomura A, Sasaki T, Yano M (2001) Hd6, a rice quantitative trait locus involved in photoperiod sensitivity, encodes the alpha subunit of protein kinase CK2. *Proc Natl Acad Sci USA* 98:7922-7927.

Takeuchi Y, Lin SY, Sasaki T, Yano M (2003) Fine linkage mapping enables dissection of closely linked quantitative trait loci for seed dormancy and heading in rice. *Theor Appl Genet* 107:1174-1180.

Tsuji H, Taoka K, Shimamoto K (2011) Regulation of flowering in rice: two florigen genes, a complex gene network, and natural variation. *Curr Opin Plant Biol* 14:45-52.

Uga Y, Nonoue Y, Liang ZW, Lin HX, Yamamoto S, Yamanouchi U, Yano M (2007) Accumulation of additive effects generates a strong photoperiod sensitivity in the extremely late-heading rice cultivar 'Nona Bokra'. *Theor Appl Genet* 114:1457-1466.

Varshney RK, Nayak SN, May GD, Jackson SA (2009) Next-generation sequencing technologies and their implications for crop genetics and breeding. *Trends Biotechnol* 27:522-530.

Wurschum T (2012) Mapping QTL for agronomic traits in breeding populations. *Theor Appl Genet* 125:201-210.

Xue WY, Xing YZ, Weng XY, Zhao Y, Tang WJ, Wang L, Zhou HJ, Yu SB, Xu CG, Li XH, Zhang QF (2008) Natural variation in Ghd7 is an important regulator of heading date and yield potential in rice. *Nat Genet* 40:761-767.

Yamamoto T, Lin HX, Sasaki T, Yano M (2000) Identification of heading date quantitative trait locus Hd6 and characterization of its epistatic interactions with Hd2 in rice using advanced backcross progeny. *Genetics* 154:885-891.

Yamamoto T, Yonemaru J, Yano M (2009) Towards the understanding of complex traits in rice: substantially or superficially? *DNA Res* 16:141-154.

Yan WH, Wang P, Chen HX, Zhou HJ, Li QP, Wang CR, Ding ZH, Zhang YS, Yu SB, Xing YZ, Zhang QF (2011) A major QTL, Ghd8, plays pleiotropic roles in regulating grain productivity, plant height, and heading date in rice. *Mol Plant* 4:319-330.

Yano M, Harushima Y, Nagamura Y, Kurata N, Minobe Y, Sasaki T (1997) Identification of quantitative trait loci controlling heading date in rice using a high-density linkage map. *Theor Appl Genet* 95:1025-1032.

Yano M, Katayose Y, Ashikari M, Yamanouchi U, Monna L, Fuse T, Baba T, Yamamoto K, Umehara Y, Nagamura Y, Sasaki T (2000) Hd1, a major photoperiod sensitivity quantitative trait locus in rice, is closely related to the arabidopsis flowering time gene CONSTANS. Plant Cell 12:2473-2483.

Figure legends

Fig. 1

Effect of genetic interactions between *Ghd7* and *Ehd1* on heading date in an F2-A58/Kasalath population. Combinations of the two loci, *Ghd7* and *Ehd1*, from A58 and Kasalath were either homozygous (A58 and Kasalath) or heterozygous (Hetero). Labels below bars refer to *Ghd7* genotypes; bar colors refer to *Ehd1* genotypes (white: A58 homozygous; gray: heterozygous; black: Kasalath homozygous). Numbers above bars correspond to number of individuals possessing the genotype combination. Error bars indicate standard errors.

Supplementary Fig. S1

Frequency and distribution of days to heading in six F2 populations. (a) F2-A58/Nipponbare ($n = 209$); (b) F2-A58/Koshihikari ($n = 212$); (c) F2-A58/T65 ($n = 195$); (d) F2-A58/Kasalath ($n = 213$); (e) F2-A58/IR36 ($n = 194$); (f) F2-A58/#108 ($n = 238$). Arrowheads indicate parental mean heading dates.

Supplementary Fig. S2

Amino acid variation in the Ghd7 protein deduced from nucleotide sequences in the seven studied rice strains. Positions correspond to those of the 257 amino acids of the Nipponbare Ghd7 polypeptide. All variable positions characterized by Lu et al. (2012) are noted. A nonsense mutation is present in the *Ghd7* allele of A58; mutations occurring in the other strains most likely do not seriously impair proper function. The CCT (CONSTANS, CO-like, and TOC1) domain is a highly conserved basic module of ~43 amino acids, and is characteristic of the Ghd7 polypeptide.

Table 1 Summary of genotyping analysis for heading date using 19 markers in the six F2 populations

F2 population (n=93) ^a	Target gene or QTL	Marker ^b	Mean of DTH ^c			Probability
			A (n)	B (n)	H (n)	
A58/Nipponbare	<i>Hd1</i>	Hd1	97 (21)	109 (25)	101.3 (47)	0.0002*
	<i>Hd2</i>	RM1306	96 (22)	110 (23)	102 (48)	0.0000*
	<i>Ghd7</i>	RM21327	95 (29)	106 (19)	105 (45)	0.0000*
	<i>Hd6</i>	RM15956	97 (17)	103 (27)	104 (49)	0.0794
	<i>Hd10</i>	RM17071	104 (25)	100 (24)	102 (44)	0.3821
	<i>Hd11</i>	RM16278	101 (28)	104 (21)	102 (44)	0.5199
	<i>Hd12</i>	RM23471	103 (30)	102 (22)	102 (41)	0.9686
	<i>Hd16</i>	87C10-17	98 (22)	104 (23)	103 (48)	0.0636
A58/Koshihikari	<i>Hd1</i>	Hd1	105 (21)	105 (31)	103 (41)	0.7492
	<i>Hd2</i>	RM1306	103 (15)	105 (21)	104 (57)	0.8553
	<i>Ghd7</i>	RM21327	100 (24)	104 (16)	107 (53)	0.0668
	<i>Hd6</i>	RM15956	106 (22)	106 (29)	102 (42)	0.3170
	<i>Hd10</i>	RM17071	108 (22)	104 (24)	103 (47)	0.2634
	<i>Hd11</i>	RM16278	108 (27)	105 (24)	102 (42)	0.0917
	<i>Hd12</i>	RM23471	105 (24)	105 (20)	104 (49)	0.8544
	<i>Hd13</i>	RM27727	105 (21)	105 (24)	104 (48)	0.9315
	<i>Hd16</i>	RM1038	105 (23)	107 (24)	103 (46)	0.3046
	<i>Hd17</i>	P548D347	104 (26)	103 (19)	105 (48)	0.8179
A58/T65	<i>Hd1</i>	Hd1	105 (22)	100 (31)	101 (40)	0.0145
	<i>Hd2</i>	RM1306	98 (25)	107 (27)	101 (41)	0.0000*
	<i>Ghd7</i>	RM7110	104 (20)	100 (20)	102 (53)	0.1167
	<i>Hd7</i>	RM3857	103 (26)	102 (25)	102 (42)	0.6281
	<i>Hd11</i>	RM16278	101.4 (25)	102.3 (21)	102.1 (47)	0.8582
	<i>Hd12</i>	RM23471	101 (23)	101 (24)	103 (46)	0.1357
	<i>Hd16</i>	87C10-17	102 (18)	103 (22)	102 (53)	0.5486
	<i>Ehd1</i>	RM25539	99 (18)	104 (30)	102 (45)	0.0263
A58/Kasalath	<i>Hd1</i>	Hd1	104 (14)	98 (37)	100 (42)	0.1366
	<i>Hd2</i>	RM1306	99 (25)	104 (26)	97 (42)	0.0163
	<i>Hd3</i>	RM204	106 (16)	97 (37)	100 (40)	0.0099*
	<i>Ghd7</i>	RM5499	87 (23)	108 (26)	101 (44)	0.0000*
	<i>Hd5</i>	GBR8001	95 (23)	99 (23)	102 (47)	0.0253
	<i>Hd6</i>	RM15956	94 (25)	100 (17)	102 (51)	0.0061*
	<i>Hd7</i>	RM8030	99 (15)	100 (29)	100 (49)	0.9222
	<i>Hd8</i>	RM5442	94 (15)	101 (31)	100 (47)	0.0602
	<i>Hd9</i>	GBR3002	93 (18)	102 (31)	100 (44)	0.0190
	<i>Hd10</i>	RM17067	99 (31)	104 (19)	98 (43)	0.1464
	<i>Hd11</i>	RM16278	101 (17)	101 (19)	99 (57)	0.7347
	<i>Hd12</i>	RM23471	101 (30)	100 (27)	98 (36)	0.5581
	<i>Hd13</i>	RM27727	97 (28)	102 (17)	100 (48)	0.3564
	<i>Hd16</i>	RM1038	93 (25)	101 (20)	102 (48)	0.0012*
	<i>Hd17</i>	P548D347	104 (18)	96 (36)	100 (39)	0.0328
	<i>Ehd1</i>	RM25539	106 (24)	96 (29)	98 (40)	0.0006*
	<i>Ehd2</i>	SSR-1	104 (22)	96 (31)	99 (40)	0.0164
	<i>Ehd3</i>	RM1381	99 (17)	99 (31)	100 (45)	0.9710
	<i>Ehd4</i>	RM6349	94 (16)	102 (34)	99 (43)	0.0466

A58/IR36	<i>Hd1</i>	Hd1	116 (19)	114 (25)	116 (47)	0.6730
	<i>Hd2</i>	RM1306	110 (23)	120 (21)	116 (47)	0.0008*
	<i>Hd3</i>	RM204	113 (28)	120 (21)	115 (42)	0.0115
	<i>Ghd7</i>	RM5499	110 (14)	119 (31)	115 (46)	0.0090*
	<i>Hd5</i>	GBR8001	117 (33)	112 (17)	116 (41)	0.1248
	<i>Hd6</i>	RM15956	117 (19)	117 (27)	114 (45)	0.4337
	<i>Hd7</i>	RM8030	114 (23)	117 (28)	116 (40)	0.3218
	<i>Hd8</i>	RM5442	107 (4)	114 (40)	118 (47)	0.0176
	<i>Hd9</i>	GBR3002	115 (18)	116 (21)	116 (52)	0.8517
	<i>Hd10</i>	RM17071	115 (19)	115 (27)	116 (45)	0.8487
	<i>Hd11</i>	RM16278	116 (17)	117 (32)	114 (42)	0.3102
	<i>Hd12</i>	RM23471	119 (25)	116 (21)	114 (45)	0.0559
	<i>Hd13</i>	RM27727	119 (7)	114 (37)	117 (47)	0.2296
	<i>Hd16</i>	RM1038	117 (17)	119 (23)	114 (51)	0.1019
	<i>Hd17</i>	0007o20	114 (33)	123 (19)	114 (39)	0.0006*
	<i>Ehd1</i>	RM25539	119 (16)	116 (26)	115 (49)	0.2862
	<i>Ehd2</i>	SSR-1	118 (18)	116 (25)	115 (48)	0.3162
	<i>Ehd3</i>	RM1381	115 (20)	117 (23)	115 (48)	0.7285
	<i>Ehd4</i>	RM6349	115 (24)	116 (17)	115 (50)	0.8740
A58/#108	<i>Hd1</i>	Hd1	97 (8)	109 (43)	104 (42)	0.0130
	<i>Hd2</i>	RM1306	104 (28)	105 (23)	107 (42)	0.5110
	<i>Hd3</i>	RM204	101 (22)	111 (29)	104 (42)	0.0044*
	<i>Ghd7</i>	RM5499	87 (9)	112 (43)	103 (41)	0.0000*
	<i>Hd5</i>	GBR8001	110 (21)	96 (25)	109 (47)	0.0000*
	<i>Hd6</i>	RM15956	103 (30)	109 (26)	105 (37)	0.2099
	<i>Hd7</i>	RM8030	106 (24)	106 (20)	105 (49)	0.9150
	<i>Hd8</i>	RM5442	104 (10)	106 (51)	105 (32)	0.8096
	<i>Hd9</i>	GBR3002	103 (17)	109 (39)	103 (37)	0.0452
	<i>Hd10</i>	RM17071	105 (31)	105 (26)	106 (36)	0.8216
	<i>Hd11</i>	RM16278	102 (19)	108 (28)	105 (46)	0.2900
	<i>Hd12</i>	RM23471	109 (18)	107 (25)	104 (50)	0.1949
	<i>Hd13</i>	RM27727	106 (19)	103 (26)	106 (48)	0.5218
	<i>Hd16</i>	RM1373	104 (27)	108 (26)	105 (40)	0.3680
	<i>Hd17</i>	P548D347	102 (24)	110 (22)	105 (47)	0.0342
	<i>Ehd1</i>	RM25539	110 (24)	106 (30)	102 (39)	0.0206
	<i>Ehd2</i>	SSR-1	108 (23)	109 (29)	107 (41)	0.1131
	<i>Ehd3</i>	RM1381	110 (22)	106 (31)	103 (40)	0.0836
	<i>Ehd4</i>	RM6349	104 (19)	108 (32)	105 (42)	0.4207

a: Paternal strains (A/B=femal/male), 93 individuals randomly selected from each of the F2 populations were examined; b: SSR markers were referred from following references; Fujino and Sekiguchi (2005a) for Hd1, GBR3002, GBR8001, McCouch et al. (2002) for all the markers with RM numbers, Matsubara et al. (2008a) for P548D347, 0007o20, Matsubara et al. (2008b) for SSR-1; c: DTH (days to heading), A represents homozygote of A58 allele, B is homozygote of the pollen parental allele, and H is heterozygote; * indicates a significance level of 1% as determined by ANOVA.

Table 2 Number of individuals expressing the extremely early heading trait categorized by genotype for 19 loci^a

Genotype/locus	<i>Hd1</i>	<i>Hd2</i>	<i>Hd3</i>	<i>Ghd7</i>	<i>Hd5</i>	<i>Hd6</i>	<i>Hd7</i>	<i>Hd8</i>	<i>Hd9</i>	<i>Hd10</i>	<i>Hd11</i>	<i>Hd12</i>	<i>Hd13</i>	<i>Hd16</i>	<i>Hd17</i>	<i>Ehd1</i>	<i>Ehd2</i>	<i>Ehd3</i>	<i>Ehd4</i>
Homozygous for A58	2	8	2	18	7	7	3	4	6	5	4	2	6	7	3	0	2	3	6
Heterozygous	8	11	5	1	5	8	7	7	6	12	13	12	7	8	5	8	5	9	6
Homozygous for pollen parent	10	1	7	1	2	5	4	3	2	3	3	6	3	5	8	6	7	2	2
No. of examined plants	20	20	14	20	14	20	14	14	14	20	20	20	16	20	16	14	14	14	14
Fit to Mendelian segregation																			
ratio ^b	0.027	0.078	0.095	2E-10	0.095	0.55	0.93	0.93	0.28	0.55	0.39	0.3	0.5	0.55	0.068	0.066	0.095	0.53	0.28

a: The 20 individuals showing extremely early heading comprised 4 in F2-A58/Nipponbare, 2 in F2-A58/Koshihikari, 12 in F2-A58/Kasalath and 2 in F2-A58/#108. b: *P*-values for the probability to Mendelian segregation of genotypes for each loci were obtained by a Chi-square test (1:2:1 ratio).

Figure 1

Supplementary Table S1 Two-way ANOVA based on Table 1 to detect additive interaction between different alleles shown by P-value.

A58/Nipponbare	Hd1	Hd2	Ghd7
Hd1			
Hd2	0.807855		
Ghd7	0.038454	0.64416	

A58/Kasalath	Hd3	Ghd7	Hd6	Hd16	Ehd1
Hd3					
Ghd7	0.411932				
Hd6	0.111289	0.329899			
Hd16	0.054606	0.055915	0.789351a		
Ehd1	0.721323	0.009632	0.592643	0.20737	

A58/IR36	Hd2	Ghd7	Hd17
Hd2			
Ghd7	0.701702a		
Hd17	0.727955	0.563473a	

A58/#108	Hd3	Ghd7	Hd5
Hd3			
Ghd7	0.803396		
Hd5	0.031798	0.182031	

a: short of degree of freedom, yellow cell $p < 5\%$, red cell $p < 1\%$

Supplementary Figure S2