



Title	Helicobacter pylori induces IL-1 protein through the inflammasome activation in differentiated macrophagic cells
Author(s)	Kameoka, Shoichiro; Kameyama, Takeshi; Hayashi, Takaya; Sato, Seiichi; Ohnishi, Naomi; Hayashi, Takeru; Murata-Kamiya, Naoko; Higashi, Hideaki; Hatakeyama, Masanori; Takaoka, Akinori
Citation	Biomedical Research, 37(1), 21-27 https://doi.org/10.2220/biomedres.37.21
Issue Date	2016-02
Doc URL	http://hdl.handle.net/2115/61351
Type	article
File Information	37_21.pdf



[Instructions for use](#)

***Helicobacter pylori* induces IL-1 β protein through the inflammasome activation in differentiated macrophagic cells**

Shoichiro KAMEOKA^{1,2,*}, Takeshi KAMEYAMA^{1,2,*}, Takaya HAYASHI^{3,*}, Seiichi SATO^{1,2}, Naomi OHNISHI⁴, Takeru HAYASHI⁵, Naoko MURATA-KAMIYA⁵, Hideaki HIGASHI⁴, Masanori HATAKEYAMA⁵, and Akinori TAKAOKA^{1,2}

¹Division of Signaling in Cancer and Immunology, Institute for Genetic Medicine, Hokkaido University, Sapporo, Hokkaido, 060-0815, Japan; ²Molecular Medical Biochemistry Unit, Biological Chemistry and Engineering Course, Graduate School of Chemical Sciences and Engineering, Hokkaido University, Sapporo, Hokkaido, 060-8628, Japan; ³Research Center for Infection-associated Cancer, Institute for Genetic Medicine, Hokkaido University, Sapporo, Hokkaido, 060-0815, Japan; ⁴Division of Infection and Immunity, Research Center for Zoonosis Control, Hokkaido University Sapporo Hokkaido, 001-0020, Japan; and ⁵Division of Microbiology, Graduate School of Medicine, University of Tokyo, Tokyo, 113-0033, Japan

(Received 15 November 2015; and accepted 30 November 2015)

ABSTRACT

More than 50% of people in the world are infected with *Helicobacter pylori* (*H. pylori*), which induces various gastric diseases. Especially, epidemiological studies have shown that *H. pylori* infection is a major risk factor for gastric cancer. It has been reported that the levels of interleukin (IL)-1 β are upregulated in gastric tissues of patients with *H. pylori* infection. In this study, we investigated the induction mechanism of IL-1 β during *H. pylori* infection. We found that IL-1 β mRNA and protein were induced in phorbol-12-myristate-13-acetate (PMA)-differentiated THP-1 cells after *H. pylori* infection. This IL-1 β production was inhibited by a caspase-1 inhibitor and a ROS inhibitor. Furthermore, K⁺ efflux and Ca²⁺ signaling were also involved in this process. These data suggest that NOD-like receptor (NLR) family, pyrin domain containing 3 (NLRP3) and its complex, known as NLRP3 inflammasome, are involved in IL-1 β production during *H. pylori* infection because it is reported that NLRP3 inflammasome is activated by ROS, K⁺ efflux and/or Ca²⁺ signaling. These findings may provide therapeutic strategy for the control of gastric cancer in *H. pylori*-infected patients.

Helicobacter pylori (*H. pylori*) is a spiral-shaped, microaerophilic, gram-negative bacterium that establishes persistent infection in the stomach and causes gastric inflammation, which contributes to the progression of various gastric diseases such as chronic gastritis, gastric ulcers, and gastric cancer (4, 24, 26, 27). Many investigators have reported the relevance of cytokine production to *H. pylori* infection

(23, 34). The inflammatory cytokines induced by *H. pylori* infection are associated with a marked infiltration of immune cells such as neutrophils, lymphocytes, monocytes/macrophages, and plasma cells in the gastric mucosa, which leads to *H. pylori*-associated gastroduodenal diseases (25). Among these cytokines induced, IL-1 β is known to be one of the key cytokines during *H. pylori* infection because the levels of IL-1 β are closely associated with gastric inflammation and carcinogenesis (5, 13). In addition, it has been described that *IL1B* polymorphisms are associated with the production levels of IL-1 β during *H. pylori* infection (6), whereas IL-1 β has a crucial role in gastric carcinogenesis *in vivo* (30, 33). According to these evidences, it is an important issue to clarify the induction mechanism of IL-1 β

*These authors contributed equally to this work.

Address correspondence to: Akinori Takaoka, MD, PhD. Division of Signaling in Cancer and Immunology, Institute for Genetic Medicine, Hokkaido University, Sapporo, Hokkaido, 060-0815, Japan

Tel: +81-11-706-5020, Fax: +81-11-706-7541

E-mail: takaoka@igm.hokudai.ac.jp

caused by bacterial and host factor(s). Indeed, recent several studies reported the signaling pathways for IL-1 β production (16, 19, 29). However, the detailed mechanism of *H. pylori*-mediated IL-1 β induction remains fully clarified.

Host innate immune system acts as a front line of host defense against the infection of pathogens. To protect from invasion of these pathogens, host cells need to detect pathogen associated molecular patterns (PAMPs) by using pattern recognition receptors (PRRs). PRR-mediated signalings result in the activation of nuclear factor-kappa B (NF- κ B) and interferon regulatory factors (IRFs) and the subsequent induction of various inflammatory cytokines such as IL-1 β , IL-6 and TNF, and interferons (IFNs). These innate immune signalings and cytokine inductions contribute to not only inducing the early host response against pathogens but also linking to adaptive immune system in the late phase of infection. The coordinated responses by innate and adaptive immune systems are essential for efficient elimination of invading pathogens.

IL-1 β production generally consists of at least two steps; induction of *IL1B* mRNA and maturation of proIL-1 β protein. The latter step is known to be mediated through the activation of inflammasome. The inflammasome is innate immune system receptors/sensors that regulate the activation of caspase-1, which is a crucial enzyme for cleavage of proIL-1 β protein (8). Several families of PRRs are known to be important components in the inflammasome complex, including NOD-like receptors (NLRs) such as NLRP3 and NLRC4. Which NLRs forms inflammasome and/or which molecule(s) trigger inflammasome response during *H. pylori* infection is an interesting issue.

In this study, we examined whether IL-1 β is induced from human cells in response to *H. pylori* infection. We used a human leukemia cell line, THP-1 cells, which are differentiated into macrophages by using phorbol-12-myristate-13-acetate (PMA) and infected with *H. pylori*. At 8 or 24 h after infection, we analyzed the induction levels of IL-1 β mRNA or proteins, respectively. Furthermore, we used several inhibitors to evaluate the inflammasome signalings.

MATERIALS AND METHODS

Cells, bacteria, and reagents. THP-1 (RCB1189) cells were provided by the RIKEN BioResource Center through the National Bio-Resource Project of the MEXT, Japan. THP-1 cells were cultured in RPMI 1640 medium (SIGMA) supplemented with 10%

heat-inactivated fetal bovine serum (FBS: GIBCO). To induce differentiation of THP-1 cells into macrophages, THP-1 cells were stimulated with 1 μ M PMA (P1585; SIGMA) for 3 h, then washed one time with PBS, and plated at 2 days prior to the experiments. An *H. pylori* (WT, NCTC11637) strain used has been reported previously (10). The inhibitors used were as follows: caspase-1 inhibitor (Ac-YVAD-CMK; Enzo Life Sciences, Farmingdale, NY, USA) used the final concentration at 50 μ M, ROS inhibitor (N-acetyl-L-cysteine; Enzo Life Sciences) at 5 mM, P2X7 inhibitor (oxidized ATP; SIGMA) at 300 μ M, inhibition of potassium efflux (KCl; Wako) at 40 mM, Ca²⁺ chelator (BAPTA/AM; AAT Bioquest, Sunnyvale, CA, USA) at 10 μ M, PLC inhibitor (U73122; Cayman CHEMICAL, Ann Arbor, MI, USA) at 10 μ M.

Bacteria culture and H. pylori infection. *H. pylori* was passaged on Trypticase Soy Agar II with 5% Sheep Blood (BD Biosciences) by incubation in microaerophilic condition (an atmosphere consisting of 8% O₂ and 6% CO₂) for 2 days at 37°C. For *H. pylori* infection with cells, bacteria were cultured in Brucella broth (Difco Laboratories, Detroit, MI, USA) supplemented with 10% FBS under the same conditions for 20–24 h at 37°C with shaking at 90–120 rpm. Bacteria were centrifuged at 6,000 rpm for one minute, resuspended in the appropriate cell culture medium, and then added to the cells at a multiplicity of infection (m.o.i.) = 100.

Quantitative-RT-PCR. PMA-differentiated THP-1 cells (2.5 \times 10⁵ cells/well, 24-well plate) were infected with *H. pylori* for 8 h. Total RNA was isolated with ISOGEN (NIPPON GENE, Tokyo, Japan), and were treated with DNaseI (Invitrogen). cDNA was synthesized with the ReverTra Ace Moloney murine leukemia virus reverse transcriptase with point mutations (TOYOBO, Osaka, Japan). Quantitative RT-PCR was analyzed with SYBR Premix Ex Taq reagent mixture (TAKARA, Shiga, Japan) and a StepOnePlus real-time PCR system (Applied Biosystems). Data were normalized to the expression of *ACTB* for each sample. Primers for quantitative RT-PCR were designed as follows; *IL1B*, 5'-GCCGCGTCAGTTGTTGTGGC-3' and 5'-TGAGTCCCGGAGCGTGCAGT-3'; *ACTB*, 5'-CATGTACGTTGCATCCAGGC-3' and 5'-CTCCTTAATGTCACGCAT-3'.

ELISA. The culture supernatant of PMA-differentiated THP-1 cells (2.5 \times 10⁵ cells/well, 24-well plate)




Fig. 1 IL-1 β mRNA and protein were induced in response to *H. pylori* infection. PMA-differentiated THP-1 cells were infected with *H. pylori* (100 m.o.i.). At 8 or 24 h after *H. pylori* infection, the induction levels of IL-1 β mRNA (A) or protein (B) were measured by quantitative RT-PCR or ELISA, respectively. Data are from one representative of at least two independent experiments (mean and SD of triplicate samples). ** $P < 0.01$, * $P < 0.05$.

was collected after 24 h of *H. pylori* infection. The levels of IL-1 β protein were measured by IL-1 β ELISA kit (R&D systems) according to manufacturer's protocol.

Statistical analysis. Values are shown as mean \pm SD. Statistical significance between two samples was determined with Student's *t*-test.

RESULTS

IL-1 β is produced during H. pylori infection in a caspase-1 dependent manner

To first examine whether IL-1 β was secreted by *H. pylori* infection, PMA-differentiated THP-1 cells were infected with *H. pylori* and analyzed. As shown in Fig. 1A, IL-1 β mRNA was induced at 8 h




Fig. 2 IL-1 β production was dependent on caspase-1. PMA-differentiated THP-1 cells treated with Ac-YVAD-CMK, were infected with *H. pylori* (100 m.o.i.) for 24 h. The induction levels of IL-1 β protein were measured by ELISA. Data are from one representative of at least two independent experiments (mean and SD of triplicate samples). ND, not detected. * $P < 0.05$.

after infection with *H. pylori*. Next we studied the protein levels of IL-1 β . IL-1 β production was also observed at 24 h after infection (Fig. 1B). For the production of IL-1 β , an intracellular cysteine protease, caspase-1, is required for processing the inactive precursors into mature form through the inflammasome complex (28). To evaluate the involvement of caspase-1 for the production of IL-1 β , we used Ac-YVAD-CMK, a caspase-1 inhibitor. *H. pylori*-mediated IL-1 β production was inhibited by treatment with Ac-YVAD-CMK (Fig. 2). These data indicated that in PMA-differentiated THP-1 cells, *H. pylori*-mediated IL-1 β production is dependent on caspase-1, possibly through the activation of inflammasome.

H. pylori-mediated IL-1 β production is dependent on reactive oxygen species, extracellular ATP and intracellular potassium

The inflammasome complex mainly consists of three members, NLRs, apoptosis-associated speck-like protein containing a CARD (ASC), and caspase-1. In many bacteria infection, the generation of reactive oxygen species (ROS) is a key signal for the activation of inflammasome (8). We examine whether ROS are important for the inflammasome activation during *H. pylori* infection. Treatment with a ROS inhibitor N-acetyl-L-cysteine significantly suppressed *H. pylori*-mediated IL-1 β production (Fig. 3), suggesting that ROS signaling is involved in this process. In order to know upstream signals for the generation of ROS, we next studied the dependency of extracellular ATP and intracellular potassium (K⁺)




Fig. 3 ROS inhibitor suppressed *H. pylori*-mediated IL-1 β production. PMA-differentiated THP-1 cells treated with N-acetyl-L-cysteine, were infected with *H. pylori* (100 m.o.i.) for 24 h. The induction levels of IL-1 β protein were measured by ELISA. Data are from one representative of at least two independent experiments (mean and SD of triplicate samples). ND, not detected. * $P < 0.05$.

signalings. As shown in Fig. 4A and 4B, *H. pylori*-mediated IL-1 β was inhibited by treatment with oxidized ATP (oxATP) and KCl, respectively. These data suggest that extracellular ATP and intracellular K⁺ signalings generate ROS, which activates inflammasome complex for the production of IL-1 β in response to *H. pylori* infection.

Intracellular calcium signaling pathway also activates H. pylori-mediated inflammasome

Recently, intracellular calcium (Ca²⁺) concentration triggers the inflammasome and subsequent producing ROS (8). We next investigated the involvement of Ca²⁺ signaling during *H. pylori* infection. Treatment with a Ca²⁺ cheator, BAPTA/AM, partially repressed *H. pylori*-mediated IL-1 β production (Fig. 5). In addition, we used U73122, an inhibitor of phospholipase C (PLC), because PLC plays an important role in Ca²⁺ signaling. As shown in Fig. 6, IL-1 β secretion during *H. pylori* infection was dependent on PLC. These data suggest that Ca²⁺ signaling also modulates the production of IL-1 β during *H. pylori* infection.

DISCUSSION

It is well known that in *H. pylori*-infected patients, the levels of IL-1 β are associated with the increased risk of gastric cancer development. Therefore, it is an important issue to reveal the detail mechanism for IL-1 β induction in response to *H. pylori* infection. In this study, we firstly show that *IL1B* mRNA




Fig. 4 Extracellular ATP and K⁺ efflux were also involved in *H. pylori*-mediated IL-1 β production. PMA-differentiated THP-1 cells treated with oxidized ATP (oxATP) (A) or KCl (B), were infected with *H. pylori* (100 m.o.i.) for 24 h. The induction levels of IL-1 β protein were measured by ELISA. Data are from one representative of at least two independent experiments (mean and SD of triplicate samples). ND, not detected. * $P < 0.05$.

is induced after infection with *H. pylori*. In general, *IL1B* transcription is mediated by transcription factor NF- κ B through PRR signalings such as TLR4 (8). In *H. pylori* infection, it is reported that TLR4-mediated NF- κ B activation was observed in gastrointestinal epithelial cells and neutrophils (1, 31), whereas CagA, one of virulence factor, which is closely related to gastric cancer (22), also activates NF- κ B (17). In relation to this, there are several reports about the relevance of *IL1B* mRNA levels to CagA positivity in *H. pylori*-infected patients (23, 32). Considering these reports, we could easily speculate that NF- κ B activation mediated by TLR4 signaling and/or CagA regulates *IL1B* mRNA induction, although further analyses are required to clarify the contribution of CagA to NF- κ B activation.

We also show that the protein levels of IL-1 β are induced after *H. pylori* infection in PMA-differenti-




Fig. 5 Intracellular Ca²⁺ signaling regulated the production of IL-1 β during *H. pylori* infection. PMA-differentiated THP-1 cells treated with BAPTA/AM, were infected with *H. pylori* (100 m.o.i.) for 24 h. The induction levels of IL-1 β protein were measured by ELISA. Data are from one representative of at least two independent experiments (mean and SD of triplicate samples). ND, not detected. * $P < 0.05$.




Fig. 6 PLC inhibitor suppressed *H. pylori*-mediated IL-1 β production. PMA-differentiated THP-1 cells treated with U73122, were infected with *H. pylori* (100 m.o.i.) for 24 h. The induction levels of IL-1 β protein were measured by ELISA. Data are from one representative of at least two independent experiments (mean and SD of triplicate samples). ND, not detected. * $P < 0.05$.

ated THP-1 cells. Consistent with our observation, a previous report demonstrated that human monocytes and macrophages secreted IL-1 β in response to *H. pylori* infection (7). We next analyzed the induction mechanism of IL-1 β . As mention above, it is well known that IL-1 β production is mediated by inflammasome activation. In this study, we show that *H. pylori*-mediated IL-1 β production is dependent on caspase-1, which is a crucial enzyme for the activation of inflammasome. To determine which signalings are involved in *H. pylori*-mediated caspase-1 activation, we used several representative inhibitors and evaluated IL-1 β production. Consequently, we also reveal that ROS, extracellular ATP, intracellular K⁺ and Ca²⁺ signalings are involved in the production of IL-1 β , suggesting that NLRP3 inflammasome plays an important role in this process in human macrophages. In this regard, two recent studies by other groups using murine bone marrow-derived dendritic cells described the involvement of NLRP3 in *H. pylori*-induced IL-1 β secretion, which is consistent with our observation.

At least a few models have been proposed for the activation of NLRP3 inflammasome. One model proposes a key role for ROS production, which is important for the priming step rather than the regulation of complex assembly, and other models implicate phagolysosomal damage and K⁺ efflux (28). K⁺ efflux is caused by external ATP, considered as a danger signal, which causes the opening of the P2X7 receptor and the subsequent recruitment of the channel pannexin-1 leading to the release of intra-

cellular K⁺ (15). It has also been reported that the bacterial toxin nigericin derived from *Streptomyces hygroscopicus* activates NLRP3 by causing K⁺ efflux in a pannexin-1-dependent manner (21). In our study, IL-1 β production is dependent on ROS, K⁺ efflux and extracellular ATP, suggesting that some factors derived from *H. pylori* affect this process. Indeed, Semper *et al.* show that *cag* pathogenicity island (*cagPAI*) and vacuolating cytotoxin A (*VacA*) are involved in the process of inflammasome activation (29). Further investigations are needed to clarify the detail mechanism for the NLRP3 activation and involvement of other bacterial factors.

We also demonstrate that intracellular Ca²⁺ signaling also regulates IL-1 β production in response to *H. pylori* infection. Recent accumulating evidences show that intracellular Ca²⁺ signaling also plays a crucial role in NLRP3 activation (12, 18). In this regard, there are interesting data that intracellular Ca²⁺ concentration is increased through *cagA* and *picB/cagE* genes after *H. pylori* infection and this Ca²⁺ upregulation is regulated by a PLC-dependent mechanism (20). In addition, another group reported that CagA was interacted with PLC γ 1 (3). CagA is known to be an effector protein that is translocated directly from bacteria into the host cells via the effector translocator system, known as a type IV secretion system (T4SS), and to interact with many host proteins, which leads to interfere with cellular signaling pathways that regulate cell growth, morphology, and motility (2, 9, 26, 27). Considering these reports, one could speculate that CagA activates not only

IL1B mRNA induction but also inflammasome assembly for IL-1 β production, although further analyses are needed to clarify the involvement of CagA and detail mechanism for the activation of NLRP3 inflammasome during *H. pylori* infection. In order to rule out the involvement of other virulence factors derived from *H. pylori*, reconstitution assay of the NLRP3 inflammasome might solve this point.

This study revealed that *H. pylori* infection induces IL-1 β mRNA and protein in human macrophages, possibly through the NLRP3 inflammasome activation. It is reported that IL-1 β increased in the stomach during *H. pylori* infection is strongly associated with the generation of gastric carcinoma (30, 33). In addition, recent evidences show that *H. pylori* and IL-1 β are closely related with not only gastric carcinoma but also diabetes and Alzheimer's disease (11, 14). Thus our findings may have clinical implications for *H. pylori*-associated diseases, and further research is desired.

Acknowledgement

The author is grateful for financial supports from the Ministry of Health, Labour and Welfare of Japan (Grant-in-Aid to A.T.), the Ministry of Education, Culture, Sports, Science and Technology of Japan (Grant-in-Aid for Scientific Research [A] [25253030] to A.T., Grant-in-Aid for Scientific Research on Innovative Areas [25115502, 23112701] to A.T., IRYO HOJIN SHADAN JIKOKAI (H. Tanaka and N. Takayanagi) to A.T., the Kato Memorial Bioscience Foundation to A.T., the Yasuda Medical Foundation to A.T., the Takeda Science Foundation to A.T., and The Waksman Foundation of Japan to A.T.

REFERENCES

1. Alvarez-Arellano L, Camorlinga-Ponce M, Maldonado-Bernal C and Torres J (2007) Activation of human neutrophils with *Helicobacter pylori* and the role of Toll-like receptors 2 and 4 in the response. *FEMS Immunol Med Microbiol* **51**, 473–479.
2. Backert S, Tegtmeyer N and Selbach M (2010) The versatility of *Helicobacter pylori* CagA effector protein functions: The master key hypothesis. *Helicobacter* **15**, 163–176.
3. Churin Y, Al-Ghoul L, Kepp O, Meyer TF, Birchmeier W and Naumann M (2003) *Helicobacter pylori* CagA protein targets the c-Met receptor and enhances the motogenic response. *J Cell Biol* **161**, 249–255.
4. Dunne C, Dolan B and Clyne M (2014) Factors that mediate colonization of the human stomach by *Helicobacter pylori*. *World J Gastroenterol* **20**, 5610–5624.
5. El-Omar EM (2001) The importance of interleukin 1 β in *Helicobacter pylori* associated disease. *Gut* **48**, 743–747.
6. El-Omar EM, Carrington M, Chow WH, McColl KE, Bream JH, Young HA, Herrera J, Lissowska J, Yuan CC, Rothman N, Lanyon G, Martin M, Fraumeni JF and Rabkin CS (2000) Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* **404**, 398–402.
7. Fehlings M, Drobbe L, Moos V, Renner Viveros P, Hagen J, Beigier-Bompadre M, Pang E, Belogolova E, Churin Y, Schneider T, Meyer TF, Aebischer T and Ignatius R (2012) Comparative analysis of the interaction of *Helicobacter pylori* with human dendritic cells, macrophages, and monocytes. *Infect Immun* **80**, 2724–2734.
8. Guo H, Callaway JB and Ting JP-Y (2015) Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nat Med* **21**, 677–687.
9. Hatakeyama M (2014) *Helicobacter pylori* CagA and gastric cancer: a paradigm for hit-and-run carcinogenesis. *Cell Host Microbe* **15**, 306–316.
10. Higashi H, Nakaya A, Tsutsumi R, Yokoyama K, Fujii Y, Ishikawa S, Higuchi M, Takahashi A, Kurashima Y, Teishikata Y, Tanaka S, Azuma T and Hatakeyama M (2004) *Helicobacter pylori* CagA induces Ras-independent morphogenetic response through SHP-2 recruitment and activation. *J Biol Chem* **279**, 17205–17216.
11. Honjo K, van Reekum R and Verhoeff NPLG (2009) Alzheimer's disease and infection: do infectious agents contribute to progression of Alzheimer's disease? *Alzheimers Dement* **5**, 348–360.
12. Horg T (2014) Calcium signaling and mitochondrial destabilization in the triggering of the NLRP3 inflammasome. *Trends Immunol* **35**, 253–261.
13. Hwang I-R, Kodama T, Kikuchi S, Sakai K, Peterson LE, Graham DY and Yamaoka Y (2002) Effect of interleukin 1 polymorphisms on gastric mucosal interleukin 1 β production in *Helicobacter pylori* infection. *Gastroenterology* **123**, 1793–1803.
14. Jeon CY, Haan MN, Cheng C, Clayton ER, Mayeda ER, Miller JW and Aiello AE (2012) *Helicobacter pylori* infection is associated with an increased rate of diabetes. *Diabetes Care* **35**, 520–525.
15. Kanneganti T-D, Lamkanfi M, Kim Y-G, Chen G, Park J-H, Franchi L, Vandenabeele P and Núñez G (2007) Pannexin-1-mediated recognition of bacterial molecules activates the cryopyrin inflammasome independent of Toll-like receptor signaling. *Immunity* **26**, 433–443.
16. Kim D-J, Park J-H, Franchi L, Backert S and Núñez G (2013) The Cag pathogenicity island and interaction between TLR2/NOD2 and NLRP3 regulate IL-1 β production in *Helicobacter pylori* infected dendritic cells. *Eur J Immunol* **43**, 2650–2658.
17. Lamb A, Yang X-D, Tsang Y-HN, Li J-D, Higashi H, Hatakeyama M, Peek RM, Blanke SR and Chen L-F (2009) *Helicobacter pylori* CagA activates NF-kappaB by targeting TAK1 for TRAF6-mediated Lys 63 ubiquitination. *EMBO Rep* **10**, 1242–1249.
18. Lee G-S, Subramanian N, Kim AI, Aksentijevich I, Goldbach-Mansky R, Sacks DB, Germain RN, Kastner DL and Chae JJ (2012) The calcium-sensing receptor regulates the NLRP3 inflammasome through Ca²⁺ and cAMP. *Nature* **492**, 123–127.
19. Li X, Liu S, Luo J, Liu A, Tang S, Liu S, Yu M and Zhang Y (2015) *Helicobacter pylori* induces IL-1 β and IL-18 production in human monocytic cell line through activation of NLRP3 inflammasome via ROS signaling pathway. *Pathog Dis* **73**, ftu024–ftu024.
20. Marlink KL, Bacon KD, Sheppard BC, Ashktorab H, Smoot DT, Cover TL, Deveney CW and Rутten MJ (2003) Effects of *Helicobacter pylori* on intracellular Ca²⁺ signaling in normal human gastric mucous epithelial cells. *Am J Physiol Gastrointest Liver Physiol* **285**, G163–G176.

21. Muñoz-Planillo R, Kuffa P, Martínez-Colón G, Smith BL, Rajendiran TM and Núñez G (2013) K⁺ efflux is the common trigger of NLRP3 inflammasome activation by bacterial toxins and particulate matter. *Immunity* **38**, 1142–1153.
22. Ohnishi N, Yuasa H, Tanaka S, Sawa H, Miura M, Matsui A, Higashi H, Musashi M, Iwabuchi K, Suzuki M, Yamada G, Azuma T and Hatakeyama M (2008) Transgenic expression of *Helicobacter pylori* CagA induces gastrointestinal and hematopoietic neoplasms in mouse. *Proc Natl Acad Sci USA* **105**, 1003–1008.
23. Peek RM, Miller GG, Tham KT, Perez-Perez GI, Zhao X, Atherton JC and Blaser MJ (1995) Heightened inflammatory response and cytokine expression in vivo to *cagA*⁺ *Helicobacter pylori* strains. *Lab Invest* **73**, 760–770.
24. Peek RM and Crabtree JE (2006) *Helicobacter* infection and gastric neoplasia. *J Pathol* **208**, 233–248.
25. Portal-Celhay C and Perez-Perez GI (2006) Immune responses to *Helicobacter pylori* colonization: mechanisms and clinical outcomes. *Clin Sci* **110**, 305–314.
26. Rieder G, Merchant JL and Haas R (2005) *Helicobacter pylori* cag-type IV secretion system facilitates corpus colonization to induce precancerous conditions in Mongolian gerbils. *Gastroenterology* **128**, 1229–1242.
27. Salama NR, Hartung ML and Müller A (2013) Life in the human stomach: persistence strategies of the bacterial pathogen *Helicobacter pylori*. *Nat Rev Microbiol* **11**, 385–399.
28. Schroder K and Tschopp J (2010) The inflammasomes. *Cell* **140**, 821–832.
29. Semper RP, Mejías-Luque R, Groß C, Anderl F, Müller A, Vieth M, Busch DH, Prazeres da Costa C, Ruland J, Gross O and Gerhard M (2014) *Helicobacter pylori*-Induced IL-1 β secretion in innate immune cells is regulated by the NLRP3 inflammasome and requires the cag pathogenicity island. *J Immunol* **193**, 3566–3576.
30. Shigematsu Y, Niwa T, Rehnberg E, Toyoda T, Yoshida S, Mori A, Wakabayashi M, Iwakura Y, Ichinose M, Kim Y-J and Ushijima T (2013) Interleukin-1 β induced by *Helicobacter pylori* infection enhances mouse gastric carcinogenesis. *Cancer Lett* **340**, 141–147.
31. Su B, Ceponis PJM, Lebel S, Huynh H and Sherman PM (2003) *Helicobacter pylori* activates Toll-like receptor 4 expression in gastrointestinal epithelial cells. *Infect Immun* **71**, 3496–3502.
32. Sugimoto M, Ohno T, Graham DY and Yamaoka Y (2009) Gastric mucosal interleukin-17 and -18 mRNA expression in *Helicobacter pylori*-induced Mongolian gerbils. *Cancer Sci* **100**, 2152–2159.
33. Tu S, Bhagat G, Cui G, Takaishi S, Kurt-Jones EA, Rickman B, Betz KS, Penz-Oesterreicher M, Bjorkdahl O, Fox JG and Wang TC (2008) Overexpression of interleukin-1 β induces gastric inflammation and cancer and mobilizes myeloid-derived suppressor cells in mice. *Cancer Cell* **14**, 408–419.
34. Yamaoka Y, Kita M, Kodama T, Sawai N, Kashima K and Imanishi J (1997) Induction of various cytokines and development of severe mucosal inflammation by *cagA* gene positive *Helicobacter pylori* strains. *Gut* **41**, 442–451.