



Title	Sulfuriferula multivorans gen. nov., sp nov., isolated from a freshwater lake, reclassification of 'Thiobacillus plumbophilus' as Sulfuriferula plumbophilus sp nov., and description of Sulfuricellaceae fam. nov and Sulfuricellales ord. nov.
Author(s)	Watanabe, Tomohiro; Kojima, Hisaya; Fukui, Manabu
Citation	International journal of systematic and evolutionary microbiology, 65(5), 1504-1508 <a href="https://doi.org/10.1099/ijs.0.000129">https://doi.org/10.1099/ijs.0.000129</a>
Issue Date	2015-05
Doc URL	<a href="http://hdl.handle.net/2115/61459">http://hdl.handle.net/2115/61459</a>
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	1-TTN_sup.pdf



[Instructions for use](#)

Supplementary material

***Sulfuriferula multivorans* gen. nov., sp. nov., isolated from a freshwater lake, reclassification of ‘*Thiobacillus plumbophilus*’ as *Sulfuriferula plumbophilus* sp. nov., and description of *Sulfuricellaceae* fam. nov. and *Sulfuricellales* ord. nov.**

Tomohiro Watanabe\*, Hisaya Kojima, Manabu Fukui

The Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan

E-mail: watanabe1986@pop.lowtem.hokudai.ac.jp




Fig. S1. Phylogenetic position of strain TTN<sup>T</sup> within the class *Betaproteobacteria*, based on SoxB amino acid sequences aligned by ClustalW (153 amino acids were used). *Allochromatium vinosum* was used as an outgroup. Tree was constructed by the neighbor-joining method with 1000 bootstrap resamplings (bootstrap values > 50% are shown at the nodes). Bar, 0.02 substitutions per nucleotide position.




Fig. S2. Phylogenetic position of strain TTN<sup>T</sup> within the class *Betaproteobacteria*, based on AprA amino acid sequences aligned by ClustalW (119 amino acids were used). *Allochromatium vinosum* was used as an outgroup. Tree was constructed by the neighbor-joining method with 1000 bootstrap resamplings (bootstrap values > 50% are shown at the nodes). Bar, 0.02 substitutions per nucleotide position.




Fig. S3. Phylogenetic position of strain TTN<sup>T</sup> within the class *Betaproteobacteria*, based on Sqr amino acid sequences aligned by ClustalW (124 amino acids were used). *Allochromatium vinosum* was used as an outgroup. Tree was constructed by the neighbor-joining method with 1000 bootstrap resamplings (bootstrap values > 50% are shown at the nodes). Bar, 0.02 substitutions per nucleotide position.




Fig. S4. Phylogenetic position of strain TTN<sup>T</sup> within the class *Betaproteobacteria*, based on DsrA amino acid sequences aligned by ClustalW (84 amino acids were used). *Allochromatium vinosum* was used as an outgroup. Tree was constructed by the neighbor-joining method with 1000 bootstrap resamplings (bootstrap values > 50% are shown at the nodes). Bar, 0.02 substitutions per nucleotide position.