



Title	Behavior of laying hens in relation to housing system and stain
Author(s)	Mohammed, Hesham H.; Said, Enas N.
Citation	Japanese Journal of Veterinary Research, 64(Supplement 2), S143-S148
Issue Date	2016-04
Doc URL	http://hdl.handle.net/2115/62013
Type	bulletin (article)
File Information	p.S143-148 Hesham H. Mohammed.pdf



[Instructions for use](#)

Behavior of laying hens in relation to housing system and strain

Hesham H. Mohammed* and Enas N. Said

Dept. of Vet. Publ. Health, Faculty of Veterinary Medicine, Zagazig University, Egypt.

*Corresponding author: heshamvet_hosny@yahoo.com

Abstract

The current study was to investigate the effect of different enriched cages and lohmann strains in behavior of layers. A total of 480 layers of lohmann breed, at 30 weeks old (240 lohmann brown "LB" & 240 lohmann selected leghorn "LSL") were reared in 2 types of enriched cages "Salmet & Big dutchman". The results revealed that the most of normal behaviors were significantly higher in salmet system, while nesting and other abnormal behaviors "feather pecking" were significantly higher in big dutchman. All normal behaviors in the present study were higher in LSL with significant difference in eating, preening, nesting and resting behavior. Nevertheless, abnormal behaviors were significantly higher in LB. In conclusion, good selection of cage system and strain of layers play role in improvement normal behavior and control of abnormal behavior.

Key Word: Layer, Enriched cage, Strain, Behavior

Introduction

There are many factors which can decrease the performance and increase abnormal behaviours of hens such as diseases, management and housing. The conditions under which laying hens are kept remain major animal welfare concern. Housing system has a significant effect on the welfare of laying hens, and one of the factors for estimating the welfare is the condition of feathers¹³. Cage system is one of the housing system and widely used for laying hens because egg production is cheaper in cages than in alternative husbandry system. It has three different categories: non-enriched cages, alternative systems and enriched cages. Non-enriched cages mean the common battery cages, alternative systems refer to non-cage system such as aviaries. While the enriched (modified) cages aim to improve hen welfare by providing with perches, nest boxes, sand bath

mates and other facilities. And the major criticisms of the cage systems are that they increase the incidence of feather damage, overgrowing claws, foot lesions and brittle bones²². The presence of apparently purposeless behavior, of high levels of aggression or redirected behaviours such as feather pecking and cannibalism are indicators that the housing system is not meeting the behavioural needs of the hens and hence is not satisfactory for bird welfare²⁶. To overcome this problem, overseas research has increasingly turned toward improving the welfare of birds in cages by modifying cage design⁷. Selective breeding for desired traits such as decreased feather pecking and cannibalism may help to improve welfare¹². The causes of variation in productivity, mortality rate and cannibalism rate revealed a strong effect of strain¹. The differences in cannibalistic and feather pecking behavior between brown and white eggs layers¹⁷. The aim of the current study was to investigate

the effect of different enriched cages and lohmann strains in behavior of layers.

Materials and Methods

This experiment was performed on 480 layers of lohmann breed, at 30 weeks old and divided into 240 Lohmann brown (LB) and 240 Lohmann selected Leghorn (LSL). Each strain was reared in 2 types of enriched cages (salmet and big dutchment) with floor area of 900 cm² for each bird and housed at the farm in the Faculty of Veterinary Medicine, Zagazig University, Egypt. Description of enriched cages was in Table (1). During the experimental period, a 14- hour lighting schedule was applied from fluorescent light. Feed and water were provided ad libitum. The basal layer diet was fed containing 18% of crude protein and 3000 Kcal/Kg of metabolized energy¹⁶⁾. Direct observations were conducted to record different behavior through scan sampling method by the same person¹⁴⁾. The observer stood directly in front

of the pen and waited ten minutes before recording to avoid any disturbance in the behavior. All cages were observed directly for 2 × 10 minutes in the morning and for 2 × 10 minutes in the afternoon on two days every week¹¹⁾. After scanning, the numbers of birds were counted & calculated the frequencies of activities¹⁵⁾. Behavioral patterns, as percentages were:

- Eating, drinking, preening, perching, nesting, resting and dust bathing.
- Resting behavior: sitting to remain dormant with the neck withdrawn¹⁸⁾.
- Dust bathing: side-rubbing, head-rubbing & wing shaking with scratching²³⁾.
- Feather pecking (Fp): only pecks to feathered parts of the body³⁾.
- Aggressive behavior (Ag): the birds counter acts toward other birds.

Data was statistically analyzed by SAS statistical system¹⁹⁾. One-factorial analysis of variance (ANOVA) was performed. Differences between means were tested by Student's t-test.

Table 1. Description of systems and birds per cage.

	Big dutchman (BD)	Salmet (S)
No. of birds in cage	60	30
No. of cage	4	8
Cage space	360 cm * 155 cm	400 cm * 70 cm
No. of nest	2	2
Nest space	90 cm * 30 cm	54.5 cm * 25 cm
No. of upper perch	2	1
Length of upper perch	175 cm	200 cm
No. of lower perch	2	5
Length of lower perch	175 cm	75 cm
No. of middle perch	1	0
Length of middle perch	175 cm	-----
No. of bath -mate	2	4
Bath-mate space	75 cm * 30 cm	35 cm * 20 cm
Area of trough / bird (cm ²)	12 cm ²	12cm ²
No. of drinking nipples	8	7

Results

Significance levels of main factors and their interaction for different behavior were shown in Table 2. It clearly showed that system of housing and strains of lohmann had significant effect on most of behavioral patterns. While, the interaction between system and strain didn't affect in the

most of behavioral patterns except dust bathing. Means \pm standard deviation of normal behaviour were presented in Table 3. It clearly revealed that these behaviors were significantly affected by housing system and strains of lohmann breed. Moreover, housing system and strains of lohmann breed had significant effect on abnormal behavior (Table 4).

Table 2. Significance levels of main factors & their interaction on behaviors of layers.

Sources of variance	Eating	Drinking	Preening	perching	Nesting	Resting	Dustbathing	Fp	Ag
System (S)	NS	***	***	**	*	***	***	*	NS
Breed (B)	*	NS	***	NS	***	***	NS	***	**
S*B	NS	NS	NS	NS	NS	NS	***	NS	NS

P < 0.05; ** P < 0.01; *** P < 0.001; NS, not significant; Fp, feather pecking; Ag, aggression.

Table 3. Means (\pm SD) of behavioral traits of hens in response to lohmann strains and housing systems (% of hens in observation period).

Behavior	Big Dutchmen		Salmet	Means
	LB	LSL		
Eating	LB	19.75 \pm 11.42	22.10 \pm 14.14	20.92 \pm 12.86 ^b
	LSL	30.15 \pm 12.03	32.6 \pm 14.95	31.38 \pm 13.57 ^a
	Means	24.98 \pm 12.81	27.32 \pm 15.42	
Drinking	LB	4.87 \pm 5.79	6.9 \pm 2.98	5.89 \pm 4.70
	LSL	6.38 \pm 7.53	7.22 \pm 3.30	6.8 \pm 5.79
	Means	5.61 \pm 6.72 ^b	7.07 \pm 3.14 ^a	
Preening	LB	0.48 \pm 1.04	6.27 \pm 368	3.37 \pm 3.96 ^b
	LSL	1.03 \pm 1.86	13.55 \pm 8.28	7.29 \pm 8.64 ^a
	Means	0.75 \pm 1.57 ^b	9.91 \pm 7.31 ^a	
Perching	LB	7.2 \pm 3.77	12.43 \pm 4.14	9.82 \pm 4.74
	LSL	8.55 \pm 6.12	13.95 \pm 5.51	11.25 \pm 6.39
	Means	7.89 \pm 5.13 ^b	13.17 \pm 4.90 ^a	
Nesting	LB	3.58 \pm 2.40	2.75 \pm 3.23	3.16 \pm 2.87 ^b
	LSL	6.29 \pm 4.34	5.10 \pm 4.95	5.69 \pm 4.69 ^a
	Means	4.95 \pm 3.74 ^a	3.91 \pm 4.34 ^b	
Resting	LB	1.67 \pm 2.08	3.14 \pm 2.79	2.4 \pm 2.56 ^b
	LSL	6.95 \pm 7.42	10.37 \pm 8.99	8.66 \pm 8.39 ^a
	Means	4.33 \pm 6.05 ^b	6.74 \pm 7.55 ^a	
Dust bathing	LB	1.84 \pm 2.60 ^b	2.26 \pm 3.64 ^{ab}	2.05 \pm 3.16
	LSL	1.53 \pm 1.84 ^b	3.47 \pm 4.93 ^a	2.5 \pm 3.88
	Means	1.63 \pm 2.27 ^b	2.97 \pm 4.37 ^a	

*Means within same column with different letters are significantly different (P \leq 0.05).

Table 4. Means (\pm SD) of abnormal behavior of layers in response to lohmann strains and housing systems (% of hens in observation period).

Behavior		Big Dutchmen	Salmet	Means
Feather pecking	LB	7.32 \pm 7.96	4.94 \pm 6.57	6.13 \pm 7.38 ^a
	LSL	5.43 \pm 3.80	2.48 \pm 3.12	3.96 \pm 3.77 ^b
	Means	6.40 \pm 6.29 ^a	3.69 \pm 5.26 ^b	
Aggression	LB	0.18 \pm 0.49	0.13 \pm 0.49	0.15 \pm 0.49 ^a
	LSL	0.09 \pm 0.36	0.07 \pm 0.35	0.08 \pm 0.37 ^b
	Means	0.13 \pm 0.44	0.10 \pm 0.43	

*Means within same column with different letters are significantly different ($P < 0.05$).

Discussion

In practice, behavioural measures are often the starting point for assessing an animal's response to its environment and hence, its welfare. The notion is that welfare assessment can provide insights into important factors when designing environments for chickens⁸. Behavior is a good indicator for the assessment of the well-being of laying hens. The data in the present study showed that eating behavior was significantly higher in salmet than big dutchman, while there was no significant difference between LSL and LB. % drinking was significantly lower in big dutchman than salmet cages. The differences in ingestive behavior (eating and drinking) under different housing system may be due to change in group size and its relation to social hierarchy.

Also, LSL had the increase in eating and drinking behaviors with significant deference in eating behavior. These results refer the effect of strain on feed intake, however the time spent eating was not affected by strains (brown-egg layers and white-egg layers), as mentioned before²⁰. While white hens spent the most time in drinking and brown hens spent the least. In other study the appetitive was not significantly affected by strain².

The type of housing had a significant effect on preening behaviour of bird⁸, where it was significantly higher in salmet than big dutchman. According to strain, LSL appears the increase of preening behaviour and this was evident in our study.

Perching the layers in salmet system was the highest with significant difference. This result may be due to the space allowance of perch/hen in salmet was higher than big dutchman⁶. The same observation for perching behavior was described by previous publication before^{4,9,21}. And, it was higher in LSL than LB, but the difference didn't reach the significance. Number of hens in nest was significantly higher in big dutchman than salmet cages. However, it was increased significantly in LSL than LB, as mentioned by. This result was in agreement with the results reported before^{10,24}. Dust bathing and resting behaviors were the highest in salmet system with significant differences. Also, it was higher in LSL than LB with significant difference in resting behaviour. There was no significant effects of strains on dust bathing²⁰, while, Hy-Line Brown layers performed more dust bathing than Hy-Line White²⁵. There was significant increase of resting behavior in LSL in compare to LB¹⁰. Regarding the influence of cage system on abnormal behavior, the results of Table (4) showed increase in big dutchman with significant difference in feather pecking. These results may be attributed to increase group size in big Dutchman, as mentioned before⁵. In conclusion, good selection of cage system and strain of layers play role in improvement normal behaviour and control of abnormal behaviours especially aggressive pecking.

References

- 1) Aerni, V., Brinkhof, M., Wechsler, B., Oester, H. and Fröhlich E. 2005. Productivity and mortality of laying hens in aviaries: a systematic review. *Worl. Poult. Sci.*, **61**: 130-142.
- 2) Anderson, K., Jones, D., Davis, G. and Jenkins, P. 2007. Effects of selection on behavioural profiles of single comb white leghorn hens through two production cycles. *Poult. Sci.*, **86**: 1814-1820.
- 3) Anja, B, Rupert, P. and Björn, F. 2006. Effect of brooder on feather pecking and cannibalism in domestic fowl (*Gallus gallus domesticus*). *Appl. Anim. Behav. Sci.*, **99**: 287-300.
- 4) Blokhuis, H., Fiks, T., Niekerk, V., Bessei, W., Elson, A., Guemene, D., Kjaer, J., Maria, L., Nicol, C., Tauson, R., Weeks, C. and Weerd, V. 2007. The laywel project: welfare implications of changes in production systems for laying hens. *Worl. Poult. Sci.*, **63**: 101-114.
- 5) Busayi, R., Channing, C. and Hocking, P. 2006. Comparisons of damaging feather pecking and time budgets in male and female turkeys of a traditional breed and a genetically selected male line. *Appl. Anim. Behav. Sci.*, **96**: 281-292.
- 6) DEFRA. 2002. Code of recommendations for the welfare of livestock: laying hens (London, defra).
- 7) Elson, H. 1990. Recent developments in laying cages designed to improve bird's welfare. *Worl. Poult. Sci.*, **46**: 34-37.
- 8) Fortomaris, P., Arsenos, G., Angeliki, T. and Yannakopoulos, A. 2007. Performance and behaviour of broiler chickens as affected by the housing system. *Arch. Geflügelk.*, **71**: 97-104.
- 9) Guinebretiere, M., Huonnic, D., De-Treglode, M., Huneau-salaun, A. and Michel, V. 2009. Furnished cages for laying hens: effect of group size and litter distribution provision on laying, feeding, perching and dust bathing behaviours. Poultry welfare symposium cervia, Italy, 18-22.
- 10) Hesham, H. M. 2012. Assessment of the behavior, plumage and feet conditions in two commercial layer breeds. *Int. Appl. Anim. Sci.*, **1**: 18-22.
- 11) Kjaer, J. and Sorensen, P. 2002. Feather pecking and cannibalism in free-range laying hens as affected by genotype, dietary level of methionine + cysteine, light intensity during rearing and age at first access to the range area. *Appl. Anim. Behav. Sci.*, **76**: 21-39.
- 12) Lay, D., Fulton, R., Hester, P., Karcher, D., Kjaer, J., Mench, J., Mullens, B., Newberry, R., Nicol, C., O'Sullivan, N. and Porter, R. 2011. Hen welfare in different housing systems. *Poult. Sci.*, **90**: 278-294.
- 13) Mirjana, D., Lidija, P. and Niko, M. 2008. Effect of housing system, age and hybrid type on the welfare of laying hens. *Lucrări Stiințif. Zootehn. Biotehnol.*, **41**: 651-654.
- 14) Mohammed, H., Badawi, M., Walaa, M., Ali, M. and Abd el-aziz, R. 2014. The influence of chromium sources on growth performance, economic efficiency, some maintenance behaviour, blood metabolites and carcass traits in broiler chickens. *Glob. Vetrin.*, **12**: 599-605.
- 15) Mohammed, H., Grashorn, M. and Bessei, W. 2010. The effects of lighting conditions on the behaviour of laying hens. *Arch.Geflügelk.*, **74**: 197-202.
- 16) NRC. 1994. Nutrient requirement of poultry. 9th rev. ed. national academy press. Washinton. D.C., USA.
- 17) Odén, K., Keeling, L. and Algers, B. 2002. Behaviour of laying hens in two types of aviary systems on 25 commercial farms in Sweden. *Br. Poult. Sci.*, **43**: 169-181.
- 18) Ruth, C., Keeling, L., Estevez, I. and Bilcik, B. 2006. Behaviour when young as a predictor of severe feather pecking in adult laying hens: the redirected foraging hypothesis revisited. *Appl. Anim. Behav. Sci.*, **107**: 262-274.
- 19) SAS. 2002. SAS/stat users guide. sas institute inc, cary, nc 27513, USA.

- 20) Singh, R. 2008. Production and behaviour of four strains of laying hens kept in conventional cages and a free run housing system. The faculty of graduate studies, the university of British Columbia, master thesis.
- 21) Struelens, E., Tuyttens, F., Duchateau, L., Leroy, T., Cox, M., Vranken, E., Buyse, J., Zoons, J., Berckmans, D., Odberg, F. and Sonck, B. 2008. Perching behaviour and perch height preference of laying hens in furnished cages varying in height. *Br. Poult. Sci.*, **49**: 381-389.
- 22) Thomas, D. and Ravindran, V. 2005. Comparison of layer performance in cage and barn systems. *Anim. Vet. Advan.*, **4**: 554-556.
- 23) Van-Liere, D. 1991. Function and organization of dust bathing behaviour in laying hens. Doctoral thesis, agricultural university, Wageningen, Netherlands.
- 24) Wall, H. 2003. Laying hens in furnished cages-use of facilities, exterior egg quality and bird health. Doctoral thesis. Swedish univ. Uppsala.
- 25) Wall, H., Tauson, R. and Elwinger, K. 2008. Effects of litter substrate and genotype on layers' use of litter, exterior appearance, and heterophil: lymphocyte ratios in furnished cages. *Poult. Sci.*, **87**: 2458-2465.
- 26) Weeks, C. and Nicol, C. 2006. Behavioural needs priorities and preferences of laying hens. *Worl. Poult. Sci.*, **62**: 296-307.