



Title	Pathological findings of cutaneous periocular habronemiasis in a horse in Egypt
Author(s)	Ali, Abdel-Moneim; Fathy, Ashraf; Ibrahem, Rania; Gomaa, Mohamed; Naif, AL-Gabri
Citation	Japanese Journal of Veterinary Research, 64(Supplement 2), S117-S122
Issue Date	2016-04
Doc URL	http://hdl.handle.net/2115/62017
Type	bulletin (article)
File Information	p.S117-122 Abdel-Moneim Ali.pdf



[Instructions for use](#)

Pathological findings of cutaneous periocular habronemiasis in a horse in Egypt

Abdel-Moneim Ali^{1,*}, Ashraf Fathy², Rania Ibrahim², Mohamed Gomaa³ and AL-Gabri Naif¹

¹Department of Pathology, Faculty of Veterinary Medicine, Zagazig University, Egypt

²Department of Pathology, Animal Health Research Institute, Egypt.

³Department of Surgery, Faculty of Veterinary Medicine, Zagazig University, Egypt

*Corresponding author: E-mail: abdelmoneim.ahmedali@yahoo.com.

Abstract

A 14 years old horse was admitted in June 2014 with an unhealed ulcer near to the medial canthus of the left eye. The ulcerated skin lesion was 3 cm in diameter reddish in color with hazy irregular borders and containing grayish and yellowish necrotic tissues with offensive odor. Histopathological examination of tissue sections showed parasitic elements embedded in multiple granulomas with caseated centers with degenerated *habronema* larvae containing masses from granulated eosinophil in the center followed by eosinophilic and basophilic necrotic debris beside numerous macrophage and giant cells followed by fibrous capsule. The covering skin revealed large deep cutaneous ulcer containing purulent or serous exudate with living and degenerated neutrophils. The inflammatory process was extended into deeper area forming pyogranuloma containing multinucleated giant cells and fibrous tissues. Also the adjacent tissues revealed aggregations of microfilaria in dermis (Onchocerciasis), and in other section from behind the granuloma we saw nonseptated mycelium (Cutaneous Phaeohyphomycosis) within the purulent exudate as an opportunistic infection. It could be concluded that complicated cutaneous Habronemiasis consider one of the parasitic cutaneous lesions which is rare among horses in Egypt. Immunological reactions of the parasitic element may be one of the etiological agents of skin ulcers and fungal invasions.

Key Words: Horses, Cutaneous *Habronema*, *microfilaria* in dermis, *Onchocerciasis*, *Phaeohyphomycosis*.

Introduction

Habronemiasis (also called summer sores, granular dermatitis, jack sores, bursati, and other terms) is a complex parasitic disease of donkeys, horses, mules, zebras, and dogs and is most commonly encountered in temperate, subtropical, and tropical regions^{9,18}. It has a seasonal distribution, with occurrences predominately in the spring and summer and coinciding with fly activity²². Habronemiasis is caused by the invasion of

Draschiamegastoma, *Habronemamajus*, and *H. muscae* nematodes which transmitted by house flies, face flies, and stable flies respectively. The adult nematodes live in the horse's stomach, laying eggs, which pass out in the horse's feces. The most common aberrant forms (Ocular and cutaneous habronemiasis) are associated with the nematode larvae being deposited in these areas, not completing its life cycle, with resultant signs probably associated with local hypersensitivity¹⁴. Diagnosis is based on gross lesions and the seasonal

nature of the disease and may be confirmed by histopathology or identification of larvae from the lesion. A rapid onset and exuberant, pruritic granuloma at the medial canthus of the eye with small yellow, calcified concretions embedded in it are considered pathognomonic for ocular habronemiasis. A definitive diagnosis is established by histopathology as a granulation tissue infiltrated by eosinophils, mast cells, neutrophils, macrophages and an occasional *habronema* larva¹²⁾. Lesions are usually reddish-brown single or multiple subcutaneous nodules, mostly bloody and purulent discharge and contained numerous caseous granules, with occasional ulceration. They often are "greasy," serosanguineous, and contain yellow calcified "rice grain-like" material and some cases reveal extensive epidermal erosion and focal ulceration with dermal necrosis and other description as a single or multiple ulcerated tumorous masses with eosinophil, epithelioid cells infiltration and scattered necrotic debris. Moreover, diffuse and heavy dermal infiltration of numerous eosinophils and fewer neutrophils were noted^{14,2,22)}. In cutaneous habronemiasis, skin lesions are ulcerated; granulomatous masses that bleed easily appear as gritty plaques as a pathognomonic sign and characteristic histological lesions consist of eosinophilic dermatitis and coagulative necrosis with or without the degeneration of nematode larvae in the center^{20,15)}. Typical signs include no healing skin lesions, ulceration of moist areas, intense itching, and formation of exuberant granulomatous tissues^{9,18)}. Ulcerated fungating growths very liable to hemorrhage they consist of exuberant fibrous tissue enclosing few or many larvae of *Habronema* spp. accompanied by dense infiltration of eosinophils, which form small eosinophilic abscesses due to immobilized and dead larvae⁸⁾. Cutaneous habronemiasis in 15 horses and 5 donkeys was described by¹¹⁾; the lesions were distributed in many parts of the body involving the medial canthus, shoulder and pectoral regions, knee and fetlock joints, abdominal wall and

prepuce. Some animals had more than one lesion. The lesions were ulcerative and filled with soft light red granulation tissue. When curetted, the deeper layers revealed a dense fibrous tissue with calcified foci.

In dogs, ulcerative, non-healing skin lesions and the formation of exuberant granulation tissue were observed as in horses. Microscopically, the epidermis was ulcerated and overlaid with a thin layer of necrotic cellular debris admixed with neutrophils. The dermis and subcutis were markedly edematous and contained extensive infiltrates of inflammatory cells, predominantly eosinophils, with fewer plasma cells, macrophages, and lymphocytes. Variably-sized tracts of necrotic cellular debris admixed with fragmented collagen and large numbers of degenerate eosinophils were also present in the dermis¹⁸⁾.

A 6-year-old castrated dromedary camel (*Camelus dromedarius*) was presented with a non-healing, severely pruritic, ulcerative fibrotic plaque located at the medial canthus. Histological examination of surgical biopsies identified degenerating nematode larvae within eosinophilic granulomas. A biopsy of this lesion revealed multiple granulomas, areas of patchy ulceration, and proliferative granulation tissue with moderate eosinophilic inflammation extending from the dermis to the subcutaneous muscle¹³⁾.

In one case with the signs of bilateral conjunctivitis on the Rhinoceros, the histological evaluation of the excised conjunctiva showed reactive fibrosis and diffuse eosinophilic infiltration throughout the substantia propria. Larger eosinophil-rich granulomas surrounding transverse and vermiform longitudinal sections of parasites were also observed⁷⁾.

Phaeohyphomycosis is a general term for disorders caused by melanized fungi with hyphal histopathology. It covers a wide variety of clinical forms, including cutaneous and subcutaneous, central nervous system and disseminated infections. The mycoses should be distinguished from mycetoma and chromoblastomycosis, which

are also caused by dematiaceous fungi¹⁷), it is considerable opportunistic infection that occurs through contamination of wounds 1). In other others the microscopically lesions described as an epithelioid macrophages with fewer multinucleated giant cells, neutrophils, lymphocytes, and plasma cells infiltrated the dermis and sub cutis in vague coalescing granulomas. Numerous intra- and extracellular brown-pigmented fungi were present which characterized by coalescing pyogranulomas involving the dermis and subcutis. Within these pyogranulomas, pigmented, septate, fungal hyphae with multiple globose dilatations were identified. The characteristic feature of this group of fungi is their pigmented appearance in section with dark colored hyphae visible on H&E stained sections⁶). Cutaneous Onchocercosis can affect horses⁴). Adult forms and larvae of *Onchocerca* are responsible for the troubles as in study in the one hundred and sixty skin biopsies on horses under the study from Poland, France and Spain in order to detect *Onchocerca* larvae. Six foreign horses were found positive (5/95 Polish and 1/18 Spanish). Equine Onchocerciasis is a parasitic infection of horses, with world-wide distribution, caused by a filarial worm of the genus *Onchocerca*. The adults of *Onchocerca cervicalis* are found in the ligamentous tissue adjacent to the nuchal attachment of the thoracic vertebral spinous processes and in and around the supra spinous bursa¹⁰). There are no literatures on cutaneous habronemiasis in equine in Egypt, so this study aimed to describe the pathological features of a complicated cutaneous granuloma in a horse.

Materials and Methods

A 14 years old mare admitted in June 2014 to the Surgery Department, Faculty of Veterinary Medicine, Zagazig University, Egypt, with an unhealed ulcer skin near to the medial canthus of the left eye. The ulcerated skin lesion was 3 cm in diameter reddish in color with hazy irregular

borders and containing grayish and yellowish necrotic tissues with offensive odor with (Fig. 1A). The animal was subjected to nasolacrimal duct lavage with normal saline 0.9% to exclude nasolacrimal duct obstruction. Biopsy of lesion was taken under tranquilization of the animal by 1.1 mg/kg I/V (Xylazine Hcl 2% (Xylaject® Adwia) and fixed in 10% neutral buffered formalin solution, dehydrated in ascending concentrations of ethanol (70-100%), cleared in xylene, and embedded in paraffin. Five-micron thick paraffin sections were prepared and then routinely stained with hematoxylin and eosin¹⁹), and then examined microscopically. Other specimens were sending for microbiological identification. An extensive debridement of the lesion around the medial canthus of the left eye was performed under deep narcosis using Chloral hydrate 10% 5 gm/ 50 kg B.wt I.V.

Results

Grossly, large single red brown protruded masse with diameter 0.5 to 1 cm was seen from unhealed ulcerated skin near to the medial canthus of the left eye. The ulcerated skin lesion was 3 cm in diameter reddish in color with hazy irregular borders and containing grayish and yellowish necrotic tissues with offensive odor (see Fig.1A). Histopathological examination of the biopsy specimen showed multiple parasitic granulomas in the dermal tissue represented by caseous centers with or without degenerated *habronema* larvae containing masses from granulated eosinophil in the center followed by eosinophils and basophils necrotic debris beside numerous macrophage and foreign body giant cells, lymphocytes and plasma cells followed by fibrous capsule on the periphery (Figs 1 B, C, D). Also the surrounding tissue of skin, subcutis and muscles had massive eosinophil infiltrations. Also dermal collagen fiber suffered from collagenolysis represented by fragmented and distorted (necrosis) or hyalinized fibers

infiltrated by numerous inflammatory cells mainly living and degranulated eosinophils. The skeletal myocytes muscle cells showed vacuolation and edema with more eosinophilic sarcoplasm beside pyknosis of their nuclei. Large deep cutaneous ulcer containing purulent or serous exudate with living and degenerated neutrophils, sometimes admixed with identified fungi elements (mycelia) (phaeohyphomycosis) over the surface of parasitic granuloma described as a septate, fungal hyphae with multiple globose dilatations were identified (Fig.1E). Moreover, the inflammatory process was extended deeper forming pyogranuloma containing multinucleated giant cells and fibrous tissues. Other biopsy specimen from the fore mentioned lesion of the skin mainly in vicinity of sebaceous glands and muscle revealed aggregations of microfilaria (*Onchocerciasis*) in dermis with or without inflammatory reactions mainly eosinophils and mononuclear cells representing *Onchocerciasis* (Fig.1F).

Discussion

In this report the case was diagnosed by the history, clinical signs and histological finding. This is a complicated case of Cutaneous Habronemiasis in horse with Phaeohyphomycosis and *Onchocerciasis*. The exact pathogenesis of Habronemiasis is unknown but it is highly probable that the disease involves as a hypersensitive reaction to dead or dying larvae^{15,21}. In our work the Cutaneous *Habronema* located in preocular region near the medial canthus of left eye. This site was a common location. This finding was in agreement with^{15,5} who mentioned that the common sites for infestation are limbs, ventral aspect of the abdomen, prepuce, external genitalia males (penis, urethral process), ocular and peri-ocular areas (conjunctiva, medial canthus, nasolacrimal duct) and commissure of the lips. Habronemiasis in this report occurred in summer season and related to fly activity and this result

was coincided with that mentioned by²².

In our histopathological findings we observed multiple parasitic granulomas in the dermal tissue represented by caseous centers with degenerated *Habronema* larvae. This lesion was observed in previous studies^{15,20,12}. Similar microscopic features of the larvae and the inflammatory response were also previously reported in Cutaneous Habronemiasis of dog¹⁸, in camel¹³, in Rhinoceros⁷ and in donkey^{11,22}.

Clinical diagnosis is unreliable as there is a range of equine skin diseases that should be considered as differential diagnoses. These include proliferative granulation tissue, sarcoids, squamous cell carcinoma, fungal granulomas, pyogranulomas and foreign body granulomas. Differential diagnoses for ophthalmic Habronemiasis include ocular neoplasia, foreign body granuloma, ocular *Onchocerciasis* and Phycomycosis^{15,16}. The exact pathogenesis of Cutaneous Habronemiasis is unknown, however, a hypersensitivity reaction to the dead or dying larvae and the marked tissue eosinophilia are presumed to be involved²². The larvae were degenerated at the time the biopsy was performed, possibly as a result of the marked tissue eosinophilia induced by the parasite and the ivermectin dose given previously by the referring veterinarian¹³. In this study we suggest that intense eosinophil's infiltration accompanying larvae due to the role of eosinophil's and their granules in the caseated center and the immunological response for focal granuloma. Also macrophages and Giant cell usually infiltrating of the granuloma formation due to the role of the detectable T-helper lymphocytes. Exuberant fibrosis and collagenolysis usually seen in the vicinity of the dermal immunological reactions of larvae.

Grossly the skin overlying of parasitic granuloma in our case revealed large single unhealed ulcerated skin with hazy irregular borders and containing grayish and yellowish necrotic tissues with offensive odor. These lesions were in accordance with^{18,9,20,14,22} also described lesion as a

reddish-brown subcutaneous nodules which were “greasy,” serosanguineous, and contain yellow calcified “rice grain-like” material also our results agree with¹⁸⁾ in dog,¹³⁾ in camel,⁷⁾ in Rhinoceros and^{11,22)} in donkey.

Microscopically skin above parasitic granuloma in our case revealed large deep cutaneous ulcer containing purulent or serous exudate with living and degenerated neutrophils, sometimes admixed with fungal elements (mycelia) (Phaeohyphomycosis). This finding is in agreement with^{3,6)} which described the Cutaneous Phaeohyphomycosis as an ulcerated skin with dark colored hyphae covered by neutrophils and dermis diffusely infiltrated by numerous epithelioid

macrophages admixed with multinucleated giant cells, neutrophils, lymphocytes, plasma cells and rare eosinophils. In this study, aggregations of microfilaria in dermis with or without inflammatory reactions mainly eosinophils and mononuclear cells were found and appeared in vicinity of sebaceous glands of skin and muscle. Similar results were reported in horse's survey studies in France by⁴⁾. It could be concluded that complicated Cutaneous Habronemiasis consider one of the parasitic cutaneous lesions which is rare among horses in Egypt. Immunological reactions of the parasitic element are one of the etiological agents of skin ulcers and fungal invasions.




Fig 1: **A)** Ulcerated lesion on the skin at medial canthus of the left eye of a horse. **B)** Skin of horse showing three multiple degenerated *habronema* larvae surrounded by granulomatous reactions. H&E (X 300). **C)** High magnification of the previous picture to show numerous macrophages and giant cell (arrow) encircled by fibrous capsule infiltrated by eosinophils (short arrow). H&E(X 1200). **D)** Skin of horse showing degenerated *habronema* larvae embedded in caseated tissue. H&E (X 800). **E)** Skin of horse showing fungal filament (arrow) with intense neutrophils infiltrations within the inflammatory exudate. H&E

References

- 1) Ajello, L. 1986. Hyalohyphomycosis and phaeohyphomycosis, two global disease entities of public health importance. *Eur. J. Epidemiol.*, **2**: 243-251.
- 2) Amininajafi, F. and Mehrara, M. R. 2014. Histopathological features of cutaneous and gastric habronemiasis in horse. *J. Parasitol.*, DOI 10.1007/s12639-014-0611-7.
- 3) Eberly, J., Wilke, M. and Singh, K. 2015. Case Report: Cutaneous phaeohyphomycosis in a horse. *Online J. Vet. Res.*, **19**: 26-31.
- 4) Franck, M. T., Colombet, C., Hugnet, J. Ducos De Lahitte, Desmaizières L. M., Delverdier, M. and Franc, M. 2006. Research of skin microfilariae on 160 horses from Poland, France and Spain. *Revue Méd. Vét.*, **157**: 323-325.
- 5) Gasthuys, F. M., Van Heerden, M. and Vercreysse, J. 2004. Conjunctival habronemiasis in a horse in Belgium. *Vet. Rec.*, **154**: 757-758.
- 6) Genovese, L. M., Whitbread, T. J. and Campbell, C. K. 2001. Cutaneous Nodular Phaeohyphomycosis in Five Horses Associated with *Alternaria alternata* infection. *Vet. Record*, **148**: 55-56.
- 7) Horowitz, I. H. and Richard, R. 2015. Case Report Conjunctival habronemiasis in a square-lipped rhinoceros (*Ceratotherium simum*). *Vet. Ophthalmol.*, **19**: 161-166.
- 8) Jubb, K. V. F., Kennedy, P. C and polmer, N. 1985. Pathology of Domestic Animals, Third Edition. Volume **2**. pp 382.
- 9) Littlewood, J. D. 1999. Equine practice: control of ectoparasites in horses. *In Pract*, **21**: 418-424.
- 10) Mellor, P. S. 1973. Studies on *Onchocerca cervicalis* (Railliet and Henry 1910): I. *Onchocerca cervicalis* in British Horses. *J. Helminthol.*, **4**: 97-110.
- 11) Mohamed, F. H, Abu Samra, M. T. and Idris, S. O. 1989. Cutaneous habronemiasis in horses and domestic donkeys (*Equus asinus asinus*). *Rev. Elev. Med. Vet. Pays. Trop.*, **42**: 535-540.
- 12) Moore, C. P. and Sarazan, R. D. 1983. Equine ocular parasites: A review. *Equine Vet. J.*, **2**: 76-84.
- 13) Myers, D. A. and Smith, C. D. 2010. Cutaneous periocular Habronema infection in a dromedary camel (*Camelus dromedaries*). *Vet. Dermatol.*, **21**: 527-530.
- 14) Pugh, D. M., David, G. M. S, Dipl, N., Dipl, T., Xing Ping, H., and Byron, B. 2014. Habronemiasis: Biology, Signs, and Diagnosis, and Treatment and Prevention of the Nematodes and Vector Flies, *J. Equine Vet. Sci.*, **34**: 241-248.
- 15) Pusterla, N., Watson, J. L, Wilson, W. D., Affolter, V. K. and Spier, S. J. 2003. Cutaneous and ocular habronemiasis in horses: 63 cases (1988-2002). *J. Am. Vet. Med. Assoc.* **222**: 978-982.
- 16) Rebhun, W. C, Mirro, E. J, Georgi, M. E. and Kern, T. 1981. Habronemic Blepharoconjunctivitis in horses. *J. Am. Vet. Med. Assoc.* **179**: 469-472.
- 17) Revankar, S. G. 2007. Dematiaceous fungi. *Mycoses*, **50**: 91-101.
- 18) Sanderson, T. P. and Nito, D. Y. 1990. Cutaneous Habronemiasis in a Dog. *Vet. Path.*, **27**: 208-209.
- 19) Suvarna, S. K, Layton, C. and Bancroft, J. D. 2013. Bancroft's Theory and Practice of Histological Techniques. 7th ed., Churchill Livingstone. Elsevier, England.
- 20) Traversa, D. and Iorio, R. 2006. Molecular cross-sectional survey of gastric habronemiasis in horses. *Vet. Parasitol.*, **141**: 285-290.
- 21) Traversa, D., Iorio, R., Petrizzi, L., De Amicis, I., Brandt, S., Meana, A., Giangaspero, A. and Otranto, D. 2007. Molecular diagnosis of equid summer sores. *Vet. Parasitol. J.*, **150**: 116-121.
- 22) Yarmut, Y., Brommer, H., Weisler, S., Shelah, M., Komarovskiy, O. and Steinman, A. 2008. Ophthalmic and cutaneous habronemiasis in a horse: case report and review of the literature. *Israel J. Vet. Med.* **63**: 87-90.