



| | |
|------------------|---|
| Title | Diversity of lactic acid bacteria isolated from raw milk in Elsharkia province, Egypt |
| Author(s) | Alnakip, Mohamed E. A.; Mohamed, Asmaa S.; Kamal, Rania M.; Elbadry, Seham |
| Citation | Japanese Journal of Veterinary Research, 64(Supplement 2), S23-S30 |
| Issue Date | 2016-04 |
| Doc URL | http://hdl.handle.net/2115/62030 |
| Type | bulletin (article) |
| File Information | p.S23-30 Mohamed E. A. Alnakip.pdf |



[Instructions for use](#)

Diversity of lactic acid bacteria isolated from raw milk in Elsharkia province, Egypt

Mohamed E. A. Alnakip^{1,*}, Asmaa S. Mohamed¹, Rania M. Kamal¹ and Seham Elbadry²

¹Food control Department, Faculty of Veterinary Medicine, Zagazig University, 44519, Egypt

²Central Laboratory, Faculty of Veterinary Medicine, Zagazig University, 44519, Egypt

*Corresponding author: Mohamed E. A. Alnakip, Email: alnakip.me@gmail.com

Abstract

A total of 50 raw cow's milk samples were collected from different areas of Elsharkia province, Egypt for characterizing lactic acid bacteria (LAB) load. Using 16S rRNA gene sequencing, a total of 41 LAB isolates have been identified corresponding to *Enterococcus* sp. (51.22 %) as the most predominant LAB genus, followed in order by *Aerococcus* (26.82 %), *Lactococcus* (7.32 %), *Lactobacillus* (7.32 %), *Leuconostoc* (4.88 %) and *Pediococcus* (2.44 %) genera. All isolates were identified to species level with exception of one strain (*Lc. lactis* subsp. *cremoris*) that has been assigned to subspecies. The phylogenetic dendrogram created has allowed good discrimination between all isolated LAB species identified with this study. Results showed a wide diversity among isolated LAB from raw milk in Elsharkia province. The impact of LAB presence in raw cow's milk on dairy safety has been discussed.

Introduction

Lactic acid bacteria (LAB) are a group of Gram-positive bacteria widely distributed in different foods. LAB were first isolated from milk and subsequently discovered that LAB are occurring naturally as indigenous microflora in raw milk. This bacterial group is united by a constellation of morphological, metabolic, and physiological characteristics; generally by being non-sporing, non-respiring cocci or rods, and ferment carbohydrates with production of lactic acid as the major end product⁹. Nearly 30% of raw milk bacterial counts is related to LAB, and production conditions, season and animal species usually affecting their numbers and diversity⁴⁵. From a practical dairy technology point of view, the following genera are considered the principal LAB: *Aerococcus*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Pediococcus* and

Streptococcus.

LAB presence in raw milk may be attributed to various origins, which can explain their diversity among seasons, animal species, etc. They can directly come from milk, but also from the surrounding animals' environment. Indeed, *Leuconostoc* sp. come from vegetation and roots but can easily propagate and persist in various niches which later on contaminate raw milk²⁷. The ubiquitous genera *Lactococcus* and *Lactobacillus* may come from plants, feces or udder skin. Meanwhile, enterococci mainly inhabit milk as a result of fecal pollution of either human or animal routes²⁹.

The discrimination of LAB into different genera is largely based on morphology, mode of glucose fermentation, growth at different temperatures, configuration of the resulted lactic acid, ability to grow at high salt concentrations, and acid or alkaline tolerance⁹. However, these methods

turned unfit nowadays due to time consuming nature, huge amount of materials and labor in addition to low diagnostic specificity and sensitivity^{3,6,40}. Chemotaxonomic markers such as fatty acids composition as well as constituents of the cell wall are also used in classification⁹. New tools for classification and identification of LAB are currently replacing and/or complementing the traditional phenotype-based methodologies such as PCR-based fingerprinting and protein fingerprinting techniques^{3,6,40,41}. In this sense and on the basis of the 16S rRNA gene sequencing (GS), the bacterium can then be identified and assigned to species or even subspecies level against phylogenetically-related strains located in different databases (e.g. NCBI GenBank). The aim of this work is to characterize the predominant LAB isolated from raw cow milk samples collected from Elsharkia province, Egypt, based on 16S rRNA GS and to discuss the impacts of isolated species on dairy safety.

Materials and Methods

Collection of milk samples: Fifty raw cow's milk samples were collected from different individual households in Elsharkia province. About 50 mL of each milk sample was aseptically collected and transported to laboratory in a 4°C vehicle-mounted refrigerator to be analyzed microbiologically within few hours.

Isolation of LAB from raw milk samples: Serial dilutions were made for each sample using 0.85% sterile physiological saline and 0.1 mL of each dilution was spread plated in duplicates of de Man, Rogosa, and Sharpe agar (MRS) (Difco Labs, Detroit, MI) adjusted to PH of 5.5¹⁴. Plates were incubated anaerobically (BBL Gas pak plus Anaerobic Sys.) at 30°C for 48 h. Colonies with distinct morphological differences were selected from each plate and further purified by re-streaking two successive times on fresh MRS plates. All isolates were maintained as frozen

cultures in MRS broth and 50% glycerol at -80 °C.

Identification of LAB isolates and Phylogenetic Analysis Based on 16S rRNA GS: All procedures were done as previously described by^{3,6,10,41}. Total genomic DNA was extracted from overnight cultures. The bacterial cells were lysed by the addition of 180 µL of lysis solution (Sigma-Aldrich) after incubation for 2 h at 37 °C. Total genomic DNA was extracted and purified using the DNeasy Tissue Mini Kit (Qiagen). A fragment of the 16S rRNA gene was amplified by PCR using the universal primer pair: p8FPL(5'-AGTTTGATCCTGGCTCAG-3') and p806R (5'-GGACTAC-CAGGGTATCTAAT-3'). All of the PCR assays were performed using a "My Cycler" Thermal Cycler (BioRad Labs, USA). Direct sequencing was performed using the "BigDye Terminator v3.1" Cycle Sequencing Kit (ABs, Perkin-Elmer, Foster city, CA) and the same primers used for PCR were also used for the sequencing. The sequencing reactions were analyzed in ABI3130 automatic GS sys. (ABs, USA). Entire 16S rRNA gene sequences were analyzed using Chromas software and aligned with Clustal-X software⁴⁴. Next, these sequences were identified by sequence homology alignment among published reference sequences using the web tool; NCBI BLAST (<http://blast.ncbi.nlm.nih.gov/>)⁸. Consensus sequences were imported into MEGA 6.0 software, with which a sequence alignment and phylogenetic trees were conducted based on the NJ method and Kimura-2 parameter model.

Rooting-out the phylogenetic dendrogram: 16S rRNA gene sequences of six strains of LAB from previous studies^{3,5,41} plus two *Lc. lactis* subsp. *lactis* strains from NCBI GenBank (gi_387286036, gi_387286035) were used to root-out the phylogenetic dendrogram (Table 1).

Table 1. Out-group strains used in the phylogenetic dendrogram.

| Strain code | Species | GenBank Accession no. |
|-------------|---|-----------------------|
| CECT4039 | <i>E. faecalis</i> | KC510231 |
| CECT410 | <i>E. faecium</i> | KC510233 |
| USC13 | <i>S. dysgalactiae</i> subsp. <i>dysgalactiae</i> | KC510218 |
| USC17 | <i>S. dysgalactiae</i> subsp. <i>dysgalactiae</i> | KC510219 |
| USC30 | <i>Lc. lactis</i> | KP001333.1 |
| USC31 | <i>Lc. lactis</i> | KP001334.1 |

Results and Discussion

During last decade, genotypic identification has been emerged as an alternative or a complement to established phenotypic methods within dairy diagnostics providing more accuracy, less labor and time saving. Among the phylogenetic marker genes used to discriminate among different species, 16S rRNA is well-established as a universal gold standard for the identification and phylogenetic classification of prokaryotic species, genera, and families³⁶. Thus, 16S rRNA GS has been applied extensively within food safety diagnostic labs proving powerful identification and discrimination potentials^{3,6,10,41}.

The partial 16S rRNA GS (800 bp) of all the strains isolated in our study were compared with related bacteria in GenBank and sequence similarities were determined using the BLAST tool. The resulted 16SrRNA gene sequences of different isolates have been submitted to be deposited in the GenBank. Based on 16S rRNA GS, forty one strains of LAB identified in our work were corresponding to Enterococcus (51.22 %) as the most predominant LAB genus, followed in order by *Aerococcus* (26.82 %), *Lactococcus* (7.32 %), *Lactobacillus* (7.32 %), *Leuconostoc* (4.88 %)

and *Pediococcus* (2.44 %) genera (Table 2).

Table 2: Isolated LAB strains in this study.

| Isolated LAB | No. | % |
|--|-----------|------------|
| <i>E. durans</i> | 1 | 2.44 |
| <i>E. faecium</i> | 8 | 19.51 |
| <i>E. hirae</i> | 2 | 4.88 |
| <i>E. faecalis</i> | 6 | 14.63 |
| <i>E. casseliflavus</i> | 2 | 4.88 |
| <i>E. saccharolyticus</i> | 2 | 4.88 |
| <i>Pediococcus pentosaceus</i> | 1 | 2.44 |
| <i>Lc. garviae</i> | 2 | 4.88 |
| <i>Lc. lactis</i> subsp. <i>cremoris</i> | 1 | 2.44 |
| <i>Lb. plantarum</i> | 1 | 2.44 |
| <i>Lb. casei</i> | 1 | 2.44 |
| <i>Lb. fermentum</i> | 1 | 2.44 |
| <i>Leuc. mesentroides</i> | 2 | 4.88 |
| <i>A. viridians</i> | 11 | 26.82 |
| Total | 41 | 100 |




Figure 1: Phylogenetic tree of LAB isolates based on 16S rRNA GS

Out of all isolates and according to identification evolved from 16S rRNA GS, 40 isolates (97.56 %) have been identified to the species levels, and only for one *Lc. lactis* subsp. *cremoris* isolate (2.44 %), the identification has been extended to the subspecies level. Nearly similar to our results, a study by²⁴⁾ showed that 94% of isolated LAB in cow's milk included lactococci, enterococci and streptococci, while the remaining 6% isolates were lactobacilli (mostly *Lb. casei*, *Lb. delbrueckii*, *Lb. paracasei* and *Lb. plantarum*), *Leuconostoc* and pediococci. Also, other study by²⁾, most LAB recovered from raw cow milk samples in Khartoum, Sudan were corresponded to *Enterococcus*, *Lactococcus* and *Lactobacillus* species. Among relevant isolates in another investigation¹⁶⁾, *Streptococcus*, *Enterococcus* and *Aerococcus* were corresponding to 52, 26 and 15 %, respectively and remaining % were lactococci. The phylogenetic dendrogram has been created using MEGA 6.0 software as shown in Figure 1. Remarkably, 16S rRNA GS has allowed for a very good discrimination among all isolated LAB species with high bootstrap values. The phylogenetic tree has been separated into 2 main branches. The first branch included *Enterococcus*, *Streptococcus* and *Lactococcus* sp., corresponded to A, B and C sub-branches respectively. Meanwhile, the second branch included *Lactobacillus*, *Pediococcus*, *Leuconostoc* and *Aerococcus* sp., which corresponded to D, E, F and G sub-branches, respectively.

Presence of LAB in raw milk could be good candidates in biopreservation of processed dairy products¹⁵⁾, especially fermented foods such as mature cheeses, cream and yoghurt. LAB have an essential role in the nutritious and organoleptic properties of fermented milk production¹¹⁾. As a result of lowered pH following sugar fermentation and acid production, the development of the desirable organoleptic properties occurs¹²⁾. Additionally, antimicrobial potentials have been linked to some LAB^{32,37)}. In this sense, *Leuc. mesenteroides* sp. *mesenteroides* FR 52, isolated

from a raw milk, produced a bacteriocin which was named Mesenterocin 52³²⁾. This bacteriocin inhibited other *Leuconostoc* strains and several strains of *Enterococcus* and *Listeria* spp. For this, most LAB are belonging to the qualified Presumption of Safety (QPS) and generally recognized as safe (GRAS) lists which insure their safety for use in food⁴²⁾. However, some LAB are excluded from these advantages as *Enterococcus* sp. because of their roles in causing certain human infections and contribution to spread of antibiotic resistance³³⁾ and more importantly, their presence in milk could indicate unsanitary production and fecal pollution of either human or animal routes or both as they are ubiquitously found in the intestinal microflora of humans and animals²⁹⁾. Unfortunately, our results indicated the unsanitary production associated with raw milk as *Enterococcus* sp. constituted alone 51.22% out of all LAB isolates. Enterococci are among predominant isolated LAB from raw milk^{1,7,16,24)}. Also, enterococci are well-known to be minor mastitis pathogens causing subclinical mastitis (SCM) in dairy animals with no apparent signs, or clinical form with abnormal milk, swelling of the udder, and fever^{4,16)}.

In one study¹⁶⁾, several enterococcal species have been isolated from subclinical intramammary infections (IMIs) in dairy cows corresponded to *E. faecalis*, *E. faecium*, *E. durans* and *E. hirae*. Also, other investigations showed isolation of enterococci as predominant LAB from raw milk, and the most isolated enterococcal species were *E. faecalis*, *E. faecium*^{1,7)} and *E. durans*²⁴⁾. Like streptococcal IMIs, those caused by enterococci may represent poor responsiveness to antibiotic therapy³⁸⁾. Biofilm-formation by enterococci is thought to contribute to this resistance³⁵⁾. Enterococci can be responsible for variety of defects in processed dairy products such as cheese, causing excessive softening, splits and cracks, off flavors and abnormal colors³⁰⁾. In human, enterococci are incriminated as direct or indirect agents of diseases^{18,25)}. Some enterococci can cause food-poisoning especially *E. faecium*

if predominated in the food. Enterococcal food-intoxication caused is greatly attributed to production of biogenic amines²⁵). In addition, *E. faecalis* has associated with a large number of gastroenteritis outbreaks and implicated in urinary tract and wound infections, intra-abdominal abscesses and endocarditis¹⁸). It is thought that enterococcal toxins behaving and producing symptoms similar to but less acute than those of staphylococcal enterotoxins²⁸).

Several *Lactococcus*, *Lactobacillus* and *Leuconostoc* species have potential technological applications within dairy industry^{21,22,24}). In other studies, *Lb. plantarum*, *Lb. fermentum*, *Lb. acidophilus*, *Lb. paracasei* and *Lb. rhamnosus* were the frequent isolated lactobacilli^{2,31}). In our study, isolated lactobacilli were only limited to 3 species in lowered incidences; *Lb. plantarum*, *Lb. fermentum* and *Lb. casei* (1 strain of each). Among isolated laccocci, *Lc. garvieae* constituted 4.88 % out of LAB. *Lc. garvieae* has been reported recently as a majority component of the autochthonous microbial populations of certain artisanal cheeses²¹) and fermented milk products¹⁷). *Lc. garvieae* cause lactococcosis in fish due to owing several virulence factors, meanwhile, *Lc. garvieae* isolates of dairy origin have shown absence of virulence determinants²²), suggesting that *Lc. garvieae* dairy strains are unrelated to the pathogenic ones²³). However, the isolation of *Lc. garvieae* from milk of bovines with SCM was reported^{16,38}). Moreover, isolates of *Lc. garvieae* of dairy origin are incriminated to carry antibiotic resistance genes⁴⁶), which might contribute to antibiotic resistance in both animal and human. Also, *Leuc. mesentroides* has been identified recently to be as a sporadic infectious agent in immune-compromised humans¹³), but further studies are needed to show if strains of dairy origin are possessing such virulence factors.

Aerococcus sp. exhibit many biochemical and physiological similarities with *Pediococcus*, *Enterococcus*, *Lactococcus* and *Leuconostoc* species, and are often confused with *Streptococcus* sp.²⁰).

A. viridans is a catalase-negative Gram-positive cocci resembling staphylococci by Gram stain, but have biochemical and growth characteristics of streptococci and enterococci¹⁹). The typing of *A. viridans* by some commercial biochemical systems may be not sufficient to achieve 100% identification accuracy⁴³), thus, the genotypic typing was recommended. In our study, accurate identification to the species level in 100% of *Aerococcus* sp. isolates based on 16S rRNA GS has been obtained, which agreed with recommendation by⁴³) for *A. viridians* identification. In current study, *A. viridians* has been isolated with incidence of 26.82%. In a study by⁴⁷), *A. viridans* was detected in 50% of 48 bulk tank milk samples from 48 dairy farms in USA. The contribution of *A. viridans* in causing clinical, subclinical or latent types of mastitis have been described^{16,43}), although the exact role of the m.o was not clear. It has reported in a study by¹⁶) that *A. viridans* constituted 15% of Gram-positive, catalase-negative, aesculin-degrading cocci isolated from clinical and subclinical bovine mastitic cases, but in a different study³⁴), only two *A. viridans* isolates were detected among 100 isolates incriminated in causing mastitis. In Humans, many infections such as endocarditis, urinary tract infections, arthritis, or meningitis have been associated with *A. viridans*^{26,39}). However and like *Lc. garvieae*, it must be demonstrated if dairy *A. viridans* isolates are possessing virulence factors of human-clinically associated strains.

Conclusion

A wide diversity in LAB isolated from raw cow milk samples collected from Elsharkia province, has been shown. Several beneficial LAB have been isolated and could be of potential application within future researches. On the other hand, the results showed presence of enterococci in high incidence which reflected bad sanitary production processes within individual households, which

reflect necessity to commitment to healthy specifications and showed that further attention of the health authorities towards individual households should be directed.

References

- 1) AbdElAziz, A., AbdElAziz, S., Hassan, N., AbdAllah, W. and Niazi, Z. 2001. Occurrence of aerobic food-poisoning microorganisms in raw milk and locally manufactured processed cheese. *Zag. Vet. J.*, **29**: 71-76.
- 2) Ali, A. A. 2011. Isolation and identification of LAB from raw cow milk in Khartoum State, Sudan. *Int. J. dairy sci.*, **6**: 66-71.
- 3) Alnakip, M. E. 2014. Development of molecular methods of bacterial identification in food products by means of genomic and proteomic tools. PhD. Fac. Vet. Med.-Zagazig Univ, Egypt.
- 4) Alnakip, M. E., Quintela-Baluja, M., Böhme, K., Fernández-, I. C., Caamaño-Antelo, S., Calo-Mata P. and Barros-Velázquez J. 2014. The immunology of mammary gland of dairy ruminants between healthy and inflammatory conditions. *J. Vet. Med.*, **2014**: 1-32. doi: 10.1155/2014/659801.
- 5) Alnakip, M. E., Quintela-Baluja, M., Böhme, K., Fernández-No, IC., Bayoumi, M. A., Caamaño-Antelo, S., Velázquez, J. B. and Calo-Mata P. 2015. Comparative discrimination of *Streptococcus* sp. incriminated in bovine mastitis by MALDI-TOF MS fingerprinting versus 16S rRNA gene sequencing-based identification. Under Publication.
- 6) Alnakip, M. E., Quintela-Baluja, M., Böhme, K., Fernández-No, I. C., Amer, I. H., Elsayed, M. S., Ayoub, M. A., Barros-Velázquez, J. and Calo-Mata, P. 2014. Comparative identification of streptococci of dairy origin by VITEK-2 system and 16S rRNA gene sequencing. In: *First Inter. Conf. of Impact of Environmental Hazards on Food Safety*, Zagazig Univ., Egypt: 30-58.
- 7) Altarazi, Y., Alzamil, A. H. and Shaltout, F. 2002. Microbiological status of raw cow milk marketed in northern Jordan. *9th Vet. Med. Zag. Conf.*, Hurgada, Egypt.
- 8) Altschul, S., Gish, W., Miller, W., Myers, E. and Lipman, D. 1990. Basic local alignment search tool. *J. Mol. Biol.*, **215**: 403-410.
- 9) Axelsson L. 2004. Lactic acid bacteria: classification and physiology. In: *Lactic Acid Bacteria: Microbiological and Functional Aspects*, 3rd Ed., pp. 1-66.
- 10) Caamaño-Antelo, S., Fernández-No, I. C., Böhme, K., Ezzat-Alnakip, M., Quintela-Baluja, M., Barros-Velázquez, J. and Calo-Mata P. 2015. Genetic discrimination of foodborne pathogenic and spoilage *Bacillus* spp. based on three housekeeping genes. *Food Microbiol.*, **46**: 288-298.
- 11) Caplice, E. and Fitzgerald, G. F. 1999. Food fermentations: role of microorganisms in food production and preservation. *Int. J. Food Microbiol.*, **50**: 131-149.
- 12) Corsetti, A., Settanni, L., López, C. C., Felis, G. E., Mastrangelo, M. and Suzzi, G. 2007. A taxonomic survey of lactic acid bacteria isolated from wheat (*Triticum durum*) kernels and non-conventional flours. *Syst. Appl. Microbiol.*, **30**: 561-571.
- 13) Cuervo, S. I., Cortés, J., Rodríguez, E., Hormaza, N. and Vargas, E. 2008. *Leuconostoc* sp. en pacientes con cáncer: *Estudio descriptivo*, **25**: 184-188.
- 14) De Man, J., Rogosa, D. and Sharpe, M. E. 1960. A medium for the cultivation of lactobacilli. *J. Appl. Bacteriol.*, **23**: 130-135.
- 15) Delavenne, E., Mounier, J., Déniel, F., Barbier, G. and Le Blay, G. 2012. Biodiversity of antifungal lactic acid bacteria isolated from raw milk samples from cow, ewe and goat over one-year period. *Int. J. Food Microbiol.*, **155**: 185-190.
- 16) Devriese, L., Homme, J., Laevens, H., Pot B., Vandamme, P. and Haesebrouck, F. 1999. Identification of aesculin-hydrolyzing streptococci, lactococci, aerococci&enterococci

- from subclinical intramammary infections in dairy cows. *Vet. Microbiol.*, **70**: 87-94.
- 17) El-Baradei, G., Delacroix-Buchet, A. and Ogier J. 2008. Bacterial biodiversity of traditional Zabady fermented milk. *Int. J. Food Microbiol.*, **121**: 295-301.
 - 18) Eley, A. R. 1996. Microbial food poisoning. Chapman & Hall.
 - 19) Facklam, R. and Elliott, J. A. 1995. Identification, classification, and clinical relevance of catalase-negative, gram-positive cocci, excluding the streptococci and enterococci. *Clin. Microbiol. Rev.*, **8**: 479-495.
 - 20) Facklam, R., Hollis, D. and Collins, M. D. 1989. Identification of gram-positive coccal and coccobacillary vancomycin-resistant bacteria. *J. Clin. Microbiol.*, **27**: 724-730.
 - 21) Fortina, M., Ricci, G., Acquati, A., Zeppa, G., Gandini, A. and Manachini P. 2003. Genetic characterization of some LAB occurring in an artisanal protected denomination origin (PDO) Italian cheese, the Toma piemontese. *Food Microbiol.*, **20**: 397-404.
 - 22) Fortina, M., Ricci, G., Foschino, R., Picozzi, C., Dolci, P., Zeppa, G., Coccolin, L. and Manachini P. 2007. Phenotypic typing, technological properties and safety aspects of *Lactococcus garvieae* strains from dairy environments. *J. Appl. Microbiol.*, **103**: 445-453.
 - 23) Foschino, R. and Nucera, D., Volponi, G., Picozzi, C., Ortoffi, M. and Bottero, M. 2008. Comparison of *Lactococcus garvieae* strains isolated in northern Italy from dairy products and fishes through molecular typing. *J. Appl. Microbiol.*, **105**: 652-662.
 - 24) Franciosi, E., Settanni, L., Cavazza, A., Poznanski, E. 2009. Biodiversity and technological potential of wild lactic acid bacteria from raw cows' milk. *Int. Dairy J.*, **19**: 3-11.
 - 25) Giraffa G. 2002. Enterococci from foods. *FEMS Microbiol. Rev.*, **26**: 163-171.
 - 26) Gopalachar, A., Akins, R. L., Davis, W. R. and Siddiqui, A. A. 2004. Urinary tract infection caused by *Aerococcus viridans*, a case report. *Med. Sci. Monit.*, **10**: CS73-75.
 - 27) Hemme, D. and Foucaud-Scheunemann, C. 2004. *Leuconostoc*, characteristics, use in dairy technology and prospects in functional foods. *Int. Dairy J.*, **14**: 467-494.
 - 28) Hobbs, B. and Roberts, D. 1993. Food poisoning and food hygiene. 6th Ed.
 - 29) Kagkli, D. M., Vancanneyt, M., Hill, C., Vandamme, P. and Cogan, T. M. 2007. *Enterococcus* and *Lactobacillus* contamination of raw milk in a farm dairy environment. *Int. J. Food Microbiol.*, **114**: 243-251.
 - 30) Marth, E. H. and Steele, J. 2001. Applied dairy microbiology. 2nd Ed., CRC Press, Marcel Dekker, Inc., Madison, New York.
 - 31) Mathara, J. M., Schillinger, U., Kutima, P. M., Mbugua, S. and Holzapfel, W. H. 2004. Isolation, identification and characterisation of the dominant microorganisms of kule naoto: the Maasai traditional fermented milk in Kenya. *Int. J. Food Microbiol.*, **94**: 269-278.
 - 32) Mathieu, F., Suwandhi, I. S., Rekhif, N., Milliere, J. and Lefebvre, G. 1993. Mesenterocin 52, a bacteriocin produced by *Leuconostoc mesenteroides ssp. mesenteroides* FR 52. *J. Appl. Bacteriol.*, **74**: 372-379.
 - 33) Mathur, S. and Singh, R. 2005. Antibiotic resistance in food lactic acid bacteria: a review. *Int. J. Food Microbiol.*, **105**: 281-295.
 - 34) McDonald, W., Fry, B. and Deighton, M. 2005. Identification of *Streptococcus* spp. causing bovine mastitis by PCR-RFLP of 16S-23S ribosomal DNA. *Vet. Microbiol.*, **111**: 241-246.
 - 35) Metzger, S. 2008. Biofilm formation by *Enterococcus* species of bovine mammary gland and environmental origins. <http://hdl.handle.net/1811/32126>
 - 36) Patel, J. B. 2001. 16S rRNA gene sequencing for bacterial pathogen identification in the clinical laboratory. *Mol. Diag.*, **6**: 313-321.
 - 37) Pfeiler, E. A. and Klaenhammer, T. R. 2007. The genomics of lactic acid bacteria. *Trends Microbiol.*, **15**: 546-553.

- 38) Pitkälä, A., Haveri, M., Pyörälä, S., Myllys, V. and Honkanen-Buzalski, T. 2004. Bovine mastitis in Finland 2001; prevalence, distribution of bacteria, and antimicrobial resistance. *J. Dairy Sci.*, **87**: 2433-2441.
- 39) Popescu, G., Benea, E., Mitache, E., Piper, C. and Horstkotte, D. 2005. An unusual bacterium, *Aerococcus viridans*, and four cases of infective endocarditis. *J. Heart Valve Dis.*, **14**: 317-319.
- 40) Quintela-Baluja, M., Böhme, K., Fernández-No, I., Alnakip, M. E., Caamaño, S., Barros-Velázquez, J. and Calo-Mata, P. 2014. MALDI-TOF Mass Spectrometry, a Rapid and Reliable Method for the Identification of Bacterial Species in Food-Microbiology Laboratories. In: *Novel Food Preservation and Microbial Assessment Techniques*: 353-385.
- 41) Quintela-Baluja, M., Böhme, K., Fernández-No, I., Morandi, S., Alnakip, M. E., Caamaño, S., Barros-Velázquez, J. and Calo-Mata P. 2013. Characterisation of different food-isolated Enterococcus strains by MALDI-TOF mass fingerprinting. *Electrophoresis*, **34**: 2240-2250.
- 42) Rossetti, L., Carminati, D., Zago, M. and Giraffa, G. 2009. A qualified presumption of safety approach for the safety assessment of Grana Padano whey starters. *Int. J. Food Microbiol.*, **130**: 70-73.
- 43) Špaková, T., Elečko, J., Vasil, M., Legáth, J., Pristaš, P. and Javorský, P. 2012. Limited genetic diversity of *Aerococcus viridans* strains isolated from clinical and subclinical cases of bovine mastitis in Slovakia. *Polish J. Vet. Sci.*, **15**: 329-335.
- 44) Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. and Higgins, D. G. 1997. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.*, **25**: 4876-4882.
- 45) Verdier-Metz, I., Michel, V., Delbes, C. and Montel, M. 2009. Do milking practices influence the bacterial diversity of raw milk?. *Food Microbiol.*, **26**: 305-310.
- 46) Walther, C., Rossano, A., Thomann, A. and Perreten, V. 2008. Antibiotic resistance in *Lactococcus* species from bovine milk: Presence of a mutated multidrug transporter mdt (A) gene in susceptible *Lact. garvieae* strains. *Vet. Microbiol.*, **131**: 348-357.
- 47) Zadoks, R., Gonzalez, R., Boor, K. and Schukken, Y. 2004. Mastitis-causing streptococci are important contributors to bacterial counts in raw bulk tank milk. *J. Food Prot.*, **67**: 2644-2650.