



Title	First total synthesis of (+)-epogymnolactam, a novel autophagy inducer
Author(s)	Okado, Yuji; Shigetomi, Kengo; Mitsuhashi, Shinya; Ubukata, Makoto
Citation	Journal of antibiotics, 68(12), 721-724 https://doi.org/10.1038/ja.2015.63
Issue Date	2015-12
Doc URL	http://hdl.handle.net/2115/62305
Type	article (author version)
File Information	71886(Ubukata).pdf



[Instructions for use](#)

First Total Synthesis of (+)-Epogymnolactam, a Novel Autophagy Inducer

Yuji Okado, Kengo Shigetomi, Shinya Mitsuhashi, and Makoto Ubukata*

Division of Applied Bioscience, Graduate School of Agriculture, Hokkaido University,
Kita-9 Nishi-9, Kita-ku, Sapporo, 060-8589, Japan.

*Correspondence: Professor M. Ubukata, Research Faculty of Agriculture, Hokkaido
University, Kita-9, Nishi-9, Kita-ku, Sapporo 060-8589, Japan.
E-mail: m-ub@for.agr.hokudai.ac.jp

Abstract

A novel autophagy inducer, (+)-epogymnolactam (**1**) was first synthesized from *cis*-4-benzyloxy-2-butene-1-ol (**2**) in 8 steps. A reliable preparation of optically pure epoxy alcohol (+)-**3** from monobenzyl derivative (**2**) was established by a tandem strategy, Sharpless epoxidation/lipase kinetic resolution.

INTRODUCTION

(+)-Epogymnolactam (**1**) was discovered as a novel autophagy inducer from a mycelial culture of *Gymnopus* sp. in our laboratory (Fig. 1).¹ Autophagy is one of the major intracellular degradation systems in eukaryotic cells, eliminating damaged organelles and protein aggregates to maintain cytoplasmic homeostasis. This degradation pathway plays important roles in such diseases as cancer, neurodegenerative and infectious diseases. Thus, the application of autophagy inducer would help to understand the regulatory roles of autophagy in human diseases, and provide insight into the development of therapeutic agents that target autophagy.²⁻⁵ As an example of the effort for the development of autophagy-inducing drug, a peptide has been reported to have benefits in the clearance of a model polyglutamine expansion protein aggregates in HeLa/htt103Q cells, inhibition of intracellular survival of the bacterium, *Listeria monocytogenes*, inhibition of HIV-1 replication in human monocyte-derived macrophages, and a reduction in the mortality of neonatal mice infected chikungunya virus and West Nile virus.⁶ Although researchers have identified different types of autophagy inducers, e.g. rapamycin, an inhibitor of mTORC1;⁷ lithium L-690330, an inhibitor of IMPase;⁸ verapamil, Ca²⁺ channel blocker;⁹ resveratrol, activator of sirtuin 1 and inhibitor of S6 kinase;¹⁰ clonidine, an imidazole-1 receptor agonist;⁹ minoxidil, a K⁺ATP channel opener;⁹ spermidine, endogenous anti-aging mediator;¹¹ α -ketoglutarate, inhibitor of ATP synthase¹² and so on, none of these compounds is similar to **1** in chemical structure.

The structure of **1** deduced by spectroscopic analysis resembled to (+)-cerulenin,¹³ a potent inhibitor of fatty acid synthesis,¹⁴⁻¹⁶ and the absolute structure of **1** was assigned by the comparison of its specific rotation with that of (+)-cerulenin.¹ To evaluate chemical and biological properties of **1** more precisely, we needed to synthesize enough amount of **1** in enantiomerically pure form. Here we report the first total synthesis, and

thus structural confirmation of **1** by direct comparison of the natural product with the synthetic compound.

RESULT and DISCUSSION

Among the total syntheses of (+)-cerulenin, the concise synthesis by Townsend group¹⁷ seemed to be most effective. Optically pure (+)-cerulenin was synthesized with use of the coupling reaction of a chiral oxiranyllithium with a side-chain aldehyde as a key step. (+)-Epogymnolactam (**1**) would be synthesized in 10 steps starting from propargyl alcohol, and the number of reaction steps in the synthetic route was shorter than any other known synthetic methods from glucose,^{18,19} tartaric acid,²⁰ or a four-carbon synthon obtained by Sharpless epoxidation.²¹ We decided, however, to develop the straightforward synthesis of (+)-**1** which could be achieved in fewer steps by using the enantiomer of Sudalai's epoxy alcohol (96% ee, as TBS-alternate of (-)-**3**)²² synthesized via Sharpless asymmetric epoxidation using (+)-DET as a chiral source. Nevertheless, we could not reproduce such a high enantioselectivity in the synthesis of TBS alternate of (+)-**3** using (-)-DET. In general, Sharpless epoxidation of *cis* allylic alcohol has been shown not to give high enantiomeric excess especially in the large-scale preparation in a reproducible fashion. Sharpless epoxidation of *cis*-4-benzyloxy-2-buten-1-ol **2** resulted in 89% ee similar to the observation by Terashima group.²³ We tried to obtain enantiopure (+)-**3** by a recrystallization of 3,5-dinitrobenzoate of **3** followed by alkaline hydrolysis,²⁴ whereas we could not obtain an acceptable result, and abandoned optimization of this procedure, because a three-step process involving dinitrobenzoylation, recrystallization, and hydrolysis was needed in any case.

Next we searched for the best conditions to obtain enantiopure (+)-**3** by a lipase-mediated kinetic resolution of the corresponding acetate prepared by acetylation of **3** (89% ee). Epoxy alcohol (+)-**3** could be obtained with up to 96% ee by hydrolysis of the acetylated precursor with porcine pancreatic lipase (PPL), unfortunately this procedure did not give reproducible results and gave mostly unsatisfactory enantioselectivity less than 90% ee.²⁵

Finally we devised the most reliable procedure to prepare enantiopure (+)-**3** (99 to 100% ee) by treating **3** (89% ee) with PPL in vinyl acetate²⁶ as shown in Scheme 1. This type of tandem strategy for preparation of epoxy alcohols could be generally useful

because Sharpless epoxidation has been applied for tremendous number of allylic alcohol but it was difficult to obtain epoxy alcohol having nearly 100% ee. We believe this tandem strategy, Sharpless epoxidation/lipase kinetic resolution for preparation of enantiopure epoxy alcohol becomes one of the standard methods in organic synthesis.

The first total synthesis of (+)-**1** was achieved in a straightforward route outlined in Scheme 2. Dess-Martin oxidation²⁷ of **3** afforded aldehyde **4** in 91% yield. Large-scale preparation of **4** was done by cost-effective TEMPO oxidation²⁸ whose yield was 85%. Grignard reaction of **4** with *n*-BuMgCl in THF at -78 °C followed by deprotection of benzyl group of **5** with hydrogen and palladium/carbon catalyst in EtOAc at room temperature gave desired epoxy diol **6** in 53% yield over two steps. TEMPO oxidation of **6** in the presence of 2.2 eq. of NaOCl²⁹ at 0 °C provided epoxy lactone **7** in 78% yield. Two diastereomers could be separated by silica gel column chromatography (EtOAc : hexane = 1 : 4). Ammonolysis of **7** with NH₃ in MeOH at 0 °C furnished desired amide alcohol **8** in 99% yield. All synthetic intermediates **5**, **6**, **7**, and **8** existed as a mixture of two diastereomers, while no inconvenience in the structure determinations of these intermediates by NMR analysis. Oxidation of the both two diastereomeric alcohols should primarily generate the open-chain form **1a**. The amide alcohol **8** was successfully converted into (+)-**1** by Dess-Martin periodinane in CH₂Cl₂ at room temperature in 76% yield. Analyses of ¹H and ¹³C NMR showed that synthetic (+)-**1** existed as a ring-chain tautomeric mixture of ketoamide (**1a**) and diastereomeric hydroxy lactams (**1b** and **1c**) in CD₃OD as in the case of natural (+)-**1**. The physicochemical properties and autophagy inducing activity of synthetic (+)-**1** were consistent with those of natural epogymnolactam. Therefore, the absolute configuration of natural epogymnolactam was unambiguously confirmed as shown in Fig. 1.

Given the enough amount of synthetic (+)-**1**, we first decided to clarify the ratio of three isomers, keto isomer **1a**, major cyclic isomer **1b**, and minor cyclic isomer **1c** in CD₃OD. A tautomeric ratio (**1a** : **1b** : **1c** = 4.7 : 4.0 : 1.3) of synthetic epogymnolactam (**1**) right after dissolving in CD₃OD changed into a different ratio (**1a** : **1b** : **1c** = 2.5 : 6.0 : 1.5) with time. This phenomenon suggests that the keto isomer **1a** is most stable in the absence of solvent. The complete NMR assignments of **1a**, **1b** and **1c** are shown in Table 1.

In conclusion, we accomplished the first total synthesis of (+)-epogymnolactam (**1**), and determined the absolute configuration of **1** unambiguously.

EXPERIMENTAL

Chemicals of the highest commercial purity were used without further purification. Thin-layer and silica gel column chromatography were performed by using Merck Silica Gel 60 F₂₅₄ and Kanto Chemical Co. Silica Gel 60N (spherical, neutral), respectively. A DAICEL Chiralpak AD-H column (ϕ 0.64 cm x 25 cm) and a Waters 600 System were used for chiral HPLC. ¹H and ¹³C NMR spectra were recorded using a JEOL JNM EX-270 FT-NMR (JEOL, Tokyo, Japan), and HSQC and HMBC spectra were measured with a Bruker AMX-500 (Bruker, MA, USA). Mass spectra were acquired with FI modes using a JMS-T100GCV (JEOL, Tokyo, Japan). ESI-MS spectra of (+)-**1** were recorded on a LTQ Orbitrap XL (Thermo Scientific, MA, USA). Optical rotations were determined on a JASCO P-2000 (JASCO, Tokyo, Japan).

(2*R*,3*S*)-4-Benzoyloxy-2,3-epoxybutane-1-ol (**3**)

To a stirred suspension of activated 4Å molecular sieves (2.29 g) in dry CH₂Cl₂ (190 ml) were sequentially added Ti(O^{*i*}Pr)₄ (7.20 ml, 24.1 mmol) and D-(–)-DIPT (5.03 ml, 24.1 mmol) under argon at -25 °C. After stirring for 30 min, **2** (4.0 g, 22.5 mmol) in dry CH₂Cl₂ (34 ml) was slowly added over 90 min and the reaction mixture was continually stirred for another 90 min at -25 °C. To the solution was added dropwise a nonane solution of *t*-BuOOH (5.5 M, 8.8 ml) and the solution was stirred for 3 days at -20 °C. After warming to room temperature, the mixture was diluted with saturated (sat.) aqueous Na₂S₂O₃ (40 ml). The resultant solution was stirred for 2 h and then filtrated. The filtrate was extracted with Et₂O and the organic layer was washed with water, dried over Na₂SO₄, concentrated, and purified by silica gel column chromatography (EtOAc : hexane = 1 : 2) to afford epoxy alcohol **3** (3.26 g, 75%) as a colorless oil. The enantiomeric excess value was determined by HPLC (DAICEL Chiralpak AD-H, 0.46 x 25 cm, hexane : EtOH = 9 : 1, 0.8 ml/min). 89% ee; [α]_D²⁵ = +23.0 (*c* 1.00, CHCl₃). ¹H NMR(270 MHz, CDCl₃): δ 2.14 (1H, s, -OH), 3.19-3.32 (2H, m, H-2 and H-3), 3.62-3.75 (4H, m, H-1 and H-4), 4.51-4.64 (2H, dd, *J* = 24.7, 11.9, benzyl), 7.28-7.39

(5H, m, aromatic).

¹³C NMR (67.5 MHz, CDCl₃): δ 54.7 (C-3), 55.6 (C-2), 60.7 (C-1), 68.0 (C-4), 73.5 (benzyl), 127.9 (aromatic), 128.0 (aromatic), 128.5 (aromatic), 137.4 (aromatic).

FI-MS: *m/z* 194.1 [M]⁺.

Kinetic resolution of **3**

To a stirred solution of **3** (1.39 g, 7.17 mmol) in vinyl acetate (73.8 ml) was added 403 mg of PPL (L3126-25G, Sigma, USA) at room temperature (rt.). The reaction mixture was stirred for 6 h, filtered with Celite pad to remove PPL, and the residue on Celite pad was washed with EtOAc. The combined filtrate and washings were concentrated *in vacuo*, and the resultant residue was purified by silica gel column chromatography (EtOAc : hexane = 1 : 2) to give 1.07 g of enantiopure (+)-**3** (1.07 g, 77%). 99% ee; [α]_D²⁵ = +24.3 (*c* 1.00, CHCl₃).

(2*R*,3*S*)-4-Benzyloxy-2,3-epoxy-1-butanal (**4**)

To a stirred solution of **3** (1.06 g, 5.47 mmol) in CH₂Cl₂ (64 ml) were added TEMPO (8.54 mg, 54.7 μmol) and 0.5 M aqueous KBr (1.09 ml) at rt., and then a mixture of 1.96 M aqueous NaOCl (3.35 ml) and sat. aqueous NaHCO₃ (3.35 ml) at 0 °C. After stirring for 4 h at 0 °C, the reaction mixture was quenched with sat. aqueous Na₂S₂O₃ and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, concentrated *in vacuo* and purified by silica gel column chromatography (EtOAc : hexane = 1 : 2) to afford aldehyde **4** (893 mg, 85%) as a colorless oil.

[α]_D²⁵ = −111.1 (*c* 1.00, CHCl₃)

¹H NMR (270 MHz, CDCl₃): δ 3.39-4.43 (1H, t, *J* = 4.5, H-2), 3.47-3.52 (1H, q, *J* = 3.1, H-3), 3.72-3.86 (2H, m, H-4), 4.55 (2H, s, benzyl), 7.29-7.38 (5H, m, aromatic), 9.42-9.44 (1H, d, *J* = 3.7, H-1).

¹³C NMR (67.5 MHz, CDCl₃): δ 57.3 (C-3), 58.0 (C-2), 66.2 (C-4), 73.5 (benzyl), 127.8 (aromatic), 128.0 (aromatic), 128.5 (aromatic), 137.1 (aromatic), 197.6 (C-1).

FI-MS: *m/z* 192.1 [M]⁺.

(2*R*,3*S*)-2,3-epoxy-1,4-octandiol (6)

To a stirred solution of **5** (44.7 mg, 0.233 mmol) in dry THF (1.0 ml) was added dropwise a solution of *n*-BuMgCl in THF (2.0 M, 129 μ l) under argon at -78 °C. The reaction mixture was stirred for 1.5 h, and quenched with MeOH. After warming to room temperature, sat. aqueous NH₄Cl was added to the solution. The mixture was stirred vigorously and extracted with Et₂O. The organic layer was dried over Na₂SO₄, concentrated *in vacuo*, and subjected to silica gel column chromatography (EtOAc : hexane = 1 : 3) to give crude alcohol **5**. To a solution of crude **5** (50.3 mg) in EtOAc (5.8 ml) was added Pd/C (66 mg) and the mixture was stirred vigorously under H₂ overnight. The resulting solution was filtered, concentrated and purified by silica gel column chromatography (EtOAc : hexane = 1 : 1) to afford a diastereomeric mixture of diol **6** (19.7 mg, 53% over 2 steps) as a colorless oil.

$[\alpha]_D^{25} = +2.4$ (*c* 1.00, CHCl₃)

¹H NMR (270 MHz, CDCl₃): δ 0.90-0.95 (3H, m, H-8), 1.31-1.77 (6H, m, H-5, H-6 and H-7), 2.93-3.30 (4H, m, H-2, H-3 and (-OH) x 2), 3.55-3.62 (1H, q, *J* = 6.7, H-4), 3.68-3.75 (1H, dd, *J* = 12.1, 3.3, H-1), 3.99-4.06 (1H, dd, *J* = 12.0, 2.8, H-1).

¹³C NMR (67.5 MHz, CDCl₃): δ 13.9 (C-8), 22.6 (C-7), 27.1 (C-6), 35.2 (C-5), 55.6 (C-2), 59.1 (C-3), 60.7 (C-1), 69.7(C-4).

FI-MS: *m/z* 161.1 [M+H]⁺.

(1*R*,5*R*)-4-Butyl-3,6-dioxabicyclo[3.1.0]hexan-2-one (7)

To a stirred solution of **6** (22.1 mg, 138 μ mol) in CH₂Cl₂ (1.8 ml) were added TEMPO (0.23 mg, 1.38 μ mol) and 0.5 M aqueous KBr (29 μ l) at rt., and then a mixture of 1.96 M aqueous NaOCl (162 μ l) and sat. aqueous NaHCO₃ (162 μ l) at 0 °C. After stirring for 4 h at 0 °C, the reaction mixture was quenched with sat. aqueous Na₂S₂O₃ and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, concentrated *in vacuo*, and purified by silica gel column chromatography (EtOAc : hexane = 1 : 3) to

afford **7** (78%), which was separable to major isomer (Rf value: 0.4, 14.5 mg, 67%) and minor isomer (Rf value: 0.3, 1.7 mg, 8%) as a colorless oil respectively.

Major isomer : $[\alpha]_{\text{D}}^{25} = +48.9$ (*c* 1.00, CHCl₃)

Minor isomer : $[\alpha]_{\text{D}}^{25} = +37.3$ (*c* 0.13, CHCl₃)

¹H NMR (270 MHz, CDCl₃): δ 0.91-0.96 (3H, t, *J* = 6.6, H-8), 1.26-1.71 (6H, m, H-5, H-6 and H-7), 3.77-3.78 (1H, d, *J* = 1.6, H-2), 3.96-3.97 (1H, d, *J* = 2.3, H-3), 4.55-4.59 (1H, t, *J* = 6.5, H-4).

¹³C NMR (67.5 MHz, CDCl₃): δ 13.8 (C-8), 22.3 (C-7), 26.3 (C-6), 31.8 (C-5), 49.8 (C-3), 58.0 (C-2), 79.8 (C-4), 170.3 (C-1).

FI-MS: *m/z* 156.1 [M+H]⁺.

(2*R*,3*R*)-2,3-epoxy-4-hydroxyoctanamide (8)

The diastereomeric mixture of **7** (14.2 mg, 91.0 μ mol) was dissolved in a solution of NH₃ in MeOH (2.0 M, 3 ml) under nitrogen atmosphere and the mixture was stirred for 2.5 h at 0 °C. The resulting solution was concentrated *in vacuo* and purified by silica gel column chromatography (MeOH : CHCl₃ = 7 : 93) to afford a diastereomeric mixture of amide **8** (15.1 mg, 99%) as a colorless oil.

$[\alpha]_{\text{D}}^{25} = +54.4$ (*c* 1.00, CHCl₃)

¹H NMR (270 MHz, CDCl₃) δ 0.87-0.95 (3H, t, *J* = 7.1, H-8), 1.32-1.69 (6H, m, H-5, H-6 and H-7), 3.07-3.21 (2H, m, H-2 and H-3), 3.45-3.58 (2H, m, H-4 and -OH), 6.28 (1H, s, -NH₂), 6.43 (1H, s, -NH₂).

¹³C NMR (67.5 MHz, CDCl₃) δ 14.0 (C-8), 22.6 (C-7), 27.0 (C-6), 34.6 (C-5), 54.3 (C-2), 60.1 (C-3), 69.0 (C-4), 170.2 (C-1).

FI-MS: *m/z* 174.1 [M+H]⁺.

(+)-Epogymnolactam (1)

To a stirred solution of **9** (7.5 mg, 43.4 μ mol) in dry CH₂Cl₂ (1.6 ml) was added Dess-Martin periodinane (25.7 mg, 60.6 μ mol) under argon at 0 °C. After stirring for 2

h, the mixture was quenched with sat. aqueous Na₂S₂O₃ and sat. aqueous NaHCO₃. The solution was extracted with EtOAc and the organic layer was washed with brine, dried over Na₂SO₄, concentrated *in vacuo*, and purified by silica gel column chromatography (EtOAc : hexane = 2 : 1) to afford (+)-epogymnolactam (**1**) (5.6 mg, 76%) as a yellow solid.

[α]_D²⁵ = +25.6 (c 0.49, MeOH)

¹H and ¹³C NMR: see Table 1.

HR-ESI-MS: *m/z* 194.07876 [M+Na]⁺ calcd. for C₈H₁₃O₃NNa, found 194.07887.

Acknowledgements

This work was supported by a grant from the Institute for Fermentation, Osaka (IFO), Japan and a Grant-in-Aid for Scientific Research (C) (No. 26450135) from the Japan Society for the Promotion of Science (JSPS). The authors are grateful to Dr. E. Fukushi (Graduate School of Agriculture, Hokkaido University) for NMR measurements of synthetic (+)-epogymnolactam.

References

1. Mitsuhashi, S., Shindo, C., Shigetomi, K., Miyamoto T. & Ubukata, M. (+)-Epogymnolactam, a novel autophagy inducer from mycelial culture of *Gymnopus* sp. *Phytochemistry* (2014), <http://dx.doi.org/10.1016/j.phytochem.2014.08.012>
2. Cuervo, A. M. Autophagy: In sickness and in health. *Trends Cell Biol.*, **14**, 70-77 (2004).
3. Nixon R. A. & Yang, D.-S. Autophagy failure in Alzheimer's disease-location the primary defect. *Neurobiol. Dis.* **43**, 38-45 (2011).
4. Jimenez, R. E., Kubli, D. A. & Gustafsson, Å. B. Autophagy and mitophagy in the myocardium: therapeutic potential and concerns. *Br. J. Pharmacol.* **171**, 1907-1916 (2014).
5. Rubinsztein, D. C., Codogno, P. & Levine, B. Autophagy modulation as a potential

- 1 therapeutic target for diverse diseases. *Nat. Rev. Drug Discov.*, **11**, 709-730 (2012).
- 2 6. Shoji-Kawata, S. *et al.* Identification of a candidate therapeutic autophagy-inducing
3 peptide. *Nature*, **494**, 201-206 (2013).
- 4 7. Ravikumar, B. *et al.* Inhibition of mTOR induces autophagy and reduces toxicity of
5 polyglutamine expansions in fly and mouse models of Huntington disease. *Nat.*
6 *Genet.* **36**, 585-595 (2004).
- 7 8. Sarkar, S. *et al.* Lithium induces autophagy by inhibiting inositol
8 monophosphatase. *J. Cell. Biol.* **170**, 1101-1111 (2005).
- 9 9. Williams, A. *et al.* Novel targets for Huntington's disease in an mTOR-independent
10 autophagy pathway. *Nat. Chem. Biol.* **4**, 295-305 (2008).
- 11 10. Jeong, J. K. *et al.* Autophagy induced by resveratrol prevents human prion
12 protein-mediated neurotoxicity. *Neurosci. Res.* **73**, 99-105 (2012).
- 13 11. Gupta, V. K. *et al.* Restoring polyamines protects from age-induced memory
14 impairment in an autophagy-dependent manner. *Nat. Neurosci.* **16**, 1453-1460
15 (2013).
- 16 12. Chin R. M. *et al.* The metabolite α -ketoglutarate extends lifespan by inhibiting
17 ATP synthase and TOR. *Nature*, **510**, 397-401 (2014).
- 18 13. D'Agnolo, G., Rosenfeld, I. S., Awaya, J., Omura, S. & Vagelos, P. R.
19 Inhibition of fatty acid synthesis by the antibiotic cerulenin. Specific inactivation
20 of beta-ketoacyl-acyl carrier protein synthetase. *Biochim. Biophys. Acta.* **326**,
21 155-156 (1973).
- 22 14. Sano, Y., Nomura, S., Kamio, Y., Omura S. & Hata, H. Studies on cerulenin, 3.
23 Isolation and physico-chemical properties of cerulenin. *J. Antibiot.* **20**, 344-348
24 (1967).
- 25 15. Funabashi, H. *et al.* Binding site of cerulenin in fatty acid synthetase. *J. Biochem.*
26 **105**, 751-755 (1989).
- 27 16. Kuhajda, F. P. *et al.* Fatty acid synthesis: a potential selective target for
28 antineoplastic therapy. *Proc. Natl. Acad. Sci. U. S. A.*, **91**, 6379-6383 (1994).
- 29 17. Mani, N. S. & Townsend, C. A. A concise synthesis of (+)-cerulenin from a chiral
30 oxiranyllithium. *J. Org. Chem.* **62**, 636-640 (1997).
- 31 18. Sueda, N., Ohru, H. & Kuzuhara, H. Stereoselective synthesis of (+)-cerulenin
32 from D-glucose. *Tetrahedron Lett.* 2039-2042 (1979).
- 33 19. Pietraszkewics, M. & Sinaý, P. Total synthesis of natural cerulenin from

- D-glucose. *Tetrahedron Lett.* 4741-4744 (1979).
20. Yoda, H., Katagiri, T. & Takabe, K. A novel stereoselective synthesis of (+)-cerulenin and (+)-tetrahydrocerulenin. *Tetrahedron Lett.* **32**, 6771-6774 (1991).
21. Furukawa, J., Funabashi, H., Morisaki, N., Iwasaki S. & Okuda, S. A new versatile synthesis of cerulenin. *Chem. Pharm. Bull.* **36**, 1229-1232 (1988).
22. Rawat, V., Dey S. & Sudalai, A. Synthesis of the anti-influenza agent (–)-oseltamivir free base and (–)-methyl 3-epi-shikimate. *Org. Biomol. Chem.* **10**, 3988-3990 (2012).
23. Yoshino, T. *et al.* Total synthesis of an enantiomeric pair of FR900482. 2. Synthesis of the aromatic and the optically active aliphatic segments. *Tetrahedron* **53**, 10239-10252 (1997).
24. Mori, K. & Seu, Y.-B. A new synthesis of (–)- α -multistriatin, the pheromone of the smaller European elm bark beetle. *Tetrahedron* **44**, 1035-1038 (1988).
25. Faigl, F. *et al.* Efficient, scalable kinetic resolution of *cis*-4-benzyloxy-2,3-epoxybutanol, *Tetrahedron: Asymmetry* **16**, 3841-3847 (2005).
26. Shen, L.-L., Wang, F., Mun, H.-S., Suh M. & Jeong, J.-H. Solvent-dependent reactivity in porcine pancreatic lipase (PPL)-catalyzed hydrolysis. *Tetrahedron: Asymmetry* **19**, 1647-1653 (2008).
27. Dess, D. B. & Martin, J. C. A useful 12-I-5 triacetoxypseudoinane (the Dess-Martin periodinane) for the selective oxidation of primary or secondary alcohols and a variety of related 12-I-5 species. *J. Am. Chem. Soc.* **113**, 7277-7287 (1991).
29. Yadav, R. N., Mondal, S. & Ghosh, S. An efficient stereoselective route to the construction of tricyclic core structure towards the synthesis of the sesquiterpens of the *seco*-prezizaane family. *Tetrahedron Lett.* **52**, 1942-1945 (2011).

Table 1. ¹H and ¹³C NMR data of (+)-epogymnolactam in CD₃OD (500 MHz for ¹H and 126 MHz for ¹³C, Bruker)

Position	1a		1b (major)		1c (minor)	
	δ_{C} , type	δ_{H} (<i>J</i> in Hz)	δ_{C} , type	δ_{H} (<i>J</i> in Hz)	δ_{C} , type	δ_{H} (<i>J</i> in Hz)
1	170.5, s	—	174.4, s	—	172.9, s	—
2	55.8, d	3.70, d (5.2)	53.1, d	3.57, d (2.6)	54.3, d	3.56, d (2.7)
3	59.4, d	3.88, d (5.2)	59.0, d	3.80, d (2.6)	58.1, d	3.84, d (2.7)
4	205.8, s	—	87.2, s	—	86.8, s	—
5	41.0, t	2.68, ddd (17.6, 8.2, 6.6) 2,56, ddd (17.6, 8.1, 6.5)	36.3, t	1.72, m	38.9, t	1.78, m
6	26.1, t	1.55, m	27.0, t	1.51, m	25.9, t	1.41, m
7	23.2, t	1.31, sext (7.4)	24.0, t	1.37, sext (7.4)	23.9, t	1.38, sext (7.4)
8	14.1, q	0.90, t (7.4)	14.3, q	0.94, t (7.4)	14.3, q	0.94, t (7.4)

Legend to figure and schemes

Fig. 1. Ring-chain tautomerism of (+)-epogymnolactam (**1**).

Scheme 1. A tandem strategy for preparation of enantiopure (+)-**3**.

Scheme 2. Total synthesis of (+)-epogymnolactam (**1**).






Fig. 1



Scheme 1



Scheme 2