Title	Photodisintegration cross section of ^9Be up to 16 MeV in the $\alpha + \alpha + n$ three-body model
Author(s)	Kikuchi, Yuma; Odsuren, Myagmarjav; Myo, Takayuki; Katayama, Kiyoshi
Citation	Physical Review C, 93(5), 054605 https://doi.org/10.1103/PhysRevC.93.054605
Issue Date	2016-05-06
Doc URL	http://hdl.handle.net/2115/62932
Rights	©2016 American Physical Society
Type	article
File Information	PhysRevC93 054605 .pdf

[Instructions for use](#)

Photodisintegration cross section of ${}^9\text{Be}$ up to 16 MeV in the $\alpha + \alpha + n$ three-body model

Yuma Kikuchi,^{1,*} Myagmarjav Odsuren,² Takayuki Myo,^{3,4} and Kiyoshi Katō⁵¹*RIKEN Nishina Center, Wako 351-0198, Japan*²*School of Engineering and Applied Sciences, National University of Mongolia, Ulaanbaatar 14200, Mongolia*³*General Education, Faculty of Engineering, Osaka Institute of Technology, Osaka 535-8585, Japan*⁴*Research Center for Nuclear Physics (RCNP), Osaka University, Ibaraki 567-0047, Japan*⁵*Nuclear Reaction Data Centre, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan*

(Received 22 February 2016; revised manuscript received 9 April 2016; published 6 May 2016)

The photodisintegration of ${}^9\text{Be}$ in the energy region lower than $E_\gamma = 16$ MeV is investigated by using the $\alpha + \alpha + n$ three-body model and the complex scaling method. The cross section exhibits two aspects in two different energy regions. In the low-energy region up to $E_\gamma = 6$ MeV, the cross section is explained by the transition strengths into the excited resonant states of ${}^9\text{Be}$, while the dipole transition into the nonresonant continuum states of ${}^8\text{Be}(2^+) + n$ dominates the cross section in the energy region of $6 \leq E_\gamma \leq 16$ MeV. Furthermore, it is shown that the dipole strength at $E_\gamma \sim 8$ MeV is understood to be caused by the single-neutron excitation from the ${}^8\text{Be}(2^+) \otimes \nu p_{3/2}$ configuration in the ground state.

DOI: [10.1103/PhysRevC.93.054605](https://doi.org/10.1103/PhysRevC.93.054605)

I. INTRODUCTION

The photodisintegration cross section of ${}^9\text{Be}$ shows various features in different energy regions. The cross section in a low-energy region has been measured to deduce a production rate of ${}^9\text{Be}$ from the astrophysical point of view [1–7]. In the low-energy region up to $E_\gamma = 6$ MeV, the cross section has been observed to come from the electromagnetic transitions into the excited states of ${}^9\text{Be}$, and theoretically, has been studied within the $\alpha + \alpha + n$ three-body model [8–11]. In particular, the first excited $1/2^+$ state, observed as a sharp peak just above the ${}^8\text{Be}(0^+) + n$ threshold, has attracted much attention, and its structure has been discussed in terms of the three-body resonance of $\alpha + \alpha + n$ [8,9,12] or the two-body virtual state of ${}^8\text{Be} + n$ [11,13,14].

In the energy region higher than $E_\gamma = 6$ MeV, the recent measurement [7] reports the photodisintegration cross section of ${}^9\text{Be}$ up to $E_\gamma = 16$ MeV, and the cross section shows a significant electric dipole strength below the giant dipole resonance (GDR). An enhancement of a low-lying dipole strength below the GDR has been observed in a wide range of the mass number in neutron-rich nuclei and is often denoted by the pygmy dipole resonance (PDR). The PDR has been discussed in neutron-rich nuclei in relation with the neutron skin thickness. On the other hand, ${}^9\text{Be}$ is a light and stable nucleus which has been found to have a cluster structure, and hence, it is interesting to investigate the mechanism of the dipole strength in ${}^9\text{Be}$ below the GDR energy region. In a recent measurement [7], Utsunomiya *et al.* showed that the enhanced dipole strength in ${}^9\text{Be}$ at the excitation energy of ~ 8 MeV exhausts 10% of Thomas-Reiche-Kuhn sum rule and almost all the cluster dipole sum rule [7]. We desire to understand the low-lying dipole strength below the GDR comprehensively from a viewpoint of the cluster structure of $\alpha + \alpha + n$ in ${}^9\text{Be}$.

In our previous work, we have investigated the structure of the $1/2^+$ state of ${}^9\text{Be}$ located just above the $\alpha + \alpha + n$ threshold energy using the $\alpha + \alpha + n$ three-body model and the complex scaling method (CSM) [11]. We calculated the photodisintegration cross section from the ground state into the $1/2^+$ state and reproduce the observed peak in the cross section just above the ${}^8\text{Be}(0^+) + n$ threshold. On the other hand, we could not find a sharp resonance corresponding to the $1/2^+$ peak in the cross section using the CSM calculation. To understand the origin of the $1/2^+$ peak in the cross section, we performed the calculation of the pole trajectory by changing the attraction of the intercluster force as an analytical continuation. From these analyses, we concluded that the excited $1/2^+$ state has a virtual-state character of the s -wave neutron in the ${}^8\text{Be} + n$ system.

In Ref. [11], we focused our discussion on the structure of the $1/2^+$ state and its contribution to the photodisintegration cross section in the low-energy region. Spin-parity states other than the $1/2^+$ state were not included in the calculation, and the photodisintegration cross section in higher energy regions was not discussed. In order to understand the feature of ${}^9\text{Be}$, it is necessary to examine the photodisintegration cross section by including all the available spin-parity states connected with the ground $3/2^-$ state via electromagnetic transitions and by taking into account transitions in higher energy regions.

The purposes of this work are the following two: One is to investigate the excited states of ${}^9\text{Be}$ in the low-energy region connected with the ground state through the electromagnetic transitions. The other is to elucidate the mechanism of the enhanced dipole transition in ${}^9\text{Be}$ below the GDR observed by the recent experiment [7]. In the present calculation, we employ the $\alpha + \alpha + n$ three-body model with complex-range Gaussian basis functions to describe the scattering states of the $\alpha + \alpha + n$ system retaining the numerical accuracy. We calculate the photodisintegration cross section by applying the CSM with Green's function to the $\alpha + \alpha + n$ three-body model, and discuss the mechanism of the photodisintegration cross section.

*yuma.kikuchi@riken.jp

This article is organized as follows: In Sec. II, we explain the $\alpha + \alpha + n$ three-body model and describe the formalism of the photodisintegration cross section using the CSM. In Sec. III, we show the results of the photodisintegration cross section, and discuss the structure of the excited states and the mechanism of the low-lying dipole strength in ${}^9\text{Be}$ up to $E_\gamma = 16$ MeV. Finally, in Sec. IV, all results and discussions are summarized.

II. THEORETICAL FRAMEWORK

A. $\alpha + \alpha + n$ three-body model for ${}^9\text{Be}$

We briefly explain the $\alpha + \alpha + n$ three-body model employed in the present work, whose details are given in Ref. [11]. We here solve the Schrödinger equation for the $\alpha + \alpha + n$ system using the orthogonality condition model [15]. The Schrödinger equation is given as

$$\hat{H}\Psi_{J^\pi}^v = E_v\Psi_{J^\pi}^v, \quad (1)$$

where J^π is the total spin and parity of the $\alpha + \alpha + n$ system and v is the state index. The energy eigenvalue E_v is measured from the $\alpha + \alpha + n$ threshold.

The Hamiltonian for the relative motion of the $\alpha + \alpha + n$ three-body system for ${}^9\text{Be}$ is given as

$$\hat{H} = \sum_{i=1}^3 t_i - T_{\text{c.m.}} + \sum_{i=1}^2 V_{\alpha n}(\xi_i) + V_{\alpha\alpha} + V_{\text{PF}} + V_3, \quad (2)$$

where t_i and $T_{\text{c.m.}}$ are kinetic operators for each particle and the center-of-mass of the system, respectively. The interaction between the neutron and the i th α particle is given as $V_{\alpha n}(\xi_i)$, where ξ_i is the relative coordinate between them. We here employ the KKNN potential [16] for $V_{\alpha n}$. For the α - α interaction $V_{\alpha\alpha}$ we employ the same potential as used in Ref. [11], which is a folding potential of the effective NN interaction [17] and the Coulomb interaction. The pseudopotential $V_{\text{PF}} = \lambda|\Phi_{\text{PF}}\rangle\langle\Phi_{\text{PF}}|$ is the projection operator to remove the Pauli forbidden states from the relative motions of α - α and α - n [18]. The Pauli forbidden state Φ_{PF} is defined as the harmonic oscillator wave functions by assuming the $(0s)^4$ configuration whose oscillator length is fixed to reproduce the observed charge radius of the α particle. In the present calculation, λ is taken as 10^6 MeV.

In the present calculation, we introduce the $\alpha + \alpha + n$ three-body potential V_3 . The explicit form of V_3 is given as

$$V_3 = v_3 \exp(-\mu\rho^2), \quad (3)$$

where ρ is the hyper-radius of the $\alpha + \alpha + n$ system. The hyper-radius is defined as

$$\rho^2 = 2r^2 + \frac{8}{9}R^2, \quad (4)$$

where r is the distance between two α 's and R is that between the neutron and the center-of-mass of the $\alpha + \alpha$ subsystem. The strength and width of the three-body potential, v_3 and μ , are determined for each spin and parity. For $3/2^-$ states, we determine the parameters to reproduce the observed binding energy and charge radii of the ground state because it is essential to reproduce the Q value and the sum rule value of the

electric dipole transition in discussing the photodisintegration of ${}^9\text{Be}$. To reproduce the ground-state properties, we take v_3 and μ as 1.10 MeV and 0.02 fm^{-2} , respectively. For other spin-parity states, we employ the same value of μ as used in $3/2^-$ states, and different strengths are used to reproduce the energy positions of the observed peaks in the photodisintegration cross section.

We solve the Schrödinger equation with the coupled-rearrangement-channel Gaussian expansion method [19]. In the present calculation, the ${}^9\text{Be}$ wave function $\Psi_{J^\pi}^v$ is described in the Jacobi coordinate system as

$$\Psi_{J^\pi}^v = \sum_{ijc} C_{ijc}^v(J^\pi) \{ [\phi_i^j(\mathbf{r}_c), \phi_\lambda^j(\mathbf{R}_c)]_L, \chi_{\frac{1}{2}} \}_{J^\pi}, \quad (5)$$

where $C_{ijc}^v(J^\pi)$ is a expansion coefficient and $\chi_{\frac{1}{2}}$ is the spin wave function. The relative coordinates $\mathbf{r}_c$ and $\mathbf{R}_c$ are those in three kinds of the Jacobi coordinate systems indexed by c ($= 1, 2, 3$), and the indices for the basis functions are represented as i and j . The spatial part of the wave function is expanded with the complex-range Gaussian basis functions [19]. The explicit forms of the complex-range Gaussian basis functions are given as

$$\phi_l^i(\mathbf{r}) = \begin{cases} N_l^S(a_i) \exp(-a_i r^2) \sin(a_i \omega r^2), \\ N_l^C(a_i) \exp(-a_i r^2) \cos(a_i \omega r^2), \end{cases} \quad (6)$$

where N_l^S and N_l^C are the normalization factors and a_i is a width of the Gaussian basis function. The basis functions in Eq. (6) enable us to treat the oscillating behavior in the relative motion and is useful to describe the photodisintegration cross section accurately.

B. Photodisintegration cross section in the complex scaling method

To calculate the photodisintegration cross section, we use the complex scaling method (CSM) [11, 20–25]. In the CSM, the relative coordinates ξ ($\mathbf{r}_c$ and $\mathbf{R}_c$) are transformed as

$$U(\theta)\xi U^{-1}(\theta) = \xi e^{i\theta}, \quad (7)$$

where $U(\theta)$ is a complex scaling operator and θ is a scaling angle being a real number. Applying this transformation to Eq. (1), we obtain the complex-scaled Schrödinger equation as

$$\hat{H}^\theta \Psi_{J^\pi}^v(\theta) = E_v^\theta \Psi_{J^\pi}^v(\theta). \quad (8)$$

The complex-scaled Hamiltonian $\hat{H}^\theta$ and the complex-scaled wave function $\Psi_{J^\pi}^v(\theta)$ are defined as in Ref. [25]. By solving the complex-scaled Schrödinger equation with the $\mathcal{L}^2$ basis function given in Eq. (6), we obtain the energy eigenvalues and eigenstates (their biorthogonal states) as $\{E_v^\theta\}$ and $\{\Psi_{J^\pi}^v(\theta)\}$ ($\{\tilde{\Psi}_{J^\pi}^v(\theta)\}$), respectively.

When we apply the CSM to the complex-range Gaussian basis functions, we need to choose the value of the scaling angle θ carefully in relation with the parameter ω in Eq. (6). Applying the CSM to the complex-range Gaussian basis

function, we obtain

$$\begin{aligned} \exp(-ar^2 e^{2i\theta}) \cos(awr^2 e^{2i\theta}) &= \frac{1}{2} [\exp\{-ar^2 e^{2i\theta}(1+i\omega)\} + \exp\{-ar^2 e^{2i\theta}(1-i\omega)\}] \\ &= \frac{1}{2} [\exp\{-a(\cos 2\theta - \omega \sin 2\theta + i\omega \cos 2\theta + i \sin 2\theta)r^2\} \\ &\quad + \exp\{-a(\cos 2\theta + \omega \sin 2\theta - i\omega \cos 2\theta + i \sin 2\theta)r^2\}] \rightarrow \infty \text{ if } \cos 2\theta - \omega \sin 2\theta < 0 \end{aligned} \quad (9)$$

for $\omega \geq 0$ and $0 \leq \theta < \pi/2$. To avoid this singularity, the complex-range Gaussian basis functions with a finite value of ω require a smaller scaling angle compared to the real-range Gaussian basis functions, in which the range of the scaling angle is $0 \leq \theta < \pi/2$. The divergent behavior in the basis function might be serious when we calculate the long-range operator, such as that of the $E1$ transition. In the present work, we take $\theta = 12$ deg and $\omega = 1$ to avoid the divergence of the basis functions and to keep the numerical accuracy in the calculation.

The energy eigenvalues $\{E_v^\theta\}$ are obtained on the complex energy plane, governed by the ABC theorem [20,21]. A schematic picture of the energy eigenvalue distribution is shown in Fig. 1. In the CSM, the energies of bound states are given by real numbers and are invariant under the complex scaling. On the other hand, resonances and continuum states are obtained as eigenstates with complex energy eigenvalues. The resonances are obtained as isolated eigenstates on the complex energy plane, whose energies are given as $E_v^\theta = E_v^r - i\Gamma_v/2$. The resonance energies E_v^r and the decay widths Γ_v are independent of the scaling angle θ . The continuum states are obtained on branch cuts rotated down by 2θ as shown in Fig. 1. The branch cuts start from the different thresholds for two- and three-body continuum states in the case of the $\alpha + \alpha + n$ system as shown in Fig. 1. This classification of the continuum states is useful in discussing the decay modes of the ${}^9\text{Be}$ photodisintegration.

Using the energy eigenvalues and eigenstates of the complex-scaled Hamiltonian $\hat{H}^\theta$, we define the complex-scaled Green's function with outgoing boundary conditions,

FIG. 1. Schematic picture of energy eigenvalue distribution on the complex energy plane for the $\alpha + \alpha + n$ system.

$\mathcal{G}^\theta(E; \xi, \xi')$, as

$$\mathcal{G}^\theta(E; \xi, \xi') = \left\langle \xi \left| \frac{1}{E - H^\theta} \right| \xi' \right\rangle = \sum_v \frac{\Psi^v(\theta) \tilde{\Psi}^v(\theta)}{E - E_v^\theta}. \quad (10)$$

In the derivation of the right-hand side of Eq. (10), we use the extended completeness relation, whose detailed explanation is given in Ref. [26].

Using the complex-scaled Green's function, we calculate the cross section of the photodisintegration of ${}^9\text{Be}(3/2^-) + \gamma \rightarrow \alpha + \alpha + n$ in terms of the multipole response. In the present calculation, we focus on the low-lying region of the photodisintegration cross section and take into account only the electromagnetic dipole responses. The photodisintegration cross section σ^γ is given by the sum of those by the $E1$ and $M1$ transitions as

$$\sigma^\gamma(E_\gamma) = \sigma_{E1}(E_\gamma) + \sigma_{M1}(E_\gamma), \quad (11)$$

where E_γ is the incident photon energy. The energy E in Eq. (10) is related to E_γ as $E = E_\gamma - E_{\text{gs}}$, where E_{gs} is the binding energy of the ${}^9\text{Be}$ ground state measured from the $\alpha + \alpha + n$ threshold. The cross sections for the electromagnetic dipole transitions σ_{EM1} is expressed as the following form:

$$\sigma_{EM1}(E_\gamma) = \frac{16\pi^3}{9} \left(\frac{E_\gamma}{\hbar c} \right) \frac{dB(EM1, E_\gamma)}{dE_\gamma}. \quad (12)$$

Using the CSM and the complex-scaled Green's function in Eq. (10), the electromagnetic dipole transition strength is given as

$$\begin{aligned} \frac{dB(EM1, E_\gamma)}{dE_\gamma} &= -\frac{1}{\pi} \frac{1}{2J_{\text{gs}} + 1} \\ &\quad \times \text{Im} \left[\sum_v \langle \tilde{\Psi}_{\text{gs}} | (\hat{O}^\theta)^\dagger(EM1) | \Psi^v(\theta) \rangle \right. \\ &\quad \left. \times \frac{1}{E - E_v^\theta} \langle \tilde{\Psi}^v(\theta) | \hat{O}^\theta(EM1) | \Psi_{\text{gs}} \rangle \right], \end{aligned} \quad (13)$$

where J_{gs} and Ψ_{gs} represent the total spin and the wave function of the ground state, respectively, and $\hat{O}^\theta(EM1)$ is an electromagnetic dipole transition operator. The $E1$ transition operator is given by

$$\hat{O}(E1) = \sum_{i=1}^3 Z_i e \mathcal{Y}_1(\mathbf{x}_i), \quad (14)$$

where Z_i is atomic number of the i th particle and $\mathbf{x}_i$ is the relative coordinate between the i th particle and the center of mass of the total system. The $M1$ transition operator is

given by

$$\hat{O}(M1) = \frac{e\hbar}{2mc} \sum_{i=1}^3 [g_l^{(i)} \mathbf{l}_i + g_s^{(i)} \mathbf{s}_i], \quad (15)$$

where m is the atomic mass unit and $g_l^{(i)}$ and $g_s^{(i)}$ are the orbital and spin g factors, respectively. For the α particle, g_l is taken as its charge and g_s is taken as 0. For the neutron, we use the free value of $g_s = -3.82$ and do not use any effective charge, namely $g_l = 0$.

We mention the properties of the strength function in Eq. (13), which is given by the sum of the contribution of each final state ν classified in the complex energy plane with the CSM as shown in Fig. 1. The total strength in Eq. (13) corresponds to the observables with positive definite for every energy. We can decompose the strength to clarify the contribution of the ν th state. It is noted that the decomposed strength for the state with complex energy is not necessarily positive definite at all energies, because the state with complex energy cannot be directly observed. In particular, the resonances generally give complex matrix elements, and their physical meanings have been discussed by Berggren [27]. Because their matrix elements are not necessarily positive definite, the matrix elements of the individual states with complex energies cannot be regarded as the physical quantities. But the quantities obtained by taking summation over all the states are physically observable because of the extended completeness relation [26]. It is also noted that the decomposed strength is useful to clarify a physical importance of each decay channel in the reaction as discussed in Ref. [28], while the decomposed strength does not correspond to the observables directly.

TABLE I. Ground-state properties in comparison with experiments. The calculated binding energies (E_{gs} , unit in MeV), charge radii (R_{ch} , unit in fm), and matter ones (R_{m} , unit in fm). The probabilities of the ${}^8\text{Be}(J^\pi) \otimes \nu l_j$ components in the ground state of ${}^9\text{Be}$ are also shown.

	Present	Exp.
E_{gs}	1.57	1.5736 ^a
R_{ch}	2.53	2.519 ± 0.012^b
R_{m}	2.42	2.38 ± 0.01^c
${}^8\text{Be}(0^+) \otimes \nu p_{3/2}$	47.06%	
${}^8\text{Be}(2^+) \otimes \nu p_{3/2}$	46.77%	
${}^8\text{Be}(2^+) \otimes \nu p_{1/2}$	2.31%	
${}^8\text{Be}(2^+) \otimes \nu f_{7/2}$	1.21%	
${}^8\text{Be}(2^+) \otimes \nu f_{5/2}$	1.20%	
${}^8\text{Be}(4^+) \otimes \nu f_{7/2}$	1.07%	
${}^8\text{Be}(4^+) \otimes \nu f_{5/2}$	0.25%	
${}^8\text{Be}(4^+) \otimes \nu h_{11/2}$	0.04%	
${}^8\text{Be}(4^+) \otimes \nu h_{9/2}$	0.04%	

^aReference [29].

^bReference [30].

^cReference [31].

TABLE II. Strength of the three-body potential v_3 for each spin-parity state.

	1/2 ⁺	3/2 ⁺	5/2 ⁺	1/2 ⁻	3/2 ⁻	5/2 ⁻
v_3 (MeV)	-0.90	-0.30	-0.30	0.30	1.10	0.35

III. RESULTS

A. Ground-state properties

The calculated ground-state properties of ${}^9\text{Be}$ are listed in Table I, in which v_3 and μ are taken as 1.10 MeV and 0.02 fm^{-2} , respectively, as mentioned in Sec. II. The binding energy and the charge radius are well reproduced by employing the three-body potential, while the calculated matter radius is slightly larger than the experimental value.

To see the ground-state structure of ${}^9\text{Be}$ more in detail, we calculate the probability of each component of the ${}^8\text{Be}(J^\pi) \otimes \nu l_j$ configurations. The calculated probabilities are also listed in Table I. The present calculation shows that the valence neutron around ${}^8\text{Be}$ occupies the $p_{3/2}$ orbit by 93.8%, while the excited 2^+ component of ${}^8\text{Be}$ has a comparable amount to the 0^+ one. This large mixture of the 2^+ component is understood using the $(\lambda, \mu) = (3, 1)$ component in the SU(3) representation, which corresponds to the π orbital of the valence neutron. In the SU(3) representation, the $(3, 1)$ component gives 53% and 47% for the ${}^8\text{Be}(0^+) \otimes \nu p$ and ${}^8\text{Be}(2^+) \otimes \nu p$ components, respectively.

B. Photodisintegration cross section and resonances in the low-energy region

We calculate the photodisintegration cross section of ${}^9\text{Be}$ using Eq. (12). Before calculating the cross section, we determine the values of v_3 for each spin-parity state as shown in Table II. It is noted that the strength for the $1/2^+$ state is slightly weakened from that in our previous work [11]. This comes from the inclusion of other spin-parity states and the improvement of the numerical accuracy in the low-energy region by using the complex-range Gaussian basis functions.

Using the CSM, we obtain the resonances corresponding to the isolated poles of the S matrix as complex energy

TABLE III. Resonance energies E^r and decay widths Γ for low-lying excited states (units in MeV). The resonance energies are measured from the $\alpha + \alpha + n$ threshold. The observed data are taken from Ref. [29].

J^π	Present (E^r, Γ)	Expt. (E^r, Γ)
1/2 ₁ ⁺		$(0.158 \pm 0.002, 0.213 \pm 0.006)^a$
5/2 ₁ ⁻	$(0.854, \sim 3 \times 10^{-4})$	$(0.8558, 7.8 \times 10^{-4})$
1/2 ₁ ⁻	$(1.11, 0.495)$	$(1.21, 1.01)$
5/2 ₁ ⁺	$(1.47, 0.323)$	$(1.475, 0.282)$
3/2 ₁ ⁺	$(3.12, 1.44)$	$(3.13, 0.743)$
3/2 ₂ ⁻	$(3.08, 1.18)$	$(4.02, 1.33)$

^aThis value is taken from Ref. [6].

FIG. 2. Energy level diagram in the present calculation in comparison with the experimental data. The levels obtained with $v_3 = 1.10$ MeV for all the spin-parity states are also presented. The experimental data are taken from Ref. [29].

eigenvalues. Our results of the resonance energies and decay widths for the excited states up to $E_\gamma = 6$ MeV are listed in Table III in comparison with the experimental data. The energy level diagram is also shown in Fig. 2. In Fig. 2, we show the levels which are obtained using $v_3 = 1.10$ MeV commonly for all the spin-parity states for reference. The calculated resonance energies and decay widths show good correspondences to the observed data, while the resonance energies of the excited states are shifted up by applying more repulsive three-body potential than those in Table II. In the present calculation, we cannot find any isolated resonance of the $1/2^+$ state similar to the result in Ref. [11].

In Fig. 3, the calculated photodisintegration cross section is shown in comparison with experimental data. Our result well reproduces the observed cross section, not only the peak positions but also the magnitudes and widths of the peaks. This agreement implies that our three-body model well describes the scattering states of ${}^9\text{Be}$ in the low-energy region. We also show the photodisintegration cross section into each spin-parity final state in Fig. 4. From the results in Fig. 4, we see that the lowest peak just above the ${}^8\text{Be}(0^+) + n$ threshold energy ($E_\gamma = 1.6654$ MeV) comes from the $E1$ transition into

FIG. 3. Calculated photodisintegration cross sections in comparison with the experimental data. The solid squares and open circles represent the experimental data taken from Refs. [6,7].

FIG. 4. Contributions of each spin-parity state in the photodisintegration cross section.

the $1/2^+$ state, and the calculated cross section shows that the strength below the ${}^8\text{Be}(0^+) + n$ threshold is negligibly small. As mentioned above, we do not find any isolated resonance of the $1/2^+$ state, while the cross section into the $1/2^+$ states has a significant peak above the ${}^8\text{Be}(0^+) + n$ threshold. These facts are consistent with our previous result [11]. It is noted that we confirm the $1/2^+$ state has the virtual-state character of the ${}^8\text{Be} + n$ system from a similar analysis to that in Ref. [11]. The second and third peaks at $E_\gamma = 2.5$ and 3.0 MeV in the cross section come from the transitions into the resonances of $5/2^-$ and $5/2^+$ states, respectively. The $M1$ transition into the $1/2^-$ resonance has a sizable contribution to the cross section at around $E_\gamma = 2.7$ MeV. The transitions into the $3/2^\pm$ states play minor roles in the photodisintegration cross section below $E_\gamma = 6$ MeV.

C. Low-lying dipole strength at $E_\gamma \sim 8$ MeV

We investigate the photodisintegration cross section above $E_\gamma = 6$ MeV and discuss the dipole strength of ${}^9\text{Be}$ below the GDR. In Fig. 5, we show the calculated photodisintegration cross section up to $E_\gamma = 16$ MeV in comparison with the experimental data. From the result in Fig. 5, we see that calculated cross section has a broad peak at $E_\gamma \sim 8$ MeV,

FIG. 5. Calculated photodisintegration cross section up to $E_\gamma = 16$ MeV. The solid squares and open circles represent experimental data taken from Refs. [6,7], respectively.

FIG. 6. Contributions of each spin-parity state in the photodisintegration cross section up to $E_\gamma = 16$ MeV.

similar to the experimental data. To investigate the origin of the peak at $E_\gamma \sim 8$ MeV, we also calculate the contribution of each spin-parity final state as shown in Fig. 6. The results in Fig. 6 show that the peak is dominated by the $E1$ transitions into $3/2^+$ and $5/2^+$. In the present CSM calculation, we do not find any isolated resonances corresponding to the peak. Therefore, the peak at $E_\gamma \sim 8$ MeV is understood to be described by the $E1$ transition from the ground state into the nonresonant continuum states of $3/2^+$ and $5/2^+$. It is also seen that our calculation slightly underestimates the observed peak in the cross section. This underestimation suggests a contribution from degrees of freedom beyond the relative motion of the $\alpha + \alpha + n$ system in the present model. For example, the GDR is known to be described by coherent $1p-1h$ excitations, but the $1p-1h$ excitations from the α particle are not taken into account in our $\alpha + \alpha + n$ three-body model.

In the recent measurement, it is reported that the energy-integrated cross section for the enhanced dipole strength measured for $4 \leq E_\gamma \leq 16$ MeV is estimated to be 11.3 mb MeV as a lower limit [7]. Moreover, it is suggested that this energy-integrated cross section exhausts 10% of the Thomas-Reiche-Kuhn (TRK) sum rule and almost all of the energy-weighted cluster dipole sum rule. In Ref. [7], the cluster dipole sum rule is defined by subtracting the contribution of an internal excitation of the α particles from the TRK sum rule, and the energy-integrated cross section is estimated by subtracting the contribution of the GDR from the total cross section. We here estimate only the energy-integrated cross section corresponding to the cluster dipole sum rule because the α particles are assumed to be inert in the present model.

We calculated the energy-integrated cross section by integrating σ_{E1} over the energy interval of $4 \leq E_\gamma \leq 16$ MeV. We obtain the energy-integrated cross section as 12.1 mb MeV for this energy interval, and the result is consistent with the experimental value (11.3 mb MeV). On the other hand, we also obtain the energy-integrated cross sections for $E_\gamma < 4$ MeV and $E_\gamma > 16$ MeV as 0.954 and 8.40 mb MeV, respectively. It is seen that the energy-integrated cross section for $4 \leq E_\gamma \leq 16$ MeV is 56.5% of the total amount in the present calculation. Our result suggests that the cluster dipole sum rule does not concentrate on the strength for

FIG. 7. Decomposed photodisintegration cross section for the $3/2^+$ states.

$4 \leq E_\gamma \leq 16$ MeV and the strength for higher energy region would have a sizable contribution to the energy-weighted cluster dipole sum rule.

In the CSM, the energy eigenvalues of unbound states are classified into those of the resonance and several families of the two- and three-body nonresonant continuum states as shown in Fig. 1. Combining the energy eigenvalues in the CSM with Eq. (13), we decompose the photodisintegration cross section into different families of the nonresonant continuum states. In Figs. 7 and 8, we show the decomposed photodisintegration cross sections for the $3/2^+$ and $5/2^+$ states, respectively. We do not show the cross sections into the ${}^5\text{He}(3/2^-) + n$ and ${}^5\text{He}(1/2^-) + n$ continuum states in Figs. 7 and 8, since their contributions are much smaller than others. In both results, we confirmed that the cross sections at $E_\gamma \sim 8$ MeV are dominated by the contributions from the ${}^8\text{Be}(2^+) + n$ nonresonant continuum states, while that from the ${}^8\text{Be}(0^+) + n$ ones is negligible in the cross section. Furthermore, we see that the three-body continuum states of $\alpha + \alpha + n$ have sizable contributions to the low-lying dipole strengths.

The decomposed photodisintegration cross sections show that the enhanced dipole strength at $E_\gamma \sim 8$ MeV is described by the single-neutron excitation from the ${}^8\text{Be}(2^+) \otimes \nu p_{3/2}$ configuration in the ground state because the $E1$ transition operator cannot directly excite the relative motion between two

FIG. 8. Same as Fig. 7 but for the $5/2^+$ states.

α 's from the 0^+ to 2^+ states. The ground-state wave function has the large (3,1) component in the SU(3) representation, and the (3,1) component gives a mixture of the ${}^8\text{Be}(0^+) \otimes \nu p$ and ${}^8\text{Be}(2^+) \otimes \nu p$ configurations. Our result shows that the ground-state structure of ${}^9\text{Be}$ is essential to reproduce the observed dipole strength at $E_\gamma \sim 8$ MeV.

On the other hand, the contribution from the nonresonant continuum states of ${}^8\text{Be}(0^+) + n$ is negligibly small in the photodisintegration cross section. It seems to be that the ${}^8\text{Be}(0^+) \otimes \nu p_{3/2}$ configuration is mainly excited into the resonances. To confirm this, it would be important to investigate the structures of the excited resonances of ${}^9\text{Be}$ and the further analysis will be performed in a forthcoming paper.

IV. SUMMARY

In this work, we investigate the photodisintegration of ${}^9\text{Be}$ in the energy region lower than $E_\gamma = 16$ MeV by using the $\alpha + \alpha + n$ three-body model and the complex scaling method (CSM). We here employ the complex-range Gaussian basis functions, which enables us to describe the oscillating behaviors of the scattering states of the $\alpha + \alpha + n$ system and to calculate the transition strength accurately. We calculate the photodisintegration cross section up to $E_\gamma = 16$ MeV and show good agreement with the observed data. From the calculated cross section, we discuss the following two points: One is the structures of the excited states of ${}^9\text{Be}$ observed in the photodisintegration in the low-energy region up to $E_\gamma = 6$ MeV. The other is the origin of the enhanced dipole strength at $E_\gamma \sim 8$ MeV below the GDR.

In addition to the photodisintegration cross section, we also show the resonance energies and decay widths of the excited states obtained in the present calculation below $E_\gamma = 6$ MeV. It is shown that the calculated resonance energies and decay widths are consistent with the observed energy levels while the $1/2^+$ state is the exceptional case. We confirm that the $1/2^+$

state has the virtual-state character of the ${}^8\text{Be} + n$ system from the similar analysis to that in our previous work. The photodisintegration cross section up to $E_\gamma = 6$ MeV can be understood by the summation of transition strengths into each excited state of ${}^9\text{Be}$.

For the energy region of $6 \leq E_\gamma \leq 16$ MeV, our calculation shows a significant dipole strength at $E_\gamma \sim 8$ MeV and a good agreement with the recent observed data of the photodisintegration. We find that this dipole strength is dominated by the transitions into the nonresonant continuum states of $3/2^+$ and $5/2^+$. To understand the origin of the dipole strength, we decompose the photodisintegration cross section into each nonresonant continuum state by the use of the energy eigenvalue distribution in the CSM. From the decomposed cross section, it is shown that the strength mainly comes from the transitions into the ${}^8\text{Be}(2^+) + n$ nonresonant continuum states. This fact can be understood by a dipole excitation of the single neutron from the ${}^8\text{Be}(2^+) \otimes \nu p_{3/2}$ configuration in the ${}^9\text{Be}$ ground state.

In the present calculation, we do not investigate the structures of the excited resonances in detail. It would be important to see the structures of the resonances to gain a comprehensive understanding of the photodisintegration of ${}^9\text{Be}$, and the detailed analysis of the resonances will be performed in a forthcoming paper.

ACKNOWLEDGMENTS

The authors are grateful to Prof. H. Utsunomiya for fruitful discussion and providing us with the experimental data for the photodisintegration cross section of ${}^9\text{Be}$. This work was supported by JSPS KAKENHI Grants No. 25400241 and No. 15K05091. One of the authors (K.K.) is also thankful for the support by the International Collaboration with Al-Farabi Kazakh National University (Grants No. 3106/GF4 and No. 1550/GF3).

-
- [1] J. H. Gibbons, R. L. Macklin, J. B. Marion, and H. W. Schmitt, *Phys. Rev.* **114**, 1319 (1959).
 - [2] W. John and J. M. Prosser, *Phys. Rev.* **127**, 231 (1962).
 - [3] M. Fujishiro, T. Tabata, K. Okamoto, and T. Tsujimoto, *Can. J. Phys.* **60**, 1672 (1982).
 - [4] H. Utsunomiya, Y. Yonezawa, H. Akimune, T. Yamagata, M. Ohta, M. Fujishiro, H. Toyokawa, and H. Ohgaki, *Phys. Rev. C* **63**, 018801 (2000).
 - [5] O. Burda, P. von Neumann-Cosel, A. Richter, C. Forssén, and B. A. Brown, *Phys. Rev. C* **82**, 015808 (2010).
 - [6] C. W. Arnold, T. B. Clegg, C. Iliadis, H. J. Karwowski, G. C. Rich, J. R. Tompkins, and C. R. Howell, *Phys. Rev. C* **85**, 044605 (2012).
 - [7] H. Utsunomiya, S. Katayama, I. Gheorghe, S. Imai, H. Yamaguchi, D. Kahl, Y. Sakaguchi, T. Shima, K. Takahisa, and S. Miyamoto, *Phys. Rev. C* **92**, 064323 (2015).
 - [8] E. Garrido, D. Fedorov, and A. Jensen, *Phys. Lett. B* **684**, 132 (2010).
 - [9] R. Álvarez-Rodríguez, A. S. Jensen, E. Garrido, and D. V. Fedorov, *Phys. Rev. C* **82**, 034001 (2010).
 - [10] J. Casal, M. Rodríguez-Gallardo, J. M. Arias, and I. J. Thompson, *Phys. Rev. C* **90**, 044304 (2014).
 - [11] M. Odsuren, Y. Kikuchi, T. Myo, M. Aikawa, and K. Katō, *Phys. Rev. C* **92**, 014322 (2015).
 - [12] V. D. Efros, P. von Neumann-Cosel, and A. Richter, *Phys. Rev. C* **89**, 027301 (2014).
 - [13] V. Efros and J. Bang, *Eur. Phys. J. A* **4**, 33 (1999).
 - [14] K. Arai, P. Descouvemont, D. Baye, and W. N. Catford, *Phys. Rev. C* **68**, 014310 (2003).
 - [15] S. Saito, *Prog. Theor. Phys., Suppl.* **62**, 11 (1977).
 - [16] H. Kanada, T. Kaneko, S. Nagata, and M. Nomoto, *Prog. Theor. Phys.* **61**, 1327 (1979).
 - [17] E. Schmid and K. Wildermuth, *Nucl. Phys.* **26**, 463 (1961).
 - [18] V. I. Kukulin, V. M. Krasnopol'sky, V. T. Voronchev, and P. B. Sazonov, *Nucl. Phys. A* **453**, 365 (1986).
 - [19] E. Hiyama, Y. Kino, and M. Kamimura, *Prog. Part. Nucl. Phys.* **51**, 223 (2003).
 - [20] J. Aguilar and J. M. Combes, *Commun. Math. Phys.* **22**, 269 (1971).

- [21] E. Balslev and J. M. Combes, [Commun. Math. Phys.](#) **22**, 280 (1971).
- [22] Y. K. Ho, [Phys. Rep.](#) **99**, 1 (1983).
- [23] N. Moiseyev, [Phys. Rep.](#) **302**, 212 (1998).
- [24] S. Aoyama, T. Myo, K. Katō, and K. Ikeda, [Prog. Theor. Phys.](#) **116**, 1 (2006).
- [25] T. Myo, Y. Kikuchi, H. Masui, and K. Katō, [Prog. Part. Nucl. Phys.](#) **79**, 1 (2014).
- [26] T. Myo and K. Katō, [Prog. Theor. Phys.](#) **98**, 1275 (1997).
- [27] T. Berggren, [Phys. Lett. B](#) **373**, 1 (1996).
- [28] T. Myo, K. Katō, S. Aoyama, and K. Ikeda, [Phys. Rev. C](#) **63**, 054313 (2001).
- [29] D. Tilley, J. Kelley, J. Godwin, D. Millener, J. Purcell, C. Sheu, and H. Weller, [Nucl. Phys. A](#) **745**, 155 (2004).
- [30] W. Nörtershäuser, D. Tiedemann, M. Žáková, Z. Andjelkovic, K. Blaum, M. L. Bissell, R. Cazan, G. W. F. Drake, C. Geppert, M. Kowalska *et al.*, [Phys. Rev. Lett.](#) **102**, 062503 (2009).
- [31] I. Tanihata, T. Kobayashi, O. Yamakawa, S. Shimoura, K. Ekuni, K. Sugimoto, N. Takahashi, T. Shimoda, and H. Sato, [Phys. Lett. B](#) **206**, 592 (1988).