

HOKKAIDO UNIVERSITY

Title	Spatiotemporal profiles of arginine vasopressin transcription in cultured suprachiasmatic nucleus
Author(s)	Yoshikawa, Tomoko; Nakajima, Yoshihiro; Yamada, Yoshiko; Enoki, Ryosuke; Watanabe, Kazuto; Yamazaki, Maya; Sakimura, Kenji; Honma, Sato; Honma, Ken-ichi
Citation	European journal of neuroscience, 42(9), 2678-2689 https://doi.org/10.1111/ejn.13061
Issue Date	2015-11
Doc URL	http://hdl.handle.net/2115/63362
Rights	This is the peer reviewed version of the following article: Yoshikawa, T., Nakajima, Y., Yamada, Y., Enoki, R., Watanabe, K., Yamazaki, M., Sakimura, K., Honma, S., Honma, Ki. (2015), Spatiotemporal profiles of arginine vasopressin transcription in cultured suprachiasmatic nucleus. European Journal of Neuroscience, 42: 2678–2689., which has been published in final form at http://dx.doi.org/10.1111/ejn.13061. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Туре	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Supplemental_information.pdf (Supporting Information)



## Supplemental figure 1



## Supplemental figure 2









В

## Supplemental figure 3



**Supplemental figure 1.** Representative double-plotted actogram of circadian rhythms in spontaneous locomotor activities of WT and AVP<sup>ELuc/+</sup> mice. Amounts of locomotor activity are plotted in 1 min bins with black columns. Black and white bars at the top of each panel indicate the dark and light phases of LD cycles. Grey bars on the right margin indicate the days in DD. Red circles in the actograms and arrows in the right margin indicate the time in ZT and day of a single light pulse.

Supplemental figure 2. A. Heat maps of parameters of the circadian AVP-ELuc rhythms on pixel level in the SCN slices (n = 4). Acrophase (far left), amplitude (left), period (right) and percent rhythm (PR) (far right) are illustrated with pseudocolor. Color scale of acrophase indicates differences from the slice mean defined as 12 h. Goodness of fit is indicated by PR: highest with 1 and lowest with 0. The limit of significance level for curve fitting is shown by a triangle. Pixels with bioluminescence intensity less than the background level are indicated with white color and aperiodic pixels are indicated with black color. Scale bars indicate 100 µm, III the third ventricle and OC the optic chiasm. B. Quantitative analyses of circadian parameters of the bioluminescence rhythm in the dorsomedial (DM), centromedial (CM) and ventromedial (VM) areas of the SCN slice. The mean differences in the acrophase between two of three areas (left), the mean period (middle) and the mean amplitude (right) of circadian AVP-ELuc rhythm in each area are expressed as the mean  $\pm$  SD (n = 5). Asterisk (\*) indicates a statistically significant difference (P<0.01, one-way ANOVA with post-hoc Tukey-Kramer test). C. Immunohistochemically identified AVP (red) and VIP (green) containing neurons in the SCN stained after ex vivo recording of bioluminescence. The SCN slices are two of the four slices as shown in A. Scale bar indicates 100 µm.

**Supplemental figure 3.** Representative circadian AVP-ELuc rhythms in the cultured slices of extra SCN brain areas and peripheral tissues from an AVP<sup>ELuc/+</sup> mouse. Only low basal levels of bioluminescence were detected except for the SCN.