



Title	Hydroxy fatty acids as fingerprint biomarkers for the atmospheric transport of soil microorganisms and terrestrial higher plant metabolites [an abstract of dissertation and a summary of dissertation review]
Author(s)	Tyagi, Poonam
Citation	北海道大学. 博士(環境科学) 甲第12415号
Issue Date	2016-09-26
Doc URL	<a href="http://hdl.handle.net/2115/63818">http://hdl.handle.net/2115/63818</a>
Rights(URL)	<a href="http://creativecommons.org/licenses/by-nc-sa/2.1/jp/">http://creativecommons.org/licenses/by-nc-sa/2.1/jp/</a>
Type	theses (doctoral - abstract and summary of review)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Tyagi_Poonam_abstract.pdf (論文内容の要旨)



[Instructions for use](#)

# 学 位 論 文 内 容 の 要 旨

博士 (環境科学)

氏 名 TYAGI Poonam

## **Hydroxy fatty acids as fingerprint biomarkers for the atmospheric transport of soil microorganisms and terrestrial higher plant metabolites**

(土壌微生物と陸上高等植物代謝物質の大気輸送バイオマーカーとしてのヒドロキシ脂肪酸)

Lipid biomarkers have long been employed as a powerful tool to deconvolute and trace the relative significance of soil microorganisms, epicuticular plant waxes and also, to some extent, for characterizing microbial communities in different environmental archives. Among the various lipid class compounds, hydroxy fatty acids (OH FAs) are one such proxy due to their ubiquitous occurrence as an essential membrane component of soil microorganisms and higher plant waxes. In particular, the positional  $\beta$ -isomers of OH FAs having carbon chain length from C10 to C18 are specific to endotoxin/lipopolysaccharides (LPS) of Gram-negative bacteria (GNB). The long-term exposure to airborne endotoxin can cause severe allergies and infections in human beings. Apart from  $\beta$ -, other positional isomers of OH FAs such as  $\alpha$ - and  $\omega$ -short chain homologous have the potential to be used as tracers of soil microorganisms. However, they are also found in microalgae, cyanobacteria, sea grasses, and plant waxes. Moreover, these OH FAs can also act as intermediate products of photochemical as well as microbial oxidation of long-chain monocarboxylic acids to dicarboxylic acids.

OH FAs from soil microbes and plants can be emitted into the atmosphere due to the wind abrasion during dust events and soil-dust re-suspension during biomass burning (forest fires/agricultural burning). Once in the atmosphere, these OH FAs can be transported to distant places and, thus, can have severe impacts on climate, human health and downwind ecosystems. The major objectives of my thesis are to better understand the source regions, atmospheric transport and deposition of soil- and plant-associated bacteria, and also, higher plant metabolites

in the East Asian outflow (Asian dust) to the North Pacific. To accomplish these targets, I have measured the OH FAs abundances in different atmospheric samples including aerosol (urban, coastal, marine), snow (fresh and snow pit) and rain water samples collected from the East Asian outflow.

Results obtained from my thesis work contribute to fill-up the gap between the sources of hydroxy FAs and their deposition into freshwater and marine sediments. A clear seasonal trend in the atmospheric abundances of OH FAs over marine (Jeju and Chichijima) islands is noteworthy with spring/winter maxima and summer/autumn minima. Furthermore, these seasonal cycles are consistent with the prevailing meteorology. The air mass back-trajectories and other soil microbial tracers (such as trehalose) indicate the influence of Asian dust outbreaks originating from Mongolian and Gobi Deserts and also from China loess plateau during winter and spring. To examine the influence of open biomass burning on soil microbes, OH FAs have also been measured in wheat residue combustion derived aerosols over Mt. Tai, North East China. These analyses have revealed significant emissions of  $\beta$ - OH FAs (biomarkers of GNB) and endotoxin in high biomass burning events. We have also found that these OH FAs can be removed from the atmosphere efficiently by snow and rain as compared to aerosols.

The overall results obtained through my thesis clearly document the long-range atmospheric transport of OH FAs, which are tracers of soil microbes (bacteria, fungi, protozoa) and higher plant waxes, from the Asian continent to North Pacific. The Asian dust outbreaks in spring can lead to increase in atmospheric abundances of OH FAs and, thereby, resulting in the occurrence of health impacts in downwind regions due to the exposure to these airborne particles. This chemical marker based approach (culture independent technique) is a suitable tool for the rapid and comparative analysis of bacterial dry mass and their endotoxin in the various environmental samples which excludes the need of biological recovery and microbial cultivation.