



Title	Toxicity of venom of Asobara and Leptopilina species to Drosophila species
Author(s)	Kohyama, Tetsuo I.; Kimura, Masahito T.
Citation	Physiological entomology, 40(4), 304-308 https://doi.org/10.1111/phen.12115
Issue Date	2015-12
Doc URL	http://hdl.handle.net/2115/63827
Rights	This is the peer reviewed version of the following article: "Toxicity of venom of Asobara and Leptopilina species to Drosophila species", which has been published in final form at DOI:10.1111/phen.12115. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	PHEN-0363_AuthorFinal.pdf (本文)



[Instructions for use](#)

1 **Toxicity of venom of *Asobara* and *Leptopilina* species to *Drosophila* species**

2

3 TETSUO I. KOHYAMA and MASAHITO T. KIMURA

4

5 Graduate School of Environmental Earth Science, Hokkaido University, Sapporo,

6 Hokkaido 060-0810, Japan

7

8 Running title: *Effects of venom of Asobara and Leptopilina*

9

10 Correspondence: Tetsuo I. Kohyama, Division of Biosphere Sciences, Graduate School

11 of Environmental Earth Science, Hokkaido University, N10W5, Kita-ku, Sapporo,

12 Hokkaido 060-0810, Japan; Tel: +81-11-706-2236; e-mail:

13 tetsuo_kohyama@ees.hokudai.ac.jp

14

15 **Abstract.** The *Drosophila* parasitoid *Asobara japonica* has highly toxic venom that
16 kills host larvae if its injection is not followed by injection of lateral oviduct
17 components along with egg-laying. In this study, the venom of seven other *Drosophila*
18 parasitoids (*Asobara rossica*, *A. rufescens*, *A. pleuralis*, *Leptopilina heterotoma*, *L.*
19 *japonica*, *L. ryukyuensis*, and *L. victoriae*) are tested against three kinds of *Drosophila*
20 species (i.e. *Drosophila* species that are suitable as host for focal parasitoids, those that
21 are resistant to the parasitoids, and a cosmopolitan species, *D. simulans*). Venom of the
22 three *Asobara* species are not toxic to any of *Drosophila* species, whereas those of the
23 four *Leptopilina* species are toxic to some *Drosophila* species. Toxicity of venom vary
24 among *Leptopilina* species, and susceptibility to venom also vary among host
25 *Drosophila* species. Furthermore, toxicity and paralytic effects of venom are not
26 correlated. Because toxicity of venom is not adaptive for parasitoids, it may be an
27 inevitable side effect of some components that play an essential role in parasitism.

28

29 **Key words.** *Asobara*, *Drosophila*, *Leptopilina*, parasitism, toxicity, venom.

30

31 **Introduction**

32

33 In oviposition, parasitoid females not only lay eggs in hosts but also inject them with
34 venom that are complex mixtures of proteins and low molecular weight compounds
35 having various functions such as induction of paralysis or regulation of host
36 development and immune responses. Some components of venom act alone, while some
37 act in combination with other factors provided from the mother's organs, parasitoid
38 embryo or teratocytes (Moreau & Guillot 2005; Asgari & Rivers, 2011; Poirié *et al.*,
39 2014). An interesting example of cooperative action of venom and oviduct products has
40 been reported for *Asobara japonica* Belokobylskij, a parasitoid of *Drosophila* species.
41 During parasitization of *Drosophila* hosts, *A. japonica* females inject venom into the
42 host immediately after the insertion of the ovipositor (Furihata & Kimura, 2009). A few
43 seconds later, parasitoid females vibrate their ovipositors for a few seconds and then
44 withdraw them. Thus, vibration of the ovipositor is an indication of egg laying (Vet &
45 Bakker, 1985; van Lenteren *et al.*, 1998; Dubuffet *et al.*, 2006). In parasitism
46 experiments using *A. japonica* and some host species such as *Drosophila melanogaster*
47 Meigen and *D. simulans* Sturtevant, interruption of oviposition after injection of venom
48 but before egg laying kills host larvae (Furihata & Kimura, 2009;
49 Mabilia-Moundougou *et al.*, 2010; Furihata *et al.*, 2013). However, this toxic action of
50 the venom is suppressed by lateral oviduct components that are injected to the host by
51 the parasitoid female along with egg (Mabilia-Moundougou *et al.*, 2010; Furihata *et al.*
52 2013). Additional studies showed that the venom of *A. japonica* is much less toxic to
53 certain *Drosophila* species that kill *A. japonica* embryos or larvae at high rates, such as
54 *D. bipectinata* Duda and *D. ficusphila* Kikkawa & Peng (Furihata & Kimura 2009;
55 Furihata *et al.*, 2013). These "resistant" *Drosophila* species may have evolved some

56 mechanisms to detoxify lethal components of *A. japonica* venom along with the
57 acquisition of capacity to kill *A. japonica* embryos and/or larvae in the course of their
58 coevolutionary interactions. On the other end of the spectrum is *A. rossica*
59 Belokobylskij, whose venom is not known to be toxic to any *Drosophila* species
60 (Furihata *et al.*, 2013). Thus, effects and functions of venom vary among different
61 parasitoid species. However, the toxicity of venom of several other *Drosophila*
62 parasitoids has not been investigated (Moreau & Guillot 2005; Asgari & Rivers, 2011;
63 Poirié *et al.* 2014).

64 In this paper, toxicity and paralytic effects of venoms of seven *Drosophila*
65 parasitoids including three *Asobara* (Braconidae) species (*A. rossica*, *A. rufescens*
66 (Förster) and *A. pleuralis* (Ashmead),) and four *Leptopilina* (Figitidae) species (*L.*
67 *heterotoma* (Thompson), *L. japonica* Novković & Kimura, *L. ryukyuensis* Novković &
68 Kimura and *L. victoriae* Nordlander) on three kinds of *Drosophila* species (suitable host,
69 resistant species and *D. simulans*) are investigated to understand how the function of
70 venom has evolved. *A. rossica*, *A. rufescens*, *L. heterotoma* and *L. japonica* are mainly
71 distributed in the temperate regions of Asia and/or Europe, whereas the other three are
72 distributed in subtropical and tropical regions of Asia and/or Africa (Allemand *et al.*
73 2002; Novković *et al.* 2011; Nomano *et al.* 2015). They show species-specific host use,
74 and therefore some *Drosophila* species are suitable as host for some parasitoid species
75 but some are resistant (Novković *et al.* 2012; Kimura & Suwito 2012, 2014; Kimura &
76 Novković 2015; Kimura unpublished data). Little is known about the virulence strategy
77 of these parasitoids, except for *L. heterotoma* and *L. victoriae* that show active
78 suppression of the host immune systems (Rizki & Rizki 1984; Morales *et al.*, 2005).

79

80 **Materials and methods**

81

82 *Laboratory strains*

83

84 Laboratory strains of *A. rossica*, *A. rufescens* and *L. heterotoma* were collected from
85 Sapporo (43.0 °N, 141.2 °E), northern Japan in August 2012, that of *L. japonica* from
86 Tokyo (35.7 °N, 139.8 °E), central Japan in June 2010, that of *L. ryukyuensis* from
87 Iriomote-jima (24.2 °N, 123.8 °E) in March 2006, southern Japan, and those of *A.*
88 *pleuralis* and *L. victoriae* from Kota-Kinabalu (5.3 °N, 117.4 °E), Malaysia in March
89 2008 (Table 1). Clumps of banana were placed in the field to allow drosophilid flies to
90 oviposit in banana and parasitoid individuals to oviposit in drosophilid larvae in banana.
91 Usually after a week, the bananas were brought back to the laboratory and placed in
92 plastic containers. When drosophilid pupae were formed, they were collected in plastic
93 Petri dishes, and examined for emergence of flies or parasitoids. When parasitoid
94 individuals emerged, they were reared on appropriate hosts.

95 At the same time of parasitoid collection, a number of *Drosophila* species were
96 collected from each locality. Suitability of these *Drosophila* species as host for
97 sympatric populations of parasitoids was examined in our previous studies (Novković *et*
98 *al.* 2012; Kimura & Suwito 2012, 2014; Kimura & Novković 2015; Kimura
99 unpublished data). From *Drosophila* species collected in each locality, two species were
100 chosen for the present experiments; one was a suitable host that allows focal parasitoid
101 to develop successfully and another was a resistant species that kills embryos or larvae
102 of focal parasitoid at high rates. In the experiments on *A. pleuralis* (from Kota
103 Kinabalu), however, *D. parabiptinata* and *D. pseudoananassae* strains that were
104 collected from Deramakot located approximately 200 km east of Kota Kinabalu in

105 March 2005 were used. In addition to *Drosophila* strains from the same or nearby
106 localities, a *D. simulans* strain collected from Tokyo in June 2007 was used.

107 These parasitoid and *Drosophila* strains originated from few to several field
108 collected females. The *Drosophila* strains were maintained with *Drosophila* medium.
109 For the maintenance of parasitoid strains, *D. simulans* was used as host with the
110 exception of *A. rossica* and *A. rufescens* strains which were maintained on *D. auraria*
111 Peng (because *D. simulans* was highly resistant to these parasitoids). Rearing and
112 experiments were performed at 23 °C under an LD 15:9 h photcycle, and experiments.

113

114 *Parasitism experiments*

115

116 Female parasitoids used in the experiments were maintained for two to five
117 days in vials containing *Drosophila* medium with host larvae after emergence, i.e., they
118 were fed, mated and had experience of oviposition. *Drosophila* larvae used in the
119 experiments were 2-3 day old (mostly second-instar). Parasitoid females were placed
120 with *Drosophila* larvae in Petri dishes (3 cm in diameter) with small amounts of food
121 medium, and were observed for oviposition behaviour under a stereoscopic microscope.
122 To obtain host larvae in which venom was injected but egg was not laid, host larvae
123 were drawn apart from parasitoid females using forceps before the females started to
124 vibrate their ovipositors ('interrupted' group). In another group, parasitoids were
125 allowed to complete oviposition without interruption ('un-interrupted' group).
126 Immediately after injection of venom or completion of oviposition, *Drosophila* larvae
127 were transferred to new vials with food and examined for emergence of adult
128 parasitoids or flies. Usually 50 *Drosophila* larvae were used for each treatment and
129 more than five parasitoid females were used to obtain 50 treated-larvae.

130 In parasitism experiments, *Drosophila* larvae showed various degrees of
131 paralysis, strong paralysis, weak paralysis and no sign of paralysis. However, it was not
132 easy to determine the boundary between weak and no paralysis, because weakly
133 paralyzed larvae were able to move more or less. On the other hand, strong paralysis
134 was distinctive; i.e., larvae did not move at all even if stimulated by forceps. In addition,
135 the occurrence of strong paralysis did not vary within strain; i.e. all larvae of a strain
136 showed the same response. Thus, *Drosophila* larvae subjected to the above experiments
137 were checked for the occurrence of strong paralysis when transferred from Petri dishes
138 to new vials.

139

140 *Statistical analysis*

141

142 The frequency of individuals from which no insect (i.e., neither parasitoid nor fly)
143 emerged was compared between the ‘un-interrupted group’ and ‘interrupted group’ by a
144 generalized linear model (GLM) with binomial error distribution and logit-link function
145 using R 3.2.1 (R Core Team 2015). The significance of the explanatory variable of the
146 model was tested with likelihood ratio test.

147

148 **Results**

149

150 Figure 1 shows the results of the experiments using previously reported data for *A.*
151 *japonica* (Furihara & Kimura, 2009) as a reference. For the *Asobara* species (*A. rossica*,
152 *A. rufescens* and *A. pleuralis*), the frequency of individuals from which no insect (i.e.,
153 neither parasitoid nor fly) emerged did not exceed 40%, even if oviposition was
154 interrupted before egg-laying. However, the frequency was significantly different

155 between the ‘interrupted’ and ‘un-interrupted’ groups for suitable hosts in the
156 experiments on *A. rossica* and *A. rufescens* and for *D. simulans* in the experiment on *A.*
157 *pleuralis* (GLM with likelihood ratio test, $P < 0.05$).

158 In *Leptopilina* species, the frequency of individuals from which no insect
159 emerged was always higher in the ‘interrupted’ group than in the ‘un-interrupted group,
160 and the difference was significant in all comparisons (GLM with likelihood ratio test, P
161 < 0.05), except for *D. simulans* in the experiment on *L. japonica* and for suitable-host
162 and resistant species in the experiment on *L. victoriae*.

163 Larvae of *D. simulans* always showed strong paralysis when parasitized by any
164 of the seven parasitoids (data not shown). In addition, both suitable-host (*D. auraria*)
165 and resistant species (*D. bifasciata*) showed strong paralysis when parasitized by *A.*
166 *rossica* (Table 1). Furthermore, a resistant species (*D. bauraria*) showed strong
167 paralysis when parasitized by *A. rufescens*, and a suitable-host species (*D.*
168 *nigromaculata*) showed strong paralysis when parasitized by *L. heterotoma* (Table 1).
169 The other parasitoids did not induce strong paralysis in suitable-host or resistant species.
170 Thus, there was no clear relation between the occurrence of strong paralysis and toxicity
171 of venom or suitability as host.

172

173 **Discussion**

174

175 *Asobara japonica* has venom that exhibits a toxic effect to host species if its injection is
176 not followed by an injection of lateral oviduct components (Mabiala-Moundougou *et*
177 *al.*, 2010; Furihata *et al.*, 2013). However, the venom of the other three *Asobara* species
178 studied here does not have such toxic effects. Four *Leptopilina* species have venom that
179 exhibits toxicity to some *Drosophila* species if egg laying is not followed; i.e., *L.*

180 *heterotoma* venom is toxic to a suitable-host species, *L. japonica* venom to both
181 suitable-host and resistant species, and *L. ryukyuensis* and *L. victoriae* venom to *D.*
182 *simulans*. However, the frequency of *Drosophila* larvae from which neither flies nor
183 parasitoids emerge is always higher when oviposition is interrupted. In this study,
184 interruption of oviposition is achieved by separating *Drosophila* larvae from ovipositing
185 parasitoid females by forceps. *Leptopilina* species have an ovipositor clip by which they
186 hold host larvae during oviposition (van Lenteren *et al.*, 1998; Buffington, 2007), and
187 *Drosophila* larvae are sometimes dragged by parasitoids when larvae and parasitoids are
188 separated by forceps (Kohyama, personal observation). It is possible that the separation
189 by forceps may injure *Drosophila* larvae and increase their mortality to some degree. In
190 the above cases of *Leptopilina*, however, the frequency of *Drosophila* larvae from
191 which no insect emerges is very high (> 84%), which suggests that their venom has at
192 least some toxicity to some *Drosophila* species. For a further understanding of the
193 toxicity of *Leptopilina* venom, artificial injection of venom is required.

194 It is not known why *A. japonica* and the four *Leptopilina* species have toxic
195 venom. In koinobiont larval parasitoids, such as *Asobara* and *Leptopilina* species, the
196 toxicity of the venom is not adaptive because parasitoids are required to allow host
197 individuals to survive and develop to the pupal stage after parasitization. The toxicity of
198 their venom may be an inevitable side effect of some components that play an essential
199 role in the regulation of host development or suppression of host immune responses. In
200 this study, the toxicity of venom varies among the parasitoid species, and each
201 parasitoid's venom shows different toxicity to different *Drosophila* species. Toxicity is
202 not related to the occurrence of strong paralysis. These results suggest that the
203 parasitoid's systems to regulate host development and immune responses are
204 complicated and have species-specific components (Dupas *et al.*, 2003, 2009;

205 Thompson, 2005; Kimura & Suwito, 2014). Biochemical and molecular biological
206 studies would provide some insight into how the virulence mechanisms of the toxicity
207 of venom has evolved and diversified.

208 *Drosophila simulans* is strongly paralysed by venom of all of the parasitoid
209 species studied. *Drosophila simulans* is an invasive species originating in Africa
210 (Lemeunier *et al.*, 1986); therefore, it has a relatively short history of interaction with
211 the Palearctic and Asian parasitoid species studied here and may not have had sufficient
212 time to adapt to these parasitoid species. In this respect, it is interesting how *D.*
213 *simulans* responds to parasitoids of the original locality of Africa.

214 In the experiments with suitable-host and resistant *Drosophila* species, strong
215 paralysis is observed only in one case (i.e., *D. nigromaculata* oviposited by *L.*
216 *heterotoma*) out of eight cases in which *Leptopilina* species were used, whereas in five
217 cases out of eight cases in which *Asobara* species were used (including *A. japonica*:
218 Furihata & Kimura 2009). Thus, *Asobara* venom induces strong paralysis in a larger
219 number of *Drosophila* species. This characteristic may be related to the absence of the
220 clip-on ovipositor in *Asobara* species. In our observation (Kohyama, personal
221 observation), the third instar larvae of *D. auraria* can easily escape from the attack by *A.*
222 *rufescens* because *A. rufescens* does not have an ovipositor clip nor the capacity to
223 paralyse *D. auraria* larvae. Therefore, many *Asobara* species may have intensified the
224 paralytic effects of venom in compensation to the absence of an ovipositor clip.

225

226 **Acknowledgements**

227

228 We thank to FY Nomano and B Novković for fruitful discussions. This study was partly
229 supported by a Grant-in-Aid to MTK (No. 23370005) from the Japan Society for
230 Promotions of Science (JSPS).

231

232 **References**

233

234 Allemand, R., Lemaître, C., Frey, F. *et al.* (2002) Phylogeny of six African *Leptopilina*
235 species (Hymenoptera : Cynipoidea, Figitidae), parasitoids of *Drosophila*, with
236 description of three new species. *Annales de la Société Entomologique de France*
237 **38**, 319–332.

238 Asgari, S., & Rivers, D.B. (2011) Venom proteins from endoparasitoid wasps and their
239 role in host-parasite interactions. *Annual Review of Entomology*, **56**, 313–335.

240 Buffington, M.L. (2007) The occurrence and phylogenetic implications of the ovipositor
241 clip within the Figitidae (Insecta: Hymenoptera: Cynipoidea). *Journal of Natural*
242 *History*, **41**, 2267–2282.

243 Dubuffet, A., Álvarez, C.I.R., Drezen, J.-M. *et al.* (2006) Do parasitoid preferences for
244 different host species match virulence? *Physiological Entomology*, **31**, 170–177.

245 Dupas, S., Carton, Y. & Poirié, M. (2003) Genetic dimension of the coevolution of
246 virulence–resistance in *Drosophila* – parasitoid wasp relationships. *Heredity*, **90**,
247 84–89.

248 Dupas, S., Dubuffet, A., Carton, Y. *et al.* (2009) Local, geographic and phylogenetic
249 scales of coevolution in *Drosophila*-parasitoid interactions. *Advance in*
250 *Parasitology*, **70**, 281-295.

251 Furihata, S.X. & Kimura, M.T. (2009) Effects of *Asobara japonica* venom on larval
252 survival of host and nonhost *Drosophila* species. *Physiological Entomology*, **34**,
253 292–295.

254 Furihata, S.X., Matsumoto, H., Kimura, M.T. *et al.* (2013) Venom components of
255 *Asobara japonica* impair cellular immune responses of host *Drosophila*
256 *melanogaster*. *Archives of Insect Biochemistry and Physiology*, **83**, 86–100.

257 Kimura, M.T. & Novković, B. (2015) Local adaptation and ecological fitting in host use
258 of the *Drosophila* parasitoid *Leptopilina japonica*. *Ecological Research*, **30**,
259 499–505.

260 Kimura, M.T. & Suwito, A. (2012) Diversity and abundance of frugivorous
261 drosophilids and their parasitoids in Bogor, Indonesia. *Journal of Natural History*,
262 **46**, 1947–1957.

263 Kimura, M.T. & Suwito, A. (2014) What determines host acceptance and suitability in
264 tropical Asian *Drosophila* parasitoids? *Environmental Entomology*, **43**, 123–130.

265 Lemeunier, F., David, J.R., Tsacas, L. *et al.* (1986) The *melanogaster* species group.
266 *The Genetics and Biology of Drosophila*, Vol. 3e (ed. by M. Ashburner, H. L.
267 Carson & J. N. Thompson Jr.), pp. 147–256. Academic Press, New York.

268 Mabilia-Moundougou, A.D.N., Doury, G., Eslin, P. *et al.* (2010) Deadly venom of
269 *Asobara japonica* parasitoid needs ovarian antidote to regulate host physiology.
270 *Journal of Insect Physiology*, **56**, 35–41.

271 Morales, J., Chiu, H., Oo, T. *et al.* (2005) Biogenesis, structure, and immune
272 suppressive effects of virus-like particles of a *Drosophila* parasitoid, *Leptopilina*
273 *victoriae*. *Journal of Insect Physiology*, **51**, 181–195.

274 Moreau, S.J.M. & Guillot, S. (2005) Advances and prospects on biosynthesis, structures
275 and functions of venom proteins from parasitic wasps. *Insect Biochemistry and*
276 *Molecular Biology*, **35**, 1209–1223.

277 Nomano, F.Y., Mitsui, H., & Kimura, M.T. (2015) Capacity of Japanese *Asobara*
278 species (Hymenoptera; Braconidae) to parasitize a fruit pest *Drosophila suzukii*
279 (Diptera; Drosophilidae). *Journal of Applied Entomology*, **139**, 105–113.

280 Novković, B., Mitsui, H., Suwito, A *et al.* (2011) Taxonomy and phylogeny of
281 *Leptopilina* species (Hymenoptera: Cynipoidea: Figitidae) attacking frugivorous
282 drosophilid flies in Japan, with description of three new species. *Entomological*
283 *Science* **14**, 333–346.

284 Novković, B., Oikawa, A., Murata, Y. *et al.* (2012) Abundance and host association of
285 parasitoids attacking frugivorous drosophilids in Iriomote-jima, a subtropical
286 island of Japan. *European Journal of Entomology*, **109**, 517–526.

287 Poirié, M., Colinet, D. & Gatti, J.-L. (2014) Insights into function and evolution of
288 parasitoid wasp venoms. *Current Opinion in Insect Science*, **6**, 52–60.

289 R Core Team (2015) *R: A language and environment for statistical computing*. R
290 Foundation for Statistical Computing, Vienna, Austria [WWW document]. URL
291 <http://www.R-project.org/> [accessed on 26 June 2015].

292 Rizki, T.M. & Rizki R.M. (1984) Selective destruction of a host blood cell type by a
293 parasitoid wasp. *Proceedings of National Academy of Science, USA*, **81**,
294 6154-6158.

295 Thompson, J.N. (2005) *The Geographic Mosaic of Coevolution*. University of Chicago
296 Press, Chicago, Illinois.

297 van Lenteren, J.C., Isidoro, N. & Bin, F. (1998) Functional anatomy of the ovipositor
298 clip in the parasitoid *Leptopilina heterotoma* (Thompson) (Hymenoptera :

299 Eucoilidae), a structure to grip escaping host larvae. *International Journal of Insect*
300 *Morphology and Embryology*, **27**, 263–268.

301 Vet, L.E.M. & Bakker, K.E. (1985) A comparative functional approach to the host
302 detection behaviour of parasitic wasps. 2. A quantitative study on eight eucoilid
303 species. *Oikos*, **44**, 487–498.

304

305 **Figure Legends**

306

307 **Fig. 1.** Percentages of fly larvae from which flies (white), parasitoids (grey) or neither
308 organism (black) emerged when *D. simulans*, suitable-host and resistant species were
309 parasitized by *Asobara* (A–D) and *Leptopilina* species (E–H) or injected with their
310 venom (I: oviposition of parasitoid females was interrupted after venom injection but
311 before egg-laying, U: oviposition was not interrupted). Suitable-host and resistant
312 species for each parasitoid species is given in Table 1. Numbers above bars indicate the
313 number of individuals used in each experiment, and symbols above bars indicate results
314 of statistical tests between ‘interrupted’ (I) and ‘un-interrupted’ (U) groups (GLM with
315 likelihood ratio test, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). Data of *A. japonica* were
316 from Furihata & Kimura (2009)

317 **Table**

Table 1. Parasitoid and *Drosophila* species used in the experiments, and the original localities of the experimental strains. Species that showed strong paralysis when parasitized by parasitoids given in the left column was shown in bold.

	Susceptible host species	Resistant species	Original locality
<i>A. japonica</i> *	<i>D. auraria</i> Peng*	<i>D. bipectinata</i> Duda**	Tokyo, Iriomote-jima
<i>A. rossica</i>	<i>D. auraria</i>	<i>D. bifasciata</i> Pomini	Sapporo
<i>A. rufescens</i>	<i>D. auraria</i>	<i>D. biauraria</i> Bock & Wheeler	Sapporo
<i>A. pleuralis</i>	<i>D. parabiptinata</i> Bock	<i>D. pseudoananassae</i> Bock	Kota Kinabalu
<i>L. heterotoma</i>	<i>D. nigromaculata</i> Kikkawa & Peng	<i>D. auraria</i>	Sapporo
<i>L. japonica</i>	<i>D. rufa</i> Kikkawa & Peng	<i>D. auraria</i>	Tokyo
<i>L. ryukyuensis</i>	<i>D. ananassae</i> Doleschall	<i>D. bipectinata</i>	Iriomote-jima
<i>L. victoriae</i>	<i>D. malekotliana</i> Parshad & Paika	<i>D. bipectinata</i>	Kota Kinabalu

*From Tokyo, **from Iriomote-jima (Furihata and Kimura 2009).

318

319