



Title	トリカブト屬數種の染色体に就て
Author(s)	酒井, 寛一
Citation	札幌博物学会会報, 13(2), 74-77
Issue Date	1933-12-15
Doc URL	http://hdl.handle.net/2115/64056
Type	article
File Information	Vol.13No.2_003.pdf



[Instructions for use](#)

A BRIEF NOTE ON THE CHROMOSOMES IN SOME *ACONITUM* SPECIES

BY

KAN-ICHI SAKAI

(With one plate)

On the chromosomes in the genus *Aconitum*, several studies have been made since the end of the last century, although most of the materials adopted were those under cultivation. E. OVERTON in 1893 was the first to deal with this problem, and following him, OSTERWALDER (1898), LANGLET (1927), G. A. LEWITSKY (1931), C. D. DARLINGTON (1932), and A. AFIFY (1933) should be mentioned.

Reviewing their reports, there can be found a polyploidal phenomenon in the chromosome number. The following list includes the plants, of which the chromosome number has already been counted.

Diploid ($n=8$, $2n=16$):

Aconitum Lycoctonum (LEWITSKY, AFIFY)

*A. barbatum*¹⁾ (DARLINGTON)

*A. vulparia*¹⁾ (DARLINGTON)

A. orientale (LEWITSKY, AFIFY)

A. luridum (AFIFY)

Triploid ($2n=24$):

A. Napellus (OVERTON, OSTERWALDER, LANGLET, and
DARLINGTON)

A. Stoekrianum (AFIFY)

Tetraploid ($n=16$, $2n=32$):

A. anglicum (DARLINGTON)

A. Anthora (LEWITSKY)

A. Napellus (LEWITSKY)

Octoploid ($n=32$, $2n=64$):

A. Wilsonii (LANGLET)

A. vshubile latisectum (DARLINGTON)

1) These species are synonyms of *A. Lycoctonum* according to BAILEY (1930).

Some detailed morphological study of their chromosomes may be found, too.

Fortunately, the writer was able to gather a few wild species belonging to this genus from Hokkaido and Karafuto, and he wishes here to describe briefly the result of his observation on the number as well as on some morphological aspects of their chromosomes.

Here the writer ought to offer his heartfelt thanks to Mr. U. KIMOTO in Konuma, through whose kindness the two species from Karafuto have been handed to him.

Materials and Methods

Six wild species have been studied. Observations were made in the mitosis of the root-tip cells. After being fixed in NAWASCHIN'S chromo-acetic-formalin, they were stained by RANDOLPH'S iodine crystal-violet method. Thickness of the paraffin sections was 20-25 μ .

Results of Observation

The chromosome numbers counted in six species are as follows:

	Jap. Name	Obtained from	2n
<i>Aconitum umbrosum</i> KOM.	Ooreijinsō	Mt. Teine	16
<i>A. yuparense</i> TAKEDA	Yezo-hosoba-torikabuto	Mt. Daisetsu	16
<i>A. Fischeri</i> RCHB.	Oobushi	Konuma	32
<i>A. sachalinense</i> FR. SCHM.	Karafuto-bushi	Konuma	32
<i>A. subcuneatum</i> NAKAI	Oku-torikabuto	Zenibako	32
<i>A. yezoense</i> NAKAI	Yezo-torikabuto	Maruyama	32

The approximate lengths of each chromosome from one specimen measured with great care under the microscope is recorded in Table I, and Fig. 7 is the chromosome-scheme founded on this table. These are made with the diploid species *A. umbrosum*.

TABLE I. The approximate length of each chromosome of *A. umbrosum* in μ .

Chromosomes	IA	IB	IIA	IIB	IIIA	IIIB	IVA	IVB	VA	VB	VIA	VIB	VIIA	VIIIB	VIIIA	VIIIB
Long arms	7.0	6.9	6.0	5.8	5.5	5.5	5.0	5.0	5.0	5.0	5.0	4.9	4.0	4.0	2.9	2.8
Short arms	6.5	6.5	3.0	3.0	1.1	1.1	1.5	1.5	1.0	1.0	1.0	1.0	2.0	2.0	1.2	1.2
Whole length	13.5	13.4	9.0	8.8	6.6	6.6	6.5	6.5	6.0	6.0	6.0	5.9	6.0	6.0	4.1	4.0

In the chromosome-scheme (Fig. 7), one can recognize eight classes from the largest I-chromosomes with approximately median constriction, to the smallest VIII-chromosomes with subterminal constriction. Among these eight types, I-chromosomes may be fairly recognized in all the species, and especially the VIII-chromosomes may be identified at a glance. The five species are compared in respect to these I- and VIII-chromosomes. Here the former is called the M, and the latter, the s chromosomes.

	M	s
<i>Aconitum umbrosum</i>	2	2
<i>A. yuparense</i>	2	2
<i>A. sachalinense</i>	(2)	$2s_1 + 2s_2$
<i>A. subcuneatum</i>	(2)	4
<i>A. yezoense</i>	(2)	4

A. Fischeri is excluded from this comparison on account of their badly fixed or stained figures.

In the above list, the number put in parentheses means that there have been found two M chromosomes at least, and consequently there might exist in reality just two or more of them. A more detailed measurement or an exact judgement with good figures is necessary in order to make this point absolutely clear.

Proportional to the chromosome number, the number of the s chromosomes is multiplied: in the diploids, we can find two of them, and in the tetraploids, four. The four s's in the latter two species are all alike in their size and form, except in the one species *A. sachalinense*, in which the 4 s's consist of 2 s_1 's and 2 s_2 's. Each length of the s_1 and s_2 chromosomes is as follows:

	Long arms	Short arms
s_1	= ca. 1.5μ	+ ca. 0.5μ
s_2	= ca. 2.0μ	+ ca. 0.5μ






The difference between the length of their long arms is ca. 0.5μ .

Conclusion

The writer, working with six wild species from Hokkaido and Karafuto, found two diploids and four tetraploids.

According to the comparison of the s-chromosomes within these species, interspecific difference in the morphology of chromosomes should exist in some species.

Plant Breeding Institute, Faculty of Agriculture,
Hokkaido Imperial University,
Sapporo, Japan



Bibliography

- AFIFY, A. (1933):—Chromosome form and behaviour in diploid and triploid *Aconitum*. J. GENET., 27, No. 2, 293-318.
- BAILEY, L. H. (1930):—Hortus.
- DARLINGTON, C. D. (1932):—Recent Advances in Cytology.
- KIHARA, H. et al. (1931):—植物染色体数の研究
- LEWITSKY, G. A. (1931):—The "karyotype" in systematics. Bull. Appl. Bot., of Genet. & Pl.-Breed., 27, No. 1, 187-239.

Explanation of Plate

Figs. 1-6. Chromosomes in root-tip cells, drawn with the aid of camera lucida under a magnification of \times ca. 2300.

Fig. 1. *Aconitum umbrosum*

Fig. 2. *A. yufarense*

Fig. 3. *A. Fischeri*

Fig. 3. *A. sachalinense*

Fig. 5. *A. subcuneatum*

Fig. 6. *A. yezoense*

Fig. 7. Scheme showing the approximate length of each chromosome in *A. umbrosum* made from Table 1.

摘 要

トリカブト属数種の染色体に就て

酒 井 寛 一

北海道及樺太に野生するトリカブト属植物6種をとりその染色体を観察した所、2種のダイプロイドと4種のテトラプロイドを發見した。トリカブト属植物の染色体の基本数は8である。ダイプロイド種なるオホレイジンサウ *Aconitum umbrosum* の各染色体の長さを測つた結果によれば(第一表及第七圖)、そこに8種の大き及び形の異なる染色体が見出される。その最小染色体なるVIII-染色体 即ち s-染色体に就て各種を比較するにこの s の數は全染色体數に比例してゐるが、4s をもつテトラプロイド種の中にはその4個が全く相同な4sより成る場合と、相同ならざる $2s_1 + 2s_2$ よりなる場合とが見出された。