



Title	満州國産蜘蛛類
Author(s)	齋藤, 三郎
Citation	札幌博物学会会報, 15(3), 191-193
Issue Date	1938-07-15
Doc URL	http://hdl.handle.net/2115/64229
Type	article
File Information	Vol.15No.3_007.pdf



[Instructions for use](#)

Arachnida of Manchoukuo III

By

Saburô SAITÔ

(齊藤三郎)

As to the Arachnida fauna of Manchoukuo the writer has reported already 69 species of spiders and 5 of harvestmen from Jehol¹⁾ and from the district along the South-Manchuria Railway²⁾. On September of 1937, Prof. Dr. T. INUKAI brought back a considerable number of spiders and harvestmen with a large collection of the other animals from his journey to Manchoukuo and Inner Mongolia as far as Kulumpeierh (呼倫貝爾) of Hei-lung-kiang (黑龍江省). These valuable materials were put at my disposal for systematical study through his courtesy. For this I wish here to express my heartiest thanks to Prof. T. INUKAI.

Of 20 species the following 13 species of spiders and 1 of harvestmen are referred again to the fauna of Manchoukuo, while the 6 listed afterwards are new;

1. *Gnaphosa sinensis* SIMON

Gnaphosa sinensis SIMON:—1880, Ann. Soc. ent. Fr., 5 ser., X, 121.

Loc. 1 ♀, the basin of the river Gan (垠河), Sept. 8.

Dist. China (Pepin), Manchoukuo (Cheng-te, Chao-yang, Harbin).

2. *Tibellus oblongus* (WALCKENAER)

Aranea oblonga WALCKENAER:—1802, Faune Par., II, 228.

Loc. 2 ♀ ♀ (subadult), Wonfu (王府), Aug. 29, 30.

Dist. Manchoukuo (Hsin-king), U.S.A., Alaska, Europe, China (Pepin), Japan (Hokkaidô, the Southern Kurile Islands).

3. *Araneus adiantus* WALCKENAER

Aranea adianta WALCKENAER:—1802, Faune Par., II, 199.

Loc. 2 ♀ ♀ (subadult), Wonfu, Aug. 29, 30.

Dist. Manchoukuo (Ai-tao-tzu-ling, Peh-ying-fang, Schuang-ta-shan, Weng-niu-te, Leo-suu-ton, Hsin-king, Kai-yuan), Japan (Hokkaidô, Southern Saghalien), Siberia, Europe.

4. *Araneus cornutus* CLERCK

Araneus cornutus CLERCK:—1757, Sv. Spindl., 39.

Loc. 1 ♀, the basin of the river Gan, Sept. 8.

1) Report of the First Sc. Exped. to Manchoukuo, Sec. V, Div. I, Pt. III, Arts. 11, 12, 1936.

2) Annot. Zool. Jap., Vol. 16, 1937.

Dist. Manchoukuo (Harbin), Japan (Southern Saghalien, Hokkaidô), U.S.A., Europe.

5. *Araneus hamatus* CLERCK

Araneus hamatus CLERCK :—1757, Sv. Spindl., 51.

Loc. 1 ♀ (subadult), Wonfu, Aug. 29, 30.

Dist. Manchoukuo (Shuang-ta-shan), Europe.

6. *Eucta chinensis* CHAMBERLIN

Eucta chinensis CHAMBERLIN :—1924, Proc. U.S. Nat. Mus., LXIII, 12.

Loc. 3 ♀ ♀ (subadult), Wonfu, Aug. 29, 30.

Dist. China, Manchoukuo (Chao-yang).

7. *Agelena labyrinthica* (LINNAEUS)

Aranea labyrinthica LINNAEUS, 1758, Syst. Nat., Ed. 10, I, 620.

Loc. 1 ♀, the basin of the river Gan, Aug. 8.

Dist. Manchoukuo (Chao-yang, Shuang-ta-shan, Leo-sun-ton, An-shan, Kai-yuan), U.S.A., Europe.

8. *Agelena opulenta* L. KOCH

Agelena opulenta L. KOCH :—1877, Verhandl. d.k.k. Zool. bot. Ges., XXVII, 757.

Loc. 1 ♀, Khailassu (哈拉蘇), Sept. 12.

Dist. Manchoukuo (Harbin, An-shan, Siong-yu-cheng, Hsin-king, Pehying-fang), Japan (Kyûsyû, Honsyû, Hokkaidô).

9. *Tegenaria Derhamii* (SCOPOLI)

Aranea Derhamii SCOPOLI, 1763, Entom. Carn., 400.

Loc. 1 ♀, Khailar (海拉爾), Sept. 10.

Dist. Manchoukuo (Leo-sun-ton, Hsin-king, Chao-yang, Shuang-ta-shan), U.S.A., Europe.

10. *Dolomedes fimbriatus* (CLERCK)

Araneus fimbriatus CLERCK :—1757, Sv. Spindl., 106.

Loc. 1 ♀, 4 subadults, Khailassu, Sept. 12.

Dist. Manchoukuo (Chang-shan-yu, Cheng-te), Europe.

11. *Lyccsa monticola* (CLERCK)

Araneus monticola CLERCK :—1757, Sv. Spindl., 91.

Loc. 2 ♀ ♀, Khailar, Sept. 9.

Dist. Manchoukuo (Hsin-king), Japan (Hokkaidô), Siberia, Europe.

12. *Lycosa T-insignita* BÖS. et STRAND

Lycosa T-insignita BÖSENBERG et STRAND :—1906, Abhandl. senckenb. Naturf. Ges., XXX,

Loc. 1 ♀, the basin of the river Gan, Aug. 8.
 Dist. Manchoukuo (Peh-wing-fang, Shuang-ta-shan, Cheng-te, Hsin-king), Japan (Hokkaidô, Honsyû, Kyûsyû, Formosa).

13. *Oxyopes sertatus* L. KOCH

Oxyopes sertatus L. KOCH:—1877, Verhandl. d.k.k. zool. bot. Ges., XXVII, 779.

Loc.; ♂, 1 ♀ (subadult), Khailassu, Sept. 11, 12.

Dist. Manchoukuo (Tieh-ling), Japan (Formosa, Honsyû, Kyûsyû).

14. *Opilio trispinifrons* ROEWER

Opilio trispinifrons ROEWER:—1911, Zool. Jahrb., Syst., V, 596.

Loc. Khailassu, Sept. 12.

Dist. Manchoukuo (Weng-niu-te), China (Pepin).

The following 6 are recorded for the first time from the Manchurian fauna;

15. *Philodromus spinitarsus* SIMON

Philodromus spinitarsus SIMON:—1895, BULL. Soc. ent. Fr., LXII.

Loc. 1 ♂ (subadult), Handshur (什殊爾), Sept. 6.

Dist. Japan (Kyûsyû), China.

16. *Xysticus ephippiatus* SIMON

Xysticus ephippiatus SIMON:—1880, Ann. Soc. ent. Fr., 5 ser., X, 107.

Loc. 5 ♀ ♀ (immature), Wonfu, Aug. 29, 30.

Dist. China (Pepin, Suifu, Soochow), Japan (Hokkaidô, Honsyû, Kyûsyû).

17. *Xysticus striatipes* L. KOCH

Xysticus striatipes L. KOCH:—1870, Beitr. Kennt. Ar.

Loc. 1 ♀, Handshur, Sept. 9, 11 ♂ ♂, 10 ♀ ♀ (subadult), Wonfu, Aug. 29, 30.

Dist. Europe.

18. *Thanatus miniaceus* SIMON

Thanatus miniaceus SIMON:—1800, Ann. Soc. ent. Fr., 5 ser., X, 110.

Loc. 11 ♀ ♀ (subadult), Wonfu, Aug. 29, 30.

Dist. China (Pepin).

19. *Larinia punctifera* BÖS. et STRAND

Larinia punctifera BÜSENBURG et STRAND:—1906, Verhandl. senckenb. Naturf. Ges., XXX, 212.

Loc. 4 ♀ ♀ (subadult), Wonfu, Aug. 29, 30.

Dist. Japan (Honsyû).

20. *Lutica* sp.

Loc. 1 ♀ (subadult), Wonfu, Aug. 29, 30.