



| | |
|------------------------|---|
| Title | Antibody detection from Middendorf ' s vole (<i>Microtus middendorffii</i>) against Tula virus captured in Mongolia |
| Author(s) | Yoshimatsu, Kumiko; Arai, Satoru; Shimizu, Kenta; Tsuda, Yoshimi; Boldgiv, Bazartseren; Boldbaatar, Bazartseren; Sergelen, Erdenebaatar; Ariunzaya, Dagvatseren; Enkhmandal, Orsoo; Tuvshintugs, Sukhbaatar; Morikawa, Shigeru; Arikawa, Jiro |
| Citation | Japanese Journal of Veterinary Research, 65(1), 39-44 |
| Issue Date | 2017-2 |
| DOI | 10.14943/jjvr.65.1.39 |
| Doc URL | http://hdl.handle.net/2115/64786 |
| Type | bulletin (article) |
| Additional Information | There are other files related to this item in HUSCAP. Check the above URL. |
| File Information | 65-1_039-044.pdf |



[Instructions for use](#)

Antibody detection from Middendorf's vole (*Microtus middendorffii*) against Tula virus captured in Mongolia

Kumiko Yoshimatsu^{1,†,*}, Satoru Arai^{2,†}, Kenta Shimizu¹, Yoshimi Tsuda¹, Bazartseren Boldgiv³, Bazartseren Boldbaatar⁴, Erdenebaatar Sergelen^{3,5}, Dagvatseren Ariunzaya³, Orsoo Enkhmandal³, Sukhbaatar Tuvshintugs³, Shigeru Morikawa⁶ and Jiro Arikawa¹

¹ Department of Microbiology, Hokkaido University Graduate School of Medicine, Sapporo 060-8638, Japan

² National Institute of Infectious Diseases, Tokyo, Japan

³ Ecology Group, Department of Biology, School of Arts and Sciences, National University of Mongolia, Ulaanbaatar 14201, Mongolia

⁴ Laboratory of Virology, Institute of Veterinary Medicine, Mongolian University of Life Sciences, Ulaanbaatar 17024, Mongolia

⁵ WWF-ASER Office in Khovd Aimag

⁶ Department of Veterinary Medicine, National Institute of Infectious Diseases, Tokyo, Japan

Received for publication, August 10, 2016; accepted, December 8, 2016

Abstract

Seroepizootiological surveys among wild rodents were carried out on the east side of Lake Khovsgol in Mongolia in 2010 and 2011. A total of 76 voles belonging to the genera *Myodes* and *Microtus* were captured. Most of the voles that were seropositive to Tula virus antigen were Middendorf's voles (*Microtus middendorffii* (6/31)). Two of the 18 *Myodes* voles were also seropositive to Tula virus antigen. On the other hand, only one vole was seropositive to Puumala virus antigen. The results suggest that Tula virus was maintained in Middendorf's vole. This is the first report of detection of anti-Tula virus antibody in the central part of the Eurasia continent.

Key Words: hantavirus, bunyavirus, rodent-borne

The hantavirus belongs to the genus *Hantavirus* of the family *Bunyaviridae*. Various hantaviruses have been isolated from rodents, shrews, moles and bats²⁷. Each hantavirus appears to have a single predominant species of mammal as a natural reservoir. Among rodent-

borne hantaviruses, subfamily Sigmodontinae-borne viruses and subfamily Neotominae-borne viruses are causative agents of hantavirus pulmonary syndrome (HPS) and subfamily Murinae-borne viruses are causative agents of hemorrhagic fever with renal syndrome (HFRS)².

†Equally contributed to this work

*Corresponding author: Kumiko Yoshimatsu, DVM, PhD., Department of Microbiology, Hokkaido University Graduate School of Medicine, Kita-15, Nishi-7, Kita-ku, Sapporo 060-8638, Japan

Phone: +81-11-706-6905. Fax: +81-11-706-7879. E-mail: yosimatu@med.hokudai.ac.jp

doi: 10.14943/jjvr.65.1.39

In addition to those viruses, various hantaviruses have been detected from rodents belonging to the subfamily Microtinae, most of which are voles. Among those viruses, Puumala virus (PUUV) carried by the bank vole, (*Myodes glareolus*) is the most important causative agent of HFRS in Europe¹. On the other hand, Tula virus (TULV) was originally isolated from the European common vole (*Microtus arvalis*)¹². TULV has been widely found from voles in Russia²², Germany¹⁷ and other European countries^{10,13,18} is thought to be apathogenic for human^{8,24}.

Voies inhabit North America, Europe, and Asia. Distributions of various vole-borne hantaviruses in North America^{11,20} European countries^{15,18,22} and eastern parts of the Eurasia continent^{3,4,7,23,19} have been reported. However, there have been few studies on hantaviruses and their reservoir animals in the central region of the Eurasia continent^{14,25}.

In this study, rodents were captured in Mongolia and serologically screened with hantavirus antigens, after species identifications were confirmed with complete cytochrome b sequences. The captured rodents included various species of voles and the occurrences of the Middendorf's vole (*Microtus middendorffii*) in Mongolia was confirmed with molecular evidence. Throughout the screening, Middendorf's voles showed anti-Tula virus antibody. This is the first report about hantavirus infection in Mongolia and in Middendorf's voles.

A total of 77 rodents were captured on the east side of Lake Hövsgöl in Mongolia in 2010 and 2011 by using Sherman traps baited with raw seeds of millet (Table 1). Blood from the rodents was absorbed on filter papers (Advantech TOYO, type-1 filter paper). After collection of sera, the filter papers were stocked in the dark and 4°C until use, approximately for 3 years. Sera were extracted from the filter paper to 1 : 50 dilutions with Dulbecco's phosphate buffered saline pH 7.3 (PBS) according to the manufacturer's instruction. Then the extracted sera were treated for 30 min at 56°C for inactivation. PUUV and TULV

immune vole sera (*Microtus monteberlli*) prepared in a previous study were used as positive controls in serological tests²¹.

PUUV strain CG1820 and TULV strain Morabia were originally provided by Dr. A. Plyusnin (Helsinki, Finland) and were cultivated with Vero E6 cells as previously described²¹. The indirect immunofluorescent antibody (IFA) assay was performed as described previously²⁶. Briefly, acetone-fixed smears of Vero E6 cells infected with PUUV or TULV were used as antigens. Alexa Fluor 488-conjugated Protein A (Thermo Fisher Scientific, Waltham, MA, USA) was used for sera of voles (*Microtus* spp. and *Myodes* spp.) and sera of control voles²¹. The IFA antibody titer was regarded as the reciprocal of the highest serum dilution that displayed characteristic fluorescence in the infected Vero cells.

To verify the identity of the host species, total DNA was extracted from lung tissues using the GC series MagDEA[®] DNA 200 (GC) Kit (Precision System Science, Matsudo, Chiba, Japan). The entire 1,140-nucleotide cytochrome *b* gene of mtDNA was amplified by PCR using modified universal primers that were previously described⁵. PCR was performed in 50- μ L reaction mixtures containing 250 μ M dNTP, 1 U of AmpliTaq 360 Gold polymerase (Thermo Fisher Scientific) and 0.25 μ M of each primer. Cycling conditions consisted of initial denaturation at 95°C for 10 min followed by 35 cycles of denaturation at 94°C for 20 sec, annealing at 55°C for 1 min, and elongation at 72°C for 2 min in a Veriti thermal cycler (Life Technologies, Foster City, CA, USA).

As shown in Table 1, 77 rodent sera were screened by the IFA test against both PUUV and TULV antigens. Among the 77 rodents, 76 were voles belonging to the genus *Myodes* or *Microtus*. The IFA patterns of sera from six Middendorf's voles (MG349R36, MG1411RR7, MG1429RR25, MG1436RR32, MG1443RR39, and MG1445RR41) were similar to the IFA pattern of the positive control serum against TULV, which was characteristic granular and perinuclear

Table 1. List of rodents used in this study and serum antibody detection against PUUV and TULV antigens in IFA

| Genus | Species | No. of tested | IFA against* | | | |
|-----------------|----------------------|---------------|--------------|---|------|---|
| | | | PUUV | | TULV | |
| | | | ± | + | ± | + |
| <i>Myodes</i> | <i>rufocanus</i> | 4 | 0 | 0 | 0 | 1 |
| | <i>rutilus</i> | 14 | 0 | 0 | 1 | 0 |
| <i>Microtus</i> | <i>gregalis</i> | 12 | 0 | 0 | 0 | 0 |
| | <i>middendorffii</i> | 31 | 1 | 0 | 3 | 3 |
| | <i>oeconomus</i> | 5 | 0 | 0 | 0 | 0 |
| <i>Apodemus</i> | <i>peninsulae</i> | 1** | 0 | 0 | 0 | 0 |
| Total | | 77 | 1 | 0 | 4 | 4 |

*+, IFA titers over 100; ±, IFA titers was 50. IFA patterns of + and ± were identical. All sera did not show granular and diffused fluorescence in IFA antigen of Vero cells without infection.

**This serum was also negative to Hantaan virus antigen.




Fig. 1. IFA profiles of Middendorff's vole sera against TULV antigens. A: Positive IFA pattern of antiserum obtained from experimentally inoculated *Microtus montebelli* (Tegshduuren *et al.*). B: Typical negative IFA pattern of Middendorff's vole serum (MG1437RR33). C, D: Typical positive IFA patterns of Middendorff's vole serum (MG1429RR25 and MG1436RR32).

fluorescence in cytoplasm. Typical IFA antibody-positive patterns obtained with MG1429RR25 and MG1436RR32 are shown in Fig. 1. A typical antibody-negative IFA pattern from Middendorff's voles is also shown in the lower left panel of Fig. 1 (MG1437RR33). Six (19.3%) of the Middendorff's voles showed IFA titers that were positive (≥ 100) or weakly positive (50) to TULV

antigen. On the other hand, 1 (3.2%) of the 31 Middendorff's voles showed a weak positive IFA profile to PUUV antigen. This vole was antibody negative to TULV antigen. No antibody-positive voles against PUUV and/or TULV antigens was not found from *M. gregalis* and *M. oeconomus*. Antibody-positive voles were captured in both 2010 and 2011. The results indicated that

Middendorf's voles in northern Mongolia might be reservoir rodents for TULV or TULV-related hantavirus.

Eighteen *Myodes* rodents were captured in the same area in Mongolia. One *Myodes rufocanus* rodent and one *Myodes rutilus* rodent showed positive and weakly positive IFA profiles against only TULV, respectively. One field mouse (*A. peninsulae*) showed negative IFA reactions to PUUV, TULV and Hantaan virus antigen in the IFA (data not shown).

Sympatric habitats of voles belonging to the genera *Myodes* and *Microtus* have been reported in European countries and the Far East^{6,16}. In Europe, *Myodes* voles with PUUV and *Microtus* voles with TULAV have been captured at the same time. In Far Eastern part of Russia, *Myodes* voles with Hokkaido virus (HOKV) and *Microtus* voles with Khabarovsk virus (KBRV) or Vladivostok virus (VLAV) have been captured in the same area. In this study, a sympatric habitat of *Myodes* voles and *Microtus* voles was found in eastern shore of Lake Hövsgöl in Mongolia.

Myodes rufocanus was reported to be a reservoir rodent for HOKV, a PUUV-related virus, in Japan and Far Eastern part of Russia. Yashina *et al.* reported HOKV from *My. rufocanus* in the central area of Russia near Lake Baikal²⁵. Plyusnina *et al.* also reported HOKV *My. rufocanus* and VLAV from *Mi. fortis* which was also PUUV-related virus¹⁴ near Lake Baikal in Buryatia. Although *My. rufocanus* was captured in Mongolia, voles with antibodies positive to PUUV antigen were not found. On the other hand, one Middendorf's vole showed an anti-PUUV antibody in serum. HOKV might be maintained in a minor population of reservoir voles in Mongolia.

In addition to these observations, two *Myodes* voles showed anti-TULV antibody in their sera. The prevalence of antibodies in Middendorf's voles against TULV was 19% (6/31), that was a higher prevalence than other species of voles. TULV-like virus maintained among Middendorf's voles in the area might be source of spillover infection from Middendorf's voles to other voles.

Similar spillover infection of PUUV was reported in Europe from *Myodes* rodents to *Microtus* rodents⁹.

Previously, we examined antibody cross-reactivities to TULV and PUUV²¹. Little cross-reactivity of antibodies was detected in the sera of experimentally inoculated mice and voles. Similarly, sera from Middendorf's voles did not show any cross-reactivity against PUUV antigen in this study. These observations indicated that both TULV and PUUV antigens should be used for serological screening of voles.

In this study, a virus genome was not detected from preserved organs of Middendorf's voles (data not shown). However, the existence of a TULV-like virus in the central region of the Eurasia continent was serologically shown (Supplement Fig. 1). The results obtained in this study might contribute to studies on radiation and evolution of *Microtus* rodents with their hantaviruses.

Acknowledgements

This manuscript was proofread by an English consultant, SES English Service, Sapporo, Japan. The research was also partly supported by the USA NSF-PIRE Grant (OISE 0729876) to B. Boldgiv.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.14943/jjvr.65.1.39>

References

- 1) Brummer-Korvenkontio, M., Vaheri, A., Hovi, T., von Bonsdorff, C. H., Vuorimies, J., Manni, T., Penttinen, K., Oker-Blom, N. and Lahdevirta, J. 1980. Nephropathia epidemica:

- detection of antigen in bank voles and serologic diagnosis of human infection. *J. Infect. Dis.*, **141**: 131-134.
- 2) de Oliveira, R. C., Guterres, A., Fernandes, J., D'Andrea, P. S., Bonvicino, C. R. and de Lemos, E. R. 2014. Hantavirus reservoirs: current status with an emphasis on data from Brazil. *Viruses*, **6**: 1929-1973.
 - 3) Ge, X. Y., Yang, W. H., Pan, H., Zhou, J. H., Han, X., Zhu, G. J., Desmond, J. S., Daszak, P., Shi, Z. L. and Zhang, Y. Z. 2016. Fugong virus, a novel hantavirus harbored by the small oriental vole (*Eothenomys eleusis*) in China. *Virol. J.*, **13**: 27.
 - 4) Horling, J., Chizhikov, V., Lundkvist, A., Jonsson, M., Ivanov, L., Dekonenko, A., Niklasson, B., Dzagurova, T., Peters, C. J., Tkachenko, E. and Nichol, S. 1996. Khabarovsk virus: a phylogenetically and serologically distinct hantavirus isolated from *Microtus fortis* trapped in far-east Russia. *J. Gen. Virol.*, **77**: 687-694.
 - 5) Irwin, D. M., Kocher, T. D. and Wilson, A. C. 1991. Evolution of the cytochrome b gene of mammals. *J. Mol. Evol.*, **32**: 128-144.
 - 6) Kariwa, H., Yoshimatsu, K., Sawabe, J., Yokota, E., Arikawa, J., Takashima, I., Fukushima, H., Lundkvist, A., Shubin, F. N., Isachkova, L. M., Slonova, R. A., Leonova, G. N. and Hashimoto, N. 1999. Genetic diversities of hantaviruses among rodents in Hokkaido, Japan and Far East Russia. *Virus Res.*, **59**: 219-228.
 - 7) Kariwa, H., Yoshizumi, S., Arikawa, J., Yoshimatsu, K., Takahashi, K., Takashima, I. and Hashimoto, N. 1995. Evidence for the existence of Puumala-related virus *Clethrionomys rufocanus* in Hokkaido, Japan. *Am. J. Trop. Med. Hyg.*, **53**: 222-227.
 - 8) Klempa, B., Meisel, H., Rath, S., Bartel, J., Ulrich, R. and Kruger, D. H. 2003. Occurrence of renal and pulmonary syndrome in a region of northeast Germany where Tula hantavirus circulates. *J. Clin. Microbiol.*, **41**: 4894-4897.
 - 9) Klingstrom, J., Heyman, P., Escutenaire, S., Sjolander, K. B., De Jaegere, F., Henttonen, H. and Lundkvist, A. 2002. Rodent host specificity of European hantaviruses: evidence of Puumala virus interspecific spillover. *J. Med. Virol.*, **68**: 581-588.
 - 10) Korva, M., Duh, D., Puterle, A., Trilar, T. and Zupanc, T. A. 2009. First molecular evidence of Tula hantavirus in *Microtus voles* in Slovenia. *Virus Res.*, **144**: 318-322.
 - 11) Lee, P. W., Amyx, H. L., Yanagihara, R., Gajdusek, D. C., Goldgaber, D. and Gibbs, C. J., Jr. 1985. Partial characterization of Prospect Hill virus isolated from meadow voles in the United States. *J. Infect. Dis.*, **152**: 826-869.
 - 12) Plyusnin, A., Vapalahti, O., Lankinen, H., Lehvaslaiho, H., Apekina, N., Myasnikov, Y., Kalliokokko, H., Henttonen, H., Lundkvist, A., Brummerkorvenkontio, M., Gavrilovskaya, I. and Vaheri, A. 1994. Tula virus: A newly detected hantavirus carried by European common voles. *J. Virol.*, **8**: 7833-7783.
 - 13) Plyusnina, A., Laakkonen, J., Niemimaa, J., Henttonen, H. and Plyusnin, A. 2008. New Genetic Lineage of Tula Hantavirus in *Microtus arvalis obscurus* in Eastern Kazakhstan. *Open Virol. J.*, **2**: 32-36.
 - 14) Plyusnina, A., Laakkonen, J., Niemimaa, J., Nemirov, K., Muruyeva, G., Pohodiev, B., Lundkvist, A., Vaheri, A., Henttonen, H., Vapalahti, O. and Plyusnin, A. 2008. Genetic analysis of hantaviruses carried by *Myodes* and *Microtus* rodents in Buryatia. *Virol. J.*, **5**: 4.
 - 15) Pounder, K. C., Begon, M., Sironen, T., Henttonen, H., Watts, P. C., Voutilainen, L., Vapalahti, O., Klempa, B., Fooks, A. R. and McElhinney, L. M. 2013. Novel hantavirus in field vole, United Kingdom. *Emerg. Infect. Dis.*, **19**: 673-675.
 - 16) Schmidt, S., Essbauer, S. S., Mayer-Scholl, A., Poppert, S., Schmidt-Chanasit, J., Klempa, B., Henning, K., Schares, G., Groschup, M. H., Spitzenberger, F., Richter, D., Heckel, G. and Ulrich, R. G. 2014. Multiple infections of rodents with zoonotic pathogens in Austria. *Vector Borne Zoonotic Dis.*, **14**: 467-475.
 - 17) Schmidt, S., Saxenhofer, M., Drewes, S., Schlegel, M., Wanka, K. M., Frank, R., Klimpel, S., von Blanckenhagen, F., Maaz, D., Herden, C., Freise, J., Wolf, R., Stubbe, M., Borkenhagen, P., Ansoerge, H., Eccard, J. A., Lang, J., Jourdain, E., Jacob, J., Marianneau, P., Heckel, G. and Ulrich, R. G. 2016. High genetic structuring of Tula hantavirus. *Arch. Virol.*, **161**: 1135-1149.
 - 18) Schmidt-Chanasit, J., Essbauer, S., Petraityte, R., Yoshimatsu, K., Tackmann, K., Conraths, F. J., Sasnauskas, K., Arikawa, J., Thomas, A., Pfeffer, M., Scharninghausen, J. J., Spletstoeser, W., Wenk, M., Heckel, G. and Ulrich, R. G. 2010. Extensive host sharing of Central European Tula virus. *J. Virol.*, **84**: 459-474.
 - 19) Song, K. J., Baek, L. J., Moon, S., Ha, S. J., Kim, S. H., Park, K. S., Klein, T. A., Sames, W., Kim, H. C., Lee, J. S., Yanagihara, R.

- and Song, J. W. 2007. Muju virus, a novel hantavirus harboured by the arvicolid rodent *Myodes regulus* in Korea. *J. Gen. Virol.*, **88**: 3121–3129.
- 20) Song, W., Torrez-Martinez, N., Irwin, W., Harrison, F. J., Davis, R., Ascher, M., Jay, M. and Hjelle, B. 1995. Isla Vista virus: a genetically novel hantavirus of the California vole *Microtus californicus*. *J. Gen. Virol.*, **76** (Pt 12): 3195–3199.
- 21) Tegshduuren, E., Yoshimatsu, K., Taruishi, M., Endo, R., Shimizu, K., Koma, T., Yasuda, S. P., Kariwa, H., Arikawa, J. and Ishihara, C. 2010. Different cross-reactivity of human and rodent sera to Tula virus and Puumala virus. *Comp. Immunol. Microbiol. Infect. Dis.*, **33**: e67–73.
- 22) Tkachenko, E. A., Witkowski, P. T., Radosa, L., Dzagurova, T. K., Okulova, N. M., Yunicheva, Y. V., Vasilenko, L., Morozov, V. G., Malkin, G. A., Kruger, D. H. and Klempa, B. 2015. Adler hantavirus, a new genetic variant of Tula virus identified in Major's pine voles (*Microtus majori*) sampled in southern European Russia. *Infect. Genet. Evol.*, **29**: 156–163.
- 23) Vapalahti, O., Lundkvist, A., Fedorov, V., Conroy, C. J., Hirvonen, S., Plyusnina, A., Nemirov, K., Fredga, K., Cook, J. A., Niemimaa, J., Kaikusalo, A., Henttonen, H., Vaheri, A. and Plyusnin, A. 1999. Isolation and characterization of a hantavirus from *Lemmus sibiricus*: Evidence for host switch during hantavirus evolution. *J. Virol.*, **73**: 5586–5592.
- 24) Vapalahti, O., Mustonen, J., Lundkvist, A., Henttonen, H., Plyusnin, A. and Vaheri, A. 2003. Hantavirus infections in Europe. *Lancet Infect. Dis.*, **3**: 653–661.
- 25) Yashina, L. N., Abramov, S. A., Dupal, T. A., Danchinova, G. A., Malyshev, B. S., Hay, J., Gu, S. H. and Yanagihara, R. 2015. Hokkaido genotype of Puumala virus in the grey red-backed vole (*Myodes rufocanus*) and northern red-backed vole (*Myodes rutilus*) in Siberia. *Infect. Genet. Evol.*, **33**: 304–313.
- 26) Yoshimatsu, K., Arikawa, J. and Kariwa, H. 1993. Application of a recombinant baculovirus expressing hantavirus nucleocapsid protein as a diagnostic antigen in IFA test: cross reactivities among 3 serotypes of hantavirus which causes hemorrhagic fever with renal syndrome (HFRS). *J. Vet. Med. Sci.*, **55**: 1047–1050.
- 27) Zhang, Y. Z. 2014. Discovery of hantaviruses in bats and insectivores and the evolution of the genus Hantavirus. *Virus Res.*, **187**: 15–21.