



Title	Caspase-dependent cleavage regulates protein levels of Epstein-Barr virus-derived latent membrane protein 1
Author(s)	Togi, Sumihito; Hatano, Yosuke; Muromoto, Ryuta; Kawanishi, Eri; Ikeda, Osamu; Hirashima, Koki; Kon, Shigeyuki; Kitai, Yuichi; Yasui, Teruhito; Oritani, Kenji; Matsuda, Tadashi
Citation	FEBS Letters, 590(6), 808-818 <a href="https://doi.org/10.1002/1873-3468.12119">https://doi.org/10.1002/1873-3468.12119</a>
Issue Date	2016-03
Doc URL	<a href="http://hdl.handle.net/2115/64895">http://hdl.handle.net/2115/64895</a>
Rights	This is the peer reviewed version of the following article: Caspase-dependent cleavage regulates protein levels of Epstein-Barr virus-derived latent membrane protein 1, FEBS Letters 590(6), pp808-818, 2016, which has been published in final form at 10.1002/1873-3468.12119. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	manuscript.pdf



[Instructions for use](#)

**Title: Caspase-dependent cleavage regulates protein levels of Epstein-Barr virus-derived latent membrane protein 1**

Running title: EBV-LMP1 is degraded by caspases

Authors: Sumihito Togi<sup>a</sup>, Yosuke Hatano<sup>a</sup>, Ryuta Muromoto<sup>a</sup>, Eri Kawanishi<sup>a</sup>, Osamu Ikeda<sup>a</sup>, Koki Hirashima<sup>a</sup>, Shigeyuki Kon<sup>a</sup>, Yuichi Kitai<sup>a</sup>, Teruhito Yasui<sup>b</sup>, Kenji Oritani<sup>c</sup>, and Tadashi Matsuda<sup>a,¶</sup>

<sup>a</sup>Department of Immunology, Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita-12 Nishi-6, Kita-Ku, Sapporo, 060-0812, Japan

<sup>b</sup>Department of Molecular Immunology, Research Institute for Microbial Diseases, Osaka University, 2-2 Yamada-oka, Suita, Osaka 565-0871, Japan

<sup>c</sup>Department of Hematology and Oncology, Graduate School of Medicine, Osaka University, 2-2 Yamada-oka, Suita, Osaka 565-0871, Japan

<sup>¶</sup>Address correspondence to: Dr. Tadashi Matsuda, Department of Immunology, Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita-Ku Kita 12 Nishi 6, Sapporo 060-0812, Japan Tel.: 81-11-706-3243, Fax: 81-11-706-4990, E-mail: [tmatsuda@pharm.hokudai.ac.jp](mailto:tmatsuda@pharm.hokudai.ac.jp)

## **Abstract**

Epstein–Barr virus (EBV)-encoded latent membrane protein-1 (LMP1) plays pathogenetic roles in EBV-related diseases. Thus, host cells employ several mechanisms to regulate LMP1 functions, and we previously reported possible regulation by signal transducing adaptor protein-2 as well as BS69. Here, we found that caspase-3 mainly degraded LMP1 proteins in HeLa cells, leading to decreased NF- $\kappa$ B and STAT3 activation. Caspase-3 cleaved the consensus DNTD sequences in the CTAR3 region of LMP1. Of importance, LMP1 expression strongly enhanced caspase-3 activity. Taken together, the reduction of LMP1 protein levels by caspases is likely to be a newly identified host defense against EBV infection.

Abbreviations: EBV, Epstein–Barr virus; LMP, latent membrane protein; CTAR, C-terminal activation region; LUC, luciferase

Keywords: EBV, LMP1, caspase, degradation, NF- $\kappa$ B activation

## 1. Introduction

Epstein–Barr virus (EBV) is linked to the development of multiple malignancies, including post-transplant lymphoproliferative disorders, Hodgkin lymphomas, and nasopharyngeal carcinomas. EBV-encoded latent membrane protein 1 (LMP1) is expressed in many EBV-associated tumor cells and is responsible for the majority of the observed altered cellular growth properties. Structurally, LMP1 is an integral membrane protein composed of 386 amino acids (aa) with a short cytoplasmic N-terminal domain (aa 1-24), six transmembrane domains (aa 25-186), and a long cytoplasmic C-terminal tail (aa 187-386) [1, 2, 3]. The cytoplasmic C-terminal tail is composed of three C-terminal activation regions (CTAR1, CTAR2, and CTAR3, respectively), which are mainly involved in LMP1-mediated signaling events [1, 4, 5]. CTAR1 contains a consensus TRAF-binding motif (PXQXT), and can bind to TRAFs 1, 2, 3, and 5 [1,6-9]. CTAR2 interacts with TNFR-associated death domain protein (TRADD), receptor-interacting protein [1, 10, 11], and a multi-domain cellular protein BS69 [12]. Signals through the CTAR domains induce p100 and p105 NF- $\kappa$ B precursors and upregulate their processing into p52 and p50, respectively [13, 14], indicating that LMP1 is involved in both canonical and non-canonical activation of the NF- $\kappa$ B pathway.

Binding of LMP1 to TRAFs and TRADD also initiates the formation of a signaling complex, which leads to the activation of mitogen-activated protein kinase, p38 [15]. LMP1 also activates c-Jun N-terminal kinase (JNK) as well as PI3K/Akt and STAT3 signaling [12, 16-19]. We recently identified an adaptor protein, STAP-2, as a new binding partner of both CTAR1 and CTAR2, and demonstrated that STAP-2 negatively regulates LMP1-mediated NF- $\kappa$ B activation (20). Moreover, we reported that BS69 regulates LMP1-mediated NF- $\kappa$ B activation by influencing binding between LMP1 and TRADD [21, 22].

LMP1 is a short-lived protein, with a reported half-life ranging from 1.5 to 7 h, depending on the cell type [23-25]. Additionally, one report has found that LMP1 is in part degraded via the ubiquitin-mediated proteasome degradation pathway in COS7/5 fibroblasts [26]. Notably, LMP1 degradation was detected in lymphomas derived from the LMP1 transgenic mice expressing LMP1 in B lymphocytes [27]. However, very little information is available on the regulation of degradation rate and stability of LMP1 proteins in individual cells.

In the present study, we have found that LMP1 is mainly degraded by the caspase cascade in HeLa cells. In this situation, caspase-3 cleaves the CTAR3 region of LMP1 at consensus DNTD sequences and is, to the best of our knowledge, a previously

undescribed host mechanism to control LMP1 protein levels. Of importance, LMP1 expression strongly enhances caspase-3 activities. We also discuss possible meanings of LMP1 degradation by caspases during EBV infection.

## 2. Materials and Methods

### 2.1. Reagents and Antibodies

Recombinant human TNF- $\alpha$ , cycloheximide (CHX) and NH<sub>4</sub>Cl were purchased from WAKO Chemicals (Osaka, Japan). MG132, Z-VAD, Z-IETD, Ac-DNLD, Ac-VEID, Ac-LEHD and Z-DEVD were purchased from Peptide Institute (Osaka, Japan). Expression vector for FLAG-tagged LMP1 was described previously [20], and its mutants were generated by PCR and cloned by using TaKaRa In-Fusion Kit (primer sequences are available upon request). The synthesized FLAG-tagged LMP1 4xANTD mutant was obtained from GenScript (Piscataway, NJ). STAT3-LUC and NF- $\kappa$ B-LUC were kindly provided by T. Hirano (Osaka University Medical School, Osaka, Japan) and T. Fujita (Kyoto University, Kyoto, Japan) [28, 29]. Anti-FLAG antibody was purchased from Sigma-Aldrich (St. Louis, MO). Anti-caspase-3, anti-caspase-7, and anti-caspase-8 antibodies were from Cell Signaling Technologies (Beverly, MA). Anti-actin antibody was from Millipore (Billerica, MA).

### 2.2. Cell Culture, Transfection, siRNA and luciferase assay

A human cervix carcinoma cell line, HeLa was maintained in DMEM containing 10 %

FCS. HeLa cells were transfected using jetPEI (Polyplus Transfection, Strasbourg, France) according to the manufacturer's instructions.. The siRNAs targeting human caspase-3, -7, or -8 used in this study were as follows: caspase-3 #1, 5'-GCGUGAUGUUUCUAAAGAAGA-3' (sense), 5'-UUCUUUAGAAACAUCACGCAU-3' (antisense) ; caspase-3 #2, 5'-CCCUGGACAACAGUUUAUAATT-3' (sense), 5'-UUAUAACUGUUGUCCAGGGTT-3' (antisense) ; caspase-7 #1, 5'-GCAUCAUAAUAAACAACAAGA-3' (sense), 5'-UUGUUGUUUAUUAUGAUGCAU-3' (antisense) ; caspase-7 #2, 5'-GGAACUCUACUUCAGUCAUA-3' (sense), 5'-UUGACUGAAGUAGAGUCCUU-3' (antisense) ; caspase-8 #1, 5'-GAGUCUGUGCCCAAUCAATT-3' (sense), 5'-UUGAUUUGGGCACAGACUCTT-3' (antisense) ; caspase-8 #2, 5'-CAGCCUACUUUCACACUAATT-3' (sense), 5'-UUAGUGUGAAAGUAGGCUGAG-3' (antisense) ; Control siRNA was obtained from Qiagen (nonsilencing; catalog No. 1022076). HeLa cells were plated on 24-well plates at  $2 \times 10^4$  cells/well and incubated with an siRNA-Lipofectamine 2000 (Invitrogen, Carlsbad, CA) mixture at 37 °C for 6 h followed by the addition of fresh medium

containing 10 % FCS [20]. Luciferase activities of HeLa cells transfected with NF- $\kappa$ B-LUC or STAT3-LUC as well as internal control, in which the  $\alpha$ 2-macroglobulin promoter drives expression of a luciferase reporter gene, were measured using a Dual-Luciferase Reporter Assay System (Promega, Madison, WI) according to the manufacturer's instructions.

### 2.3. Immunoblotting

Western blotting assays were performed, as described previously [30]. The cell lysates were resolved on SDS-PAGE and transferred to polyvinylidene difluoride transfer membrane (PerkinElmer, Boston, MA). The filters were then immunoblotted with each antibody. Immunoreactive proteins were visualized using an ECL detection system (Millipore).

### 2.4. RNA Isolation and Quantitative Real-time (qRT)-PCR

Cells were harvested and total RNAs were prepared by using TRI Reagent (Molecular Research Center, Cincinnati, OH, USA). First-strand cDNA was synthesized from 1  $\mu$ g of total RNA with ReverTra Ace (TOYOBO, Osaka, Japan). Quantitative real-time PCR analysis of mRNA transcripts was carried out using a combination of a KAPA

SYBR FAST qPCR master mix (KAPA Biosystems, Woburn, MA, USA) with an Mx3005P real-time PCR system (Stratagene, Santa Clara, CA, USA). The primers used for qPCR were: *IL-6*, 5'- AACTGGATATAATCAGGAAAT-3' (sense), 5'- CTAGGTTTGCCGAGTAGATCTC-3' (antisense); *ACTIN*, 5'- TGTTACCAACTGGGACGACA-3' (sense), 5'-GGGGTGTGGAAGGTCTCAAA-3' (antisense).

## 2.5. Analysis of caspase activity

Caspase-3 activity was measured using the Caspase-Glo 3/7 Assay kits essentially according to the manufacturer's instructions (Promega) [31].

### 3. Results

#### 3.1. LMP1 is degraded in a caspase-dependent manner in HeLa cells

To confirm the degradation of LMP1 protein, we first assessed the stability of LMP1 proteins in LMP1-transfected HeLa cells. HeLa cells ectopically expressing LMP1 were cultured in the presence of CHX to block protein synthesis. As shown in Fig. 1A and B, the treatment of LMP1-transfected HeLa cells with CHX induced a rapid reduction of full-length LMP1 (FL-LMP1) protein levels; in addition, multiple smaller LMP1-derived bands were clearly visible. This result likely indicates that LMP1 protein was comprehensively degraded in HeLa cells. We next examined effects of various protease inhibitors on LMP1 protein degradation using this experimental system. As shown in Fig. 1C and D, a pan-caspase inhibitor, Z-VAD, significantly blocked the degradation of LMP1 protein in HeLa cells while a proteasomal inhibitor, MG132, or a lysosomal inhibitor, NH<sub>4</sub>Cl, did not. In the same situation, the treatment of HeLa cells with MG132 blocked TNF- $\alpha$ -induced I $\kappa$ B $\alpha$  degradation in HeLa cells (Fig. 1S).

Therefore, in HeLa cells, ectopically expressed LMP1 is mostly degraded in a caspase-dependent manner.

### 3.2. Caspase-3 is responsible for LMP1 protein degradation in HeLa cells

To identify which caspase is responsible for LMP1 protein degradation in HeLa cells, LMP1-transfected HeLa cells were cultured in the presence of a caspase-3 specific inhibitor, Ac-DNLD, a caspase-3/7 specific inhibitor, Z-DEVD, a caspase-6 specific inhibitor, Ac-VEID, a caspase-8-specific inhibitor, IETD, a caspase-9 specific inhibitor, Ac-LEHD or a pan-caspase inhibitor, Z-VAD. As shown in Fig. 2A and B, the treatment with Z-DEVD, and Ac-VEID as well as Z-VAD significantly blocked LMP1 protein degradation in HeLa cells. Although these results indicate the involvement of caspases, the evaluation of each caspase in LMP1 degradation is difficult. This might be due to the limited selectivity of the caspase inhibitors because they are designed based on the substrate specificity profiles and have similar constructs. To directly clarify this issue, we employed siRNA techniques to specifically knockdown each caspase. We transfected specific siRNAs for caspase-3 (#1 and #2), caspase-7 (#1 and #2), or caspase-8 (#1 and #2) into HeLa cells expressing LMP1. The reduction of caspase-3, but not caspase-7 or caspase-8 expression, resulted in a significant increase of FL-LMP1 protein levels in LMP1-transfected HeLa cells (Fig. 2C and D). Therefore, caspase-3 is the main caspase responsible for degradation of LMP1 in HeLa cells. However, caspase-3 siRNA treatment of HeLa cells also increased accumulation of the

partially degraded fragment of LMP1 as shown in the case of the treatment with Ac-DNLD (Fig. 2A). Thus, we could not exclude the involvement of other caspases in LMP1 protein degradation in HeLa cells, although caspase-3 knockdown in HeLa cells by siRNA treatment was not complete.

### 3.3. LMP1 expression induces caspase-3 activation in HeLa cells

To investigate the relationship between LMP1 and caspase-3, we analyzed the caspase enzymatic activities of caspase-3 after LMP1-transfection into HeLa cells. As shown in Fig. 3A, the activities of these caspases markedly increased in correlation with LMP1 protein degradation in LMP1-transfected HeLa cells, whereas they were completely blocked by the addition of Z-VAD. This was also confirmed by western blot analyses (Fig. 3B) showing that the cleavage of caspase-3 and -8 was significantly increased in LMP1-transfected HeLa cells compared with vector-transfected HeLa cells. However, we could not detect the cleavage of caspase-7 in these cells (data not shown).

### 3.4. Caspase inhibition enhances LMP1-mediated signals in HeLa cells

To analyze the involvement of caspases on LMP1-mediated signals, we examined the effects of Z-VAD on LMP1-induced NF- $\kappa$ B or STAT3 activation using transient

reporter assays with NF- $\kappa$ B-LUC or STAT3-LUC. Both luciferase activities were markedly induced by the expression of LMP1 in HeLa cells (Fig. 4A and B). Importantly, LMP1-induced NF- $\kappa$ B-LUC or STAT3-LUC activities were significantly enhanced by Z-VAD treatment in parallel with the increase in LMP1 protein levels, suggesting that caspase activation negatively regulates LMP1-induced NF- $\kappa$ B and/or STAT3 activation.

NF- $\kappa$ B activation induces the expression of inflammatory cytokines, such as IL-6. To confirm the regulation of LMP1-induced NF- $\kappa$ B activation by caspases, LMP1-induced IL-6 mRNA expression was analyzed in the Z-VAD-treated or untreated cells. LMP1-induced IL-6 mRNA expression was markedly augmented by the treatment of LMP1-transfected HeLa cells with Z-VAD in (Fig. 4C). Interestingly, the treatment with Z-VAD had no effect on TNF- $\alpha$ -induced NF- $\kappa$ B activation (Fig. 4D).

Therefore, caspase activity negatively regulates NF- $\kappa$ B and/or STAT3 activation mediated by LMP1, but not TNF- $\alpha$ .

### 3.5. Caspase-3 regulates LMP1-mediated NF- $\kappa$ B activation in HeLa cells

To further explore whether caspase-3 regulates LMP1-induced NF- $\kappa$ B activation, we used siRNAs to reduce endogenous expression of caspase-3 in HeLa cells. HeLa

cells were transfected with specific siRNAs for caspase-3 (#1 and #2) or a control siRNA. We then determined the effects of caspase-3 siRNAs on LMP1-induced NF- $\kappa$ B activation in HeLa cells. As shown in Fig. 5A, siRNA-mediated knock down of caspase-3 resulted in a significant enhancement of LMP1-induced NF- $\kappa$ B-LUC activities in parallel with the increase in LMP1 protein levels. We also determined the effects of caspase-3 siRNAs on LMP1-induced IL-6 mRNA expression. LMP1-induced IL-6 mRNA expression was also significantly augmented by reduced expression of caspase-3 in LMP1-transfected HeLa cells (Fig. 5B).

Therefore, caspase-3 is responsible for LMP1 protein degradation and LMP1-induced NF- $\kappa$ B activation in HeLa cells.

### 3.6. Determination of caspase cleavage sites in LMP1 in HeLa cells

To determine which domains of LMP1 were involved in caspase-dependent degradation, HeLa cells were transfected with a series of FLAG-tagged LMP1 deletion mutants (Fig. 6A). The transfectants were treated with Z-VAD or left untreated, lysed, and immunoblotted with an anti-FLAG antibody. As shown in Fig. 6B, the protein levels of both FL-LMP1 and LMP1(1-349) were sensitive to Z-VAD treatment. However, no significant alteration of protein levels of both LMP1(1-192) and LMP1(1-231) was

observed in samples with Z-VAD. These results indicate that the caspase cleavage sites in LMP1 may exist in the CTAR3 region. We found four putative consensus casapase-3 cleavage sequences (DNTD) at the positions of aa 257–260, aa 268–271, aa 284–287, and aa 295–298 in that region (data not shown). We then examined these four repeated DNTD motifs using a  $\Delta$ 4xDNTD mutant with a deletion of aa 249–307 in LMP1(1-349) (Fig. 6C). As shown in Fig. 6D, LMP1(1-349/ $\Delta$ 4xDNTD) protein levels did not change before or after the Z-VAD treatment. In addition, the multiple degraded bands observed in control LMP1(1-349) samples were not seen in LMP1(1-349/ $\Delta$ 4xDNTD). Of note, LMP1(1-349/ $\Delta$ 4xDNTD) protein yielded two bands, which migrated a little faster, in the absence of Z-VAD, indicating that other cleavage sites may exist in LMP1(1-349/ $\Delta$ 4xDNTD). Upon investigation, another potential caspase cleavage site was identified as LMTD (aa338–341). To further evaluate the potential cleavage of LMP1 at this site, we generated a LMP1 mutant with substitution of glutamic acid for aspartic acid at residue 341 (D341E) in LMP1(1-349/ $\Delta$ 4xDNTD). The LMP1(1-349/ $\Delta$ 4xDNTD/D341E) protein showed a single protein band in the absence or presence of Z-VAD, indicating that the LMP1(1-349/ $\Delta$ 4xDNTD/D341E) protein gained complete resistance to caspase-dependent proteolysis. We also introduced these mutations into FL-LMP1 (FL-LMP1 ( $\Delta$ 4xDNTD/D341E)) (Fig. 6E).

As shown in Fig. 6F, the FL-LMP1 ( $\Delta$ 4xDNTD/D341E) protein was completely resistant to caspase-dependent proteolysis. We finally generated a LMP1 4xANTD mutant with substitution of alanine for aspartic acid at residues 257, 268, 284 and 295 in LMP1, and a LMP1 4xANTD/D341E mutant with substitution of glutamic acid for aspartic acid at residue 341 in LMP1 4xANTD. As shown in Fig. 6G, multiple smaller LMP1-derived degraded bands disappeared in the LMP1 4xANTD mutant, although LMP1 4xANTD protein yielded two bands as well as LMP1(1-349/ $\Delta$ 4xDNTD). More importantly, LMP1 4xANTD/D341E was completely resistant to degradation in HeLa cells (Fig. 6G). The treatment of Z-VAD tended to slightly increase the protein levels in LMP1 mutants, which do not contain the suggested caspase cleavage sites. However, there is no statistically significant difference, when we normalized the protein level LMP1 to the actin protein of the same sample (data not shown). Thus, we have identified four repeated DNTD motifs and a LMTD sequence in CTAR3 as consensus caspase cleavage sites responsible for degradation by caspase-3 and another caspase, respectively, although we could not exclude the possibility that the caspase-8 cleavage motif LMTD may be newly exposed by deletion of the four repeated DNTD motifs in the CTAR3 region.

#### 4. Discussion

EBV, among other tumor-associated viruses, can affect host cell growth and survival, resulting in malignant transformation [32, 33]. Some virus proteins function to mediate escape from immune surveillance, while others directly regulate growth and/or apoptotic signals [34, 35]. In the case of EBV, a nuclear antigen, EBNA1, has anti-apoptotic effects, while EBNA3 binds to Rb and promotes cell-cycle progression in Burkitt lymphoma cells. The small nuclear RNA, EBER1, binds to and inhibits protein kinase RNA, which is an effector of the apoptotic signaling pathway. In addition, EBERs can induce resistance to apoptosis via increased expression of an anti-apoptotic factor BCL-2 [1, 33]. LMP1 also contributes to survival and/or tumor progression by providing CD40/CD40L-like signals or modifying the TNF receptor pathway [1, 2, 3, 33]. Here, we describe a new caspase cascade-dependent regulatory mechanism controlling LMP1. In HeLa cells, caspase-3 mainly degrades LMP1 proteins, resulting in decreased LMP1-induced NF- $\kappa$ B activation and IL-6 production. We have found that LMP1 expression induces caspase-3 activation, and activated caspase-3 in turn cleaves the CTAR3 region of LMP1. Therefore, caspases negatively regulate LMP1 functions by controlling protein levels.

Caspases are often targeted by a variety of viruses in infected cells because these host enzymes play a central role in controlling cell death machinery [33, 34]. Thus, many viruses can interfere with caspase activation or inhibit caspase activities. Indeed, caspase-8 activities are inhibited by various virus-encoded proteins, including the baculovirus p35 protein, the adenovirus E3 14.7 kDa protein, several poxvirus serpins and vFLIPs (viral FLIPs), the gammaherpesvirus vFLIPs, and the cytomegalovirus viral inhibitor of caspase 8 activation protein. In contrast, we have shown that caspase-3 activities are upregulated in response to LMP1 expression and that activated caspase-3 degrades LMP1 protein. As mentioned above, EBV-derived LMP1 provides strong signals for cell growth and survival. Thus, the relationship between caspase-3 and LMP1 suggests a likely target for the host defense to block EBV infection. However, our experimental model was restricted to LMP1, although EBV produces a number of viral proteins. Further analysis will clarify the extent to which caspases act to inhibit EBV infection in host cells. Of note, an interesting report indicates that the inhibition of caspase activities enhances virus replication in a HIV-infected T-cell line [36]. Virus replication was caused by sensitization of TNF-R1 towards endogenously produced TNF- $\alpha$ . The activation of Fas also triggered HIV replication, which was further enhanced in the presence of Z-VAD. The authors concluded that caspase inhibitors

sensitize both Fas and TNF-R1 to mediate the activation of HIV in latently infected cells. Therefore, caspases function to protect host cells from virus infection at multiple intracellular events, and the degradation of EBV LMP1 by caspase-3 is likely to be a potential mechanism of host defense against EBV.

With regard to LMP1 protein degradation, Aviel et al. reported that LMP1 protein was degraded by the ubiquitin-proteasome pathway in a fibroblast line COS7/5 [26]. LMP1 protein levels were significantly increased by treatment with a proteasome inhibitor MG132. As we show here, treatment of HeLa cells with Z-VAD, but not MG132, increased LMP1 protein levels. In addition, our preliminary experiments also indicated that LMP1 protein was degraded via both ubiquitin-proteasome and caspase pathways in B lymphocyte lines (data not shown). Therefore, at least two mechanisms exist for LMP1 protein degradation, and the use of these two pathways to degrade LMP1 protein is likely to be dependent on cell type.

The CTAR3 region of LMP1 contains 4xDNTD and LMTD, which are the consensus caspase cleavage sequences. CTAR3 has been shown to bind JAK3 via PXXPP motifs (aa 275–280 and aa 302–307) that activate DNA binding of STAT1 [5]. CTAR3, but not CTAR1 or CTAR2, has been shown to interact with SUMO-conjugating enzyme Ubc9 [37]. Interactions between Ubc9 and CTAR3 mediate LMP1-induced protein

sumoylation, in turn influencing cellular migration. Furthermore, LMP1 expression induced sumoylation of IRF7 in a CTAR3-dependent manner, followed by decreasing its turnover, increasing nuclear retention, decreasing DNA binding, and limiting transcriptional activation [38]. Our results indicate that the cleavage of LMP1 protein at the CTAR3 region significantly reduces the ability to induce NF- $\kappa$ B activation and IL-6 production. Although the precise mechanisms are unknown, the CTAR3 region may be required for full LMP1 functions.

Here, we provide evidence that caspases can inhibit EBV LMP1-mediated NF- $\kappa$ B activation through protein degradation of LMP1. Our data suggest the possibility that controlling LMP1 protein levels may be a novel strategy for anti-viral therapy to regulate EBV LMP1-induced NF- $\kappa$ B activation.

## **Acknowledgments**

This study was supported in part by Grant-in-Aid for scientific research from Ministry of Education, Culture, Sports, Science and Technology of Japan.

## **Disclosures**

The authors have no financial conflicts of interest.

## **Author contributions**

ST, YH, RM, EK, OI and KH performed experiments. SK and YK analyzed data. TY distributed materials. KO wrote paper and TM designed experiments, supervised the project and wrote the paper.

## References

- [1] Kieff, E. (1996) Epstein-Barr virus and its replication. *In Fields virology*: 3rd Ed, 2 (Field, D. N., Knipe, D. M., Howley, P. M., eds), 2343-2396., Raven Press, NY
- [2] Liebowitz, D., Wang, D., and Kieff, E. (1989) Orientation and patching of the latent infection membrane protein encoded by Epstein-Barr virus. *J. Virol.* 58, 233-237
- [3] Li, H.P., and Y.S. Chang. 2003. Epstein-Barr virus latent membrane protein 1: structure and functions. *J. Biomed. Sci.* 10, 490-504
- [4] Huen, D. S., Henderson, S. A., Croom-Carter, D., and Rowe, M. (1995) The Epstein-Barr virus latent membrane protein-1 (LMP1) mediates activation of NF-kappa B and cell surface phenotype via two effector regions in its carboxy-terminal cytoplasmic domain. *Oncogene* 10, 549-560
- [5] Gires, O., Kohlhuber, F., Kilger, E., Baumann, M., Kieser, A., Kaiser, C., Zeidler, R., Scheffer, B., Ueffing, M., and Hammerschmidt, W. (1999) Latent membrane protein 1 of Epstein-Barr virus interacts with JAK3 and activates STAT proteins. *EMBO. J.* 18, 3064-73
- [6] Brodeur, S. R., Cheng, G., Baltimore, D., and Thorley-Lawson, D. A. (1997) Localization of the major NF-kappaB-activating site and the sole TRAF3 binding site of

LMP-1 defines two distinct signaling motifs. *J. Biol. Chem.* 272, 19777-19784

[7] Devergne, O. E., Hatzivassiliou, K. M., Izumi, K. M., Kaye, K. M., Kleijnen, K. F., Kieff, E., and Mosialos, G. (1996) Association of TRAF1, TRAF2, and TRAF3 with an Epstein-Barr virus LMP1 domain important for B-lymphocyte transformation: role in NF-kappaB activation. *Mol Cell Biol.* 16, 7098-7108

[8] Kaye, K. M., Devergne, O., Harada, J. N., Izumi, K. M., Yalamanchili, R., Kieff, E., and Mosialos, G. (1996) Tumor necrosis factor receptor associated factor 2 is a mediator of NF-kappa B activation by latent infection membrane protein 1, the Epstein-Barr virus transforming protein. *Proc. Natl. Acad. Sci. USA.* 93, 11085-11090

[9] Mosialos, G., Birkenbach, M., Yalamanchili, R., VanArsdale, T., Ware, C., and Kieff, E. (1995) The Epstein-Barr virus transforming protein LMP1 engages signaling proteins for the tumor necrosis factor receptor family. *Cell* 80, 389-399

[10] Izumi, K. M., and Kieff, E. (1997) The Epstein-Barr virus oncogene product latent membrane protein 1 engages the tumor necrosis factor receptor-associated death domain protein to mediate B lymphocyte growth transformation and activate NF-kappaB. *Proc. Natl. Acad. Sci. USA.* 94, 12592-12597

[11] Izumi, K. M., Cahir-McFarland, E. D., Ting, A. T., Riley, E. A., Seed, B., and Kieff, E. (1999) The Epstein-Barr virus oncoprotein latent membrane protein 1 engages

the tumor necrosis factor receptor-associated proteins TRADD and receptor-interacting protein (RIP) but does not induce apoptosis or require RIP for NF-kappaB activation.

*Mol Cell Biol.* 19, 5759-5767

[12] Wan, J., Zhang, W., Wu, L., Bai, T., Zhang, M., Lo, K. W., Chui, Y. L., Cui, Y., Tao, Q., Yamamoto, M., Akira, S., and Wu, Z. (2006) BS69, a specific adaptor in the latent membrane protein 1-mediated c-Jun N-terminal kinase pathway. *Mol Cell Biol.*

26, 448-456

[13] Laherty, C. D., Hu, H. M., Opipari, A. W., Wang, F., and Dixit, V. M. (1992) The Epstein-Barr virus LMP1 gene product induces A20 zinc finger protein expression by activating nuclear factor kappa B. *J. Biol. Chem.* 267, 24157-24160

[14] Luftig, M., Yasui, T., Soni, V., Kang, M. S., Jacobson, N., Cahir-McFarland, E., Seed, B., and Kieff, E. (2004) Epstein-Barr virus latent infection membrane protein 1 TRAF-binding site induces NIK/IKK alpha-dependent noncanonical NF-kappaB activation. *Proc. Natl. Acad. Sci. USA.* 101, 141-146

[15] Eliopoulos, A. G., Gallagher, N. J., Blake, S. M., Dawson, C. W., and Young, L. S. (1999) Activation of the p38 mitogen-activated protein kinase pathway by Epstein-Barr virus-encoded latent membrane protein 1 coregulates interleukin-6 and interleukin-8 production. *J. Biol. Chem.* 274, 16085-16096

- [16] Eliopoulos, A. G., and Young, L. S. (1998) Activation of the cJun N-terminal kinase (JNK) pathway by the Epstein-Barr virus-encoded latent membrane protein 1 (LMP1). *Oncogene* 16, 1731-1742
- [17] Dawson, C.W., Tramontanis, G., Eliopoulos, A.G., and Young, L.S. (2003) Epstein-Barr virus latent membrane protein 1 (LMP1) activates the phosphatidylinositol 3-kinase/Akt pathway to promote cell survival and induce actin filament remodeling. *J. Biol. Chem.* 278, 3694-704
- [18] Mainou, B.A., Everly, D.N. Jr., and Raab-Traub, N. (2005) Epstein-Barr virus latent membrane protein 1 CTAR1 mediates rodent and human fibroblast transformation through activation of PI3K. *Oncogene* 24, 6917-24
- [19] Chen, H., Hutt-Fletcher, L., Cao, L., and Hayward, S.D. (2003) A positive autoregulatory loop of LMP1 expression and STAT activation in epithelial cells latently infected with Epstein-Barr virus. *J. Virol.* 77, 4139-48
- [20] Ikeda, O., Sekine, Y., Yasui, T., Oritani, K., Sugiyama, K., Muromoto, R., Ohbayashi, N., Yoshimura, A., and Matsuda, T. (2008) STAP-2 negatively regulates both canonical and non-canonical NF-kappaB activation induced by Epstein-Barr virus-derived LMP1. *Mol Cell Biol.* 28, 5027-5042
- [21] Ikeda, O., Sekine, Y., Mizushima, A., Oritani, K., Yasui, T., Fujimuro, M.,

Muromoto, R., Nanbo, A., and T. Matsuda. (2009) BS69 negatively regulates the canonical NF- $\kappa$ B activation induced by Epstein-Barr virus-derived LMP1. *FEBS Lett.* 583, 1567-1574

[22] Ikeda, O., Miyasaka, Y., Yoshida, R., Mizushima, A., Oritani, K., Sekine, Y., Kuroda, M., Yasui, T., Fujimuro, M., Muromoto, R., Nanbo, A., and T. Matsuda. (2010) BS69 cooperates with TRAF3 in the regulation of Epstein-Barr virus-derived LMP1/CTAR1-induced NF- $\kappa$ B activation. *FEBS Lett.* 584, 865-872

[23] Baichwal, V. R., Sugden, B. (1987) Posttranslational processing of an Epstein-Barr virus-encoded membrane protein expressed in cells transformed by Epstein-Barr virus. *J. Virol.* 61, 866-875

[24] Mann, K. P., and Thorley-Lawson, D. (1987) Posttranslational processing of the Epstein-Barr virus-encoded p63/LMP protein. *J. Virol.* 61, 2100-2108

[25] Martin, J., and Sugden, B. (1991) Transformation by the oncogenic latent membrane protein correlates with its rapid turnover, membrane localization, and cytoskeletal association. *J. Virol.* 65, 3246-3258

[26] Aviel, S., Winberg, G., Massucci, M., and Ciechanover, A. (2000) Degradation of the Epstein-Barr virus latent membrane protein 1 (LMP1) by the ubiquitin-proteasome pathway. Targeting via ubiquitination of the N-terminal residue. *J. Biol Chem.* 275, 23491-23499

- [27] Shair, K. H., Bendt, K. M., Edwards, R. H., Bedford, E. C., Nielsen, J. N., Raab-Traub N. (2007) EBV latent membrane protein 1 activates Akt, NFkappaB, and Stat3 in B cell lymphomas. *PLoS Pathog.* 3, e166
- [27] Muromoto, R., Nakao, K., Watanabe, T., Sato, N., Sekine, Y., Sugiyama, K., Oritani, K., Shimoda, K., and Matsuda, T. (2006) Physical and functional interactions between Daxx and STAT3. *Oncogene* 25, 2131-2136
- [28] Togi, S., Shiga, K., Muromoto, R., Kato, M., Souma, Y., Sekine, Y., Kon, S., Oritani, K., and Matsuda, T. (2013) Y14 positively regulates TNF- $\alpha$ -induced NF- $\kappa$ B transcriptional activity via interacting RIP1 and TRADD beyond an exon-junction complex protein. *J. Immunol.* 191, 1436-1444
- [29] Matsuda, T., Yamamoto, T., Muraguchi, A., and Saatcioglu, F. (2001) Cross-talk between TGF-beta and estrogen receptor signaling through Smad3. *J. Biol. Chem.* 276, 42907-42914
- [30] Sekine, Y., Yamamoto, C., M. Kakisaka, M., Muromoto, R., Kon, S., Ashitomi, D., Fujita, N., Yoshimura, A., K. Oritani, K., and T. Matsuda (2012) Signal-transducing adaptor protein-2 modulates Fas-mediated T-cell apoptosis by interacting with caspase-8 *J. Immunol.* 188, 6194-6204
- [31] Butel, J. S. (2000) Viral carcinogenesis: revelation of molecular mechanisms and etiology of human disease. *Carcinogenesis* 21, 405-426.

- [32] McLaughlin-Drubin, M. E., and Munger, K. (2009) Viruses associated with human cancer. *Biochim Biophys Acta*. 1782, 127-150.
- [33] Lamkanfi, M., and Dixit, V.M. (2010) Manipulation of host cell death pathways during microbial infections. *Cell. Host. Microbe*. 8, 44-54
- [34] Mocarski, E.S., Upton, J.W., and Kaiser, W.J. (2011) Viral infection and the evolution of caspase 8-regulated apoptotic and necrotic death pathways. *Nat. Rev. Immunol.* 12, 79-88
- [35] Scheller, C., Sopper, S., Chen, P., Flory, E., Koutsilieri, E., Racek, T., Ludwig, S., ter Meulen, V., and Jassoy, C. (2002) Caspase inhibition activates HIV in latently infected cells. Role of tumor necrosis factor receptor 1 and CD95. *J. Biol. Chem.* 277, 15459-64
- [36] Bentz, G.L., Whitehurst, C.B., and Pagano, J.S. (2011) Epstein-Barr virus latent membrane protein 1 (LMP1) C-terminal-activating region 3 contributes to LMP1-mediated cellular migration via its interaction with Ubc9. *J. Virol.* 85, 10144-53
- [37] Bentz, G.L., Shackelford, J., and Pagano, J.S. (2012) Epstein-Barr virus latent membrane protein 1 regulates the function of interferon regulatory factor 7 by inducing its sumoylation. *J. Virol.* 86, 12251-61

## Figure Legends

**FIGURE 1. LMP1 is degraded in a caspase-dependent manner in HeLa cells.** A, HeLa cells in a 24 well plate were transfected with FLAG-tagged LMP1 (500 ng). At 24 h after transfection, the cells were treated with cycloheximide (5  $\mu\text{g/ml}$ ) for the indicated periods. Cells were then lysed, and an aliquot of total cell lysate (TCL) (1%) was blotted with anti-FLAG or anti-actin antibody. B, Densitometric quantification of the above results was also shown. The relative intensity of FLAG-LMP1 was normalized to the actin protein of the same sample. Results are representative of three independent experiments, and the error bars represent the SD. \*,  $p<0.05$ . \*\*,  $p<0.01$ . C, HeLa cells in a 24 well plate were transfected with FLAG-tagged LMP1 (500 ng). At 12 h after transfection, the cells were treated with MG132 (10  $\mu\text{M}$ ), Z-VAD (10  $\mu\text{M}$ ) or  $\text{NH}_4\text{Cl}$  (10  $\mu\text{M}$ ) for an additional 12h, with or without cycloheximide (5  $\mu\text{g/ml}$ ). Cells were then lysed, and TCL (1%) was blotted with anti-FLAG or anti-actin antibody. D, Densitometric quantification of the above results was also shown. The relative intensity of FLAG-LMP1 was normalized to the actin protein of the same sample. Results are representative of three independent experiments, and the error bars represent the SD. \*\*,  $p<0.01$ , \*\*\*,  $p<0.001$ .

**FIGURE 2. Caspase-3 is responsible for LMP1 protein degradation in HeLa cells.**

A, HeLa cells in a 24 well plate were transfected with FLAG-tagged LMP1 (500 ng). At 12 h after transfection, the cells were treated with Ac-DNLD (10  $\mu$  M), Z-DEVD (10  $\mu$ M), Ac-VEID (10  $\mu$  M), Z-IETD (10  $\mu$ M), Ac-LEHD (10  $\mu$  M), or Z-VAD (10  $\mu$ M) for an additional 12h. Cells were then lysed, and an aliquot of total cell lysate (TCL) (1%) was blotted with anti-FLAG or anti-actin antibody. B, Densitometric quantification of the above results was also shown. The relative intensity of FLAG-LMP1 was normalized to the actin protein of the same sample. Results are representative of three independent experiments, and the error bars represent the SD. \*,  $p < 0.05$ . \*\*,  $p < 0.01$ . C, HeLa cells in a 24-well plate were transfected with FLAG-tagged LMP1 (500 ng) together with control, caspase-3, -7, or -8 siRNA (20 pmol). At 36 h after transfection, the cells were lysed, and TCL (1%) was blotted with anti-FLAG or anti-actin antibody. D, Densitometric quantification of the above results was also shown. The relative intensity of FLAG-LMP1 was normalized to the actin protein of the same sample. Results are representative of three independent experiments, and the error bars represent the SD. \*,  $p < 0.05$ . \*\*,  $p < 0.01$ .

**FIGURE 3. LMP1 expression induces caspase-3/8 activation in HeLa cells.** A, HeLa

cells in a 24 well plate were transfected with FLAG-tagged LMP1 or empty vector (500 ng). At 12 h after transfection, the cells were treated with DMSO or Z-VAD (10  $\mu$ M) for an additional 12h. Cells were then lysed and assessed for caspase-3/7 activities (Top panel). The results are indicated as -fold induction of caspase activity from triplicate experiments, and the error bars represent S.D. \*,  $p < 0.05$ . \*\*,  $p < 0.01$ . An aliquot of total cell lysate (TCL) (1%) was blotted with anti-FLAG or anti-actin antibody (Bottom panel). B, HeLa cells in a 24 well plate were transfected with FLAG-tagged LMP1 or empty vector (500 ng). At 24 h after transfection, the cells were lysed, and then TCL (1%) was blotted with anti- caspase-3, anti-caspase-8, anti-FLAG or anti-actin antibody.

**FIGURE 4. Caspase inhibition enhances LMP1-mediated signals in HeLa cells.** A, HeLa cells in a 24-well plate were transfected with NF- $\kappa$ B-LUC (250 ng), together with FLAG-tagged LMP1 or empty vector. At 12 h after transfection, the cells were treated with DMSO or Z-VAD (10  $\mu$ M) for an additional 12 h. And then the cells were harvested and assayed for luciferase activity. The results are indicated as -fold induction of luciferase activity from triplicate experiments, and the error bars represent S.D. \*\*,  $p < 0.01$ . An aliquot of total cell lysate (TCL) (1%) was blotted with an anti-FLAG or anti-actin antibody (IB). B, HeLa cells in a 24-well plate were transfected with


STAT3-LUC (250 ng), together with FLAG-tagged LMP1 or empty vector. At 12 h after transfection, the cells were treated with DMSO or Z-VAD (10  $\mu$ M) for an additional 12 h. And then the cells were harvested and assayed for luciferase activity as described the above. \*\*,  $p < 0.01$ . TCL (1%) was blotted with an anti-FLAG or anti-actin antibody. C, HeLa cells in a 24-well plate were transfected with FLAG-tagged LMP1 or empty vector. At 12 h after transfection, the cells were treated with DMSO or Z-VAD (10  $\mu$ M) for an additional 12 h. Total RNA samples isolated from these cells were also quantified by reverse transcription and quantitative real-time PCR analysis. Data represent the levels of *IL-6* mRNA normalized to that of a *ACTIN* internal control and are expressed relative to the value of control sample that was transfected with empty vector and treated with DMSO. The results are representative of three independent experiments, and the error bars represent the SD. \*\*,  $p < 0.01$ . D, HeLa cells in a 24-well plate were treated with DMSO or Z-VAD (10  $\mu$ M) for 12 h, followed by TNF- $\alpha$  (10 ng/ml) stimulation for 30 min. Total RNA samples isolated from these cells were also quantified by reverse transcription and quantitative real-time PCR analysis as described the above.

**FIGURE 5. Caspase-3 regulates LMP1-mediated NF- $\kappa$ B activation in HeLa cells. A,**

HeLa cells in a 24 well plate were transfected with NF- $\kappa$ B-LUC (250 ng), together with FLAG-tagged LMP1 or empty vector. At 12 h after transfection, the cells were transfected with control or caspase-3 siRNA (20 pmol). 36 h later, the cells were harvested and assayed for luciferase activity as described the above. \*\*,  $p < 0.01$ . An aliquot of total cell lysate (TCL) (1%) was blotted with an anti-FLAG, anti-caspase-3 or anti-actin antibody. B, HeLa cells in a 24 well plate were transfected with FLAG-tagged LMP1 or empty vector. At 12 h after transfection, the cells were transfected with control or caspase-3 siRNA (20 pmol). 36 h later, the cells were harvested and total RNA samples isolated from these cells were also quantified by reverse transcription and quantitative real-time PCR analysis as described the above.

**FIGURE 6. Determination of caspase cleavage sites in LMP1 in HeLa cells.** A, C, E, Domain structure of LMP1 and its mutant fragments are shown schematically. B, D, F, G, HeLa cells in a 24 well plate were transfected with FLAG-tagged LMP1 or its mutants (500 ng). At 12 h after transfection, the cells were treated with DMSO or Z-VAD (10  $\mu$ M) for an additional 12h. Cells were then lysed, and an aliquot of total cell lysate (TCL) (1%) was blotted with anti-FLAG or anti-actin antibody.

**Figure 1**



**Figure 2**




Figure 3






Figure 4




**Figure 5**




**Figure 6**


**A**




**B**




**C**




**D**



**E**



**F**



**G**

