Title	Impact of elevated CO ₂ on root traits of a sapling community of three birches and an oak : a free-air-CO ₂ enrichment (FACE) in northern Japan
Author(s)	Agathokleous, Evgenios; Watanabe, Makoto; Nakaji, Tatsuro; Wang, Xiaona; Satoh, Fuyuki; Koike, Takayoshi
Citation	Trees : structure and function, 30(2), 353-362 https://doi.org/10.1007/s00468-015-1272-6
Issue Date	2016-04
Doc URL	http://hdl.handle.net/2115/64926
Rights	The final publication is available at link.springer.com .
Type	article (author version)
File Information	Trees SF30-2_353-362.pdf

[Instructions for use](#)

**Impact of elevated CO₂ on root traits of a sapling community of three birches and an oak:
A free-air-CO₂ enrichment (FACE) in northern Japan**

Evgenios Agathokleous¹, Makoto Watanabe², Tatsuro Nakaji³, Xiaona Wang¹, Fuyuki Satoh³,
Takayoshi Koike^{1*}

¹Silviculture and Forest Ecological Studies, Hokkaido University, Sapporo, Hokkaido, 060-8589, Japan

²Institute of Agriculture, Tokyo University of Agriculture and Technology, Fuchu, Tokyo 183-8509, Japan

³Hokkaido University Forests, Sapporo, Hokkaido, 060-0809, Japan

*e-mail address: tkoike@for.agr.hokudai.ac.jp [T.K.]

ABSTRACT

We evaluated the root response to elevated CO₂ fumigation of 3 birches (*Betula* sp.) and 1 deciduous oak (*Quercus* sp.) grown in immature volcanic ash soil (VA) or brown forest soil (BF). VA is a nutrient-poor, phosphorus impoverished soil, broadly distributed in northern Japan. Each species had been exposed to either ambient (375-395 μmol mol⁻¹) (aCO₂) or elevated (500 μmol mol⁻¹) (eCO₂) CO₂ during the daytime (more than 70 μmol m⁻²s⁻¹) over 4 growing seasons. The results suggest that eCO₂ did not cause an increase in total root production when the community had grown in fertile BF soil, however it did cause a large increase when the community was grown in infertile VA soil. Yet, carbon allocation to plant roots was not affected by eCO₂ in either the BF and VA soils. Rhizo-morphogenesis appeared to occur to a greater extent under eCO₂. It seems that the saplings developed a massive amount of fine roots under the VA and eCO₂ conditions. Unexpectedly, eCO₂ resulted in a larger total root mass when the community was grown in VA soil than when grown in BF soil (eCO₂×VA vs. eCO₂×BF). These results may hint to a site-specific potential of communities to sequester future atmospheric carbon. The

growing substance of plants is an important factor which root response to eCO₂ depends on, however further studies are needed for a better understanding.

Key words: birch, elevated CO₂, FACE, oak, root, volcanic ash soil

Key message: The CO₂ effect on the root production of a broadleaved community was insignificant when grown in brown forest soil, however it was positively large when grown in volcanic ash soil.

Author contribution statement:

Evgenios Agathokleous: Root measurements, data analysis and interpretation, and synthesis of the manuscript

Makoto Watanabe: Data collection and article discussion

Tatsuro Nakaji: Guidance on the root research and article discussion

Xiaona Wang: Article discussion

Fuyuki Satoh: Management of the FACE system

Takayoshi Koike: Funds, excavation of roots, collection of data and management of all technical procedures

INTRODUCTION

Roots are the hidden half of a plant and account for a high proportion of the biomass production (Eshel and Beeckman 2013). They also act as an interface between soil and the plant body (e.g. Fitter and Hay 2002). Most roots of woody plants have high plasticity to soil properties, such as soil moisture content (e.g. Koike et al. 2003), nitrogen availability (Qu et al. 2003), temperature of rhizosphere (Korotkii et al. 2002; Qu et al. 2009; Kajimoto et al. 2010) and other factors. The

growth dynamic of roots is strongly affected by atmospheric conditions (Wang et al. 2015a), as in the case of the Aspen FACE (Pregitzer and Tlhelm 2013). CO₂ levels have already hit a record of ~400 $\mu\text{mol mol}^{-1}$ (NOAA 2014), and are projected to rise to 500 $\mu\text{mol mol}^{-1}$ with the possibility to elevate even further until 2040 (IPCC 2007). It is known that elevated CO₂ (eCO₂) alters above- and below-ground functions of plants (Koike et al. 1995; Beerling 1996; Karnosky 2003; Ozanne et al. 2003; Heath et al. 2005; Cheng et al. 2012; Slattery et al. 2013; Guo et al. 2015; Koike et al. 2015).

Soil fertility is one of the factors affecting the biomass production of the loblolly pine (Oren et al. 2001) and several other tree species (e.g. Norby and Zak 2011; Pregitzer and Tlhelm 2013) under elevated CO₂ conditions. According to estimations, more than 50% of the global potential arable lands are acidic (von Uexkull and Mutert 1995) and usually P-deficient (Zheng 2010; Cordell and Neset 2014; Reijnders 2014; Ulrich and Frossard 2014). Specifically, the pH in Japanese forests ranges from 4.8 to 5.2, except for red soil and podzolic soil (e.g. Hashimoto et al. 2012). Soils of Northern Japan are partly dominated by nutrient-poor immature volcanic ash or pumice soils, where phosphorous (P) is usually impoverished (e.g. Eguchi et al. 2005; Kim et al. 2010; Hashimoto et al. 2012). Many plants grown under such nutrient limited conditions are usually suffering from nutrient deficiency, and are consequently forced to develop fine roots (Hermans et al. 2006) in order to establish symbioses with ectomycorrhizae (e.g. Smith and Read 2008; Qu et al. 2010).

In general, the allocation pattern of photosynthates to above- and below-ground differs broadly among deciduous tree species (e.g. Schulze et al. 2005). This pattern strongly depends on the successional traits of tree species; for example, early successional species (i.e. birch) versus mid-late successional species (i.e. oak) (Sato 1995). Among birches, this tendency is

typically found in mountain birch (Koike et al. 2003). The majority of the early successional species, such as birch, grow above- and belowground simultaneously. Yet, late successional species, usually, first allocate photosynthates to their roots which store them, and then prepare for quick growth in the succeeding year (e.g. Sato 1995). Consequently, these woody plants flourish aboveground in the early part of the growing season and then gradually develop their belowground structure. Under elevated CO₂, root growth is expected to increase, boosting the absorption of mineral nutrients and water which support an increased aboveground body - if increased.

A review of current literature revealed that several studies have dealt with responses of individual species to elevated CO₂, while other studies tested the responses of communities to elevated CO₂. However, there is a lack of evidence on the response of communities to elevated CO₂ as a whole unit (e.g. Pregitzer et al. 2008).

We aimed to test if and how elevated CO₂ and soil infertility could impact the belowground functioning of a community of trees in terms of carbon (C) allocation within plant body and atmospheric C sequestration capacity. To this purpose, we established a free-air-CO₂ enrichment (FACE) system with 4 deciduous, broad-leaved species, which are known to be physiologically (aboveground) affected by elevated CO₂ (Koike et al. 2015). We hypothesized that the root impact of elevated CO₂ on the communities would depend on soil nutrient status due to shifts in the photosynthetic function and the nutritive demands. We were expecting the soil nutrient deficiency to induce a differential impact of elevated CO₂ on the belowground functioning of the community.

In this study, we focused on a community of species – consisting of 2 ecological groups (3 early successional: birch, and 1 late successional: oak) - because these broad-leaved trees play an integral role in the above-mentioned forest functions (Jensen and Anderson 1995).

MATERIALS & METHODS

Experimental Design: This study was conducted in the FACE system located in the Sapporo Experimental Forest of Hokkaido University, Japan (43°06' N, 141°20' E, 60 m a.s.l.). The main-plot treatment consisted of ambient (375–395 $\mu\text{mol mol}^{-1}$) and elevated (500 $\mu\text{mol mol}^{-1}$) CO₂ treatments, with three site replications for each treatment (Filion et al. 2000). The sub-plot treatments were brown forest (BF) soil, and BF mixed with immature volcanic ash plus pumice (VA) soil. The latter is nutrient poor soil (Masyagina et al. 2006) which was excavated and brought from Tomakomai Experimental Forest of Hokkaido University (42°40' N, 141°37' E, 30 m a.s.l.). It is widespread in Hokkaido and ideal for experiments mimicking soil-nutrient deficiency scenarios.

Since BF soil is native to Sapporo Experimental Forest, half of each FACE ring was excavated and refilled with a mixture of 65 % BF soil and 35 % VA soil. Chemical and physical properties of BF and VA soils used in the present study are illustrated in Eguchi et al. (2008) and Watanabe et al. (2013). The VA soil had about 17 % lower P₂O₅ content than the BF soil.

Plant Materials: The materials we used included saplings of 3 birch species (*Betula ermanii* (Be: mountain birch), (*B. maximowicziana* (Bm: Monarch birch), *B. platyphylla* var. *japonica* (Bp: white birch)), and a deciduous oak (*Quercus mongolica* var. *crispula* (Qm: oak). All the saplings were obtained from Hokkaido Hort-green Co. Ltd. near Sapporo, and planted in the

FACE rings with a distance of 50 cm among them, in 2010 (after the snow had melted). Equal numbers of individuals among species were planted in each research condition.

The mean values ($\pm$ SD) of the height and trunk basal diameter (at 2 cm from the starting point of root system) of the juveniles were measured for the 3 birches and the Oak; the variability between the two gas treatments within each species was minimal (Table 1). The oak had been suppressed by the birches (i.e. restricted relative light conditions) after 1.5 years with CO₂ fumigation (data not shown). Similarly, data from leaf area per ground area have shown that the mountain birch was suppressed by the white and monarch birches (Hara et al. 2014). We assumed that it was a phenomenon that could have occurred in pristine conditions and thus can be considered as a realistic growth of a community. Nonetheless, the obtained growth data could have been different if the trees had grown independently or in an uncompetitive environment.

CO₂ treatment: Fumigation of CO₂ was carried out in 4 consecutive Julian years (2010-2013). Saplings were fumigated in each growing season during daytime, when the photosynthetic photon flux (PPF) exceeded the 70 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (i.e. light compensation point of photosynthesis; Koike 1988), from leaf emergence to leaf senescence (from May until late November). This FACE system consists of 6 rings, 3 of which were fumigated by additional CO₂ in order to reach the targeted concentration of 500 $\mu\text{mol mol}^{-1}$ (hereafter “eCO₂”). The diameter and height of each FACE ring are 5 m. The 3 ring plots remained under ambient CO₂ (ca. 370-390 $\mu\text{mol mol}^{-1}$) (hereafter “aCO₂”) as noted in researches by Eguchi et al. (2008) and Watanabe et al. (2013).

Climatic & meteorological conditions: The snow-free period lasts usually from early April to early November. The monthly mean air temperature, wind speed, and relative humidity and the monthly total sunshine duration and precipitation were averaged per year, and the means for the

entire experiment duration were calculated. The mean values for the years 2010-2013 were 9.40 (± 0.27 CI) °C, 3.45 (± 0.13 CI) m s⁻¹, 69.75 (± 0.94 CI) %, 1687 (± 126 CI) h, and 1301 (± 42 CI) mm, for each variable, respectively. The meteorological data were collected from a station located in Sapporo (WMO, ID: 47412) at 43°03.6'N 141°19.7'E (Japan Meteorological Agency, 2015).

Sampling and measurements: At the end of the final growing season (2013), the trunk basal diameter was measured for each grown sapling, and all the roots were excavated using a small bulldozer. This mechanistic method was previously compared with the manual method (by hand) and it was revealed that excavation by this method causes destruction of the roots during the excavation process. Although the target species were different from the present 4 species, the final excavated root system had less mass; approximately 70~90 % of the whole roots (Matsunami 2008). Nevertheless, we assume there could be equal amounts of error across all the subjects.

After the excavation, the following procedure was carried out: a) the roots of small classes were sampled, and dried at 75 °C, for more than 5 days, to constant mass. The dry mass of intermediate roots ($\varnothing$ 2 - 4 mm) and fine roots ($\varnothing$ < 2 mm) – including hyphae of ectomycorrhizae – were determined; b) the whole root systems were left on the field to physically dry; this was unavoidable because of the large size of the root systems. The next summer, dry mass measurements were taken for the total root systems (TDM). Due to harsh natural conditions, the identities of some root systems were lost, therefore these samples were not measured. Nonetheless, the sample size was sufficient to perform parametric tests.

Statistics: The accepted threshold of significance was defined at $\alpha=0.05$. Trunk basal diameter data were first converted to trunk basal area (Area), and then the Trunk-basal-area:Total-root-dry-mass rate (Area:TDM) was calculated. Area:TDM is a significant indicator of the plant health, and is based on the Pipe Model theory of tree form (Shinozaki et al. 1964a, 1964b). Prior to the analysis, the data of each variable were standardized (Saitanis et al. 2014, 2015). For the standardization, T-scores ($T = z - score \times 10 + 50$) were used, with a mean equal to 50 and a standard deviation equal to 10. Therefore, the mean T-score of each species in each soil and each experimental unit, constituted the real-replicates in the overall analysis. As previously suggested by Saitanis et al. (2014, 2015), this process also allows a fair intercomparison of measured variables and removes any heterogeneity. In order to utilize the overall information and generalize the results at a community level, two steps were followed: First, in relation to the biomasses, data for each community (including all the species) in each treatment (i.e. soil and CO₂) and block (FACE ring) were calculated by averaging the values of individuals of each species in each treatment and then summing the returned values for all the species in each treatment (this process treats any differences in ratios among species and gives better estimates). In order to increase accuracy of the ratio and rates, the values for each individual were calculated for each separately, and averaged per species in each of the research conditions for each block. The values were averaged again for all the species (community) in each of the research conditions for each block.

Second, the data from all the experimental units were pooled (resulting to n=3) and analyzed. Prior to this analysis, the data were analyzed using individual species in order to test for interactions with species; there were no significant ($p>0.05$) species-specific differences. All the data were subjected to split-plot general linear model randomized by block (GLM), based on

Kuehl (1999), and, if there were significant effects (only for factors with more than two levels), a Tukey range post-hoc test was performed. For the presentation of the data, the non-standardized values were used.

Finally, in order to quantify the size of the difference between the community of plants grown under ambient CO₂ and elevated CO₂, under the different soils, and for each dependent variable (Table 2), the effect size *ESi* was calculated according to Cohen (1988). The formula was:

$$ESi = \left(\frac{[C_{i(EA)} - C_{i(CTRL)}]}{\sqrt{[(n_{i(EA)} - 1) * (SD_{i(EA)})^2 + (n_{i(CTRL)} - 1) * (SD_{i(CTRL)})^2] / (n_{i(EA)} + n_{i(CTRL)} - 2)}} \right),$$

C_i , n_i , SD_i , stands for the mean score, the number of observations, and the standard deviation respectively, in the plants under CO₂ enriched air (*EA*) or ambient air (*CTRL*; control group), of the *i* case. The denominator stands for the pooled standard deviation. Absolute values of the effect size within the arbitrary segments 0.00-0.20, 0.20-0.50, 0.50-0.80, and 0.80+ indicate neutral, small, moderate, and large effects, respectively. *ES* was calculated only for the pairs with statistically significant difference in order to quantify the size of the difference. Data processing and statistical analysis were conducted using MS EXCEL 2010 (© Microsoft) and STATISTICA v.10 (© StatSoft Inc.) software.

RESULTS

CO₂ appeared to have a large main effect on all variables with the exception of Area:TDM. Besides, soil had only a large effect on Fine:Area and a medium effect on fine roots (Tables 2 and 3). Concerning the CO₂ × soil interactions, only for Area:TDM and intermediate roots were not significant.

Regarding the Area, eCO₂ was responsible for a large effect, namely a 17 % smaller Area, cf. aCO₂. There was a convergent interaction in Area due to a higher (ns) Area of the communities under eCO₂×VA, compared to eCO₂×BF, and a 22 % significantly smaller (*ES* = 3.69) Area of the community grown under aCO₂×VA, compared to aCO₂×BF (Table 3). Under eCO₂×VA, there was not such a reduction. Instead, eCO₂×VA led to a 32 % larger (*ES* = 6.97) Area than that of plants in aCO₂×VA. There was no significant effect of CO₂ on the communities grown in BF soil.

Similarly, the large 20 % greater TDM induced by eCO₂ can be attributed to the communities grown in VA soil and not in BF soil. Under VA soil and eCO₂, the TDM was 35 % higher (*ES* = 4.65) than under aCO₂×VA, and, unexpectedly, it was also 25 % higher than that under eCO₂×BF (*ES* = 3.55).

Likely to TDM, the coarse root biomass was largely boosted (19 %) by eCO₂, regardless of the soil treatment. This caused by a large increase of coarse root biomass in eCO₂×VA, which also led to a divergent interaction. Under eCO₂×VA, coarse root biomass was largely higher than under aCO₂×VA (35 %) and eCO₂×VA (25 %). It is revealed that there was no effect of eCO₂ when the communities were grown in BF soil. On the other hand, none of the factors had a significant effect on Area:TDM (Table 2).

Fine root biomass was 27 % superior (*ES* = 2.90) in eCO₂ than in aCO₂, as a main effect. Furthermore, it was 11 % suppressed (*ES* = 0.63) by VA soil, compared to BF soil. The interaction of the two factors was significant and could be characterized as divergent with differences in sizes as well. When the communities had grown in BF and VA soils, eCO₂ caused large increases of fine root biomass by 14 and 39 % respectively. The moderate main effect of

the VA soil was produced by a 27 % lower ($ES = 5.09$) fine root biomass under $aCO_2 \times VA$, compared to $aCO_2 \times BF$.

Intermediate root biomass experienced a large positive effect (+32 %) by eCO_2 , independent of soil treatment, however it was not affected by the main effect of soil. There was also no interaction between the 2 factors.

As a result, the Fine: Intermediate root dry mass ratio (Fine:Intermediate) was largely negatively affected (-10 %) by eCO_2 as a main effect, and there was a significant interaction with soil that tended to mimic a directional interaction. It seems that both the eCO_2 and VA soil alone or in combination caused a higher increase of intermediate root than fine root biomass. Communities under $aCO_2 \times BF$ had 22, 21, and 14 % higher Fine:Intermediate than under $eCO_2 \times BF$, $aCO_2 \times VA$, and $eCO_2 \times VA$, respectively. Among the latter three treatments, there was no statistical difference.

Finally, Fine:Area was largely increased by eCO_2 (27 %) and VA (42 %) factors, regardless of each other factor. The factor of the interaction of these 2 variables was also significant. eCO_2 had no significant effect on the Fine:Area of the community grown in BF, however it had a large effect (+34 %) on the Fine:Area of the community grown in VA.

DISCUSSION

We predicted that root growth and root architecture of the saplings would be affected by eCO_2 and soil infertility; thus, the response of the community to eCO_2 would depend on the soil fertility and species seral traits. Based on the results, and in contrast to our expectations, this 4-year fumigation with eCO_2 resulted in neither a greater Area nor an increased TDM (or coarse

root biomass) when the community had grown in the fertile BF soil. eCO₂ also drastically increased Area, coarse root biomass and TDM when the community had grown in the infertile VA soil. Moreover, no specific seral traits were found in final production of root biomass (p>0.05; data not shown), which may be due to only one final assessment of the production of root biomass. In addition, root growth may be restricted after canopy closure (Sato 1995; Hara et al. 2014; Wang et al. 2015b).

Area:TDM and TDM were assessed in order to evaluate the root responses to eCO₂. The former is involved in the tree functional traits, i.e. C allocation, and the latter is important for C sequestration capacity of the ground. Although VA suppressed the Area, it did not suppress the TDM under aCO₂. Meanwhile, it appeared that eCO₂ had a surprisingly large hostile effect on the VA soil, and thus the negative effects of the VA soil on Area were compensated. Although neither eCO₂ nor VA soil affected the Area:TDM, VA soil in combination with eCO₂ led to a marginally (p>0.05) higher belowground allocation than aboveground (i.e., trunk area) compared to aCO₂×BF (-14%), aCO₂×VA (-9%) and eCO₂×VA (-11%). Despite this, under ambient CO₂, soil infertile conditions are usually expected to alter the C partitioning to favor root growth (Hermans et al. 2006), yet this was not the case under aCO₂.

Our results agree with the findings of quantitative reviews which state that, on average, eCO₂ does not alter the allometry of plants (Poorter and Nagel 2000; Poorter et al. 2012). However, the results of our statistical analyses differ to the general finding which notes that soil infertility does alter the allometry (Poorter and Nagel 2000; Poorter et al. 2012) as explained by the *functional equilibrium* theory (Brouwer 1962; Poorter and Nagel 2000). Our findings pointed out that C allocation within a plant body was not altered in this community of trees by a long-term experimental fumigation with eCO₂, while the sequestration of atmospheric C, in eCO₂,

increased only in the community grown in VA soil but not BF, which may be due to physical structure of BF (e.g. Eguchi et al. 2005).

The usual response of trees to elevated CO₂ is an increased root growth and root turnover of fine roots (e.g. Karnosky 2003; Pregitzer et al. 2008; de Oliveira et al. 2012; King et al. 2013; Cao et al. 2008). Our results support this response with the large effect of eCO₂ on TDM when the community was grown in VA soil. On the other hand, there was no effect of eCO₂ when the community was grown in BF. Insignificant increases of root biomass due to 2-growing-season fumigation with eCO₂ were also found in *Larix gmelinii* var. *japonica* × *L. kaempferi* seedlings grown in well-homogenized BF soil in FACE (Watanabe et al. 2013) or OTCs (Wang et al. 2015a), in the same region. Furthermore, short-term fumigation with elevated CO₂ had no significant effects on the root biomass of neither the *Betula papyrifera* seedlings grown in slight fertilization (Zhang et al. 2006) nor the *Fagus crenata* seedlings (Hirano et al. 2012). In general, root longevity is positively correlated with poor soil-nutrient status and well aeration of soils (Fitter and Hay 2002), which in our case may be attributed to high porosity of VA soil with pumice (e.g. Eguchi et al. 2005; Masyagina et al. 2006).

The eCO₂ increased fine roots, of which the majority naturally colonizes A horizon (e.g. Jaloviar et al. 2009) in both the BF and VA soils. It seems fine roots are ecologically important components as they account for some 13 % of net primary production (NPP) (Ostonen et al. 2005) and a high proportion of total litter (e.g. Leppalammi-Kujansuu et al. 2014). Thus, increased production of fine roots may have an ecologically important benefit in nutrient cycling in a forest. As observed in the Area, although VA soil unexpectedly suppressed fine roots under aCO₂, with the interaction of VA×eCO₂, the fine roots still reached levels similar to that of BF×eCO₂. We expected that under the influence of either eCO₂ or VA soil, the saplings would

undergo a higher need to seek nutrients thereby increasing the fine root area/biomass in order to access more nutrients. Instead, this was the case only for eCO₂ since VA soil caused a reduction of fine root biomass, until the canopy closed (Hara et al. 2014). At present, however, we cannot give a plausible reason for this phenomenon.

There was an unexplained decrease in fine root biomass in VA soil under aCO₂. Yet the Fine:Area, which is an appropriate index to assess the belowground responses in relation to nutrient cycling, showed something different. Namely, VA soil induced a large ($ES = 3.54$) increase when the community had been exposed to aCO₂ (cf. aCO₂×BF). Among the interaction levels, the largest Fine:Area was observed in eCO₂×VA: it was 34 % higher than aCO₂×VA and 49 % ($ES = 8.13$) higher than eCO₂×BF. A different foraging strategy of fine roots among the research conditions is clear, in accordance with the findings of Ostonen et al. (2011) and Leppalammi-Kujansuu et al. (2014). Particularly, fine roots underwent extensive foraging strategy (Ostonen et al. 2011), i.e. higher Fine:Area, in VA soil, in both CO₂ conditions. Moreover, eCO₂ had a synergistic effect with VA soil leading to the highest Fine:Area. These results coincide with the results of Area:TDM, albeit insignificant, and indicating greater responses to eCO₂ in relation to nutrient cycling when the community was grown in VA soil.

Differences of intermediate root biomass were attributed only to eCO₂ which increased the biomasses of the communities grown in both the BF and VA soils, compared to the pairwise aCO₂. Consequently, there was a higher amount of intermediate roots than fine roots both in VA soil and eCO₂. The ratio of fine roots to intermediate roots indicates the balance of the two root classes, and its modification is dependent on the needs of the plants. For instance, plants in the need of increasing nutrient uptake are expected to modify a higher proportion of fine roots compared to intermediate roots. However, this could be misleading in such cases which take into

account single observations of only standing biomasses. A single observation, as in our case, of the living fine roots, does not stand for the total production of fine roots, but only a partial production of that given time. In our study, this ratio should be taken with a degree of caution as fine roots were increased by eCO₂ as well. One explanation for the higher amount of intermediate roots than fine roots could be the mortality and regeneration of fine roots that occur (Pritchard et al. 2008) more frequently throughout the growing season which deposit a significant amount of C. As such, we can only speculate that there was an accelerated turnover of fine roots in all research conditions except aCO₂×BF.

Despite our expectations that soil infertility might lead to a maximum fine root production independent of the CO₂ treatment, CO₂ treatment did appear to have a combination effect with soil infertility, resulting in higher production of fine roots. Could this phenomenon be an ecological growth method of saplings in order to mitigate the negative effects via plasticity? In other terms, this phenomenon could be a kind of “*rhizo-morphogenesis*” as the root architecture is changing under these external stressors. This might be induced by a better communication between above- and below-ground under eCO₂, thus resulting in mitigation of the VA soil separated effects.

In general, eCO₂ increases root biomass and the demand for N, P, and K (Norby et al. 1986; Li et al. 2013), however, even if the concentrations of elements in the soil are adequate, a proportionally smaller root system accompanied by less microbial activity is quite hard to support the dissimilation organs. On the other hand, eCO₂ acts as an external force for plants in order to extract more nutrients from the soil: eCO₂ increased NaHCO₃- and NaOH-extractable organic P in the rhizosphere of wheat by 160 and 53 %, respectively (Jin et al. 2014).

It seems that the growing substance of plants is an important factor to which the plant response to CO₂ usually depends on (e.g. Conroy et al. 1990), and makes it difficult to draw general conclusions. It is also critical that nutrient dynamics between different elements may change under elevated CO₂ (Cong et al. 2015; Li et al. 2015; Wang et al. 2015a). However, the responses are still not well understood since many factors such as soil temperature, light condition, nitrogen deposition, O₃, etc. (Cao et al. 2008; Danyagri et al. 2014; Cong et al. 2015; Wang et al. 2015a) may confound the nutrient allocation under elevated CO₂.

We should mention that increased microbial activity contributes to P and root-derived materials immobilization in rhizosphere under elevated CO₂ (Jin et al. 2014; Wang et al. 2015a). Considering fine root dynamics, we should include the interaction of soil microbial activity (e.g. ectomycorrhizae) that could be a critical and determinant factor in the modulation of nutritional starvation and elevated CO₂ interaction. Unfortunately, we did not find any significant species-specific difference in root biomasses and structure of fine roots. In order to understand this, we should trace the seasonal trends of above- and belowground (Sato 1995). Moreover, future studies should consider further investigations on soil microbial activity under the interaction of nutrient-poor soils and elevated CO₂.

Summarizing, in agreement with similar studies (e.g. de Oliveira et al. 2012), our results show that soil type has the potential to modify the response of trees to eCO₂. However, in contrast to our expectations, the data support the hypothesis that the root growth of the community in BF soil was not altered by eCO₂ while in the infertile VA soil it was positively altered. Elevated CO₂ led to largely higher coarse root biomass (*ES* = 3.55), TDM (*ES* = 4.08) and Fine:Area (*ES* = 8.13) when the community was grown in VA soil than when the community was grown in BF soil. This indicates the magnitude of response varies considerably between

communities growing in different regions and that models dealing with the estimation of C allocation and storage under elevated CO₂ scenarios may overestimate the potential in some regions and underestimate it in others.

CONCLUSIONS

Elevated CO₂ did not cause an increase in trunk basal area or total root mass when the community had grown in fertile BF soil after canopy closure, but it did cause a large increase in both characteristics when the community had grown in infertile VA soil. Furthermore, eCO₂ caused an evident rhizo-morphogenesis through a large increase in fine root production in both soils. On the other hand, eCO₂ had no effect on C allocation to roots even in the case where eCO₂ increased Area and TDM in VA soil. Similarly, VA soil did not cause any significant disturbance to C allocation under ambient or elevated CO₂, albeit there was an insignificant trend for higher allocation to belowground when combined with eCO₂. Nevertheless, our results do not support the drawing of general conclusions because the adaptability of plants to VA soil as well as to eCO₂ may differ among species of different growth traits and soil physicochemical structure.

We conclude that eCO₂ fumigation of 4 growing seasons has no potential to alter the total root production of a community of the 4 deciduous broad-leaved trees when it is growing in brown forest soil. Tree species growing in less fertile soils, such as VA with high porosity, may respond more vigorously to eCO₂ if they are well adapted to the single effects of such soils. New directions should also deal with more infertile soil than the one used in our study. Moreover, ectomycorrhizae research under similar conditions would be essential as their symbiosis may directly or indirectly help plants to mediate the VA- or CO₂-induced suppressing stressors.

ACKNOWLEDGEMENTS

The authors appreciate Mr. Tatsushiro Ueda of Dalton Co. (Hokkaido Branch) for his continuous engineering assistance in the FACE system, as well as Mr. K. Ichikawa of Hokkaido University Forests for the operation of the bulldozer. The authors also recognize Dr. Rhett Loban of the Queensland University of Technology, Australia, for proofreading the manuscript. Two anonymous reviewers are acknowledged for their positive comments, especially related to above- and below-ground balance. E.A. extends his sincere thanks to JSPS for funding (scholarship no: 140539). This study was financially supported by JSPS via research funds to T.K. (Innovation type: 21114008, Type B: 26292075).

Conflict of interest: The financial source support of this study (Japan Society for the Promotion of Science) is a non-profit organization. We declare that our research has no conflict of interest.

LITERATURE CITED

- Beerling D J (1996) Ecophysiological responses of woody plants to past CO₂ concentrations. *Tree Physiol* Netherlands
- Brouwer R (1962) Distribution of dry matter in the plant. *Neth J Agr Sci* 10: 399–408
- Cao et al (2008) Effects of [CO₂] and nitrogen on morphological and biomass traits of white birch (*Betula papyrifera*) seedlings. *Forest Ecol Manag* 254: 217–224
- Cheng L, Booker F L, Tu C, et al (2012) Arbuscular mycorrhizal fungi increase organic carbon decomposition under elevated CO₂. *Science* 337: 1084–1087
- Cohen J (1988) Statistical power analysis for the behavioral sciences, 2nd edition. Lawrence Erlbaum Associates, USA, pp. 590

395 Cong S, Kitao M, Agathokleous E, et al. (2015) Chemical composition of two oak seedlings
 396 grown under elevated O₃ and CO₂ with a free-air fumigation system. J Agr Meteo. Accepted

397 Conroy J P, Milham P J, Bevege D I, Barlow E W R (1990) Influence of phosphorus deficiency
 398 on the growth response of four families of *Pinus radiata* seedlings to CO₂-enriched atmospheres.
 399 Forest Ecol Manag 30: 175-188

400 Cordell D, Neset T-SS (2014) Phosphorus vulnerability: a qualitative framework for assessing
 401 the vulnerability of national and regional food systems to the multidimensional stressors of
 402 phosphorus scarcity. Global Environ Chang 24: 108–122

403 Danyagri G, Dang Q-L (2014) Effects of elevated [CO₂] and soil temperature on photosynthetic
 404 responses of mountain maple (*Acer spicatum* L.) seedlings to light. Environ Exp Bot 107: 64–70

405 De Oliveira E A D, Approbato A U, Legracie J R Jr, Martinez C A (2012) Soil-nutrient
 406 availability modifies the response of young pioneer and late successional trees to elevated carbon
 407 dioxide in a Brazilian tropical environment. Environ Exp Bot 77: 53–62

408 Eguchi N, Funada R, Ueda T, et al (2005) Soil moisture condition and growth of deciduous tree
 409 seedlings native to northern Japan grown under elevated CO₂ with a FACE system. Phytom
 410 (Austria) 45: 133-138

411 Eguchi N, Karatsu K, Ueda T, Funada R, et al (2008) Photosynthetic responses of birch and alder
 412 saplings grown in a free air CO₂ enrichment system in northern Japan. Trees 22: 437-447

413 Eshel A, Beeckman T (2013) Plant Roots: The Hidden Half. 4th edition. CRC Press, Boca
 414 Raton. 784 pp.

415 Filion M, Dutilleul P, Potvin C (2000) Optimum experimental design for Free-Air-Carbon
 416 dioxide Enrichment (FACE) studies. *Glob Change Biol* 6: 843-854

417 Fitter A H, Hay R K M (2002) *Environmental physiology of plants*, 3rd edition. Academic Press,
 418 San Diego, CA. 367 pp.

419 Guo B, Dai S, Wang R, Guo J, Ding Y, Xu Y (2015) Combined effects of elevated CO₂ and Cd-
 420 contaminated soil on the growth, gas exchange, antioxidant defense, and Cd accumulation of
 421 poplars and willows. *Environ Exp Bot* 115: 1-10

422 Hara Y, Watanabe M, Takagi K, Sato F, Mao Q Z, Koike T (2014) Dynamics of Leaf Area Index
 423 of 3 kinds of birch species grown under free air CO₂ enrichment (FACE) system –Temporal
 424 changes of CO₂ exposure for 4 years. *Boreal For Res* 62: 57-58 (in Japanese)

425 Hashimoto S, Ugawa S, Morisada K, Wattenbach M, Smith P, Matsuura Y (2012) Potential
 426 carbon stock in Japanese forest soils – simulated impact of forest management and climate
 427 change using the CENTURY model. *Soil Use Manage* 28: 45-53

428 Heath J, Ayres E, Possell M, Bardgett R D, et al. (2005) Rising atmospheric CO₂ reduces
 429 sequestration of root-derived soil carbon. *Science* 309: 1711

430 Hermans C, Hammond P J, White J P, Verbruggen N (2006) How do plants respond to nutrient
 431 shortage by biomass allocation? *Trends Plant Sci* 11: 610-617

432 Hirano A, Hongo I, Koike T (2012) Morphological and physiological responses of Siebold's
 433 beech (*Fagus crenata*) seedlings grown under CO₂ concentrations ranging from pre-industrial to
 434 expected future levels. *Landscape Ecol Eng* 8: 59–67

435 IPCC (Intergovernmental Panel on Climate Change) (2007) *Climate Change 2007: Synthesis*
 436 *Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the*
 437 *Intergovernmental Panel on Climate Change* (ed Pachauri RK, Reisinger A). Geneva:IPCC
 438 Jaloviar P, Bakosova L, Kucbel S, Vencurik K (2009) Quantity and distribution of fine root
 439 biomass in the intermediate stage of beech virgin forest Badínsky prales. J Forest Sci 55: 502-
 440 510
 441 Japan Meteorological Agency (2012) <http://www.jma.go.jp/jma/indexe.html> Website accessed
 442 on 29th January 2015
 443 Jensen E C, Anderson D J (1995) The reproductive ecology of broadleaved trees and shrubs: an
 444 overview. Forest Research Laboratory, Oregon State University, Corvallis. Research Publication
 445 9a. 10 p.
 446 Jin J, Tang C, Robertson A, et al (2014) Increased microbial activity contributes to phosphorus
 447 immobilization in the rhizosphere of wheat under elevated CO₂. Soil Biol Biochem 75: 292-299
 448 Kajimoto T (2010) Root system development of larch trees growing on Siberian permafrost. *In*:
 449 Osawa A et al (eds.) Permafrost Ecosystems. Springer Ecol Studies 209: 303-330
 450 Karnosky D F (2003) Impacts of elevated atmospheric CO₂ on forest trees and forest ecosystems:
 451 knowledge gaps. Environ Int 29: 161-169
 452 Kim YS (2013) Soil-Atmosphere Exchange of CO₂, CH₄ and N₂O in Northern Temperate
 453 Forests: Effects of Elevated CO₂ Concentration, N Deposition and Forest Fire. Eurasian J For
 454 Res 16: 1-43

455 King J, Liu L, Aspinwall M (2013) Tree and forest responses to interacting elevated atmospheric
 456 CO₂ and tropospheric O₃: A synthesis of experimental evidence. In: Matyssek R, Clarke N,
 457 Cudlin P et al. (eds.) Climate Change, Air Pollution and Global Challenges: Understanding and
 458 Solutions from Forest Research, A COST action. Elsevier, Oxford, pp. 179-208

459 Koike T (1988) Leaf structure and photosynthetic performance as related to forest succession of
 460 deciduous broad-leaved trees. Plant Spec Biol 3: 77-87

461 Koike T, Kohda H, Mori S, et al (1995) Growth responses of the cuttings of two willow species
 462 to elevated CO₂ and Temperature. Plant Spec Biol 10:95-101

463 Koike, T., Kato, S., Shimamoto, Y, et al (1998) Mitochondrial DNA variation follows a
 464 geographic pattern in Japanese beech species. Bot Acta 11: 87-91

465 Koike T, Kitao M, Quoreshi A M, Matsuura Y (2003) Growth characteristics of root-shoot
 466 relations of three birch seedlings raised under different water regimes. Plant Soil 255: 303-310

467 Koike T, Watanabe M, Watanabe Y, Agathokleous E, et al (2015) Ecophysiology of deciduous
 468 trees native to Northeast Asia grown under FACE (Free Air CO₂ Enrichment). J Agr Meteo.
 469 (accepted)

470 Korotkii T, Prokushkin S G, Matsuura Y, Koike T (2002) Effects of soil temperature on the
 471 contents of nitrogen compounds in seedlings of *Larix gmelinii* regenerated on permafrost in
 472 central Siberia. Eurasian J For Res 5: 39-48

473 Kuehl R O (1999) *Design of experiments: Statistical principles of research design and analysis*,
 474 2nd ed. Duxbury Press, Boston. pp. 688

475 Leppalammi-Kujansuu J, Aro L, Salemaa M, et al (2014) Fine root longevity and carbon input
 476 into soil from below- and aboveground litter in climatically contrasting forests. *Forest Ecol*
 477 *Manag* 326: 79–90

478 Li J, Dang Q-L, Man R, Marfo J (2013) Elevated CO₂ alters N-growth relationship in spruce and
 479 causes unequal increases in N, P and K demands. *Forest Ecol Manag* 298: 19-26

480 Li P, Han X, Zong Y, et al. (2015) Effects of free-air CO₂ enrichment (FACE) on the uptake and
 481 utilization of N, P and K in *Vigna radiata*. *Agriculture, Ecosystems Environ* 202: 120-125

482 Masyagina O V, Hirano T, Ji D H, Choi D S et al. (2006) Effect of spatial variation of soil
 483 respiration rates following disturbance by timber harvesting in a larch plantation in Northern
 484 Japan. *Forest Sci Technol* 2: 80-91

485 Matsunami S (2008) Ecophysiological survey on the dispersal capacity of root sucker of Black
 486 locust and its application for the management, Ms thesis, Agr School Hokkaido Univ. pp.36.

487 NOAA (2014) Dataset accessed on August 4th, 2014, from <http://www.esrl.noaa.gov>

488 Norby R J, O'Neil E G, Luxmoore R J (1986) Effects of atmospheric CO₂ enrichment on the
 489 growth and mineral nutrition of *Quercus alba* seedlings in nutrient-poor soil. *Plant Physiol* 82:
 490 83-89

491 Norby R J, Zak D R (2011) Ecological lessons from Free-Air CO₂ Enrichment (FACE)
 492 Experiments. *Annu Rev Ecol Evol Syst* 42: 181-203

493 Oren R, Ellsworth D S, Johness K H, Phillips N, et al (2001) Soil fertility limits carbon
 494 sequestration by forest ecosystems in a CO₂-enriched atmosphere. *Nature* 411: 469-472

495 Ostonen I, Lohmus K, Pajuste K (2005) Fine root biomass, production and its proportion of NPP
 496 in a fertile middle-aged Norway spruce forest: Comparison of soil core and ingrowth core
 497 methods. *Forest Ecol Manag* 212: 264–277

498 Ostonen I, Helmisaari H-S, Borken W, et al (2011) Fine root foraging strategies in Norway
 499 spruce forests across a European climate gradient. *Glob Change Biol* 17: 3620-3632

500 Ozanne C M P, Anhuf D, Boulter S L, et al (2003) Biodiversity meets the atmosphere: a global
 501 view of forest canopies. *Science* 301: 183-186

502 Poorter H, Nagel O (2000) The role of biomass allocation in the growth response of plants to
 503 different levels of light, CO₂, nutrients and water: a quantitative review. *Aust J Plant Physiol* 27:
 504 595–607

505 Poorter H, Niklas K J, Reich P B, Oleksyn J, Poot P, Mommer L (2012) Biomass allocation to
 506 leaves, stems and roots: meta-analyses of interspecific variation and environmental control. *New*
 507 *Phytol* 193: 30-50

508 Pregitzer K S, Burton A J, King J S, Zak D R (2008) Soil respiration, root biomass, and root
 509 turnover following long-term exposure of northern forests to elevated atmospheric CO₂ and
 510 tropospheric O₃. *New Phytol* 180: 153-161

511 Pregitzer KS, Tlhelm AK (2013) Belowground carbon cycling at Aspen FACE: dynamic
 512 responses to CO₂ and O₃ in developing forests. *In: Matyssek, R. et al (eds), Climate Change, Air*
 513 *Pollution and Global Challenges: Understanding and Solutions from Forest Research. Elsevier,*
 514 *Oxford, pp. 209-226*

515 Pritchard S G, Strand A E, McCormack M L, Davis M A, et al (2008) Fine root dynamics in a
 516 loblolly pine forest are influenced by free-air-CO₂-enrichment: a six-year-minirhizotron study.
 517 Glob Change Biol 14: 588-602

518 Qu L Y, Makoto K, Choi D S, et al. (2010) The role of ectomycorrhiza in boreal forest
 519 ecosystem. *In*: Osawa, A., et al (eds), Permafrost Ecosystem: Siberian Larch Forests. Springer
 520 Ecological Studies, 209: 413-426

521 Qu L, Quoreshi A M, Koike T (2003) Root growth characteristics, biomass and nutrient
 522 dynamics of seedlings of two larch species raised under different fertilization regimes. Plant Soil
 523 255: 293-302

524 Qu L Y, Kitaoka S, Makoto K, Kuromaru M, et al (2009) Root–shoot communication of the
 525 seedlings of Japanese larch and a hybrid species grown in different soil-temperature regimes.
 526 Landscape Ecol Eng 5:115–123

527 Reijnders L (2014) Phosphorus resources, their depletion and conservation, a review. Res
 528 Conserv Recy 93: 32–49

529 Saitanis C, Bari S M, Burkey K O, Stamatelopoulos D, Agathokleous E (2014) Screening of
 530 Bangladeshi winter wheat (*Triticum aestivum* L.) cultivars for sensitivity to ozone. Environ Sci
 531 Pollut Res 21:13560-13571

532 Saitanis C, Lekkas D V, Agathokleous E, Flouri F (2015) Screening agrochemicals as potential
 533 protectants of plants against ozone phytotoxicity. Environ Pollut 197: 247-256

534 Sato T. (1995) Fundamental studies on the root growth of trees in Hokkaido. Bull Hokkaido
 535 Forest Res Inst 31: 22-38. (in Japanese with English summary).

536 Shinozaki K, Yoda K, Hozumi K, Kira T (1964a) A quantitative analysis of plant form-the pipe
537 model theory. I. Basic analyses. Jap J Ecol 14: 97-105

538 Shinozaki K, Yoda K, Hozumi K, Kira T (1964b) A quantitative analysis of plant form - the pipe
539 model theory II. Further evidence of the theory and its application in forest ecology. Jap J Ecol
540 14: 133-139

541 Schulze E-D, Beck E, Muller-Hohenstein K (2005) *Plant ecology*. Springer, Berlin-Heidelberg.
542 702pp

543 Slattery R A, Ainsworth E A, Ort D R (2013) A meta-analysis of responses of canopy
544 photosynthetic conversion efficiency to environmental factors reveals major causes of yield gap.
545 J Exp Bot 64: 3723-3733

546 Smith D E, Read D J (2008) *Mycorrhizal Symbiosis, Third Edition*. Academic Press, Great
547 Britain, pp. 787

548 Ulrich A E, Frossard E (2014) On the history of a reoccurring concept: phosphorus scarcity. Sci
549 Total Environ 490: 694-707

550 Von Uexkull H R, Mutert E (1995) Global extent, development and economic impact of acid
551 soils. In: Date RA, Grundon NJ, Raymet GE, Probert ME, (ed) *Plant–Soil Interactions at Low*
552 *pH: Principles and Management*. Dordrecht, The Netherlands: Kluwer Academic Publishers, 5–
553 19

554 Wang X, Qu L, Mao Q, et al. (2015a) Ectomycorrhizal colonization and growth of the hybrid
555 larch F₁ under elevated CO₂ and O₃. Environ Pollut 197: 116-126

556 Wang X, Fujita S, Nakaji T, et al. (2015b) Fine root turnover of Japanese white birch (*Betula*
557 *platyphylla* var. *japonica*) grown under elevated CO₂ in northern Japan. *Trees* (accepted)

558 Watanabe M, Mao Q, Novriyanti E, et al (2013) Elevated CO₂ enhances the growth of hybrid
559 larch F₁ (*Larix gmelinii* var. *japonica* x *L. kaempferi*) seedlings and changes its biomass
560 allocation. *Trees* 27: 1647-1655

561 Zhang S, Dang Q-L, Yu X (2006) Nutrient and [CO₂] elevation had synergistic effects on
562 biomass production but not on biomass allocation of white birch seedlings. *Forest Ecol Manag*
563 234: 238–244

564 Zheng S J (2010) Crop production on acidic soils: overcoming aluminum toxicity and
565 phosphorus deficiency. *Ann Bot* 106: 183-184

566

567

568

569

570

571

572

573

574

Captions

Table 1 The mean ($\pm$ SD) values of the initial height and diameter (at 2 cm) of the juveniles

Table 2 Summary of the GLM results for the measured variables trunk basal area (Area), total belowground dry mass (TDM), coarse root dry mass, Area:TDM rate, fine and intermediate roots dry masses, Fine:Intermediate dry mass ratio and Fine:Area rate. Text in bold indicates statistical significance. Data obtained from communities (*Betula ermanii*, *B. maximowicziana*, *B. platyphylla* var. *japonica*, and *Quercus mongolica* var. *crispula*) of saplings exposed to (370–390 $\mu\text{mol mol}^{-1}$) or elevated (500 $\mu\text{mol mol}^{-1}$) CO_2 and grown either in brown forest soil or immature volcanic ash plus pumice soil for 4 consecutive years. Three real replications were used for each treatment

Table 3 Mean real values ($\pm$ SD) of the measured variables trunk basal area (Area), total belowground dry mass (TDM), coarse root dry mass, Area:TDM rate, fine and intermediate roots dry masses, Fine:Intermediate dry mass ratio and Fine:Area rate. The lowercase letters above the mean values indicate the significant differences among the 4 combination treatments while the uppercase indicate the differences within the 2 main treatments. The results of each variable obtained by one GLM analysis or a Tukey range, post-hoc test, after significant results of the GLM analysis, based on standardized data. Means within each variable marked with different letter differ statistically significantly at a level of significance $\alpha=0.05$. Uppercase letters stand for the differences within each main factor and lowercase letters stand for the differences within the interactive effects of the two main factors. In case of no significant effects post-hoc test was not followed, rather means marked with the same letter. ES_v indicates the Cohen effect size of the v case for each pair with statistically significant difference and *n/a* means not applicable. Data obtained from communities (*Betula ermanii*, *B. maximowicziana*, *B. platyphylla* var. *japonica*,

598 and *Quercus mongolica* var. *crispula*) of saplings exposed to (370–390 $\mu\text{mol mol}^{-1}$) or elevated
599 (500 $\mu\text{mol mol}^{-1}$) CO_2 and grown either in brown forest soil or immature volcanic ash soil for 4
600 consecutive years. Each value is the product of three replications

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

Table 1

	Ambient CO ₂		Elevated CO ₂	
	Height (cm)	Diameter (mm)	Height (cm)	Diameter (mm)
<i>Betula ermanii</i>	30.6 (0.8)	4.2 (0.1)	29.4 (1.2)	4.1 (0.2)
<i>B. maximowicziana</i>	64.4 (1.1)	6.1 (0.2)	61.3 (1.4)	6.3 (0.3)
<i>B. platyphylla</i> var. <i>japonica</i>	45.1 (0.9)	4.1 (0.2)	44.1 (1.0)	3.9 (0.3)
<i>Quercus mongolica</i> var. <i>crispula</i>	55.1 (0.9)	9.3 (0.4)	54.2 (1.1)	9.8 (0.5)

631 **Table 2**

	CO₂	Soil	CO₂ × Soil
Area	F = 22.56, p < 0.010	F = 01.83, p = 0.213	F = 21.75, p < 0.010
TDM	F = 14.67, p < 0.010	F = 01.76, p = 0.221	F = 14.68, p < 0.010
Coarse root	F = 09.72, p < 0.050	F = 01.64, p = 0.236	F = 11.27, p < 0.010
Area:TDM	F = 00.18, p = 0.681	F = 00.75, p = 0.413	F = 04.42, p = 0.069
Fine roots	F = 79.15, p < 0.001	F = 10.65, p < 0.050	F = 18.91, p < 0.010
Intermediate roots	F = 73.00, p < 0.001	F = 04.19, p = 0.075	F = 01.23, p = 0.300
Fine:Intermediate	F = 05.75, p < 0.050	F = 04.62, p = 0.064	F = 16.75, p < 0.010
Fine:Area	F = 22.04, p < 0.010	F = 66.02, p < 0.001	F = 10.71, p < 0.050

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646 **Table 3**

	CO ₂		Soil				CO ₂ × Soil					
	aCO ₂	eCO ₂	ES _a	BF	VA	ES _b	aCO ₂ ×BF	eCO ₂ ×BF	ES _c	aCO ₂ ×VA	eCO ₂ ×VA	ES _d
Area (cm²)	5.438 ^A (0.864)	6.545 ^B (0.533)	1.69	6.149 ^A (0.377)	5.834 ^A (1.251)	<i>n/a</i>	6.064 ^a (0.501)	6.159 ^a (0.323)	<i>n/a</i>	4.736 ^b (0.371)	6.931 ^a (0.400)	6.97
TDM (kg)	0.737 ^A (0.096)	0.918 ^B (0.154)	1.55	0.796 ^A (0.059)	0.859 ^A (0.216)	<i>n/a</i>	0.774 ^a (0.085)	0.796 ^a (0.039)	<i>n/a</i>	0.678 ^a (0.074)	1.040 ^b (0.113)	4.65
Coarse root (kg)	0.702 ^A (0.097)	0.867 ^B (0.160)	1.36	0.751 ^A (0.071)	0.818 ^A (0.209)	<i>n/a</i>	0.740 ^{ab} (0.089)	0.744 ^a (0.067)	<i>n/a</i>	0.647 ^a (0.080)	0.989 ^b (0.121)	4.08
Area:TDM (cm² g⁻¹ ×10)	0.344 ^A (0.021)	0.338 ^A (0.031)	<i>n/a</i>	0.347 ^A (0.022)	0.335 ^A (0.029)	<i>n/a</i>	0.337 ^a (0.020)	0.358 ^a (0.021)	<i>n/a</i>	0.352 ^a (0.021)	0.319 ^a (0.029)	<i>n/a</i>
Fine roots (g)	12.824 ^A (2.368)	17.529 ^B (0.835)	2.90	16.040 ^A (1.520)	14.314 ^B (3.935)	0.63	14.822 ^a (0.807)	17.242 ^b (0.885)	3.50	10.812 ^c (1.099)	17.816 ^b (0.844)	8.75
Intermediate roots (g)	23.892 ^A (3.196)	35.384 ^B (2.087)	4.66	31.015 ^A (5.999)	28.261 ^A (7.299)	<i>n/a</i>	24.855 ^a (3.217)	36.015 ^b (2.146)	5.00	21.769 ^a (1.291)	34.754 ^b (2.257)	8.65
Fine:Intermediate	0.557 ^A (0.086)	0.502 ^B (0.045)	0.88	0.554 ^A (0.090)	0.505 ^A (0.041)	<i>n/a</i>	0.613 ^a (0.045)	0.479 ^b (0.036)	-	0.485 ^b (0.029)	0.524 ^b (0.047)	<i>n/a</i>
Fine:Area (g cm⁻² ×10)	0.285 ^A (0.068)	0.388 ^B (0.141)	1.02	0.247 ^A (0.036)	0.425 ^B (0.103)	2.55	0.227 ^a (0.038)	0.263 ^{ab} (0.032)	<i>n/a</i>	0.338 ^b (0.039)	0.513 ^c (0.043)	5.25

647