Title	Combined effect of astaxanthin and squalene on oxidative stress in vivo
Author(s)	Kumar, Sangeetha Ravi; Narayan, Bhaskar; Sawada, Yuki; Hosokawa, Masashi; Miyashita, Kazuo
Citation	Molecular and cellular biochemistry, 417(1-2), 57-65 https://doi.org/10.1007/s11010-016-2713-2
Issue Date	2016-06
Doc URL	http://hdl.handle.net/2115/65792
Rights	The final publication is available at link.springer.com
Type	article (author version)
File Information	Sangeetha et al (2016)-MCB-MS Revised-2.pdf

[Instructions for use](#)

**COMBINED EFFECT OF ASTAXANTHIN AND SQUALENE ON OXIDATIVE
STRESS IN VIVO**

Sangeetha Ravi Kumar^{1,2*}, Bhaskar Narayan^{1,3}, Yuki Sawada¹, Masashi Hosokawa¹ & Kazuo Miyashita^{1*}

¹Faculty of Fisheries Sciences, Hokkaido University,
3-1-1, Minato Cho, Hakodate 041-8611, Hokkaido, Japan.

²Presently, Post Doctoral Researcher at Laboratory of Lung Biology, Department of Pathobiological Sciences & Center for Experimental Infectious Disease Research, Louisiana State University, Baton Rouge 70803, Louisiana, USA.

³Department of Meat and Marine Sciences, CSIR-Central Food Technological Research Institute, Mysore 570 020, Karnataka, India.

***Author(s) for Correspondence**

Email: sangeetha.rk@gmail.com (Dr Sangeetha)/kmiya@fish.hokudai.ac.jp (Prof. Miyashita)
Faculty of Fisheries Sciences, Hokkaido University, 3-1-1, Minato Cho, Hakodate 041-8611,
Hokkaido, Japan
Tel: +81 138 408804; Fax: +81 138 408804

Abstract

Obesity and diabetes, risk factors for metabolic syndrome, are characterized by oxidative stress and inflammatory responses. Marine biofunctionals, astaxanthin (Ax) and squalene (SQ), were evaluated for their combined effect. Groups of male *KK-A^y* mice were fed high fat/sucrose diet for 4 weeks, supplemented with either 0.1% Ax, 2% SQ or 0.1 % Ax+2% SQ. In comparison to control, Sod was elevated in only Ax+SQ. while Gpx was highest in Ax+SQ, indicating the combined antioxidant effect of Ax and SQ. This was supported by elevated mRNA expression of Sod1 and Gpx1. Except adiponectin (elevated in Ax and Ax+SQ), expression of other inflammatory markers was not altered. Blood glucose levels were decreased in SQ and Ax+SQ while liver triglycerides decreased in SQ group. This is the first *in-vivo* study demonstrating combined effects of Ax and SQ resulting in antioxidant effects and modulation of glucose/cholesterol levels. This study highlights the benefit of utilizing Ax and SQ together for management of obesity/diabetes.

Keywords: Antioxidant enzymes; astaxanthin; *KK-A^y* mice; lipid hydroperoxides; squalene

37

38 **1 Introduction**

39 The metabolic syndrome - a complex condition involving obesity, insulin resistance,
40 hypertension, diabetes, hyperinsulinemia and dyslipidemia - is a major risk factor for
41 atherosclerosis and coronary artery disease. The conditions involved in metabolic syndrome
42 are believed to lead to insulin resistance and clinical and biochemical changes [1]. Further,
43 obesity is a common risk factor for diabetes along with advanced age, undesirable diet and
44 lifestyle changes. These contribute to an environment of increased oxidative stress which is
45 believed to be a major risk factor for the onset of diabetes as well as progression of the
46 disease. Diabetes is characterized by hyperglycemia which contributes to the progression and
47 maintenance of an oxidative environment [2]. Hence, compounds exerting antioxidant,
48 hypoglycemic, anti-obesity or anti-inflammatory effect and/or attenuate lipid parameters
49 would be beneficial in these conditions.

50 Astaxanthin (Ax) is a red-colored xanthophyll that is primarily found in the marine
51 environment. Ax is biosynthesized by microalgae/ phytoplankton, accumulates in
52 zooplankton and crustaceans and subsequently in fish from where they are added to the
53 higher levels in the food chain. Ax is shown to exert several benefits to human health such as
54 antioxidant, anti-inflammatory, anti-cancer, anti-diabetic effects as well as protection against
55 cardiovascular, neurological, gastro-intestinal disorders and in ocular health and drug toxicity
56 [3]. The potential of Ax to act as an antioxidant has been established *in-vitro* and *in-vivo* by
57 several researchers [4-9].

58 Squalene (SQ) is a triterpenoid that is ubiquitously found in the plant and animal kingdoms
59 by virtue of its importance as an intermediate in the sterol pathway. Although the most
60 abundant source of squalene is shark liver oil, it is also found in terrestrial oil sources such as
61 olive, amaranth seed, rice bran etc. In humans, the main storage organ is the skin where it is

62 believed to protect against the oxidative stress caused by exposure to ionizing rays [10]. Sq is
63 reported to act as an antioxidant, anti-inflammatory, anti-cancer, hypoglycemic and
64 hypocholesterolemic agent [11,12]. However, several researchers have reported the negative
65 effect of feeding squalene, viz. hypercholesterolemia [13,14].

66 Saw et al [15] carried out a combination study involving Ax and PUFAs (EPA/DHA) and
67 reported their synergistic effect *in-vitro*. Dhandapani et al [16] have reported the synergistic
68 effect of SQ and PUFA (fish oil concentrate with ~65% EPA/DHA) on lipid peroxidation and
69 increase in activities of antioxidant enzymes such as superoxide dismutase (Sod), catalase
70 (Cat), glutathione peroxidase (Gpx), and glutathione transferase. However, apart from an *in-*
71 *vitro* report by our group very recently [17], there are no studies available detailing the effect
72 of Ax and SQ when present together. The results of the *in-vitro* study by our group had
73 revealed the combined effect of Ax and SQ in suppressing lipid hydroperoxides in RAW
74 264.7 cells stimulated by lipopolysaccharide [17]. It is evident from the reports in literature
75 that both Ax and SQ exert antioxidant and anti-inflammatory effects and this may involve
76 common factors such as NF-kB, ERK, MAPK, iNOS, Cox-2, P13/Akt and related
77 downstream molecules like interleukin-1 β (IL-1 β), interleukin-6 (IL-6), tumor necrosis factor-
78 α (TNF- α), interferon- γ (INF), monocyte chemoattractant protein-1 (MCP-1), peroxisome
79 proliferator activated receptor-alpha (PPAR- α) and PPAR- γ [11,18-21]. With this
80 background, the present study was undertaken to study the effect of Ax and SQ, individually
81 and in combination, in obese/diabetic *KK-A^y* mice fed a high fat (20%) diet. The activity of
82 antioxidant enzymes, cholesterol and triglyceride levels was measured in liver and white
83 adipose tissue (WAT). Serum glucose levels as well as blood cholesterol profile was
84 determined. Molecules pertaining to antioxidant activity of enzymes as well as markers of
85 inflammation such as TNF- α , PPAR- α , PPAR- γ , IL-6, IL-1 β , INF, Cox-1, Cox-2 and
86 adiponectin were studied.

2 Materials and Methods

2.1 Materials

Squalene (SQ) and dietary lipids were purchased from Wako Pure Chemicals, Ltd., Osaka (Japan). Astaxanthin (Ax) (5%) was obtained from AstaReal, Japan. Soybean oil and lard were from Summit oil Mill Co. Ltd., Chiba (Japan) and Junsei Chemical Co. Inc., Tokyo, (Japan), respectively. All the other chemicals and solvents used in the study were of analytical grade, unless mentioned otherwise and were purchased from Wako Pure Chemicals, Ltd., Osaka (Japan).

2.2 Animals and diets

Male obese/diabetic model, *KK-A^y* mice (4 weeks old, n=28) were acclimatized with basal diet for a week following which they were divided into 4 groups and fed a high fat (20%) diet based on the the recommendations of American Institute of Nutrition (AIN-93G), for 4 weeks [22]. The different groups (n=7/group) included control, astaxanthin (Ax-0.1%), squalene (SQ-2%) and combination of astaxanthin and squalene (Ax 0.1% + SQ 2%). The dosage of SQ was fixed at the mid level of dosages reported (1-4%) in previous works [12, 23-25]. Ax dosage was based on the work carried out in our laboratory (unpublished). The composition of the control and experimental diets is given in Table 1. The animals were maintained at 23 ± 2°C temperature and 60 ± 5 % humidity with a 12 hour light/dark cycle. All the animals had free access to food and water. All the procedures and protocols for the use and care of animals were approved by the Ethical Committee for Experimental Animal Care of the Hokkaido University, Japan (Approval no. 113 14-0072).

Daily measurements of body weight and food intake were recorded. At the end of the 4 weeks feeding trial, the animals were sacrificed and blood and organs were sampled. . The blood was collected in vacuettes for separation of serum. Organs were excised, weighed, divided and stored in liquid nitrogen, RNA later solution (Life Technologies, USA) and

stored in deep freezer (-40°C) till further analyses. Analyses were carried out with serum, liver and epididymal white adipose tissue (WAT) tissues. Hereafter, reference to WAT would indicate epididymal WAT unless mentioned otherwise.

2.3 Lipid Hydroperoxides

Total lipid extracts were obtained from liver and WAT by the method of Folch et al [26] and the lipid hydroperoxides present were allowed to react with non-fluorescent diphenyl-1-pyrenylphosphine (DPPP) to give fluorescent DPPP oxide which could be measured by the fluorescence detector of the HPLC system [17]. In brief, the weighed lipid extract was dissolved in chloroform: methanol (2:1, v/v) (containing 10 mg butylhydroxytoluene per mL chloroform) and 100 µl of this sample solution was allowed to react with 50 µL of DPPP solution (1 mg/10 ml chloroform) in a water bath for 60 min at 60°C. The reaction mixture was cooled on ice and isopropyl alcohol (3 ml) was added to it. An aliquot was drawn, diluted with mobile phase and injected to the HPLC system (reverse phase). The HPLC system used was Hitachi L-2350 HPLC system (Hitachi, Tokyo, Japan) with a pump (L-2130), an auto-sampler (L-2200) and a fluorescence detector (L-2485). The DPPP oxide was measured at 40°C with a Develosil-ODS column (UG-5, Nomura Chemicals, Aichi, Japan), protected by a ODS guard column (10 x 4.0 mm i.d.). The mobile phase contained HPLC grade butanol and methanol (10:90, v/v), and the flow rate was 1.0 ml/min. The fluorescence detector was set at Ex. 352 nm and Em. 380 nm. The lipid hydroperoxide concentration in the samples were calculated using the DPPP standard curve and expressed as pmol/g lipid.

2.4 Antioxidant Enzymes and Molecules

Antioxidant enzymes, superoxide dismutase (Sod), catalase (Cat), glutathione peroxidase (Gpx) as well as reduced glutathione (GSH) and oxidised glutathione levels (GSSG) were estimated in liver tissue while Sod and Gpx were measured in WAT, with the help of research kits. Kits for estimation of SOD and GSH/GSSG were obtained from Dojindo

Molecular Technologies Inc (MD, USA), protocols as defined by the supplier were followed and results were obtained by reading at 450 and 405/415 nm, respectively. Cat and Gpx kits were procured from Cayman Chemical Company (MI, USA), instructions of the supplier were followed and final readings were recorded at 540 and 340 nm, respectively.

2.5 Profiling of lipids and glucose in tissues and serum

Lipids extracted from liver and WAT by the method of Folch et al [26] were subjected to cholesterol and triglyceride (TG) analysis by LabAssay Kits (Wako Pure Chemical Industries, Osaka, Japan) by spectrophotometric method using a plate reader. Serum samples were outsourced to Hakodate Medical Association Health Examination and Clinical Laboratory Centre for estimation of cholesterol, LDL-C and HDL-C. Blood glucose levels (fasting) were measured in the live animals by using tail blood on a glucose monitor, the Glutest Neo Sensor (Sanwa Kagaku Kenkyusho Co. Ltd., Aichi, Japan). This sensor is an amperometric sensor with flavin adenine dinucleotide (FAD)-dependent glucose dehydrogenase and ferricyanide ($\text{Fe}(\text{CN})_6^{3-}$).

2.6 Quantitative real time PCR

Total RNA was extracted from liver and WAT of mice using the Rneasy Lipid Tissue Mini Kit (Qiagen, Tokyo, Japan) according to the manufacturer's protocol. The cDNA was then synthesized from total RNA using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems Japan Ltd., Tokyo, Japan). Quantitative real time PCR analysis of individual cDNA was performed with ABI Prism 7500 (Applied Biosystems Japan Ltd., Tokyo, Japan) using primers for TaqMan Gene Expression Assays (Applied Biosystems Japan Ltd., Tokyo, Japan; [Sod1: Mm01344233_g1, Sod2: Mm01313000_m1, Cat: Mm00437992_m1, GPX1: Mm00656767_g1, Adiponectin: Mm00456425_m1, PPAR α : Mm00440939_m1, PPAR γ : Mm00440940_m1, Cox1: Mm00477214_m1, Cox2: Mm00478374_m1, Il-1 β : Mm00434228_m1, Il-6: Mm00446190_m1, MCP-1:

Mm0041242_m1, TNF α : Mm00443258_m1, IFN: Mm01168134_m1, GAPDH: Mm99999915_g1]. PCR cycling conditions were 50°C for 2 min, 95°C for 10 min, followed by 40 cycles 15s at 95°C and 60°C for 1 min.

2.7 Statistical analysis

All data were analysed by one way analysis of variance (ANOVA) and wherever significant, mean separation was accomplished using Student's t-test at 95% confidence level.

3 Results

3.1 Body and organ weights

Body weight of the animals was measured daily. A slight decrease ($p>0.05$) was observed in the average body weight of the SQ and Ax+SQ groups as compared to the Ax and Con groups. A significant increase ($p<0.05$) was observed in the weight of liver in SQ group and brown adipose tissue (BAT) in Ax group, while BAT and retroperitoneal WAT was significantly decreased ($p<0.05$) in SQ group, compared to control. Weights of other organs such as spleen, kidney, brain and epididymal WAT were unaffected ($p>0.05$) by feeding Ax and/or SQ in the diet. Body weight and weight of the various organs in different groups is given in Table 2.

3.2 Lipid hydroperoxides and anti-oxidant molecules

Lipid hydroperoxide levels, measured as pmol DPPP/g lipid, were significantly suppressed ($p<0.05$) in liver of all the experimental groups, whereas, in WAT, the decrease was not significant ($p>0.05$), when compared to control. No significant difference ($p>0.05$) in lipid hydroperoxide levels was observed amongst the experimental groups in liver and WAT. Lipid hydroperoxide levels as well as activities of antioxidant enzymes in are shown in Figure 1. Alteration of antioxidant molecules/activity was observed in liver of mice, but not in WAT. Sod activity was significantly ($p<0.05$) increased in liver of Ax+SQ groups while there was only slight ($p>0.05$) increase in the case of Ax group. No significant change

($p>0.05$) was observed for Sod activity in WAT. Liver Cat activity was significantly elevated ($p<0.05$) in Ax, SQ and Ax+SQ groups when compared to control, and this rise was higher ($p<0.05$) in Ax group than SQ and Ax+SQ groups. Liver Gpx activity was also significantly increased ($p<0.05$) in the experimental groups when compared to control and this was in the order of $Ax<SQ<Ax+SQ$. GSSG/GSH, considered to be an indicator of oxidative stress status, was significantly ($p<0.05$) lower in the experimental groups as compared to control but were not significantly different ($p>0.05$) amongst the test groups.

3.3 Expression of mRNA relevant to antioxidant and immunomodulatory molecules

Quantitative RT-PCR analysis results for liver mRNA expression of genes relevant to antioxidant activity and immunomodulatory pathways are shown in Figures 2 and 3 respectively. The results demonstrated an increase in mRNA expression of liver Sod1 and adiponectin in the experimental groups when compared to control, though significant ($p<0.05$) only for Ax and Ax+SQ groups. In addition, the mRNA expression of liver Gpx1 was significantly elevated for Ax+SQ over control ($p<0.05$). Whereas, only a slight ($p>0.05$) increase in mRNA expression was observed for liver Sod2 and Cat; no significant changes ($p>0.05$) were observed in expression of genes involved in the antioxidant and immunomodulatory pathways in epididymal WAT tissue (data not shown). No significant change ($p>0.05$) was observed for immunomodulatory factors such as MCP-1, IL-6, IL-1 β , TNF- α , IFN, PPAR- α , PPAR- γ , Cox-1 and Cox-2 on feeding Ax, SQ or Ax+SQ to mice while adiponectin expression was at increased levels ($p<0.05$) in Ax and Ax+SQ groups as compared to control.

3.4 Lipids and glucose profile of serum and tissue

The lipid profile of blood and tissues of the different groups as well as fasting blood glucose levels are presented in Table 3. Increase in serum cholesterol and HDL-C was observed in mice fed either Ax or SQ as compared to control ($p<0.05$), whereas, the cholesterol and

212 HDL-C levels of Ax+SQ as well as LDL-c levels in all groups were comparable with control.
213 In addition, blood glucose levels saw a significant ($p<0.05$) decrease in the SQ and Ax+SQ
214 groups while the decrease in Ax group was not significant ($p>0.05$). Although statistically
215 significant ($p<0.05$), changes in liver and WAT cholesterol were not apparent on feeding SQ
216 and Ax+SQ. With regard to TG, levels in liver were significantly decreased ($p<0.05$) in SQ
217 and Ax+SQ groups while in Ax, it was significantly increased ($p<0.05$), when compared to
218 control. TG levels in WAT were unaffected ($p<0.05$) by feeding Ax and/or SQ.

219 **4. Discussion**

220 Obese/diabetic *KK-A^y* mice were fed 20% fat diet supplemented with either Ax, SQ or
221 Ax+SQ. Feeding Ax and/or SQ did not have any significant effect ($p>0.05$) on weight gain
222 although average weight of SQ and Ax+SQ fed mice appeared to be lower than control
223 ($p>0.05$). Organs weights were also not affected except for liver and BAT which were
224 significantly ($p<0.05$) increased in SQ and Ax respectively, while BAT and retro-peritoneal
225 WAT were decreased ($p<0.05$) in Ax+SQ group. Gabas-Rivera et al [27] have reported a
226 similar effect with no change in weight and increase in liver weight of mice fed SQ at a dose
227 of 1g/kg while a decrease in body weight of rats was observed with SQ feeding by Liu et al
228 [28].

229 Lipids, present in the cells are prone to peroxidation by free radicals. The effects of these are
230 neutralized by antioxidant molecules including endogenous enzymes and molecules as well
231 as externally supplied antioxidants. Feeding Ax, SQ and Ax+SQ resulted in decrease in lipid
232 hydroperoxides in liver. Activity of antioxidant enzymes like Sod, Cat and Gpx in liver were
233 elevated on feeding Ax, SQ and Ax+SQ. Hashimoto et al [5] and Choi et al [6] have also
234 reported Ax to be a potent antioxidant with increase in Sod activity and other antioxidant
235 enzymes and decrease in hydroperoxides in human subjects. Similarly, Tripathi and Jena [7]
236 have reported the protective effect of Ax resulting in increased Sod activity and GSH levels

237 in murine model. Whereas, Augusti et al [8] have reported Ax to increase in Sod and decrease
 238 in Cat and Gpx activities in mercuric chloride treated rats. The present results also
 239 corroborate with the results of Buddhan et al [29] and Farvin et al [30], who have also
 240 reported suppression of lipid peroxidation and improvement in the antioxidant enzymes as a
 241 result of feeding 2% SQ to rats, while Moreno [31] reported the lack of antioxidant activity
 242 for SQ in murine cell lines. Das et al [10] have also demonstrated increased GSH levels and
 243 increased activity of Gpx and glutathione transferase antioxidant enzymes with SQ in mice.
 244 In the present study, antioxidant enzymes and the ratio of the indigenous antioxidant
 245 molecules, GSSG/GSH was also decreased in the experimental groups. The GSSG/GSH ratio
 246 is accepted as a marker of oxidative stress status [32]. Collectively, this implies that feeding
 247 Ax, SQ or Ax+SQ protected the cells from oxidative stress. In particular, Ax+SQ was more
 248 potent ($p<0.05$), than Ax and SQ individually, in elevating the activity of Sod and Gpx while
 249 increase in activity of Cat was significantly more in Ax. The clearly indicates that Ax and SQ
 250 are more potent in combination than when present alone. This surmise is further supported by
 251 the mRNA expression of Sod1, Gpx1 and adiponectin in liver, where Ax and Ax+SQ showed
 252 a significant elevation over control and in addition, Ax +SQ had a markedly higher
 253 expression of Gpx1. This is the first study reporting the combined effect of carotenoids and
 254 SQ *in-vivo*. Previously, Saw et al [15] have reported that the combination of Ax with
 255 EPA/DHA resulted in synergistic antioxidant effects in Hep-G2-C8 cells and Dhandapani et
 256 al [16] have reported on the synergistic effect of SQ and PUFA as antioxidants in rats. We
 257 had previously reported the combined effect of Ax and SQ on lipid peroxidation in
 258 macrophages [17] and speculated that it may involve regulation of inflammatory markers
 259 common to Ax and SQ. However, feeding Ax, SQ or Ax+SQ did not significantly influence
 260 mRNA expression of IL-1 β , IL-6, MCP-1, TNF- α , IFN, Cox-1, Cox-2, PPAR- α and PPAR- γ .
 261 A significant increase ($p<0.05$) was observed in the mRNA expression of adiponectin for Ax

262 and Ax+SQ. Previously, Cardeno et al [11] have reported SQ to decrease the mRNA
 263 expression of TNF- α , IL-1 β , IL-6, IFN, Cox-2 and other markers of inflammatory processes
 264 in neutrophils and monocytes stimulated with lipopolysaccharide. Similarly, Ax was reported
 265 to suppress mRNA expression of NF- κ B, Cox-1, Cox-2, MCP-1, TNF- α , IL-1 β *in-vitro* and
 266 *in-vivo* [18-21]. Jia et al [33] have reported that Ax activated PPAR- α but suppressed PPAR-
 267 γ in mice. Park et al [34] have reported increase in IFN and decrease in IL-6 in human
 268 subjects fed high dose of Ax (8mg/day) but not for those fed lower dose (2 mg/day). A
 269 possible reason for the discrepancy in our results from those of others may be due to the use
 270 of a very low dosage of Ax (0.01%) in the diet of the mice. A longer duration study may be
 271 considered for confirming the true effect of Ax and SQ in the mice.

272 In addition to the protective effects of feeding Ax, SQ and Ax+SQ on oxidative stress related
 273 parameters, the experimental diets also had an effect on the cholesterol and TG in the blood
 274 and tissues as well as fasting blood glucose levels. While Ax and SQ groups had increased
 275 ($p < 0.05$) levels of cholesterol and HDL-C compared to control, the levels in Ax+SQ group
 276 was comparable ($p > 0.05$) with control. An interesting observation made from Table 3 is that
 277 while cholesterol levels increased in Ax and SQ fed groups, there was a corresponding rise in
 278 HDL-C and LDL-C in Ax but a rise in HDL-C and fall in LDL-c levels in SQ. The
 279 statistically significant decrease observed for cholesterol in liver and WAT of SQ and Ax+SQ
 280 could be misleading as it was not apparent when viewed holistically. Liver TG decreased
 281 ($p < 0.05$) in SQ group, whereas, Ax was found to result in increased liver TG. Aguilera et al
 282 [35] have reported increased cholesterol and TG levels in chick embryos treated with alcohol
 283 and this effect was attenuated by feeding SQ indicating its hypocholesterolemic and
 284 hypotriglyceridemic effect. Farvin et al [12] have also reported the ability of SQ to suppress
 285 increase in cholesterol and TG levels in rats. de Castro et al [23] whereas, have reported no
 286 change in the lipid profile as a result of feeding SQ to hamsters. Contrary results have been

287 reported by Relas et al [13] and Zhang et al [14] who have reported increase in cholesterol
288 and TG levels as a result of SQ feeding, *in-vivo*. The decreased serum cholesterol and liver
289 TG observed in our study may have been due to a squalene-induced suppression of the HMG-
290 CoA reductase activity, by a feedback mechanism [35]. In addition to lipids, blood glucose
291 levels were significantly decreased in SQ and Ax+SQ groups. Liu et al [28] have also
292 reported lowering of plasma cholesterol, TG and glucose levels on feeding SQ to rats.
293 Tsuchiya et al [36] have suggested that this may be due to glucose stimulated insulin
294 secretion and increased insulin levels. Overall, the results indicate that while Ax, SQ and
295 Ax+SQ protected the cells from oxidative stress, the effect was more pronounced when Ax
296 and SQ were administered in combination. Moreover, the combination Ax+SQ resulted in a
297 significant decrease in tissue cholesterol and blood glucose levels, indicating its multifaceted
298 beneficial effect in diabetes/obesity.

299 This is the first study *in-vivo* to highlight the advantage of using them in combination as
300 exhibited by their superior antioxidant effects and ability to modulate glucose and cholesterol
301 levels in blood and tissues. Dietary supplementation with Ax or/and SQ resulted in decreased
302 lipid hydroperoxide levels as well as increased activity of antioxidant enzymes and molecules
303 in *KK-A^y* mice fed a high fat/sucrose diet. In combination, the elevation in the activities of
304 Sod and Gpx antioxidant enzymes was greater than when Ax and SQ were present
305 individually and this was supported by the increased mRNA gene expression of Sod1, Gpx1
306 and adiponectin. In addition, tissue cholesterol and triglyceride levels as well as serum
307 glucose levels were lowered in SQ and Ax+SQ groups. This study thus confirms the
308 antioxidant effects of Ax and SQ. Given that Ax and SQ are obtained naturally from marine
309 and other food sources and have multifarious effects, especially when used in combination,
310 they may be considered for improving the outcome of obesity/diabetes management regimes.

Further studies are required for establishing the combined effect of Ax and SQ in normal mice models.

List of abbreviations

Ax : Astaxanthin, BAT: Brown Adipose Tissue, Cat : catalase, Cox : Cyclooxygenase, DHA : Docosahexaenoic acid, DPPP: diphenyl-1-pyrenylphosphine, EPA : Eicosapentaenoic acid, ERK : Extracellular-signal regulated kinase, Gpx : glutathione peroxidase, GSH : Reduced glutathione, GSSG : Oxidised glutathione, HDL-C: High density cholesterol, iNOS : Inducible nitric oxide synthase, MAPK : Mitogen activated protein kinase, MCP : Monocyte chemoattractant protein, NF : Nuclear factor, PPAR : Peroxisome proliferator activated receptor, Sod : Superoxide dismutase, SQ : Squalene, TG: Triglycerides, TNF : Tumor necrosis factor, WAT : White adipose tissue

Acknowledgements

This work was supported by “Scientific technique research promotion program for agriculture, forestry, fisheries and food industry” and partially supported by a National Project for the Formation of Tohoku Marine Science Center from MEXT (Ministry of Education, Culture, Sports, Science & Technology in Japan). SRK and BN are grateful to Japan Society for Promotion of Science (JSPS) for Post-Doctoral Fellowship and Invitation Fellowship, respectively.

Conflict Of Interest

The authors have no conflict of interest to report.

336

337 **REFERENCES**

- 338 [1] Dandona P, Aljada A, Chaudhuri A, Mohanty P, Garg R (2005) Metabolic
339 syndrome: A comprehensive perspective based on interactions between obesity,
340 diabetes, and inflammation. *Circulation* 111: 1448-1454.
- 341 [2] Rains JL, Jain SK (2011) Oxidative stress, insulin signalling, and diabetes. *Free*
342 *Rad Biol Med* 50: 567–575.
- 343 [3] Yuan J-P, Peng J, Yin K, Wang J-H (2011) Potential health-promoting effects of
344 astaxanthin: A high-value carotenoid mostly from microalgae. *Mol Nutr Food Res*
345 55: 150–165.
- 346 [4] Otsuka T, Shimazawa M, Nakanishi T, Ohno Y, Inoue Y, Tsuruma K, Ishibashi T,
347 Hara H (2013) The protective effects of a dietary carotenoid, astaxanthin, against
348 light-induced retinal damage. *J Pharmacol Sci* 123:209 – 218.
- 349 [5] Hashimoto H, Arai K, Hayashi S, Okamoto H, Takahashi J, Chikuda M, Obara Y
350 (2013) Effects of astaxanthin on antioxidation in human aqueous humour. *J Clin*
351 *Biochem Nutr* 53: 1-7.
- 352 [6] Choi HD, Kim JH, Chang MJ, Kyu-Youn Y, Shin WG (2011) Effects of
353 astaxanthin on oxidative stress in overweight and obese adults. *Phytother Res* 25:
354 1813–1818.
- 355 [7] Tripathi DN, Jena GB (2009) Intervention of astaxanthin against
356 cyclophosphamide-induced oxidative stress and DNA damage: A study in mice.
357 *Chem Biol Interact* 180: 398–406.
- 358 [8] Augusti PR, Conterato GMM, Somacal S, Sobieski R, Spohr PR; Torres JV,
359 Charao MF, Moro AM, Rocha MP, Garcia SC, Emanuelli T (2008) Effect of

astaxanthin on kidney function impairment and oxidative stress induced by
mercuric chloride in rats. *Food Chem Toxicol* 46: 212–219.

[9] Nakajima Y, Inokuchi Y, Shimazawa M, Otsubo K, Ishibashi T, Hara H (2008)

Astaxanthin, a dietary carotenoid, protects retinal cells against oxidative stress in-
vitro and in mice in-vivo. *J Pharm Pharmacol* 60: 1365–1374.

[10] Das B, Antoon R, Tsuchida R, Lotfi S, Morozova O, Farhat W, Malkin D, Koren

G, Yeager H, Baruchel S (2008) Squalene selectively protects mouse bone marrow
progenitors against cisplatin and carboplatin-induced cytotoxicity in vivo without
protecting tumor growth. *Neoplasia* 10: 1105–1119.

[11] Cardeno A, Aparicio-Soto M, Montserrat-de laPaz S, Bermudez B, Muriana FJG,

Alarcon-de-la-Lastra C (2015) Squalene targets pro and anti-inflammatory
mediators and pathways to modulate over activation of neutrophils, monocytes and
macrophages. *J Func Foods* 14:779–790.

[12] Farvin KHS, Anandan R, Kumar SHS, Shiny KS, Mathew S, Sankar TV, Nair

PGV (2006) Cardioprotective effect of squalene on lipid profile in isoprenaline-
induced myocardial infarction in rats. *J Med Food* 9: 531–536.

[13] Relas H, Gylling H, Miettinen TA (2000) Dietary squalene increases cholesterol

synthesis measured with serum noncholesterol sterols after a single oral dose in
humans. *Atherosclerosis* 152: 377–383.

[14] Zhang Z, Yeung WK, Huang Y, Chen Z-Y (2002) Effect of squalene and shark

liver oil on serum cholesterol level in hamsters. *Int J Food Sci Nutr* 53: 411–418.

[15] Saw CLL, Yang AY, Guo Y, Kong A-NT (2013) Astaxanthin and omega-3 fatty

acids individually and in combination protect against oxidative stress via the Nrf2–
ARE pathway. *Food Chem Toxicol* 62:869–875.

- [16] Dhandapani N, Ganesan B, Anandan R, Jeyakumar R, Rajaprabhu D, Ezhilan RA (2007) Synergistic effects of squalene and polyunsaturated fatty acid concentrate on lipid peroxidation and antioxidant status in isoprenaline-induced myocardial infarction in rats. *Afr J Biotechnol* 6:1021-1027.
- [17] Sangeetha RK, Bhaskar N, Kizawa Y, Hosokawa M, Miyashita K (2015) Does squalene alter the antioxidant potential of astaxanthin and fucoxanthinol? In vitro evidence in RAW264.7 cells, a murine macrophage. *J Food Sci Tech* DOI: 10.1007/s13197-015-2077-8 (Available online).
- [18] Yasui Y, Hosokawa M, Mikami N, Miyashita K, Tanaka T (2011) Dietary astaxanthin inhibits colitis and colitis-associated colon carcinogenesis in mice via modulation of the inflammatory cytokines. *Chem Biol Interact* 193: 79–87.
- [19] Macedo RC, Bolin AP, Marin DP, Otton R (2010) Astaxanthin addition improves human neutrophils function: in vitro study. *Eur J Nutr* 49:447-457.
- [20] Manabe E, Handa O, Naito Y, Mizushima K, Akagiri S, Adachi S, Takagi T, Kokura S, Maoka T, Yoshikawa T (2008) Astaxanthin protects mesangial cells from hyperglycemia-induced oxidative signaling. *J Cell Biochem* 103: 1925–1937.
- [21] Pashkow FJ, Watumull DG, Campbell CL (2008) Astaxanthin: A novel potential treatment for oxidative stress and inflammation in cardiovascular disease. *Am J Cardio* 101S:58D– 68D.
- [22] Reeves PG, Nielsen FH, Fahey GC (1993) AIN-93 purified diets for laboratory rodents: Final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet. *J Nutr* 123: 1939-1951.

- [23] deCastro LIA, Soares RAM, Saldiva PHN, Ferrari RA, Miguel AMRO, Almeida CAS, Are^as JAG (2013) Amaranth oil increased fecal excretion of bile acid but had no effect in reducing plasma cholesterol in hamsters. *Lipids* 48: 609–618.
- [24] Farvin KHS, Anandan R, Kumar SHS, Shiny KS, Mathew S, Sankar TV, Nair PGV (2006) Cardioprotective Effect of Squalene on Lipid Profile in Isoprenaline-Induced Myocardial Infarction in Rats. *J Med Food* 9: 531–536.
- [25] Smith D, Espino-Montoro A, Perez-Jimenez F, Pedro-Botet J, Pereperez JJ, Ordovas JM (2000) Effect of a high saturated fat and cholesterol diet supplemented with squalene or β -sitosterol on lipoprotein profile in F1B hamsters. *Nutr Res* 20: 1309–1318.
- [26] Folch J, Lees M, Stanley GHS (1957) A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 26: 497–509.
- [27] Gabas-Rivera C, Barranquero C, Martinez-Beamonte R, Navarro MA, Surra JC, Osada J (2014) Dietary squalene increases high density lipoprotein-cholesterol and paraoxonase 1 and decreases oxidative stress in mice. *PLoS ONE* 9: e104224.
- [28] Liu Y, Xu X, Bi D, Wang X, Zhang X, Dai H, Chen S, Zhang W (2009) Influence of squalene feeding on plasma leptin, testosterone and blood pressure in rats. *Indian J Med Res* 129: 150-153.
- [29] Buddhan S, Sivakumar R, Dhandapani N, Ganesan B, Anandan R (2007) Protective effect of dietary squalene supplementation on mitochondrial function in liver of aged rats. *Prost Leuk Essen Fatty Acids* 76: 349–355.
- [30] Farvin KHS, Anandan R, Kumar SHS, Shiny KS, Sankar TV, Thankappan TK (2004) Effect of squalene on tissue defence system in isoproterenol-induced myocardial infarction in rats. *Pharmacol Res* 50: 231–236.

- [31] Moreno JJ (2003) Effect of olive oil minor components on oxidative stress and arachidonic acid mobilization and metabolism by macrophages raw 264.7. *Free Rad Biol Med* 35:1073–1081.
- [32] Baeza I, Fdez-TresguerresJ, Ariznavarreta C, De la Fuente M (2010) Effects of growth hormone, melatonin, oestrogens and phytoestrogens on the oxidized glutathione (GSSG)/reduced glutathione (GSH) ratio and lipid peroxidation in aged ovariectomized rats. *Biogerontology* 11: 687–701.
- [33] Jia Y, Wu C, Kim J, Kim B, Lee S-J (2015) Astaxanthin reduces hepatic lipid accumulations in high-fat fed C57BL/6J mice via activation of peroxisome proliferator-activated receptor (PPAR) alpha and inhibition of PPAR gamma and Akt. *J Nutr Biochem* DOI: 10.1016/j.jnutbio.2015.09.015 (Available online).
- [34] Park JS, Chyun JH, Kim YK, Line LL, Chew BP (2010) Astaxanthin decreased oxidative stress and inflammation and enhanced immune response in humans. *Nutr Metab* 7:18-24.
- [35] Aguilera Y, Dorado ME, Prada FA, Martí'nez JJ, Quesada A, Ruiz-Gutie'rrrezV (2005) The protective role of squalene in alcohol damage in the chick embryo retina. *Exp Eye Res* 80: 535–543.
- [36] Tsuchiya M, Hosaka M, Moriguchi T, Zhang S, Suda M, Yokota-Hashimoto H, Shinozuka K, Takeuchi T (2010) Cholesterol biosynthesis pathway intermediates and inhibitors regulate glucose-stimulated insulin secretion and secretory granule formation in pancreatic β -cells. *Endocrinol* 151: 4705–4716.

Table 1. Composition of control and experimental diets

Ingredient (g/kg)	Control	Ax[*]	SQ[*]	Ax+SQ
β-Corn starch	224.6	224.6	224.6	224.6
α-Corn starch	84.9	84.9	84.9	84.9
Milk casein	260	260	260	260
Sucrose	130	130	130	130
Cellulose	50	50	50	50
Soybean oil	70	50	50	30
Lard	130	130	130	130
Ax oil [#]	-	20	-	20
Squalene	-	-	20	20
L-cystine	3	3	3	3
AIN93G mineral mix	35	35	35	35
AIN93G vitamin mix	10	10	10	10
Choline bitartrate	2.5	2.5	2.5	2.5
Tert-butyl hydroquinone	0.014	0.014	0.014	0.014

^{*}Ax - Astaxanthin, SQ - Squalene; [#]Ax oil contained 5% Ax

Table 2. Effect of feeding Astaxanthin (Ax) or/and Squalene (SQ) on body and organ weights.

	Control	Ax	SQ	Ax+SQ
<i>Body weight, g</i>				
Initial	26.12±0.99 ^a	26.02±1.56 ^a	25.78±1.45 ^a	25.70±1.02 ^a
Final	42.77±1.83 ^a	43.13±1.59 ^a	42.01±1.40 ^a	40.94±1.63 ^a
<i>Organ weight, g</i>				
Liver	2.24±0.35 ^a	2.00±0.18 ^{a,b}	2.64±0.29 ^{b,c}	2.39±0.34 ^{a,c}
Spleen	0.12±0.03 ^a	0.10±0.01 ^a	0.10±0.03 ^a	0.12±0.05 ^a
BAT [*]	0.23±0.03 ^a	0.26±0.03 ^b	0.20±0.04 ^a	0.20±0.04 ^c
Epididymal WAT [#]	1.92±0.29 ^a	2.05±0.24 ^a	1.96±0.25 ^a	1.94±0.47 ^a
Retroperitoneal WAT [#]	0.81±0.14 ^a	0.81±0.18 ^a	0.78±0.17 ^a	0.64±0.08 ^b
Kidney	0.59±0.05 ^a	0.64±0.05 ^a	0.60±0.07 ^a	0.57±0.04 ^a
Brain	0.35±0.05 ^a	0.37±0.02 ^a	0.38±0.03 ^a	0.35±0.02 ^a

^{*}BAT – Brown Adipose Tissue; [#]WAT – White Adipose Tissue; Columns not sharing common alphabets are significantly different (p<0.01)

Table 3. Effect of feeding astaxanthin (Ax) or/and squalene (SQ) on lipid profile and fasting blood glucose in serum, liver and epididymal white adipose tissue (WAT).

Parameter	Control	Ax	SQ	Ax+SQ
Serum (all parameters in mg/dl)				
Cholesterol	146.71±10.7 ^a	173.00±23.5 ^b	164.29±16.4 ^b	152.57±24.0 ^a
LDL-C	12.57±6.3 ^a	14.00±1.6 ^{a,b}	10.14±4.5 ^{a,c}	9.57±2.8 ^{a,c}
HDL-C	72.29±7.9 ^a	85.14±7.4 ^b	82.00±11.9 ^b	77.00±11.4 ^a
HDL:LDL	6.7±2.4 ^a	6.2±0.5 ^{a,b}	9.3±3.2 ^{a,c}	8.8±3.3 ^{a,c}
Glucose	372.86±112.4 ^a	299±77.9 ^a	249.4±89.9 ^b	226.7±120.6 ^b
Liver (all parameters in mg/g)				
Cholesterol	2.1312±0.5×10 ⁻³ ^a	2.1320±0.9×10 ⁻³ ^a	2.1297±0.3×10 ⁻³ ^b	2.1299±0.8×10 ⁻³ ^b
Triglycerides	0.444±0.05 ^a	0.716±0.17 ^b	0.343±0.09 ^c	0.450±0.08 ^{a,c}
Epididymal WAT (all parameters in mg/g)				
Cholesterol	2.1306±0.9×10 ⁻³ ^a	2.1298±0.5×10 ⁻³ ^a	2.1282±0.2×10 ⁻⁶ ^b	2.1282±0.3×10 ⁻⁶ ^b
Triglycerides	0.035±0.0 ^a	0.029±0.0 ^a	0.045±0.02 ^a	0.047±0.02 ^a

Columns not sharing common alphabets are significantly different (p<0.01)

List of Figures

Figure 1. Effect of Ax or/and SQ on antioxidant enzymes and molecules in liver (L) and epididymal white adipose tissue (WAT). *LHpx-Lipid hydroperoxides (pmol DPPP/g lipid), Sod-Superoxide dismutase activity (inhibition rate %), Cat-Catalase (Formaldehyde nmol/min/g), Gpx-Glutathione peroxidase (nmol/min/ml), GSSG/GSH-Ratio of oxidised glutathione and reduced glutathione. Groups not sharing a common alphabet within a parameter are significantly different ($p < 0.05$).

Figure 2. Effect of Ax or/and SQ on mRNA expression of genes related to antioxidant enzymes in liver. Sod-Superoxide dismutase, Cat-catalase, Gpx-Glutathione peroxidase. Groups not sharing a common alphabet within a parameter are significantly different ($p < 0.05$).

Figure 3. Effect of Ax or/and SQ on mRNA expression of genes related to immunomodulatory pathways in liver. TNF α -Tumor necrosis factor α , PPAR α -Peroxisome proliferator-activated receptor α , Cox-1-Cyclooxygenase 1, Cox-2-Cyclooxygenase 2, MCP-1- Monocyte chemoattractant protein 1, IFN-Interferon γ , IL-1 β -Interleukin 1 β , IL-6-Interleukin 6, PPAR γ -Peroxisome proliferator-activated receptor γ . Groups not sharing a common alphabet within a parameter are significantly different ($p < 0.05$).

Figure 1.

Figure 2.

Figure 3.