Title	The mitochondrial genomes of <i>Pecten albicans</i> and <i>Pecten maximus</i> (Bivalvia: Pectinidae) reveal a novel gene arrangement with low genetic differentiation
Author(s)	Marín, Alan; Fujimoto, Takafumi; Arai, Katsutoshi
Citation	Biochemical Systematics and Ecology, 61, 208-217 https://doi.org/10.1016/j.bse.2015.06.015
Issue Date	2015-08
Doc URL	http://hdl.handle.net/2115/66898
Rights	© 2015. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	all.pdf

[Instructions for use](#)

1 **The mitochondrial genomes of *Pecten albicans* and**
2 ***Pecten maximus* (Bivalvia: Pectinidae) reveal a novel**
3 **gene arrangement with low genetic differentiation**

4

5 Alan Marín* • Takafumi Fujimoto² • Katsutoshi Arai³

6 *Hokkaido University, Graduate School of Fisheries Sciences, 3-1-1 Minato,*
7 *Hakodate, Hokkaido, 041-8611, Japan.*

8

9 *Correspondent author

10 e-mail: marin@fish.hokudai.ac.jp

11 Tel.: +81(90)6444 1955

12 ²fujimoto@fish.hokudai.ac.jp

13 ³arai@fish.hokudai.ac.jp

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28 **Abstract**

29 Scallops (Bivalvia: Pectinidae) comprise more than 350 extant taxa including
30 species of high economic importance. However, its phylogenetic classification
31 remains unclear and so far only 7 scallop mitogenomes have been determined. In
32 this study, the mitochondrial genomes of two congeneric scallop species *Pecten*
33 *albicans* and *Pecten maximus* were determined. Both mitogenomes contain 12
34 protein-coding genes (*atp8* is missing), two ribosomal genes, 20-22 transfer *RNA*
35 genes, all encoded in the same strand. Overall, both *Pecten* mitogenomes are
36 highly similar with high nucleotide (93 %) and amino acid (98.7 %) sequence
37 identity. Both mitogenomes also contain the same gene order arrangement,
38 which is a novel contribution within Pectinidae. The highly similar mitochondrial
39 organization between both *Pecten* species suggested a recent speciation event.
40 Phylogenetic analysis based on complete protein-coding gene information as
41 well as large syntenic gene blocks confirmed the sister group relationship
42 between the genera *Pecten* (subfamily Pectininae) and *Argopecten* (tribe
43 Aequipectini).

44

45 **Keywords** • Mitochondrial genome • Pectinidae • Synteny • Phylogenetics •
46 Amino acids • Scallops

47

48

49

50

1. Introduction

Metazoan mitochondrial genome is typically built of double-stranded circular DNA molecules of about 15-17 kb and usually encodes for 37 genes: 13 protein-coding genes (PCGs), 2 ribosomal RNA genes and 22 transfer *RNAs* that are indispensable for protein translation (Boore, 1999; Liu et al., 2013). Mitochondrial genes are considered important tools for evolutionary and phylogenetic studies because of its high mutation rates, maternal transmission (with exceptions in some animal groups) and the abundance of mitochondrial copies in cells. In most metazoan taxa, gene order is usually well conserved. However, marine bivalves are a notable exception to the common metazoan mitochondrial constitution. Marine bivalve mitogenomes are known for their high levels of gene rearrangement (Dreyer and Steiner, 2006; Yuan et al., 2012), even among species of the same family (i.e. Pectinidae; Xu et al., 2010). An extra large mitogenome size is another unusual characteristic of marine bivalves. From all Pectinidae species with available mitogenome, *Placopecten magellanicus* has the largest size (32115 bp; GenBank access number DQ088274). The presence of tandem repeats elements and transposition involving *tRNAs* or *tRNA*-like structures are responsible for the large size and length variation (30-40 kb) among individuals (La Roche et al., 1990; Smith and Snyder, 2007). Another common characteristic in most bivalve mitogenomes is the lack of *atp8* gene and certain *tRNAs* (Smith and Snyder, 2007).

The family Pectinidae comprises about 350 living species (Biscotti et al., 2007). In spite of the large fossil record, the phylogenetic relationships among scallop species are still unresolved. With 15 recognized species (World Register

75 of Marine Species, WORMS; <http://www.marinespecies.org>), scallops of the
76 genus *Pecten* represent commercially important species distributed throughout
77 temperate and sub-tropical seas in distant geographic areas such as Europe,
78 Africa, Asia and Australia (Duncan and Wilson, 2012; Saavedra and Peña, 2004).
79 *Pecten albicans* and *Pecten maximus* constitute important fisheries (Saavedra and
80 Peña, 2004) as well as interesting evolutionary models due to their unknown
81 divergence times. The Japanese baking scallop *P. albicans* occurs from southern
82 Hokkaido to Kyushu in Japan (Okutani, 2000). On the other hand, *P. maximus* is
83 distributed along the Northeast Atlantic from Norway down to North Africa
84 (Beaumont and Gjedrem, 2007). To date, little is known about the mitochondrial
85 genome composition of *Pecten* species and few studies have addressed the
86 extremely low genetic differentiation among species of this genus, using only
87 partial mitochondrial gene fragments (Saavedra and Peña, 2004). It has been
88 shown that phylogenetic analysis based on complete mitogenome sequence data
89 have proved to produce much robust phylogenetic reconstructions enhancing
90 resolution and statistical confidence of inferred phylogenetic trees when
91 compared with analyses based only on smaller portions of the mtDNA (Ingman
92 et al., 2000 and Yuan et al., 2012).

93 Before this study, only seven complete mitogenomes from the family
94 Pectinidae were available in GenBank database including 5 species from 2
95 subfamilies named Chlamydinae (*Mizuhopecten yessoensis*, *Chlamys farreri*,
96 *Mimachlamys nobilis* and *Mimachlamys senatoria*) and Palliolinae: (*P.*
97 *magellanicus*), and two species from the tribe Aequipectini (*Argopecten irradians*
98 and *Argopecten purpuratus*). This is the first report on the complete

mitochondrial coding region and gene rearrangement of two Pectininae species: *P. albicans* and *P. maximus*. The results of the present study will help to get further insights into the very complex taxonomy and evolutionary relationships among Pectinidae species.

2. Materials and Methods

2.1. Sample collection and genomic DNA extraction

One individual of *P. albicans* and *P. maximus* were collected from Oki Island in Japan and Rye Bay in England, respectively. The adductor muscles were dissected and preserved in 95% ethanol and stored at -20 °C. Genomic DNA was extracted following the standard phenol-chloroform protocol and adjusted to a concentration of 50 ng/μl and used for PCR.

2.2. Long PCR amplification and sequencing

Two long PCR primer sets were designed to amplify the entire mitogenome in two fragments with overlapping segments, covering the complete circular mitochondrial genome. The primer sets 12SpectUF and PecnaXX (amplified fragments of 9922 and 9932 bp in *P. albicans* and *P. maximus*, respectively) and Pectco3F and Pect12SR (amplified fragments of approx. 13 and 15 kb in *P. albicans* and *P. maximus*, respectively) were designed based on multi-alignment comparisons of the mitochondrial 12S rRNA, *cox3* and *nad2* genes from Pectinidae species retrieved from GenBank database (primer sequences are listed in Table 1). Large PCR products were obtained using TaKaRa LA *Taq* polymerase (TaKaRa, Japan). To quicken the sequencing process, additional shorter overlapping PCR fragments (2-6 kb) were amplified (see Table 1 for

primer list) using the large PCR fragments (10-15kb) as DNA templates. Full length of both scallop mitogenomes were estimated based in the nucleotide size content of the first large PCR product amplified by primers 12SpectUF and PecnaXX (which was fully sequenced) plus the combination of overlapping large and small PCR fragments from the "second half" of the mitogenome, which were estimated based on comparison with molecular ladders of different sizes by agarose electrophoresis. All PCR amplifications were performed on an ASTEC PC816 thermal cycler in a total volume of 50 µl, containing 26.5 µl sterile H₂O, 5 µl 10 × LA PCR Taq buffer (Mg²⁺ free, TaKaRa), 5 µl MgCl₂ (25mM), 8 µl dNTPs (2.5mM each), 2 µl each primer (10 µM), 0.5 µl LA *Taq* polymerase (5 units/µl, TaKaRa) and 1 µl of template DNA. Thermal cycling conditions were as follows: initial denaturation for 2 min at 98 °C, followed by 35 cycles of denaturation for 20 s at 94 °C, annealing for 35 s at 60 °C, and extension for 10 min at 68 °C, followed by a final extension for 10 min at 72 °C. Amplicons were electrophoresed in 1% agarose gels, stained with ethidium-bromide, and visualized under UV light. Subsequently, PCR products were extracted and purified using the QIAquick Gel Extraction kit (Quiagen, Hilden, Germany), directly sequenced in both directions by primer walking method on an ABI PRISM 3130XL Genetic Analyzer (Applied Biosystems, Hitachi, Japan).

The largest non-coding region of the *P. albicans* and *P. maximus* mitogenomes is characterized for the presence of long tandems repeats (1.6 kb approx; Rigaa et al., 1995) and homopolymer runs, which affected the PCR reaction and DNA sequencing process by polymerase slippage. Regrettably, a fragment containing the control region was not determined (approx. 6 and 7.5 kb

in size in *P. albicans* and *P. maximus*, respectively). Thus, only the complete mitochondrial coding region is reported for both species.

Additionally, to identify the gene cluster *nad1-rrnL-cox1* in more scallop species, a highly conserved fragment including partial sequence of genes *nad1* and *cox1* flanking the complete genes *rrnL*, *trnR* and *trnM* were amplified and sequenced in six additional scallop species including *Amusium pleuronectes*, *Ylistrum japonicum*, *Annachlamys macassarensis*, *Bractechlamys vexillum*, *Decatopecten radula* and *Mimachlamys sanguinea*. We followed same strategy as Marín et al. (2015), using a combination of degenerated and specific primers.

2.3. Sequence assembling and gene identification

The mitogenome sequences were assembled manually using MEGA 5.2 software (Tamura et al., 2011). The mitochondrial genomes were annotated using MITOS web service (Bernt et al., 2013). PCGs were identified with ORF Finder (NCBI website) using invertebrate mitochondrial code. PCGs denoted by putative ORFs were then identified by sequence similarity with BLASTP (NCBI website). Ribosomal RNA genes were identified by multi-alignment with GenBank sequences of bivalve mitogenomes. The *tRNA* genes were identified by tRNAScan-SE 1.21 Search Server (Lowe and Eddy 1997), using the invertebrate mitochondrial codon sequence, MITOS annotation web service or using secondary structures and anticodons when necessary. The whole mitochondrial genome sequences were tested for potentially tandem repeats by TANDEM REPEAT FINDER, Version 4.0 (Benson 1999). Mitochondrial genomes were drawn using GenomeVx (Conant and Wolfe 2008). Nucleotide and amino acid sequence identities were estimated as *p*-distances using MEGA 5.2.

2.4. Phylogenetic analysis

The complete PCGs from the available mitogenomes of 7 scallop species along with the two newly developed mitogenomes were used to infer their phylogenetic relationships. The oysters *Pinctada maxima* (NC_018752) and *Crassostrea gigas* (EU672831) were used as outgroups in all the analyses. The translated amino acid sequences of 12 PCGs were aligned separately with ClustalW. Poorly aligned positions and gaps were excluded using Gblocks (Castresana, 2000) as implemented in SeaView v. 4.5.3 (Gouy et al., 2010) and concatenated to construct phylogenetic trees. The nucleotide sequence was substituted from the concatenated amino acid alignment. Substitution saturations in single codon positions from each PCGs were analyzed using DAMBE 5 (Xia, 2013). The third codon positions of the genes *atp6*, *cytb*, *cox2*, *nad1*, *nad2*, *nad4L*, *nad5*, and *nad6* showed clear signs of saturation. Thus, two dataset were used for phylogenetic analyses: nucleotide sequence of 12 PCGs including all codon positions (PCG123) and nucleotide sequence of 12 PCGs excluding the third codon positions (PCG12). Two phylogenetic methods were performed including maximum likelihood (ML) with MEGA 5.2 and Bayesian inference (BI) with MRBAYES 3 (Ronquist and Huelsenbeck, 2003). For the ML analysis, the best fit of substitution model was determined in MEGA 5.2 using Bayesian Information Criterion (BIC). The GTR+G model was selected for the PCG123 and PCG12 datasets. Node reliability was tested using 1000 bootstrap replicates. For the Bayesian approach, we used jModelTest 2 (Darriba et al., 2012) under the Akaike Information Criterion (AIC) to find the best-fit model of evolution. For both datasets (PCG123 and PCG12) BI analysis was performed

under the GTR model of nucleotide substitution with a gamma-distributed rate variation across sites and a proportion of invariable sites. Two runs were performed simultaneously, each with four Markov chains. The analyses were run for 1,000,000 generations with sampling every 100 generations. The first 25 % of the sampled trees were discarded as burn-in. All phylogenetic trees were drawn using Figtree 1.4.2 program (<http://tree.bio.ed.ac.uk/software/figtree/>).

3. Results

3.1. Genome composition

The mitochondrial genomes of *P. albicans* (GenBank accession KP900974) and *P. maximus* (GenBank accession KP900975) are 22.6 and 24.8 kb in length, respectively (Figure 1). As aforementioned, due to the presence of long tandem repeats within the control region of both *Pecten* species, partial mitogenomes containing the complete protein coding regions were obtained in *P. albicans* (16653 bp) and *P. maximus* (17252 bp). Excluding the main control region, the overall nucleotide sequence identity between both *Pecten* species is 93 %, with 1,089 variable sites. The structure and organization of *P. albicans* and *P. maximus* mitogenomes showed characteristics of a typical marine bivalve mitogenome: 12 PCGs (*atp6*, *cox1* to *cox3*, *cytb*, *nad1* to *nad6*, *nad4L*), two ribosomal RNA genes, 20 (*P. albicans*) to 22 (*P. maximus*) transfer RNAs and the missing *atp8* gene. In both *Pecten* species all genes are encoded in the same strand with two overlapping genes (17 nt between *nad4L* and *cox3*). In *P. albicans* the genes *nad2* and *nad3* are also overlapping by one nucleotide.

3.2. Protein-coding genes

Twelve PCGs were identified in both *Pecten* mitogenomes (Table 2). Overall, the complete length of PCGs in *P. albicans* (11403 nt) and *P. maximus* (11397 nt) accounted for 50.5% and 46.0% of their whole genome, respectively. Total A+T content from complete PCG sequences was 59.4 and 59.8 in *P. albicans* and *P. maximus*, respectively. Excluding the termination codons, a total of 3780 and 3787 amino acids are encoded by *P. albicans* and *P. maximus* mitogenomes, respectively. The amino acid frequencies are quite similar between both *Pecten* species. For both species, the upmost used amino acids are Leucine (12.9%), Valine (11.0%), Serine (10.3%), Glycine (9.9-10.0%) and Phenylalanine (9.7-9.8%). The overall amino acid sequence identity between *P. albicans* and *P. maximus* is 98.7%. The pairwise amino acid identity for homologous genes ranged from 97.2% for *cox3* to 100% for *nad4* and *nad4L* genes (Table 2). The most common eukaryotic genome start codon ATN was found in more than half of all PCGs of both species (*atp6*, *cox1*, *cox2*, *cox3*, *nad4*, *nad5*, *nad4L*, and *nad3* only in *P. albicans*), whereas the alternative triplets GTG and TTG are used as start codons for the genes *nad1* and *nad2* and the genes *nad6* and *cytb*, respectively. The triplet TTG was also used by *P. maximus* as start codon of the gene *nad3*. Fifty percent of the open reading frames in *P. albicans* and *P. maximus* mitogenomes end with a TAG stop codon. The other 50% ends with a TAA stop codon. No incomplete (T-, TA-) stop codons were found.

3.3. rRNA and tRNA genes

In *P. albicans* and *P. maximus*, the length of the *rrnS* and *rrnL* genes were 965 and 1423 bp and 965 and 1421 bp, respectively. In both species, the *rrnS* gene is located between *trnA* and *trnQ* genes, and the *rrnL* gene between *trnR*

and *cox1* genes. Sequence identity for *rrnS* and *rrnL* genes between *P. albicans* and *P. maximus* is 98 % and 96.7 %, respectively. In *P. albicans* only 20 transfer RNAs were identified. Sixteen *tRNAs* were identified with tRNAScan-SE and 3 *tRNAs* (*trnR*, *trnS1* and *trnS2*) using MITOS annotation web service. The *trnL1* and *trnM* were not found because they might be located in the beginning of the control region, which was not sequenced. The mitogenome of *P. maximus* encodes a total of 22 *tRNA* genes, 19 *tRNAs* were identified with tRNAScan-SE and 3 *tRNAs* (*trnM*, *trnS1* and *trnS2*) using MITOS annotation web service. The *tRNAs* in both *Pecten* species ranged from 63 (*trnL2*) to 71 nucleotides (*trnQ* and *trnR*). In both scallop species two *trnS*s were detected. On the other hand, a second *trnL* was identified only in the beginning of the control region of *P. maximus*. The inferred secondary structures for the two mitogenomes showed that most *tRNAs* formed typical cloverleaf secondary structures (Additional file S1). However, in some *tRNAs* non Watson-Crick matches, unpaired nucleotides and aberrant loops were found. One non Watson-Crick base pair (C-C) was detected in the D-arm of the *trnR* of *P. maximus*. Unpaired nucleotides were detected in the C-arm of *trnD* from both *P. albicans* and *P. maximus*, as well as the acceptor stem of *trnM* in *P. maximus*. Finally, the genes *trnL2* and *trnQ* from both *Pecten* species formed a non-canonical oversized 9 nt anticodon loop.

3.4. Gene arrangement

The mitochondrial gene order of *P. albicans* and *P. maximus* displayed the same gene order arrangement, except for last two *tRNAs* identified in *P. maximus* that could not be found in *P. albicans* due to unconcluded sequencing of the region downstream the gene *trnT*. The mitochondrial gene order revealed in this

study constitutes a novel gene arrangement among metazoans. Two highly conserved gene clusters (*nad6-trnL-cytb* and *nad1-rrnL-cox1*) were detected in most scallop species with available mitogenomes (see Fig. 2), except for *P. magellanicus* (subfamily Palliolinae). The sequences spanning the 3' end of the *nad1* through 5' end of the *cox1* gene were determined in 6 different scallop species (GenBank access numbers from KP900976 to KP900981). Depending on species, the insertion of *trnR* between the genes *nad1* and *rrnL*, and *trnM* between the genes *rrnL* and *cox1* were detected (Fig. 2). Overall, gene cluster *nad1-rrnL-cox1* is present in the mitogenomes of 13 scallop species, whereas the gene cluster *nad6-trnL-cytb* was detected in a total of 8 scallop species. The mitogenomes of the genera *Argopecten* (tribe Aequipectini) and *Pecten* (subfamily Pectininae) shared two large similar gene clusters *rrnS-trnQ-trnV-nad1-trnR-rrnL-cox1-trnaF-trnE-nad6-trnL-cytb-cox2-nad4L-cox3-nad3-nad4* and *trnH-trnW-trnP-nad5-atp6-trnC-trnY-trnT-trnL-trnM*, with just few gene insertion and deletion differences (Fig. 3). At least 9 rearrangement events were identified between the mitogenomes of *A. purpuratus* and *P. maximus* including two *trna* gene insertions, one *trna* gene deletion, and the translocation of four *trna* genes, one PCG and the control region (Fig. 3).

3.5. Phylogenetic analysis

Results based on nucleotide sequences deduced from amino acid information from all 12 PCGs are shown in Fig. 4. Both phylogeny estimation approaches (ML and BI) inferred with two datasets (PCG123 and PCG12) showed very similar topologies, similar branch lengths and high bootstrap support and posterior probabilities. The 9 scallop species were recovered in 3 clear

monophyletic groups. The sister taxa *Pecten* (subfamily Pectininae) and *Argopecten* (tribe Aequipectini) scattered in two subclades within a larger basal clade. Within the subfamily Chlamydinae, two discrete subclades (*M. nobilis* and *M. senatoria*) + (*C. farreri* and *M. yessoensis*) were recovered. Finally, *P. magellanicus* (subfamily Palliolinae) split in a single branch after the Chlamydinae clade.

4. Discussion

In this study, two new mitogenomes belonging to the subfamily Pectininae were revealed, including a novel gene arrangement. The complete mitochondrial genome lengths of both *Pecten* species were estimated based on their PCR amplicon sizes. Previously, Rigaa et al. (1995) described the presence of a tandem repeat unit of 1.6 kb occurring from two to five times within the control region of *P. maximus*. Accordingly, a mitogenome of 20 and 24.8 kb should contain two and five repeat units, respectively. Taking into consideration the results by Rigaa et al. (1995), the complete mitogenome of the *P. maximus* analyzed in this study (24.8 kb in length), should contain five repeat units, which explains the large mitochondrial molecule size. The increased mitochondrial size caused by the presence of large tandem repeats within the control region is a common feature among scallops (Gjetvaj et al., 1992; Rigaa et al., 1995). Large mitochondrial genomes have been found in *M. yessoensis* (20964 bp; Wu et al., 2009), *P. magellanicus* (32115 bp; Smith and Snyder, 2007), *Crassodoma gigantea* 22.8-24.8 kb, *Aequipecten opercularis* 21-28.2 kb, *Chlamys hastata* 23.9-27.2 kb, *C. islandica* 22.2-25 kb (Gjetvaj et al., 1992). To date, all described scallop mitogenomes encode all PCGs in one strand. Placement of all coding

genes on the same strand and lacking of *atp8* gene are one of the most distinctive characteristics of marine bivalve mitogenomes (He et al., 2011). Besides, it is believed that coding on both strands may inhibit mitogenome rearrangement, and marine bivalve mitogenomes are known for showing high levels of gene rearrangement (Ren et al., 2010a). Lack of the *atp8* gene is another common signature in marine bivalve mitogenomes. Indeed, the *atp8* gene is absent or highly modified in different animal groups including Chaetognatha, Rotifera, most Mollusca Bivalvia, Nematoda and Platyhelminthes (Gissi et al., 2008). Its high nucleotide variability together with its short and variable length might have hampered the annotation of the *atp8* gene in many studies (Gissi et al., 2008; Smietanka et al., 2010). Thus, "putative" *atp8* gene has been reported in some bivalve species such as *Mytilus* spp (Breton et al., 2010; Smietanka et al., 2010)

Overall, both *Pecten* mitogenomes showed similar start codons. About 40 % of the PCGs from *Pecten* mitogenomes used alternative start codons GTG or TTG. The non-conventional start codons GTG and TTG are not usual in molluscan mitogenomes, however their use have been reported in other scallop species (Marín et al., 2014; Smith and Snyder, 2007; Wu et al., 2009) and in many gastropods (Yuan et al., 2012). Duplication of transfer RNAs is common in mollusks (Ren et al., 2010b) and total *tRNAs* number identified in Pectinidae varies depending on species, ranging from 16 in *M. yessoensis* (Wu et al., 2009) to 32 in *P. magellanicus* (Smith and Snyder 2007). In *P. maximus* and *P. albicans* two *trnS*s were detected. On the contrary, lack of *trnS* has been reported in some scallop species including *C. farreri*, *M. nobilis* and *M. yessoensis* (Wu et al., 2009),

337 *A. irradians irradians* (Ren et al., 2010b), and *A. purpuratus* (Marín et al., 2014).
338 Previous mollusk mitogenome studies have revealed that among the *tRNA* genes,
339 *trnS* is the least conserved (Ren et al., 2010b).

340 The small conserved gene block *nad4-trnH-trnW* was observed in *C.*
341 *farreri*, *Mimachlamys* spp. (Chlamydinae), *P. magellanicus* (Palliolinae) and
342 *Pecten* spp. (Pectininae), and the gene block *trnL-cytb-cox2* in *P. magellanicus*
343 (Palliolinae), *Pecten* spp. and *Argopecten* spp. (Pectininae) (Fig. 2). Additionally,
344 as described by Ren et al. (2010b), the gene order rearrangement of *M.*
345 *yessoensis* closely resembles those of *Chlamys farreri* and *Mimachlamys* spp., all
346 belonging to the subfamily Chlamydinae (Fig. 2). Three conserved gene clusters
347 (*nad4L-nad6-trnL-cytb*, *trnD-cox3-trnK-trnF-trnQ-trnE-atp6-cox2-nad2*, and
348 *nad1-trnR-rrnL-trnM*) are shared among these three genera. Similarly, the
349 synapomorphy of two large gene clusters identified in our results (*rrnS-trnQ-*
350 *trnV-nad1-trnR-rrnL-cox1-trnaF-trnE-nad6-trnL-cytb-cox2-nad4L-cox3-nad3-*
351 *nad4* and *trnH-trnW-trnP-nad5-atp6-trnC-trnY-trnT-trnL-trnM*) between genera
352 *Argopecten* and *Pecten*, supports the sister relationship between tribe
353 Aequipectini (genus *Argopecten*) and subfamily Pectininae (genus *Pecten*) (Fig.
354 3). The usefulness of gene rearrangements for phylogenetic reconstructions of
355 distant Pectinidae groups might be limited due to the drastic variability of
356 mitochondrial gene order in this family. Nevertheless, our results confirm that
357 Pectinidae gene rearrangement can be used to determine and support
358 phylogenetic relationships among closer groups. In this study, additional scallop
359 species mitogenomes were found to contain the highly conserved gene clusters
360 reported by Ren et al. (2010b): *nad1-rrnL-cox1* (now detected in 13 species; Fig.

2) and *nad6-trnL-cytb* (now detected in 8 species, Fig. 2). These conserved blocks may have been inherited from a common ancient scallop ancestor (Ren et al., 2010b) and therefore could be used as molecular markers for a deeper understanding of mitochondrial genome evolution in members of Pectinidae.

At least 9 rearrangement events occurred between the mitogenomes of *A. purpuratus* and *P. maximus*. Because of the relatively complex rearrangement events, their evolutionary pathways of gene rearrangements cannot be explained by one rearrangement mechanism alone, instead it might be a result of a combination of more mechanisms such as tandem duplication and random loss (TDRL) model (Boore, 1999), transposition (Macey et al., 1997) and intramolecular recombination (Lunt and Hyman 1997). Comparisons among mitogenomes at lower taxonomic levels (e.g., within genera) are highly useful for further interpretation and understanding of mitochondrial gene rearrangement mechanisms (Gissi et al., 2008). Thus, the analysis of additional mitogenomes from species with intermediate stages can be useful to trace missing mitochondrial evolutionary pathways in Pectinidae. Interestingly, most of the rearrangement events occurred in genes flanking the control region of *Argopecten* and *Pecten* species (Fig. 3). This result suggests that the control region in the family Pectinidae is a potential hotspot for gene rearrangement. In metazoan mitogenomes, the control region neighborhood has been recognized as one of the gene rearrangement hotspots (Kurabayashi et al., 2008).

Phylogenetic relationships in Pectinidae are still a matter of debate, due to the incongruent results between genetic and morphological studies. In this study, the concatenation of 12 PCGs allowed the recovery of strong supported

phylogeny based on two different phylogenetic approaches (ML and BI). Overall, our results strongly supported the basal position and sister group relationship between *Argopecten* and *Pecten*, and the apical position of Chlamydinae and Palliolinae subfamilies. The groups formed by Aequipectini + Pectinidae and Chlamys + Palliolinae have been inferred as the most ancient and recently derived groups from the family Pectinidae, respectively (Puslednik et al., 2008). Our results strongly supported the existence of the subfamily Palliolinae, which is in agreement with previous studies (Puslednik et al., 2008; Saavedra and Peña, 2006; Waller, 2006). However, our phylogenetic results recovered the subfamily Palliolinae in the tip of the tree, just after the subfamily Chlamydinae. In this regard, our results slightly differed from previous studies. For example, Puslednik et al. (2008) used partial mitochondrial and nuclear genes to infer the phylogenetic relationships of 46 scallop species. In the same study, members of the subfamily Palliolinae (covered by 3 species) were recovered in a monophyletic group between Chlamydinae and Pectininae clades. In this study, we used the only available mitochondrial genome from Palliolinae (*P. magellanicus*). It has been shown that the inclusion of dense taxa dataset results in more accurate estimation of evolutionary relationships (Heath et al., 2008). Thus, the inclusion of complete mitogenomes from more representatives of the subfamily Palliolinae will provide an accurate estimation of the phylogenetic position of this subfamily within Pectinidae.

Pecten species are characterized for a broad geographic range distribution, inhabiting the coasts of four continents. However, in spite of the clear allopatric populations, very low mitochondrial genetic distances (ranging

from 0.006 to 0.042) have been reported for members of this genus (Peña and Saavedra, 2012). The estimated genetic distance between *P. albicans* and *P. maximus* mitogenomes are in agreement with previous results by Peña and Saavedra (2012). Relatively low genetic distance (7 %) and quite similar amino acid sequence identity (98.7 %) found between the mitogenomes of *P. albicans* and *P. maximus* suggest a recent divergence event. Further studies using complete mitochondrial genome information from other *Pecten* species are needed to corroborate the recent diversification within this genus. Recently diverged species are challenging for identification (van Velzen et al., 2012). Previous studies have demonstrated the great utility of the mitochondrial 16S gene for scallop species identification (Marín et al., 2013a, 2015). Interestingly, our results showed that the genes with the highest pairwise sequence diversities were *nad1* ($p=0.079$) and *nad6* ($p=0.081$). Thus, the *nad1* and *nad6* genes can be also considered as alternative potential markers in further identification and population studies of *Pecten* species, which are characterized for showing extremely low genetic variation. Finally, our results will also be of great utility to clarify the possible conspecificity between *P. maximus* and *P. jacobaeus* (Canapa et al., 2000).

Acknowledgments

We want to thank Prof. Kazuo Inaba (University of Tsukuba, Japan), Dr. Kenta Suda (Okabe Co. Ltd., Japan), Ms. Shelah Buen-Ursua (SEAFDEC/AQD, Philippines) and Ms. Mika Yoshimura (Hokkaido University, Japan) for providing scallop samples. A.M. was supported by the Japanese Government Scholarship-Ministry of Education, Culture, Sports, Science and Technology (MEXT, Monbukagakusho) for doctoral studies.

Figures

Fig. 1. Circular and linearized mitochondrial genome organization of *Pecten albicans* and *Pecten maximus*. All genes are encoded in the same strand. Protein-coding genes are marked in blue (gene name abbreviations: *atp6*, ATP synthase subunit 6; *cytb*, apocytochrome *b*; *cox1-3*, cytochrome *c* oxidase subunits; *nad1-4*, *4L*, *5-6*, NADH dehydrogenase subunits), ribosomal genes in red (gene name abbreviations: *rrnL*, *rrnS*, large and small subunit rRNAs), *tRNAs* (abbreviated using universal amino acid code) in yellow, and control region in grey.

Fig. 2. Synteny among the linear-mapping mitogenomes of 12 scallop species, including the novel *Pecten maximus* mitogenome developed in this study. All mitogenomes are opened downstream of control region for easier comparison. Species inside pink box belong to the subfamily Chlamydinae (*Mimachlamys senatoria* is omitted because of its high similarity with *Mimachlamys nobilis*), white box represents subfamily Palliolinae, purple box represents subfamily

455 Pectininae (*Pecten albicans* is not included because of its high similarity with
 456 *Pecten maximus*), and green box the tribe Aequipectini (*Argopecten irradians* is
 457 not included because of its high similarity with *Argopecten purpuratus*).
 458 Conserved gene cluster *nad1-rrnL-cox1* is represented in 6 additional species
 459 *Amusium pleuronectes*, *Ylistrum japonicum*, *Annachlamys macassarensis*,
 460 *Brachtechlamys vexillum*, *Decatopecten radula* and *Mimachlamys sanguinea*. Black
 461 horizontal bars indicate the conserved gene cluster “*nad1-rrnL-cox1*”. Blue bars
 462 indicate the conserved gene cluster “*nad6-L-cytb*”. Orange and red bars represent
 463 the conserved gene clusters “*L2-cytb-cox2*” and “*nad4-H-W*”, respectively.

464 **Fig. 3.** Linear-mapping mitogenomes rearrangement comparison between
 465 *Pecten maximus* and *Argopecten purpuratus*. Protein-coding genes are marked in
 466 blue, ribosomal genes in red, *tRNAs* (abbreviated using universal amino acid
 467 code) in yellow, and control region in grey. Large conserved gene blocks are
 468 located inside black line rectangles. The blue stars represent gene insertions, red
 469 stars represent gene deletions and black lines indicate six gene translocations.

470 **Fig. 4.** Maximum likelihood (A) and Bayesian Inference (B) phylogenetic
 471 consensus trees constructed based on the concatenation of 12 mitochondrial
 472 protein-coding genes in 9 scallop species. The two species with newly
 473 determined mitogenomes developed in this study (*Pecten albicans* and *Pecten*
 474 *maximus*, subfamily Pectininae) are highlighted in bold. The conserved gene
 475 cluster *nad1-trnR-rrnL-cox1* is represented by a black circle, *nad1-trnR-rrnL-*
 476 *trnM-cox1* is represented by a black triangle, *nad1-trnR-rrnL-trnM-rrnS-cox1* is
 477 represented by a black square (indicated in all species except for *Placopecten*
 478 *magellanicus*, where the gene cluster is absent). Bootstrap values (maximum

likelihood) and posterior probabilities (Bayesian inference) at correspondent nodes are shown in percentages. The oysters *Pinctada maxima* (NC_018752) and *Crassostrea gigas* (EU672831) were used as outgroups.

Table 1 Primer list

Table 2 Mitochondrial gene annotation

Supplemental Material Inferred secondary structure of *tRNAs* in *Pecten albicans* and *Pecten maximus*.

References

- Beaumont, A., Gjedrem, T., 2007. Scallops—*Pecten maximus* and *P. Jacobeus*. In: Svåsand, T., Crosetti, D., García-Vázquez, E., Verspoor, E., (eds) Genetic impact of aquaculture activities on native populations. Genimpact final scientific report (EU contract RICA-CT-2005-022802), pp 83–90.
URL <http://www.imr.no/genimpact/filarkiv/2007/07/scallops.pdf/en>
Accesed 15.02.19
- Benson, G., 1999. Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res. 27, 573-580.
- Bernt, M., Donath, A., Jühling, F., Externbrink, F., Florentz, C., Fritzsche, G., Pütz, J., Middendorf, M., Stadler, P.F., 2013. MITOS: improved *de novo* metazoan mitochondrial genome annotation. Mol. Phylogenet. Evol. 69, 313-319.
- Boore, J.L., 1999. Animal mitochondrial genomes. Nucleic Acids Res. 27, 1767-1780.
- Breton S., Stewart, D.T., Hoeh, W.R., 2010. Characterization of a mitochondrial ORF from the gender-associated mtDNAs of *Mytilus* spp. (Bivalvia: Mytilidae): identification of the “missing” ATPase 8 gene. Mar Genomics 3, 11–18.
- Canapa, A., Barucca, M., Marinelli, A., Olmo, E., 2000. Molecular data from the 16S rRNA gene for the phylogeny of Pectinidae (Mollusca: Bivalvia). J. Mol. Evol. 50, 93–97.
- Castresana, J., 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Mol. Biol. Evol. 17, 540-552.
- Conant, G.C., Wolfe, K.H., 2008. GenomeVx: simple web-based creation of editable circular chromosome maps. Bioinformatics 24, 861-862.

522 Biscotti, M.A., Canapa, A., Olmo, E., Barucca, M., Teo, C.H., Schwarzacher, T.,
523 Dennerlein, S., Richter, R., Heslop-Harrison, J.S., 2007. Repetitive DNA,
524 molecular cytogenetics and genome organization in the King scallop (*Pecten*
525 *maximus*). *Gene* 406, 91–98.

526 Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2012. jModelTest 2: more models,
527 new heuristics and parallel computing. *Nature Methods* 9, 772.

528 Dreyer, H., Steiner, G., 2006. The complete sequences and gene organization of
529 the mitochondrial genomes of the heterodont bivalves *Acanthocardia*
530 *tuberculata* and *Hiatella arctica* and the first record for a putative Atpase
531 subunit 8 gene in marine bivalves. *Front. Zool.* 3, 13.

532 Duncan, P., Wilson, G., 2012. A new species of *Pecten* (Mollusca: Bivalvia:
533 Pectinidae) from northern Western Australia. *Molluscan Res.* 31, 21–26.

534 Gissi, C., Iannelli, F., Pesole, G., 2008. Evolution of the mitochondrial genome of
535 Metazoa as exemplified by comparison of congeneric species. *Heredity* 101,
536 301–320.

537 Gjetvåg, B., Cook, D.I., Zouros, E., 1992. Repeated sequences and large-scale size
538 variation of mitochondrial DNA: a common feature among scallops (Bivalvia:
539 Pectinidae). *Mol. Biol. Evol.* 9, 106–124.

540 Gouy, M., Guindon, S., Gascuel, O., 2010. SeaView version 4: a multiplatform
541 graphical user interface for sequence alignment and phylogenetic tree
542 building. *Mol. Biol. Evol.* 27, 221–224.

543 He, C.B., Wang, J., Gao, X.G., Song, W.T., Li, H.J., Li, Y.F., Liu, W.D., Su, H. 2011. The
544 complete mitochondrial genome of the hard clam *Meretrix meretrix*. Mol. Biol.
545 Rep. 38, 3401–3409.

546 Heath, T.A., Hedtke, S.M., Hillis, D.M., 2008. Taxon sampling and the accuracy of
547 phylogenetic analyses. J. Syst. Evol., 46, 239–257.

548 Ingman, M., Kaessmann, H., Paabo, S., Gyllensten, U., 2000. Mitochondrial genome
549 variation and the origin of modern humans. Nature 408, 708–712.

550 Kurabayashi, A., Sumida, M., Yonekawa, H., Glaw, F., Vences, M., Hasegawa, M.,
551 2008. Phylogeny, recombination, and mechanisms of stepwise mitochondrial
552 genome reorganization in mantellid frogs from Madagascar. Mol. Biol. Evol. 25,
553 874–891.

554 La Roche, J., Snyder, M., Cook, D.I., Fuller, K., Zouros, E., 1990. Molecular
555 characterization of a repeat element causing large-scale size variation in the
556 mitochondrial DNA of the sea scallop *Placopecten magellanicus*. Mol. Biol. Evol.
557 7, 45–64.

558 Liu, Y.G., Kurokawa, T., Sekino, M., Tanabe T., Watanabe, K., 2013. Complete
559 mitochondrial DNA sequence of the ark shell *Scapharca broughtonii*: An ultra-
560 large metazoan mitochondrial genome. Comp. Biochem. Physiol. D. 8, 72–81.

561 Lowe, T.M., Eddy, S.R., 1997. A program for improved detection of transfer RNA
562 genes in genomic sequence. Nucleic Acids Res. 25, 955–964.

563 Lunt, D.H., Hyman, B.C., 1997. Animal mitochondrial DNA recombination.
564 Nature 387, 247.

565 Macey, J.R, Larson, A., Ananjeva, N.B., Fang, Z., Papenfuss, T.J., 1997. Two novel
566 gene orders and the role of light-strand replication in rearrangement of the
567 vertebrate mitochondrial genome. *Mol. Biol. Evol.* 14, 91-104.

568 Marín, A., Fujimoto, T., Arai, K., 2013a. Rapid species identification of fresh and
569 processed scallops by multiplex PCR. *Food Control* 32, 472-476.

570 Marín, A., Fujimoto, T., Arai, K., 2013b. Genetic structure of the Peruvian scallop
571 *Argopecten purpuratus* inferred from mitochondrial and nuclear DNA
572 variation. *Mar. Genomics*, 9, 1-8.

573 Marín, A., Alfaro, R., Fujimoto, T., Arai, K. 2014. Mitochondrial genome of the
574 Peruvian scallop *Argopecten purpuratus* (Bivalvia: Pectinidae). Mitochondrial
575 DNA. DOI: 10.3109/19401736.2013.845760

576 Marín, A., Fujimoto, T., Arai, K., 2015. The variable 5' end of the 16S rRNA gene as
577 a novel barcoding tool for scallops (Bivalvia, Pectinidae). *Fish. Sci.* 81, 73-81.

578 Okutani, T., (ed.) 2000. Marine mollusks in Japan. Tokai University
579 Press, Tokyo, 1175 pp.

580 Peña, J.B., Saavedra, C., 2012. Pectínidos de las aguas de Castellón: faunística,
581 taxonomía y evolución. *Nemus. Revista de l'Ateneu de Natura* 2, 71-82.

582 Puslednik, L., Serb, J.M., 2008. Molecular phylogenetics of the Pectinidae
583 (Mollusca: Bivalvia) and effect of increased taxon sampling and outgroup
584 selection on tree topology. *Mol. Phylogenet. Evol.* 48, 1178–1188.

585 Ren, J.F., Liu, X., Zhang, G.F., Liu, B., Guo, X.M., 2010a. Unusual conservation of
586 mitochondrial gene order in *Crassostrea* oysters: evidence for recent
587 speciation in Asia. *BMC Evol. Biol.* 10, 394.

588 Ren, J., Shen, X., Jiang, F., Liu, B., 2010b. The mitochondrial genomes of two
589 scallops, *Argopecten irradians* and *Chlamys farreri* (Mollusca: Bivalvia): the
590 most highly rearranged gene order in the family Pectinidae. J. Mol. Evol., 70,
591 57–68.

592 Rigaa, A., Monnerot, M., Sellos, D., 1995. Molecular cloning and complete
593 nucleotide sequence of the repeated unit and flanking gene of the scallop
594 *Pecten maximus* mitochondrial DNA: putative replication origin features. J. Mol.
595 Evol. 41, 189–195.

596 Ronquist, F., Huelsenbeck, J.P., 2003. MRBAYES 3: Bayesian phylogenetic
597 inference under mixed models. Bioinformatics 19, 1572-1574.

598 Saavedra, C., Peña, J.B., 2004. Phylogenetic relationship of commercial European
599 and Australasian king scallops (*Pecten* spp.) based on partial 16S ribosomal
600 RNA gene sequences. Aquaculture 235, 153–166.

601 Saavedra, C., Peña, J.B., 2006. Phylogenetics of American scallops (Bivalvia:
602 Pectinidae) based on partial 16S and 12S ribosomal RNA gene sequences. Mar.
603 Biol. 150, 111–119.

604 Smietanka, B., Burzynski, A., Wenne, R., 2010. Comparative genomics of marine
605 mussels (*Mytilus* spp.) gender associated mtDNA: Rapidly evolving atp8. J. Mol.
606 Evol. 71, 385–400.

607 Smith, D.R., Snyder, M., 2007. Complete mitochondrial DNA sequence of the
608 scallop *Placopecten magellanicus*: Evidence of transposition leading to an
609 uncharacteristically large mitochondrial genome. J. Mol. Evol. 65, 380-391.

610 Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011.
611 MEGA5: molecular evolutionary genetics analysis using maximum likelihood,
612 evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28,
613 2731-2739.

614 Van Velzen, R., Weitschek, E., Felici, G., Bakker, F.T., 2012. DNA barcoding of
615 recently diverged species: relative performance of matching methods. *PLoS*
616 *One* 7:e30490.

617 Waller, T.R., 2006. New phylogenies of the Pectinidae (Mollusca: Bivalvia):
618 reconciling morphological and molecular approaches. In: Shumway SE,
619 Parsons GJ (eds) *Scallops: biology, ecology and aquaculture*. Elsevier BV,
620 Amsterdam, The Netherlands, pp 1–44.

621 Wu, X., Xu, X., Yu, Z., Kong, X., 2009. Comparative mitogenomic analyses of three
622 scallops(Bivalvia: Pectinidae) reveal high level variation of genomic
623 organization and a diversity of transfer RNA gene sets. *BMC Res. Notes* 2, 69.

624 Xia, X., 2013. DAMBE5: A comprehensive software package for data analysis in
625 molecular biology and evolution. *Mol. Biol. Evol.* 30, 1720-1728.

626 Xu, K.F., Kanno, M., Yu, H., Li, Q., Kijima, A., 2010. Complete mitochondrial DNA
627 sequence and phylogenetic analysis of Zhikong scallop *Chlamys farreri*
628 (Bivalvia: Pectinidae). *Mol. Biol. Rep.* 38, 3067–3074.

629 Yuan, Y., Li, Q., Yu, H., Kong, L., 2012. The complete mitochondrial genomes of six
630 heterodont bivalves (Tellinoidea and Solenoidea): variable gene
631 arrangements and phylogenetic implications. *PLoS ONE* 7, e32353.

Pecten maximus

Argopecten purpuratus

★ Gene insertion

★ Gene deletion

└┐ Gene translocation

A

B

Table 1

List of universal and *Pecten*-specific primers used to amplify the complete mitochondrial genomes of *Pecten albicans* (PA) and *Pecten maximus* (PM). Long PCR products were used as templates in all subsequent amplifications of short PCR products.

Primer name	Sequence 5'-3'	Direction	Species	Gene	Source
LONG PCR					
12SpectUF	AGTCCAACCAGGTGCCAGCA	Forward	PA/PM	<i>rrnS</i>	Marín et al. (2013a)
PecnaXX	CCCCTGCAAACATAAAGAATTCAAC	Reverse	PA/PM	<i>nad2</i>	This study
Pectco3F	TGGCCTCAGGAGATTTATACAATACC	Forward	PA/PM	<i>cox3</i>	This study
Pect12SR	CAACAAGCTGACAACGGCAATACACC	Reverse	PA/PM	<i>rrnS</i>	This study
SHORT PCR					
12SpectUF	AGTCCAACCAGGTGCCAGCA	Forward	PA/PM	<i>rrnS</i>	Marín et al. (2013a)
CO1AB	GGTGCTGGGCAGCcayatncngg	Reverse	PA/PM	<i>cox1</i>	Marín et al. (2013b)
16SCC	GCGTAATCCGTCTTGACAGT	Forward	PA/PM	<i>rrnL</i>	Marín et al. (2013b)
PecnaXX	CCCCTGCAAACATAAAGAATTCAAC	Reverse	PA/PM	<i>nad2</i>	This study
Cox3F1	TGGCCTCAGGAGATTTATACAATACC	Forward	PA/PM	<i>cox3</i>	This study
Nad4R1	CCAGTATGGGCAATAGAAGAGTA	Reverse	PA	<i>nad4</i>	This study
Nad5R1	AAGAACTGTGGCAAGGATAGAACAA	Reverse	PM	<i>nad5</i>	This study
Nad3F2	GAGGGTTTTTTAGTGGTTCTTG	Forward	PA	<i>nad3</i>	This study
TrnpR2	CCCAATGTACTAACTTATACTACCC	Reverse	PA	<i>trnP</i>	This study
Nad5F2	GCGGTGGCTTTGGCTGGTGATACTT	Forward	PM	<i>nad5</i>	This study
Atp6R2	CCCCTCATTGTTCAATCAAACCTG	Reverse	PM	<i>atp6</i>	This study
TrnHF3	TGGTGTAGAAAGGCACGAAAGGTTG	Forward	PA	<i>trnH</i>	This study
CRpa3	ATACTACCCTGCTCAAACAAGTGG	Reverse	PA	<i>CR</i>	This study
Nad5F3	TTCCGTGCCATGTTAGGAAGGCATT	Forward	PM	<i>nad5</i>	This study
CRpm3	AGGCTCCACTGACCTACCTATAAAC	Reverse	PM	<i>CR</i>	This study
CRpm4	GGTCAGTGGAGCCTCTTTAGAGCTA	Forward	PA/PM	<i>CR</i>	This study
Pect12SR	CAACAAGCTGACAACGGCAATACACC	Reverse	PA/PM	<i>rrnS</i>	This study

Table 2

Complete mitochondrial annotation, number of amino acids in each gene, pairwise amino acid and nucleotide identity based on the mitochondrial genomes of *Pecten albicans* (PA) and *Pecten maximus* (PM).

Gene	Location		Size (bp)		Start codon		Stop codon		Anticodon		Intergenic nucleotides		No. Amino acid		% Amino acid identity PA/PM	% Nucleotide identity PA/PM
	PA	PM	PA	PM	PA	PM	PA	PM	PA	PM	PA	PM	PA	PM		
<i>trnG</i>	1-66	1-66	66	66					TCC	TCC	-	-				
<i>trnN</i>	75-141	75-141	67	67					GTT	GTT	8	8				
<i>trnI</i>	170-239	168-237	70	70					GAU	GAU	28	26				
<i>trnK</i>	269-337	267-335	69	69					TTT	TTT	29	29				
<i>trnA</i>	349-414	348-413	66	66					TGC	TGC	11	12				
<i>rrnS</i>	450-1414	447-1411	965	965							35	33				98
<i>trnQ</i>	1430-1500	1427-1497	71	71					TTG	TTG	16	15				
<i>trnV</i>	1527-1593	1524-1590	67	67					TAC	TAC	26	26				
<i>nad1</i>	1597-2544	1594-2541	948	948	GTG	GTG	TAA	TAA			3	3	315	315	99	92.1
<i>trnR</i>	2551-2621	2544-2615	71	72					TCG	TCG	6	2				
<i>rrnL</i>	2665-4087	2662-4082	1423	1421							43	46				96.7
<i>cox1</i>	4106-5716	4101-5711	1611	1611	ATG	ATG	TAA	TAA			18	18	536	536	99.1	93.2
<i>trnF</i>	5749-5812	5743-5806	64	64					GAA	GAA	32	31				
<i>trnS2</i>	5830-5895	5823-5888	66	66					TGA	TGA	17	16				
<i>trnE</i>	5913-5978	5906-5971	66	66					TTC	TTC	17	17				
<i>nad6</i>	6204-6731	6188-6715	528	528	TTG	TTG	TAG	TAG			225	216	175	175	97.7	91.9
<i>trnL2</i>	6734-6796	6718-6780	63	63					TAA	TAA	2	2				
<i>cob</i>	6800-8110	6784-8094	1311	1311	TTG	TTG	TAA	TAA			3	3	436	436	99.1	93.3
<i>cox2</i>	8224-8919	8208-8903	696	696	ATG	ATG	TAG	TAG			113	113	231	231	98.3	93.7
<i>trnD</i>	8944-9011	8927-8994	68	68					GTC	GTC	24	23				
<i>nad4L</i>	9051-9344	9034-9327	294	294	ATA	ATG	TAA	TAA			39	39	97	97	100	92.2
<i>cox3</i>	9328-10194	9311-10177	867	867	ATG	ATG	TAG	TAG			-17	-17	288	288	97.2	93.3
<i>nad2</i>	10200-11162	10183-11145	963	963	GTG	GTG	TAA	TAA			5	5	320	320	98.4	93.3
<i>nad3</i>	11162-11521	11151-11504	360	354	ATA	TTG	TAG	TAG			-1	5	119	117	98.3	95.8
<i>trnS1</i>	11615-11681	11511-11577	67	67					TCT	GCT	93	6				
<i>nad4</i>	11722-12972	11707-12957	1251	1251	ATG	ATG	TAG	TAG			40	129	416	416	100	93.4
<i>trnH</i>	12985-13050	12970-13035	66	66					GTG	GTG	12	12				
<i>trnW</i>	13061-13127	13046-13112	67	67					CCA	CCA	10	10				
<i>trnP</i>	13146-13210	13131-13195	65	65					TGG	TGG	18	18				
<i>nad5</i>	13226-15013	13212-14999	1788	1788	ATG	ATG	TAG	TAG			15	16	595	595	98.7	94.2
<i>atp6</i>	15020-15805	15006-15791	786	786	ATA	ATA	TAA	TAA			6	6	261	261	98.1	95.3
<i>trnC</i>	15824-15889	15810-15875	66	66					GCA	GCA	18	18				
<i>trnY</i>	15900-15965	15886-15951	66	66					GTA	GTA	10	10				
<i>trnT</i>	15979-16046	15965-16032	68	68					TGT	TGT	13	13				
<i>trnL1</i>	-	17051-17120	-	70					-	TAG		1018				
<i>trnM</i>	-	17122-17193	-	72						CAU		1				

bp: base pairs.

**The mitochondrial genomes of *Pecten albicans* and *P. maximus* (Bivalvia: Pectinidae)
reveal a novel gene arrangement with low genetic differentiation**

Biochemical Systematics and Ecology

Alan Marín* • Takafumi Fujimoto² • Katsutoshi Arai³

Hokkaido University, Graduate School of Fisheries Sciences, 3-1-1 Minato, Hakodate, Hokkaido, 041-8611, Japan.

*Correspondent author

e-mail: marin@fish.hokudai.ac.jp

Supplementary 1a. Inferred secondary structure of the identified transfer RNAs in *Pecten albicans*. Blue and red circles represent U-A or U-G and C-G pairings, respectively

Supplementary 1a. (Continued)

Tyr
Y

Val
V

Supplementary 1b. (Continued)