Title	Genetic and antigenic characterization of H5, H6 and H9 avian influenza viruses circulating in live bird markets with intervention in the center part of Vietnam
Author(s)	Duc-Huy Chu; Okamatsu, Masatoshi; Matsuno, Keita; Hiono, Takahiro; Ogasawara, Kohei; Lam Thanh Nguyen; Long Van Nguyen; Tien Ngoc Nguyen; Thuy Thu Nguyen; Dong Van Pham; Dang Hoang Nguyen; Tho Dang Nguyen; Thanh Long To; Hung Van Nguyen; Kida, Hiroshi; Sakoda, Yoshihiro
Citation	Veterinary Microbiology, 192, 194-203 https://doi.org/10.1016/j.vetmic.2016.07.016
Issue Date	2016-08-31
Doc URL	http://hdl.handle.net/2115/67102
Rights	©2016, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Manuscript_Table1-5_Fig.1-3.pdf

[Instructions for use](#)

1 **Genetic and antigenic characterization of H5, H6 and H9 avian influenza viruses circulating**
2 **in live bird markets with intervention in the center part of Vietnam**

3
4 Duc-Huy Chu^{a,b}, Masatoshi Okamatsu^a, Keita Matsuno^{a,c}, Takahiro Hiono^a, Kohei Ogasawara^a,
5 Lam Thanh Nguyen^a, Long Van Nguyen^b, Tien Ngoc Nguyen^b, Thuy Thu Nguyen^b, Dong Van
6 Pham^b, Dang Hoang Nguyen^b, Tho Dang Nguyen^b, Thanh Long To^b, Hung Van Nguyen^e, Hiroshi
7 Kida^{c,d}, Yoshihiro Sakoda^{a,c,*}

8
9 ^a*Laboratory of Microbiology, Department of Disease Control, Graduate School of Veterinary*
10 *Medicine, Hokkaido University, Sapporo, Hokkaido 060-0818, Japan*

11 ^b*Department of Animal Health, Ministry of Agriculture and Rural Development, 15/78 Giai Phong,*
12 *Phuong Mai, Dong Da, Hanoi, Vietnam*

13 ^c*Global Station for Zoonosis Control, Global Institution for Collaborative Research and Education*
14 *(GI-CoRE), Hokkaido University, Sapporo, Hokkaido 001-0020, Japan*

15 ^d*Research Center for Zoonosis Control, Hokkaido University, Sapporo, Hokkaido 001-0020, Japan*

16 ^e*Sub-Department of Animal Health, 62 Nguyen Chi Dieu, Hue city, Vietnam*

17
18 *Corresponding author: Yoshihiro Sakoda

19 *Laboratory of Microbiology, Department of Disease Control, Graduate School of Veterinary*
20 *Medicine, Hokkaido University, Sapporo, Hokkaido 060-0818, Japan*

21 Tel: +81-11-706-5207; Fax: +81-11-706-5273

22 E-mail: sakoda@vetmed.hokudai.ac.jp

23

24 **Abstract**

25 A total of 3,045 environmental samples and oropharyngeal and cloacal swabs from apparently
26 healthy poultry have been collected at three live bird markets (LBMs) at which practices were
27 applied to reduce avian influenza (AI) virus transmission (intervention LBMs) and six conventional
28 LBMs (non-intervention LBMs) in Thua Thien Hue province in 2014 to evaluate the efficacy of the
29 intervention LBMs. The 178 AI viruses, including H3 (19 viruses), H4 (2), H5 (8), H6 (30), H9
30 (114), and H11 (5), were isolated from domestic ducks, muscovy ducks, chickens, and the
31 environment. The prevalence of AI viruses in intervention LBMs (6.1%; 95% CI: 5.0 to 7.5) was
32 similar to that in non-intervention LBMs (5.6%; 95% CI: 4.5 to 6.8; $\chi^2 = 0.532$; $df = 1$; $P = 0.53$) in
33 the study area. Eight H5N6 highly pathogenic avian influenza (HPAI) viruses were isolated from
34 apparently healthy ducks, muscovy ducks, and an environmental sample in an intervention LBM.
35 The hemagglutinin genes of the H5N6 HPAI viruses belonged to the genetic clade 2.3.4.4, and the
36 antigenicity of the H5N6 HPAI viruses differed from the H5N1 HPAI viruses previously circulating
37 in Vietnam. Phylogenetic and antigenic analyses of the H6 and H9 viruses isolated in both types of
38 LBMs revealed that they were closely related to the viruses isolated from domestic birds in China,
39 Group II of H6 viruses and Y280 lineage of H9 viruses. These results indicate that the interventions
40 currently applied in LBMs are insufficient to control AI. A risk analysis should be conducted to
41 identify the key factors contributing to AI virus prevalence in intervention LBMs.

42

43 **Key words:** avian influenza; antigenic analysis; phylogenetic analysis; live bird market;
44 surveillance

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63 **1. Introduction**

64 Transmission of avian influenza (AI) viruses among wild and domestic birds is an important
65 target of control measures aiming to minimize the risk to both human and animal health worldwide
66 (Kilpatrick et al., 2006; Nomura et al., 2012; Okamatsu et al., 2013). Vietnam has a large
67 population of poultry (approximately 308 million), and a majority of these animals (approximately
68 80%) are raised under backyard conditions at households in rural areas without biosecurity
69 application (General Statistics Office of Vietnam. Results of the 2012 Rural, Agricultural and
70 Fishery Census. https://www.gso.gov.vn/default_en.aspx?tabid=778; Hanh et al., 2007). In Vietnam,
71 H5N1 highly pathogenic avian influenza (HPAI) viruses have caused a large number of outbreaks
72 in poultry since 2003 (Minh et al., 2009). In our previous study, we conducted surveillance of AI
73 viruses in live bird markets (LBMs) and households that raise poultry in a large number of
74 Vietnamese provinces in 2009. We found that the prevalence of AI viruses was substantially higher
75 in LBMs than in backyard farms, and H5N1 viruses were isolated from apparently healthy domestic
76 ducks and chickens in LBMs (Okamatsu et al., 2013; Nomura et al., 2012). The findings suggest
77 that in Vietnam, LBMs are more responsible for the amplification, maintenance, circulation, and
78 transmission of AI viruses than backyard farms. Therefore, continuous surveillance of AI viruses in
79 LBMs in Vietnam is essential to understand the distribution of AI viruses and to minimize the risk
80 to public and animal health.

81 LBMs are ubiquitous and integral parts of the poultry industry in Vietnam and other
82 developing countries in Asia (Wan et al., 2011; Indriani et al., 2010). In China, LBMs have become

83 a major source of human infection with H5 HPAI viruses and H7 low pathogenic AI viruses (He et
84 al., 2014; Yu et al., 2014; Wan et al., 2011). When keeping live birds overnight in LBMs was
85 banned, the rate of virus isolation declined (Leung et al., 2012). In addition, it was reported that
86 closing LBMs effectively reduced the risk of virus transmission to the public (Fournié et al., 2011;
87 He et al., 2014; Yu et al., 2014). However, banning LBMs in developing countries remains
88 challenging because changing the traditional market style should take a long time. Therefore,
89 government intervention to improve the biosecurity measures employed at LBMs has been thought
90 to represent a promising strategy to minimize the transmission of viruses in Asian countries
91 including Vietnam. To support for the evaluation of local authority on the current intervention, we
92 surveyed the prevalence of AI viruses at nine LBMs with or without the intervention of Vietnam
93 Avian and Human Influenza Control and Preparedness Project (VAHIP) in Thua Thien Hue
94 province, located in the central region of Vietnam. The VAHIP was a project funded by the World
95 Bank aiming to reduce the risk of AI virus transmission to humans (The World Bank, Projects and
96 Operations. Vietnam - Avian and Human Influenza Control and Preparedness Project.
97 [http://documents.worldbank.org/curated/en/2011/03/14026433/vietnam-additional-financing-](http://documents.worldbank.org/curated/en/2011/03/14026433/vietnam-additional-financing-vietnam-avian-human-influenza-control-preparedness-project-environmental-impact-assessment)
98 [vietnam-avian-human-influenza-control-preparedness-project-environmental-impact-assessment.](http://documents.worldbank.org/curated/en/2011/03/14026433/vietnam-additional-financing-vietnam-avian-human-influenza-control-preparedness-project-environmental-impact-assessment)).
99 The project developed the new model of LBM in selected provinces in Vietnam by deploying
100 infrastructure for the poultry markets and operating periodic disinfection in these intervention
101 LBMs. In this study, various subtypes of AI viruses were isolated during surveillance of LBMs with
102 or without intervention in August and December, 2014. The representative isolates were

103 phylogenetically and antigenically analyzed to characterize the genetic and antigenic variation of
104 the AI viruses currently circulating in LBMs in Vietnam.

105

106 **2. Materials and methods**

107 *2.1. Sample collection*

108 The surveillance was conducted in August and December, 2014. Oropharyngeal, cloacal swabs
109 and fecal samples from domestic birds and water troughs (environmental samples) were collected at
110 nine LBMs (Supplemental Fig. 1). At three of the nine LBMs, a biosecurity infrastructure had been
111 established by the VAHIP program in Thua Thien Hue province, Vietnam (intervention LBMs;
112 Table 1; Supplemental Fig. 1). The other six LBMs were conventional markets at which no
113 particular biosecurity infrastructure was established and at which poultry and other animals were
114 usually mixed together in low biosecurity conditions (non-intervention LBMs; Table 1;
115 Supplemental Fig. 1).

116 All collected samples were stored in sterile tubes with transport medium (minimum essential
117 medium, Nissui, Japan) containing 10,000 U/ml penicillin G (Meiji Seika, Japan), 10 mg/ml
118 streptomycin (Meiji Seika, Japan), 0.3 mg/ml gentamicin (Schering Plough, USA), 250 U/ml
119 nystatin (Sigma, USA), and 0.5% bovine serum albumin fraction V (Roche, Switzerland) at -80°C
120 until use.

121

122 *2.2. Isolation and identification of AI viruses*

123 Samples were resuspended in virus transport medium and centrifuged at 2,000 rpm for 5 min.
124 The supernatant was inoculated into the allantoic cavity of a 10-day-old chicken embryo. After
125 incubation at 35°C for 30–48 h, allantoic fluids exhibiting hemagglutination activity were collected
126 for subtyping of influenza viruses by hemagglutination-inhibition (HI) and neuraminidase-
127 inhibition tests with antisera to the reference influenza virus strains (Kida et al., 1979).

128

129 *2.3. Sequencing and phylogenetic analysis*

130 Viral RNA was extracted from the 250 µl of allantoic fluids by TRIzol LS Reagent (Life
131 Technologies, USA) following the manufacturer's protocol and reverse transcribed with the Uni12
132 primer (Hoffmann et al., 2001) and M-MLV Reverse Transcriptase (Life Technologies, USA). Full-
133 length cDNAs of the eight gene segments were amplified by polymerase chain reaction with Ex-
134 Taq (TaKaRa, Shiga, Japan) and gene-specific primer sets (Hoffmann et al., 2001). Direct
135 sequencing of each gene segment was performed using 3500 Genetic Analyzer (Life Technologies,
136 USA).

137 For phylogenetic analysis, nucleotide sequences of the isolates, together with those from a
138 public database, were aligned using Clustal W. Phylogenetic trees were constructed using the
139 maximum-likelihood (ML) method with 1,000 bootstrap replicates using MEGA 5.0 software
140 (Tamura et al., 2011). The genome sequences identified in this study have been registered in
141 GenBank/EMBL/DDBJ (Table 2).

142

143 2.4. Antigenic analysis

144 Polyclonal antisera were prepared from chickens immunized with reference AI virus strains
145 that had been inactivated with formalin (Kida et al., 1979). Antigenic analysis of H5, H6 and H9
146 viruses was performed using polyclonal antisera by HI test.

147

148 2.5. Pathogenicity of an H5N6 AI virus in chickens

149 To assess the pathogenicity of the representative H5N6 virus in chickens, 0.2 ml of the 1:10-
150 diluted fresh allantoic fluid of chicken embryos infected with A/duck/Vietnam/HU1-1151/2014
151 (H5N6) was inoculated intravenously into four 7-week-old chickens (*Gallus gallus*). Each chicken
152 was housed in a self-contained isolator unit (Tokiwa Kagaku, Japan) in a BSL3 biosafety facility in
153 the Graduate School of Veterinary Medicine, Hokkaido University, Japan.

154

155 2.6. Knowledge, attitude, and practices survey

156 The KAP survey was designed to investigate the knowledge, attitude, and practice of
157 individual poultry sellers that associated with AI in LBMs using questionnaires written in
158 Vietnamese. A self-designed questionnaire set comprised 47 questions about the general
159 background of the sellers, the source of poultry, and application of any biosecurity measures for the
160 identification of possible risk factors contributing to AI virus circulation in LBMs.

161

162 2.7. Ethics statements

163 Animal experiments were authorized by the Institutional Animal Care and Use Committee of
164 Hokkaido University (approval number: 13-0108), and all experiments were performed according to
165 the guidelines of the committee.

166

167 **3. Results**

168 *3.1. Identification of AI viruses circulating in intervention and non-intervention LBMs*

169 A total of 178 viruses were identified from 3,045 cloacal and oropharyngeal samples of
170 domestic birds and environmental samples (Table 1). In total, 19 H3, 2 H4, 8 H5, 30 H6, 114 H9,
171 and 5 H11 AI viruses were isolated from samples collected at nine LBMs. At the individual market
172 level, the prevalence of AI viruses in No market was the highest (12.2%; 95% CI: 9.6 to 15.4) in the
173 group of intervention LBMs as well as that of Phu Bai market (13.3%; 95% CI: 9.9 to 17.6) in the
174 group of non-intervention LBMs. At the type of LBMs level, the prevalence of AI viruses in
175 intervention LBMs (6.1%; 95% CI: 5.0 to 7.5) was similar to that in non-intervention LBMs (5.6%;
176 95% CI: 4.5 to 6.8; $\chi^2 = 0.532$; $df = 1$; $P = 0.53$). The subtypes of AI viruses isolated in LBMs with
177 intervention were H3N2 (7), H3N6 (1), H4N6 (2), H6N2 (10), H6N6 (9), H9N2 (49), H9N6 (2),
178 H11N6 (1), and H11N7 (3). The subtypes of AI viruses isolated in non-intervention LBMs were
179 H3N2 (11), H6N2 (4), H6N6 (7), H9N2 (60), H9N6 (3), and H11N7 (1). Eight H5N6 viruses were
180 isolated from apparently healthy ducks (5), muscovy ducks (2), and an environmental sample at the
181 “No” market with intervention. To assess the pathogenicity of the H5N6 viruses, a representative

182 H5N6 virus, A/duck/Vietnam/HU1-1151/2014 (H5N6) was inoculated intravenously into four 7-
183 week-old chickens. All chickens died within 1 day of infection.

184

185 3.2. Genetic and antigenic analysis of H5 viruses

186 The full-length nucleotide sequences of the HA genes of H5 representative isolates were
187 phylogenetically analyzed and divided into two lineages: Eurasian and North American. The H5
188 viruses in the Eurasian lineage were identified on the basis of the nomenclature defined by
189 WHO/OIE/FAO H5N1 Evolution Working Group in 2014 (Smith et al., 2015). The HA genes of
190 six representative H5N6 Vietnam isolates were classified into the clade 2.3.4.4, which was newly
191 established in 2014 (Smith et al., 2015) and is closely related to the
192 A/environment/Zhenjiang/C13/2013 (H5N6) virus isolated in China (Qi et al., 2014) (Fig. 1). All
193 the H5N6 isolates contain multiple basic amino acids, R and K, at the proteolytic cleavage site of
194 the hemagglutinin protein, PLREKRRKR/GLF, identical to the typical cleavage site motif of H5
195 HPAI viruses. The residues at position 190, 225, 226, and 228 (H3 numbering) are associated with
196 influenza virus receptor specificity. In these viruses, those residues were E, G, Q, and G,
197 respectively; all of them are of avian-type motif (Paulson et al., 2013).

198 Four representative strains of the H5 isolate were antigenically analyzed by cross HI test (Table
199 3). Chicken antiserum against the reference virus of the clade 2.3.4.4, A/chicken/Kumamoto/1-
200 7/2014 (H5N8) (Kanehira et al., 2015), effectively inhibited hemagglutination of A/muscovy
201 duck/Vietnam/HU1-1144/2014 (H5N6), A/duck/Vietnam/HU1-1151/2014 (H5N6), and A/muscovy

202 duck/Vietnam/HU2-26/2014 (H5N6) and partly inhibited hemagglutination of
203 A/environment/Vietnam/HU1-1434/2014 (H5N6) causing a 4-fold reduction in HI titer. In contrast,
204 these viruses exhibited low HI titers in comparison with homologous titers in reactions with the
205 antisera of different clades (HI titer from 20 to 80).

206

207 *3.3. Genetic and antigenic analysis of H6 viruses*

208 The H6 HA genes were phylogenetically divided into two lineages: Eurasian and North
209 American. The H6 viruses in the Eurasian lineage were clustered into five different sublineages:
210 Group II, W312, Group III, Early and Group I, as described in our previous study (Okamatsu et al.,
211 2013). All H6 viruses isolated in Vietnam in 2014 were found to belong to the Group II sublineage
212 (Fig. 2), as were the viruses previously isolated in Vietnam between 2010 and 2012 and those
213 isolated in China between 2003 and 2011. However, the H6 viruses isolated in this study and in our
214 previous study occupied different clusters.

215 Four strains representative of the H6 isolates were antigenically analyzed by cross HI test
216 (Table 4) using a panel of chicken antisera against four viruses of different sublineages. These H6
217 viruses weakly reacted with A/duck/Vietnam/OIE-4429/2010 (H6N2) antiserum, a virus of the
218 same genetic sublineage, Group II. In reaction with viruses of other sublineages, these H6 viruses
219 exhibited HI titers 4-fold lower than the homologous groups. In addition, these H6 viruses share
220 equivalent HI titers with homologous hyperimmune serum against A/duck/Vietnam/HU1-637/2014
221 (H6N6) virus, while A/duck/Vietnam/OIE-4429/2010 (H6N2) virus reacted weakly with the

222 antiserum with 16-fold lower HI titers. These results suggest that the antigenicity of the H6 viruses
223 we isolated differed from viruses isolated in Vietnam between 2010 and 2012 (Okamatsu et al.,
224 2013).

225

226 *3.4. Genetic and antigenic analysis of H9 viruses*

227 The HA genes of H9 isolates were phylogenetically divided into two lineages: Eurasian and
228 North American lineages. All H9 viruses isolated at LBMs in Thua Thien Hue province in 2014
229 were classified as Y280 sublineage of the Eurasian lineage (Okamatsu et al., 2013; Nomura et al.,
230 2012). These viruses were genetically related to the A/chicken/Vietnam/OIE-1611/2012 (H9N2)
231 virus isolated in the North Vietnam in 2012 and other viruses isolated from poultry in China in 1997
232 and 2012, which were also classified into the Y280 sublineage (Fig. 3).

233 Representative strains of the H9 isolates were antigenically analyzed by cross HI test (Table 5).
234 All viruses from Vietnam reacted with the antiserum of A/duck/Hong Kong/Y280/1997 (H9N2)
235 virus, which belonged to the Y280 sublineage. These Vietnamese H9N2 viruses reacted weakly
236 with antisera of other H9N2 viruses classified into different sublineages, such as the G1 or North
237 American lineages, and reacted moderately with antisera to Y349 sublineage virus. This suggests
238 that the antigenicities of the H9N2 viruses isolated in Vietnam have been stable in the poultry
239 population.

240

241 *3.5. The neuraminidase genes of AI viruses isolated in LBMs*

242 For neuraminidase (NA) gene segments, the names of groups were defined based on previous
243 studies (Kim et al., 2013; Okamatsu et al., 2013). The N2 NA genes of H6 and H9 AI viruses were
244 phylogenetically categorized into two groups, Group II and Y280 (Supplemental Fig. 2a). All N6
245 NA genes of the H5 viruses belonged to Group II and of H6 viruses belonged to Group I
246 (Supplemental Fig. 2b).

247

248 3.6. Genetic diversity of AI viruses isolated in LBMs

249 The six internal gene segments of the AI viruses were then phylogenetically analyzed to
250 investigate the genetic diversity of AI viruses circulating in Vietnamese LBMs (Supplemental Fig.
251 3). For each internal gene segment, the names of groups were defined as previously described (Kim
252 et al., 2013; Okamatsu et al., 2013). In this study, the six internal gene segments were also classified
253 into H6 Group I, H6 Group II, H6 Group III, JX8264 like, and Gs/Gd like, indicating that these
254 viruses were closely related to the viruses isolated in China. For instance, the PB2 and PA genes of
255 the A/muscovy duck/Vietnam/HU1-1144/2014 (H5N6) were classified as JX8264 like. The NP, M,
256 and NS genes of this H5N6 virus were grouped into Gs/Gd like and H6 Group I where Chinese
257 viruses were also grouped into. Phylogenetic analysis also indicated that the PB2, PB1, PA, and NS
258 internal genes of H9N2 viruses, A/chicken/Vietnam/HU1-1050/2014 and A/chicken/Vietnam/HU1-
259 976/2014, were classified into the same groups: Vietnam II, Vietnam, and JX8286 like. However,
260 other gene segments, NP and M genes, were classified into different groups (Supplemental Fig. 3).

261

262 3.7. *Knowledge, attitude, and practices*

263 A total of 52 sellers working at LBMs were interviewed during the first sampling collection, 29
264 sellers working at intervention LBMs and 23 working at non-intervention LBMs. All respondents
265 working at intervention LBMs declared that their shops were cleaned by chemical disinfection
266 twice per day before the LBM was opened and after closing their business. In contrast, only 17.4%
267 of respondents working at non-intervention LBMs made this claim. In intervention LBMs, 52.0% of
268 sellers acquired their poultry from within Thua Thien Hue province, within a 5 km radius. The other
269 48.0% of sellers imported their poultry from neighboring provinces, from within a radius of 50 km.
270 In contrast, 100% of the sellers working at non-intervention LBMs sell poultry that originated from
271 the local areas surrounding the LBMs.

272

273 4. Discussion

274 LBMs provide an ideal environment for genetic reassortment events and interspecies
275 transmission of AI viruses (He et al., 2014; Yu et al., 2014; Okamatsu et al., 2013; Wan et al., 2011).
276 In this study, 178 AI viruses were isolated from poultry in both intervention and non-intervention
277 LBMs. The H5N6 viruses were isolated from apparently healthy domestic ducks, muscovy ducks,
278 and an environmental sample at an intervention LBM, indicating that subclinical HPAI virus
279 infections are endemic in domestic birds at LBMs. These results remind us that LBMs play an
280 important role as a hotspot for the transmission of AI viruses, including HPAI viruses, in Vietnam.
281 Although closure of LBMs or banning the storage of poultry overnight at the markets represents a

282 highly effective method to reduce amplification and persistence of viruses in LBMs (He et al.,
283 2014; Yu et al., 2014; Leung et al., 2012; Fournié et al., 2011), it is hard to change the LBMs
284 system in Vietnam due to well-established traditional cultures and the behavior of customers.
285 Therefore, biosecurity measures were developed with the intention of reducing AI transmission. In
286 this study, we conducted surveillance of AI viruses in both intervention and non-intervention LBMs
287 in a central province of Vietnam. We sampled viruses on two occasions and found the AI virus
288 prevalence at intervention LBMs to be 6.1%. The prevalence of AI viruses at non-intervention
289 LBMs did not differ significantly from that at non-intervention LBMs, which was 5.6%. All the
290 viruses identified in this study were isolated from apparent healthy LBM poultry. In addition,
291 various subtypes of AI viruses were identified at intervention LBMs. The H6 and H9 viruses
292 isolated in intervention LBMs were genetically similar to H6 and H9 viruses isolated at non-
293 intervention LBMs. Although disinfection was reported to be performed twice per day at
294 intervention LBMs, AI viruses still contaminated the environment. These results indicate that the
295 current intervention LBMs remain several limitations for the control of AI contamination such as
296 problems of periodic disinfection procedure and concentration of chemical disinfection. The
297 intervention LBMs are generally larger capability to hold poultry than these of non-intervention
298 LBMs meaning that intervention LBMs can contain poultry originating from many different sources,
299 including neighboring provinces. In addition, KAP surveys indicated that the sources of poultry in
300 intervention LBMs in Thua Thien Hue province are widely distributed from local area to other
301 provinces. Therefore, the source of poultry may play an important role in AI transmission at LBMs,

302 and this factor should be further studied. As a next step, a risk analysis will be conducted to identify
303 the risk factors contributing to the appearance of various subtypes of AI viruses at intervention
304 LBMs.

305 The H5N1 HPAI viruses were first detected in Vietnam in 2001, and outbreaks have been
306 regularly reported since the end of 2003 (Minh et al., 2009). The H5 viruses circulating in Vietnam
307 were classified into seven major genetic clades: 1.1, 2.3.2.1, 2.3.4.1, 2.3.4.2, 2.3.4.3, 7.1, and 7.2
308 (Nguyen et al., 2014; Okamatsu et al., 2013; Nguyen et al., 2012). In this study, we report for the
309 first time the characterization of H5N6 viruses isolated in an intervention LBM in a center province
310 of Vietnam. The HA genes of the H5N6 viruses isolated from poultry and an environmental sample
311 at an intervention LBM in Thua Thien Hue province were classified as clade 2.3.4.4. Genetic
312 analysis indicated that these H5N6 viruses were closely related to an H5N6 virus isolated in
313 Zenjiang, China and differed slightly from the H5N6 viruses isolated in Laos, indicating an existing
314 genetic diversity within this new clade 2.3.4.4 (Food and Agriculture Organization of the United
315 Nations. Avian influenza A (H5N6): the latest addition to emerging zoonotic avian influenza threats
316 in East and Southeast Asia, 2014; www.fao.org/ag/empres.html). The H5 viruses found in northern
317 Vietnam may have been introduced from China. In April, 2014, the first outbreak of H5N6 was
318 found in a flock of chickens in Lang Son province, located near the border with China. Then
319 another outbreak was reported in a flock of ducks in Ha Tinh province, located in central part of
320 Vietnam and China (World Animal Health Information Database Interface, Disease Information,
321 2015; http://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?reportid=15802). In

322 Vietnam and China, AI vaccines are used to control H5 HPAI virus infections in poultry (Hu et al.,
323 2015; Le et al., 2014). However, immunological selection pressure has driven development of
324 antigenic variants of H5 HPAI viruses (Grund et al., 2011; Lee et al., 2004). The H5N6 HPAI
325 viruses we identified were also antigenically distinct from the clade 2.3.4 virus used in the Re-5
326 vaccine, which has been applied in China and Vietnam (Hu et al., 2015; Le et al., 2014). Recently,
327 several human H5N6 AI virus infections were reported, and avian-originated H5N6 viruses were
328 also isolated from healthy pigs in China, although the H5N6 AI virus has not adapted to the swine
329 population yet (Pan et al., 2016; Li et al., 2015). In Thua Thien Hue province, Vietnam, piglets, and
330 poultry are often housed together in non-intervention LBMs. This LBM system may facilitate
331 influenza virus reassortment and transmission of influenza viruses between poultry and from
332 poultry to pigs or humans. Therefore, it will be important to monitor swine influenza, and strict
333 controls should be applied to limit interspecies transmission of influenza viruses.

334 H6 and H9 AI viruses are widely distributed among poultry and wild birds in Asia (Huang et
335 al., 2010; Moon et al., 2010). H9N2 viruses have become the most prevalent subtype detected in
336 poultry populations in China since 2004 (Choi et al., 2004) and in Vietnam since 2009 (Okamatsu
337 et al., 2013; Nomura et al., 2012). In this study, H6 and H9 viruses isolated from domestic ducks,
338 chickens, and environment samples were genetically identical in both types of LBMs in Thua Thien
339 Hue province, Vietnam. These viruses were phylogenetically closely related to viruses previously
340 isolated in poultry in China. The HA genes of H9 viruses did not differ from those previously
341 isolated in 2011 and 2012, indicating that these viruses were maintained in poultry in Vietnam.

342 Antigenic analysis of H6 viruses indicated that all the Vietnamese H6 viruses we isolated in 2014
343 exhibited low cross reactivity with antiserum of A/duck/Vietnam/OIE-4429/2010 (H6N2), another
344 Group II virus. In addition, genetic analysis also indicated that the H6 viruses isolated in this study
345 differed from other Vietnamese H6 viruses isolated in our previous study (Okamatsu et al., 2013).
346 These results suggest that the antigenicity of these Vietnamese H6 viruses isolated in 2014 differed
347 from those isolated in Vietnam between 2010 and 2012. Perhaps the H6 viruses circulating in
348 Vietnam in 2014 exhibited altered antigenicity as a result of repetitive infections of the poultry
349 population. Further studies should be conducted to characterize the antigenic variation of H6 viruses
350 circulating in Vietnam.

351 In Vietnam, domestic birds are mainly raised in households in a free-range manner and are
352 transported to LBMs by their owners or poultry sellers. AI viruses are transmitted and spread within
353 the poultry population. Although domestic birds are vaccinated against H5N1 in Vietnam (Le et al.,
354 2014; Soares et al., 2010; Minh et al., 2009), HPAI viruses have silently spread in the poultry
355 population. Our study provides initial evidences for the improvement of intervention strategies in
356 study LBMs. The intervention with supporting a good infrastructure as “a good hardware” need to
357 comprise well with “a good software” like increasing of education level of people involved in
358 poultry trade and improvement of hygiene procedures by local authority at LBMs. Active
359 surveillance program of AI and hygiene practice performance monitoring in LBMs will be essential
360 to eradicate HPAI from Vietnam as well as Asian countries.

361

362 **Conflict of interest**

363 None.

364

365 **Acknowledgments**

366 We thank Dr. T. Saito, National Institute of Animal Health, Japan for kindly providing viruses
367 A/chicken/Ibaraki/1/2005 (H5N2), A/chicken/Yamaguchi/7/2004 (H5N1), and
368 A/chicken/Kumamoto/1-7/2014 (H5N8). We also thank Dr. Luk S.M. Geraldine of the Tai Lung
369 Veterinary Laboratory, Hong Kong SAR, China for providing A/peregrine falcon/Hong
370 Kong/810/2009 (H5N1) virus. This research was supported by Japan Society for the Promotion of
371 Science (JSPS) KAKENHI (grant number 16J06369 to D-H. Chu) by the Program for Leading
372 Graduate Schools from JSPS (grant number F01) and was partially funded by Japan Initiative for
373 Global Research Network on Infectious Diseases (J-GRID) from Japan Agency for Medical
374 Research and Development (AMED). D-H. Chu is supported by JSPS Research Fellowships for
375 young scientists.

376

377

378

379

380

381

382 **References**

- 383 Choi, Y.K., Ozaki, H., Webby, R.J., Webster, R.G., Peiris, J.S., Poon, L., Butt, C., Leung, Y.H.,
384 Guan, Y., 2004. Continuing evolution of H9N2 influenza viruses in Southeastern China. J.
385 Virol 78, 8609-8614.
- 386 Food and Agriculture Organization of the United Nations. Avian influenza A (H5N6): the latest
387 addition to emerging zoonotic avian influenza threats in East and Southeast Asia.
388 <http://www.fao.org/3/a-i4199e.pdf> (accessed 25.09.15).
- 389 Fournié, G., Guitian, F.J., Mangtani, P., Ghani, A.C., 2011. Impact of the implementation of rest
390 days in live bird markets on the dynamics of H5N1 highly pathogenic avian influenza. J. R.
391 Soc. Interface 8, 1079-1089.
- 392 General Statistics Office of Vietnam. Results of the 2012 Rural, Agricultural and Fishery Census.
393 https://www.gso.gov.vn/default_en.aspx?tabid=778 (accessed 25.09.15).
- 394 Grund, C., Abdelwhab, e.-S., Arafa, A.S., Ziller, M., Hassan, M.K., Aly, M.M., Hafez, H.M.,
395 Harder, T.C., Beer, M., 2011. Highly pathogenic avian influenza virus H5N1 from Egypt
396 escapes vaccine-induced immunity but confers clinical protection against a heterologous
397 clade 2.2.1 Egyptian isolate. Vaccine 29, 5567-5573.
- 398 Hanh PTH., Burgos S., Ronald-Holst D., The poultry sector in Viet Nam: prospects for smallholder
399 producers in the aftermath of the HPAI crisis. Pro-Poor Livestock Policy Initiative Research
400 Report 2007. [http://www.fao.org/ag/againfo/programmes/en/pplpi/docarc/rep-](http://www.fao.org/ag/againfo/programmes/en/pplpi/docarc/rep-hpai_vietnampoultryquality.pdf)
401 [hpai_vietnampoultryquality.pdf](http://www.fao.org/ag/againfo/programmes/en/pplpi/docarc/rep-hpai_vietnampoultryquality.pdf) (accessed 25.09.15).

402 He, Y., Liu, P., Tang, S., Chen, Y., Pei, E., Zhao, B., Ren, H., Li, J., Zhu, Y., Zhao, H., Pan, Q., Gu,
 403 B., Yuan, Z., Wu, F., 2014. Live poultry market closure and control of avian influenza
 404 A(H7N9), Shanghai, China. *Emerg. Infect. Dis.* 20, 1565-1566.

405 Hoffmann, E., Stech, J., Guan, Y., Webster, R.G., Perez, D.R., 2001. Universal primer set for the
 406 full-length amplification of all influenza A viruses. *Arch. Virol* 146, 2275-2289.

407 Hu T., Song J., Zhang W., Zhao H., Duan B., Liu Q., Zeng W., Qiu W., Chen G., Zhang Y., Fan Q.,
 408 Zhang F., 2015. Emergence of novel clade 2.3.4 influenza A (H5N1) virus subgroups in
 409 Yunnan Province, China. *Infect. Genet. Evol.* 33, 95-100.

410 Huang, K., Bahl, J., Fan, X.H., Vijaykrishna, D., Cheung, C.L., Webby, R.J., Webster, R.G., Chen,
 411 H., Smith, G.J., Peiris, J.S., Guan, Y., 2010. Establishment of an H6N2 influenza virus
 412 lineage in domestic ducks in southern China. *J. Virol.* 84, 6978-6986.

413 Indriani, R., Samaan, G., Gultom, A., Loth, L., Irianti, S., Indryani, S., Adjid, R., Dharmayanti,
 414 N.L., Weaver, J., Mumford, E., Lokuge, K., Kelly, P.M., Darminto, 2010. Environmental
 415 sampling for avian influenza virus A (H5N1) in live-bird markets, Indonesia. *Emerg. Infect.*
 416 *Dis.* 16, 1889-1895.

417 Kanehira, K., Uchida, Y., Takemae, N., Hikono, H., Tsunekuni, R., Saito, T., 2015.
 418 Characterization of an H5N8 influenza A virus isolated from chickens during an outbreak of
 419 severe avian influenza in Japan in April 2014. *Arch. Virol* 160, 1629-1643.

420 Kida, H., Yanagawa, R., 1979. Isolation and characterization of influenza a viruses from wild free-
 421 flying ducks in Hokkaido, Japan. *Zentralbl. Bakteriол. Orig. A* 244, 135-143.

422 Kilpatrick, A.M., Chmura, A.A., Gibbons, D.W., Fleischer, R.C., Marra, P.P., Daszak, P., 2006.
 423 Predicting the global spread of H5N1 avian influenza. *Proc. Natl. Acad Sci. U.S.A.* 103,
 424 19368-19373.

425 Kim, K.I., Choi, J.G., Kang, H.M., To, T.L., Nguyen, T.D., Song, B.M., Hong, M.S., Choi, K.S.,
 426 Kye, S.J., Kim, J.Y., Lee, H.S., Lee, Y.J., 2013. Geographical distribution of low pathogenic
 427 avian influenza viruses of domestic poultry in Vietnam and their genetic relevance with
 428 Asian isolates. *Poult. Sci.* 92, 2012-2023.

429 Le, T.H., Nguyen, N.T., 2014. Evolutionary dynamics of highly pathogenic avian influenza
 430 A/H5N1 HA clades and vaccine implementation in Vietnam. *Clin. Exp. Vaccine Res.* 3,
 431 117-127.

432 Lee, C.W., Senne, D.A., Suarez, D.L., 2004. Effect of vaccine use in the evolution of Mexican
 433 lineage H5N2 avian influenza virus. *J. Virol.* 78, 8372-8381.

434 Leung, Y.H., Lau, E.H., Zhang, L.J., Guan, Y., Cowling, B.J., Peiris, J.S., 2012. Avian influenza
 435 and ban on overnight poultry storage in live poultry markets, Hong Kong. *Emerg. Infect.*
 436 *Dis.* 18, 1339-1341.

437 Li, X., Fu, Y., Yang, J., Guo, J., He, J., Weng, S., Jia, Y., Liu, B., Zhu, Q., Chen, H., 2015. Genetic
 438 and biological characterization of two novel reassortant H5N6 swine influenza viruses in
 439 mice and chickens. *Infect. Genet. Evol.* 36, 462-466.

440 Minh, P.Q., Morris, R.S., Schauer, B., Stevenson, M., Benschop, J., Nam, H.V., Jackson, R., 2009.
 441 Spatio-temporal epidemiology of highly pathogenic avian influenza outbreaks in the two

442 deltas of Vietnam during 2003-2007. *Prev. Vet. Med.* 89, 16-24.

443 Moon, H.J., Song, M.S., Cruz, D.J., Park, K.J., Pascua, P.N., Lee, J.H., Baek, Y.H., Choi, D.H.,

444 Choi, Y.K., Kim, C.J., 2010. Active reassortment of H9 influenza viruses between wild

445 birds and live-poultry markets in Korea. *Arch. Virol* 155, 229-241.

446 Nguyen, D.T., Bryant, J.E., Davis, C.T., Nguyen, L.V., Pham, L.T., Loth, L., Inui, K., Nguyen, T.,

447 Jang, Y., To, T.L., Nguyen, T.D., Hoang, D.T., Do, H.T., Nguyen, T.T., Newman, S.,

448 Jennifer Siembieda, Pham, D.V., 2014. Prevalence and distribution of avian influenza

449 A(H5N1) virus clade variants in live bird markets of Vietnam, 2011-2013. *Avian Dis.* 58,

450 599-608.

451 Nguyen, T., Rivailler, P., Davis, C.T., Hoa, D.T., Balish, A., Dang, N.H., Jones, J., Vui, D.T.,

452 Simpson, N., Huong, N.T., Shu, B., Loughlin, R., Ferdinand, K., Lindstrom, S.E., York, I.A.,

453 Klimov, A., Donis, R.O., 2012. Evolution of highly pathogenic avian influenza (H5N1)

454 virus populations in Vietnam between 2007 and 2010. *Virology* 432, 405-416.

455 Nomura, N., Sakoda, Y., Endo, M., Yoshida, H., Yamamoto, N., Okamatsu, M., Sakurai, K., Hoang,

456 N.V., Nguyen, L.V., Chu, H.D., Tien, T.N., Kida, H., 2012. Characterization of avian

457 influenza viruses isolated from domestic ducks in Vietnam in 2009 and 2010. *Arch. Virol*

458 157, 247-257.

459 Okamatsu, M., Nishi, T., Nomura, N., Yamamoto, N., Sakoda, Y., Sakurai, K., Chu, H.D., Thanh,

460 L.P., Van Nguyen, L., Van Hoang, N., Tien, T.N., Yoshida, R., Takada, A., Kida, H., 2013.

461 The genetic and antigenic diversity of avian influenza viruses isolated from domestic ducks,

462 muscovy ducks, and chickens in northern and southern Vietnam, 2010-2012. *Virus Genes*
463 47, 317-329.

464 Pan, M., Gao, R., Lv, Q., Huang, S., Zhou, Z., Yang, L., Li, X., Zhao, X., Zou, X., Tong, W., Mao,
465 S., Zou, S., Bo, H., Zhu, X., Liu, L., Yuan, H., Zhang, M., Wang, D., Li, Z., Zhao, W., Ma,
466 M., Li, Y., Li, T., Yang, H., Xu, J., Zhou, L., Zhou, X., Tang, W., Song, Y., Chen, T., Bai,
467 T., Zhou, J., Wu, G., Li, D., Feng, Z., Gao, G.F., Wang, Y., He, S., Shu, Y., 2016. Human
468 infection with a novel, highly pathogenic avian influenza A (H5N6) virus: Virological and
469 clinical findings. *J. Infect.* 72 (1):52-9.

470 Paulson, J.C., de Vries, R.P., 2013. H5N1 receptor specificity as a factor in pandemic risk. *Virus*
471 Res. 178, 99-113.

472 Qi, X., Cui, L., Yu, H., Ge, Y., Tang, F., 2014. Whole-Genome Sequence of a Reassortant H5N6
473 Avian Influenza Virus Isolated from a Live Poultry Market in China, 2013. *Genome*
474 Announc. 2 (5), e00706-14

475 Smith, G.J., Donis, R.O., World Health Organization/World Organisation for Animal Health/Food
476 and Agriculture Organization (WHO/OIE/FAO), H5N1 Evolution Working Group, 2015.
477 Nomenclature updates resulting from the evolution of avian influenza A(H5) virus clades
478 2.1.3.2a, 2.2.1, and 2.3.4 during 2013-2014. *Influenza Other Respir. Viruses* 9, 271-276.

479 Soares Magalhães, R.J., Pfeiffer, D.U., Otte, J., 2010. Evaluating the control of HPAIV H5N1 in
480 Vietnam: virus transmission within infected flocks reported before and after vaccination.
481 *BMC. Vet. Res.* 6, 31.

482 Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: molecular
483 evolutionary genetics analysis using maximum likelihood, evolutionary distance, and
484 maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731-2739.

485 The World Bank, Projects and Operations. Vietnam - Avian and Human Influenza Control and
486 Preparedness Project.
487 <http://documents.worldbank.org/curated/en/2011/03/14026433/vietnam-additional->
488 [financing-vietnam-avian-human-influenza-control-preparedness-project-environmental-](http://documents.worldbank.org/curated/en/2011/03/14026433/vietnam-additional-financing-vietnam-avian-human-influenza-control-preparedness-project-environmental-impact-assessment)
489 [impact-assessment](http://documents.worldbank.org/curated/en/2011/03/14026433/vietnam-additional-financing-vietnam-avian-human-influenza-control-preparedness-project-environmental-impact-assessment) (accessed 25.09.15).

490 Wan, X.F., Dong, L., Lan, Y., Long, L.P., Xu, C., Zou, S., Li, Z., Wen, L., Cai, Z., Wang, W., Li,
491 X., Yuan, F., Sui, H., Zhang, Y., Dong, J., Sun, S., Gao, Y., Wang, M., Bai, T., Yang, L., Li,
492 D., Yang, W., Yu, H., Wang, S., Feng, Z., Wang, Y., Guo, Y., Webby, R.J., Shu, Y., 2011.
493 Indications that live poultry markets are a major source of human H5N1 influenza virus
494 infection in China. *J. Virol* 85, 13432-13438.

495 World Animal Health Information Database (WAHID) Interface, Disease Information.
496 http://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?reportid=15802
497 (accessed 25.09.15).

498 Yu, H., Wu, J.T., Cowling, B.J., Liao, Q., Fang, V.J., Zhou, S., Wu, P., Zhou, H., Lau, E.H., Guo,
499 D., Ni, M.Y., Peng, Z., Feng, L., Jiang, H., Luo, H., Li, Q., Feng, Z., Wang, Y., Yang, W.,
500 Leung, G.M., 2014. Effect of closure of live poultry markets on poultry-to-person
501 transmission of avian influenza A H7N9 virus: an ecological study. *Lancet* 383, 541-548.

502

503 **Figure captions**

504 **Fig. 1.** Phylogenetic tree for the influenza virus H5 HA genes. Full-length HA genes of six H5
505 subtype viruses, and reference strains were analyzed using the maximum-likelihood method with
506 MEGA 5.0 software and divided into two lineages: Eurasian and North American. The Eurasian H5
507 viruses were identified on the basis of the nomenclature defined by WHO/OIE/FAO H5N1
508 Evolution Working Group, 2014 (Smith et al., 2015). Horizontal distances are proportional to the
509 minimum number of nucleotide differences required to join nodes and sequences. Digits at the
510 nodes indicate the probability of confidence levels in a bootstrap analysis with 1,000 replications.
511 The viruses isolated in this study are highlighted in gray. The viruses isolated in our previous study
512 are underlined. The black circle indicates the viruses isolated in an intervention LBM.

513

514 **Fig. 2.** Phylogenetic tree for the influenza virus H6 HA genes. Nucleotides 51–943 (893 bp) of the
515 HA genes of four viruses of H6 subtype and reference strains were analyzed using the maximum-
516 likelihood method with MEGA 5.0 software and divided into two lineages: Eurasian and North
517 American. The Eurasian H6 viruses were clustered into five different sublineages: Group II, W312,
518 Group III, Early, and Group I (Okamatsu et al., 2013). Horizontal distances are proportional to the
519 minimum number of nucleotide differences required to join nodes and sequences. Digits at the
520 nodes indicate the probability of confidence levels in a bootstrap analysis with 1,000 replications.
521 The viruses isolated in this study are highlighted in gray, and the representative of each sublineage

522 is indicated in bold. The virus isolated in a previous study is underlined. The viruses isolated at
523 intervention LBMs are indicated by black circles, and the virus isolated at a non-intervention LBM
524 is indicated by a white circle.

525

526 **Fig. 3.** Phylogenetic tree for the influenza virus H9 HA genes. Full-length HA genes of eight
527 viruses of H9N2 subtype, and reference strains were analyzed using the maximum-likelihood
528 method with MEGA 5.0 software and divided into two lineages: Eurasian and North American
529 lineages. The Eurasian H9 viruses were clustered into three different sublineages: Y280, G1, and
530 Y439 (Nomura et al., 2012; Okamatsu et al., 2013). Horizontal distances are proportional to the
531 minimum number of nucleotide differences required to join nodes and sequences. Digits at the
532 nodes indicate the probability of confidence levels in a bootstrap analysis with 1,000 replications.
533 The viruses isolated in this study are highlighted in gray, and the representative of each sublineage
534 is indicated in bold. The virus isolated in a previous study is underlined. The viruses isolated in
535 intervention LBMs are indicated by black circles, and the viruses isolated in non-intervention LBMs
536 are indicated by white circles. The H9N2 subtype of each virus strain is omitted.

Table 1 Viruses isolated from LBMs in Vietnam in 2014

Type of LBMs	Name of LBMs	Latitude/Longitude	Number of samples	Number of isolates	% Positive (95% CI)	Subtype of isolates (number of isolates)
Intervention	An Lo	16.545962/107.452388	500	18	3.6 (2.3–5.6)	H3N2 (1) H4N6 (1) H6N6 (1) H9N2 (10) H9N6 (2) H11N7 (3)
	No	16.511793/107.600336	500	61	12.2 (9.6–15.4)	H3N2 (4) H3N6 (1) H5N6 (8) H6N2 (9) H6N6 (3) H9N2 (36)
	Thuy Phuong	16.433257/107.635256	496	13	2.6 (1.5–4.4)	H3N2 (2) H4N6 (1) H6N2 (1) H6N6 (5) H9N2 (3) H11N6 (1)
	Total		1,496	92	6.1 (5.0–7.5)	
Non-intervention	Phu Bai	16.408326/107.677989	300	40	13.3 (9.9–17.6)	H6N2 (3) H6N6 (1) H9N2 (35) H11N7 (1)
	Phu Da Than Phu	16.434797/107.710014	200	1	0.5 (0.1–2.8)	H3N2 (1)
		16.424635/107.656268	199	12	6.0 (3.5–10.2)	H3N2 (1) H6N6 (5) H9N2 (3) H9N6 (3)
	Quang Phuoc Tay Ba	16.57431/107.51984	212	6	2.8 (1.3–6.0)	H9N2 (6)
		16.537018/107.560679	338	25	7.4 (5.1–10.7)	H3N2 (9) H6N2 (1) H9N2 (15)
	Vinh Thanh	16.431796/107.783475	300	2	0.7 (0.2–2.4)	H6N6 (1) H9N2 (1)
	Total		1,549	86	5.6 (4.5–6.8)	

CI, confidence interval

Table 2 Representative H5, H6, and H9 viruses isolated in live bird markets in Vietnam in 2014

HA subtypes	Type of LBMs	Name of LBMs	Isolates	Accession numbers							
				PB2	PB1	PA	HA	NP	NA	M	NS
H5	Intervention	No	A/muscovy duck/Vietnam/HU1-1144/2014 (H5N6)	LC069835	LC069867	LC069898	LC041323	LC069953	LC041324	LC070009	LC070040
			A/duck/Vietnam/HU1-1151/2014 (H5N6)	LC041310	LC041311	LC041312	LC041313	LC086330	LC041314	LC041315	LC041316
			A/environment/Vietnam/HU1-1434/2014 (H5N6)	LC069834	LC069866	LC069897	LC041321	LC069952	LC041322	LC070008	LC070039
			A/duck/Vietnam/HU1-1507/2014 (H5N6)	LC069832	LC069864	LC069895	LC041317	LC069950	LC041318	LC070006	LC070037
			A/duck/Vietnam/HU1-1511/2014 (H5N6)	LC069833	LC069865	LC069896	LC041319	LC069951	LC041320	LC070007	LC070038
H6	Intervention	An Lo	A/duck/Vietnam/HU1-1245/2014 (H6N2)	LC069813	LC069846	LC069877	LC069908	LC069932	LC069963	LC069988	LC070019
		No	A/environment/Vietnam/HU1-1423/2014 (H6N2)	LC069836	LC041334	LC041335	LC041336	LC041337	LC041338	LC041339	LC041340
			A/environment/Vietnam/HU1-1426/2014 (H6N2)	LC069837	LC069868	LC069899	LC041341	LC069954	LC041342	LC070010	LC070041
	Non-intervention	Than Phu	A/duck/Vietnam/HU1-637/2014 (H6N6)	LC069817	LC069850	LC069881	LC069912	LC069936	LC069967	LC069992	LC070023
H9	Intervention	An Lo	A/chicken/Vietnam/HU1-3/2014 (H9N2)	LC069838	LC069869	LC069900	LC041350	LC069955	LC069980	LC070011	LC070042
			A/duck/Vietnam/HU1-225/2014 (H9N2)	LC069842	LC069873	LC069904	LC069928	LC069959	LC069984	LC070015	LC070046
		No	A/chicken/Vietnam/HU1-1286/2014 (H9N2)	LC069840	LC069871	LC069902	LC069926	LC069957	LC069982	LC070013	LC070044
			A/environment/Vietnam/HU1-1424/2014 (H9N2)	LC069841	LC069872	LC069903	LC069927	LC069958	LC069983	LC070014	LC070045
			A/duck/Vietnam/HU1-1512/2014 (H9N2)	LC041343	LC041344	LC041345	LC041346	LC086331	LC041347	LC041348	LC041349
	Non-intervention	Phu Bai	A/chicken/Vietnam/HU1-381/2014 (H9N2)	LC069839	LC069870	LC069901	LC069925	LC069956	LC069981	LC070012	LC070043
		Than Phu	A/duck/Vietnam/HU1-675/2014 (H9N2)	LC069820	LC069853	LC069884	LC069915	LC069939	LC069970	LC069995	LC070026
		Quang Phuoc	A/chicken/Vietnam/HU1-786/2014 (H9N2)	LC069843	LC069874	LC069905	LC069929	LC069960	LC069985	LC070016	LC070047

Table 3 Cross-reactivity of H5 influenza viruses with antisera by HI test

Lineage	Viruses	Clade	Antiserum to						
			Vac-3	Mon/05	Hok/08	HK/09	Km/14	Yama/04	Ibr/05
Eurasia	A/duck/Hokkaido/Vac-3/2007 (H5N1)	–	<u>10,240</u>	10,240	640	1,280	1,280	5,120	2,560
	A/whooper swan/Mongolia/3/2005 (H5N1)	2.2	80	<u>640</u>	80	40	20	1,280	80
	A/whooper swan/Hokkaido/1/2008 (H5N1)	2.3.2.1	80	1,280	<u>1,280</u>	80	80	1,280	40
	A/peregrine falcon/Hong Kong/810/2009 (H5N1)	2.3.4	40	20	40	<u>2,560</u>	40	160	20
	A/muscovy duck/Vietnam/HU1-1144/2014 (H5N6)	2.3.4.4	80	40	20	640	640	80	<20
	A/duck/Vietnam/HU1-1151/2014 (H5N6)	2.3.4.4	40	20	20	640	640	40	<20
	A/environment/Vietnam/HU1-1434/2014 (H5N6)	2.3.4.4	20	20	20	160	160	20	<20
	A/muscovy duck/Vietnam/HU2-26/2014 (H5N6)	2.3.4.4	40	40	20	640	640	80	<20
	A/chicken/Kumamoto/1-7/2014 (H5N8)	2.3.4.4	20	20	20	320	<u>640</u>	80	<20
	A/chicken/Yamaguchi/7/2004 (H5N1)	2.5	640	10,240	640	160	160	<u>10,240</u>	320
North America	A/chicken/Ibaraki/1/2005 (H5N2)	–	160	640	20	20	<20	2,560	<u>10,240</u>

The H5 isolates identified in the present study are shown in bold.

Vac-3, A/duck/Hokkaido/Vac-3/2007; Mon/05, A/whooper swan/Mongolia/3/2005; Hok/08, A/whooper swan/Hokkaido/1/2008; HK/09, A/peregrine falcon/Hong Kong/810/2009; Km/14, A/chicken/Kumamoto/1-7/2014; Yama/04, A/chicken/Yamaguchi/7/2004; Ibr/05, A/chicken/Ibaraki/1/2005

Homologous titers are underlined.

"–" indicates that the virus does not belong to clade 0–9

Table 4 Cross-reactivity of H6 influenza viruses with antisera by HI test

Lineage	Sublinage	Viruses	Antiserum to				
			HK/960/80	VN/OIE-4429/10	VN/HU1-637/14	Aus/1/72	Mas/3740/65
Eurasia	Early	A/duck/Hong Kong/960/1980 (H6N2)	<u>5,120</u>	640	640	640	640
	Group II	A/duck/Vietnam/OIE-4429/2010 (H6N2)	640	<u>5,120</u>	640	160	2,560
		A/duck/Vietnam/HU1-637/2014 (H6N6)	640	1,280	<u>10,240</u>	80	640
		A/duck/Vietnam/HU1-1245/2014 (H6N2)	160	160	2,560	20	160
		A/environment/Vietnam/HU1-1423/2014 (H6N2)	320	640	2,560	40	320
—		A/environment/Vietnam/HU1-1426/2014 (H6N2)	320	640	2,560	80	320
		A/shearwater/South Australia/1/72 (H6N5)	2,560	40	80	<u>1,280</u>	640
North American		A/turkey/Masachusetts/3740/65 (H6N2)	2,560	1,280	40	1,280	<u>5,120</u>

The H6 isolates identified in the present study are shown in bold.

HK/960/80 , A/duck/Hong Kong/960/1980 (H6N2); VN/OIE-4429/10, A/duck/Vietnam/OIE-4429/2010 (H6N2); VN/HU1-637/14, A/duck/Vietnam/HU1-637/2014 (H6N6); Aus/1/72, A/shearwater/South Australia/1/72 (H6N5); Mas/3740/65 , A/turkey/Masachusetts/3740/65 (H6N2)

Homologous titers are underlined.

"—" indicates that the virus does not belong to any lineage.

Table 5 Cross-reactivity of H9N2 influenza viruses with antisera by HI test

Lineage	Sublinage	Viruses	Antiserum to			
			HK/Y280/97	HK/G1/97	Hok/49	Wis/66
Eurasia	Y280	A/duck/Hong Kong/Y280/1997	<u>10,240</u>	1,280	2,560	80
		A/chicken/Vietnam/HU1-3/2014	10,240	1,280	1,280	80
		A/duck/Vietnam/HU1-225/2014	2,560	1,280	640	80
		A/chicken/Vietnam/HU1-381/2014	10,240	640	1,280	40
		A/chicken/Vietnam/HU1-786/2014	20,480	1,280	2,560	80
		A/chicken/Vietnam/HU1-1286/2014	10,240	640	640	80
		A/environment/Vietnam/HU1-1424/2014	10,240	320	1,280	40
		A/duck/Vietnam/HU1-1512/2014	10,240	640	2,560	40
	G1	A/quail/Hong Kong/G1/1997	1,280	<u>5,120</u>	640	80
	Y439	A/duck/Hokkaido/49/1998	640	80	<u>2,560</u>	320
North America	—	A/turkey/Wisconsin/1/1966	80	40	320	<u>320</u>

The H9 isolates identified in the present study are shown in bold.

HK/Y280/97, A/duck/Hong Kong/Y280/1997; HK/G1/97, A/quail/Hong Kong/G1/1997; Hok/49, A/duck/Hokkaido/49/1998; Wis/66, A/turkey/Wisconsin/1/1966

Homologous titers are underlined.

"—" indicates that the virus does not belong to any sublinage.

Fig. 1. Chu *et al.*

Fig. 2. Chu *et al.*

0.02

Fig. 3. Chu *et al.*