



Title	Elucidation of the biosynthetic pathway of cis-jasmone in <i>Lasiodiplodia theobromae</i>
Author(s)	Matsui, Ryo; Amano, Naruki; Takahashi, Kosaku; Taguchi, Yodai; Saburi, Wataru; Mori, Hideharu; Kondo, Norio; Matsuda, Kazuhiko; Matsuura, Hideyuki
Citation	Scientific reports, 7(1), 6688 https://doi.org/10.1038/s41598-017-05851-7
Issue Date	2017-07-27
Doc URL	http://hdl.handle.net/2115/67110
Rights(URL)	http://creativecommons.org/licenses/by/4.0/
Type	article
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	s41598-017-05851-7.pdf



[Instructions for use](#)

SCIENTIFIC REPORTS

OPEN

Elucidation of the biosynthetic pathway of *cis*-jasmone in *Lasiodiplodia theobromae*

Ryo Matsui¹, Naruki Amano¹, Kosaku Takahashi¹, Yodai Taguchi¹, Wataru Saburi¹, Hideharu Mori¹, Norio Kondo¹, Kazuhiko Matsuda^{1,2} & Hideyuki Matsuura¹

In plants, *cis*-jasmone (CJ) is synthesized from α -linolenic acid (LA) via two biosynthetic pathways using jasmonic acid (JA) and *iso*-12-oxo-phytodienoic acid (*iso*-OPDA) as key intermediates. However, there have been no reports documenting CJ production by microorganisms. In the present study, the production of fungal-derived CJ by *Lasiodiplodia theobromae* was observed for the first time, although this production was not observed for *Botrytis cinerea*, *Verticillium longisporum*, *Fusarium oxysporum*, *Gibberella fujikuroi*, and *Cochliobolus heterostrophus*. To investigate the biosynthetic pathway of CJ in *L. theobromae*, administration experiments using [18,18,18-²H₃], [17,17-²H₂]LA (LA-d5), [18,18,18-²H₃, 17,17-²H₂]12-oxo-phytodienoic acid (*cis*-OPDA-d5), [5',5',5'-²H₃, 4',4'-²H₂, 3'-²H₁]OPC 8:0 (OPC8-d6), [5',5',5'-²H₃, 4',4'-²H₂, 3'-²H₁]OPC 6:0 (OPC6-d6), [5',5',5'-²H₃, 4',4'-²H₂, 3'-²H₁]OPC 4:0 (OPC4-d6), and [11,11-²H₂, 10,10-²H₂, 8,8-²H₂, 2,2-²H₂]methyl *iso*-12-oxo-phytodienoate (*iso*-MeOPDA-d8) were carried out, revealing that the fungus produced CJ through a single biosynthetic pathway via *iso*-OPDA. Interestingly, it was suggested that the previously predicted decarboxylation step of 3,7-didehydroJA to afford CJ might not be involved in CJ biosynthesis in *L. theobromae*.

Jasmonic acid (JA) is a phytohormone and key mediator in plant wound responses to insects, necrotrophic pathogens^{1–3} and other environmental stresses. The biosynthesis of JA begins with the oxygenation of α -linolenic acid (LA) in the chloroplast to give (+)-7-*iso*-JA in the peroxisome⁴ via 12-oxo-phytodienoic acid (*cis*-OPDA), 3-oxo-2-(2'-[Z]-pentenyl)-cyclopentane-1-octanoic acid (OPC 8:0), 3-oxo-2-(2'-[Z]-pentenyl)-cyclopentane-1-hexanoic acid (OPC 6:0), and 3-oxo-2-(2'-[Z]-pentenyl)-cyclopentane-1-butanoic acid (OPC 4:0). (+)-7-*iso*-JA is readily epimerized to afford JA, which has an absolute configuration of (3R, 7R) (Fig. 1). The synthesized (+)-7-*iso*-JA is metabolized to afford other jasmonates, including MeJA, 12-hydroxy-JA (12-OH-JA), and JA-amino acid conjugates. It has been generally accepted that the compounds derived from oxygenated polyunsaturated fatty acids are categorized as oxylipins, and thus, JA and its derivatives are members of this family. JA-related compounds are well known, and their biological functions have been reported^{5,6}. Among of them, jasmonoyl isoleucine (JA-Ile) is an important player due to its curtail biological roles to interact with its receptor, COI1^{7–10}, which needs to induce JA dependent biological phenomena. However, Yan *et al.*¹¹ reported that JA-Ile and other four JA conjugates are the ligand of the JA receptor. Methyl jasmonate (MeJA) is a volatile compound involved in the response to wound stress, and volatized MeJA is thought to induce defensive proteins in plants¹².

cis-Jasmone (CJ) is one of oxylipins, and it was reported that the biosynthetic pathway of CJ in plant was proceeded using JA as a biosynthetic intermediate (pathway A, Fig. 1)¹³. In this report, [7-²H₁, 5,5-²H₂, 2,2-²H₂]JA and [9-²H₁, 8-²H₁]methyl 3,7-didehydrojasmonate were converted into deuterium-labeled CJs in jasmine flowers and the leaves of six higher plants, including lima beans (*Phaseolus lunatus*) and willow (*Salix alba*). It was proposed that the conversion of 3,7-didehydroJA to CJ is achieved by decarboxylation¹³. Moreover, a recent study has shown that *iso*-12-oxo-phytodienoic acid (*iso*-OPDA) is an early precursor of CJ¹⁴, suggesting biosynthesis via pathway B (Fig. 1). *iso*-OPDA is an isomer of *cis*-OPDA, an important intermediate in the JA biosynthetic pathway (Fig. 1). Interestingly, the successful conversion of labelled *iso*-OPDA into CJ by the yeast *Saccharomyces cerevisiae* has also been reported, although the yeast had no operative pathway for CJ. Several reports have demonstrated the biological activities of *cis*-OPDA^{15,16}. The existence of OPDA-Ile in *Arabidopsis thaliana* was proven¹⁷, the related biological activity was evaluated by Arnold *et al.*¹⁸. However, no similar reports

¹Research Faculty of Agriculture, Hokkaido University, Sapporo, 060-8589, Japan. ²Graduate School of Agriculture, Faculty of Agriculture, Kinki University, Nakamachi, Nara, 631-8505, Japan. Correspondence and requests for materials should be addressed to H.M. (email: matsuura@chem.agr.hokudai.ac.jp)




Figure 1. Biosynthetic pathway to give CJ and JA. LA: α -linolenic acid, OPDA: 12-oxo-phytodienoic acid, OPC 8:0: 3-oxo-2-(2'-[Z]-pentenyl)-cyclopentane-1-octanoic acid, OPC 6:0: 3-oxo-2-(2'-[Z]-pentenyl)-cyclopentane-1-hexanoic acid, OPC 4:0: 3-oxo-2-(2'-[Z]-pentenyl)-cyclopentane-1-butanoic acid (OPC 4:0). 13-LOX: 13-lipoxygenase, AOS: alene oxide synthase, AOC: alene oxide cyclase, OPR3: 12-oxophytodienoate reductase 3.

exist for *iso*-OPDA. There is an interesting report about *iso*-OPDA as a metabolite, which revealed that *cis*-OPDA isomerase is involved in phytohormone detoxification in the insect gut, a process by which *cis*-OPDA is converted to *iso*-OPDA^{19–23}. It has been generally accepted that CJ is involved in plant defense systems, similar to JA. For example, CJ is released from wounded leaves, attracts aphid parasitoids and acts as a repellent of the pest^{24, 25}. CJ is also released from the flowers of many plants, such as jasmine, neroli (*Citrus bigaradia*), jonquil (*Narcissus jonquilla* L.), bergamot (*Citrus bergamia*), and the *Pittosporum* family. On the other hand, some insects such as butterflies²⁶ and silk worms²⁷ use CJ as a sex pheromone. Therefore, potential agricultural use of CJ in applications such as aphid control has been studied²⁸.

Plant pathogens produce various secondary metabolites, and some fungi produce phytohormones, such as JA, auxin, and abscisic acid²⁹. It was initially thought that fungi produce phytohormones to disturb signaling systems of plants and enable facile invasion of plants. The fungus, *Verticillium longisporum* uses CORONATINE INSENSITIVE1 (COI1), which is an essential protein for establishing the JA-dependent wound response, to invade plants, and thus, it was hypothesized that the fungus might synthesize JA-related compounds. *Fusarium oxysporum* causes vascular wilt disease in more than 100 plant species and reportedly produces 22 kinds of JA analogues³⁰. Furthermore, Thatcher and coworkers discovered that *F. oxysporum* also hijacks COI1-mediated jasmonate signaling to promote disease development in *Arabidopsis*³¹. The fungus *Botrytis cinerea*, *Gibberella fujikuroi* and *Lasiodiplodia theobromae* also synthesize JA, and the JA biosynthetic pathway in *L. theobromae* has been reported by Tsukada *et al.*³². *Cochliobolus heterostrophus* can cause corn leaf blight in maize and produces a death acid that is an analogue of JA³³. The biosynthetic pathway of this death acid is similar to that of JA, and the fungus reportedly uses the plant's JA biosynthetic pathway to synthesize the death acid. However, fungal production of CJ has not been reported. In this study, we discovered that *L. theobromae* produces CJ. Furthermore, we used a combined approach involving feeding the fungus deuterium-labeled compounds and gas chromatography–mass spectrometry (GC-MS) analysis to elucidate the CJ biosynthetic pathway in *L. theobromae*.

Results and Discussion

Screening of fungal producers of CJ. The similarities in the chemical structures of CJ and JA suggest that some species of plant pathogens that synthesize JA or its related compounds could produce CJ. Thus, the ability of six species of plant pathogens, *L. theobromae*, *B. cinerea*, *V. longisporum*, *F. oxysporum*, *G. fujikuroi*, and *C. heterostrophus*, to synthesize CJ was evaluated using GC-MS. To ensure accuracy in the experiment, [4,4-²H₂,




Figure 2. Representative GC MS chromatograms analyzing fungal derived *cis*-jasmone. (A, B, C, D, E and F) measuring fungal derived *cis*-jasmone using selected ion monitoring at m/z 164 (lower) and 171 (upper, internal standard) for *L. theobromae*, *B. cinerea*, *V. longisporum*, *F. oxysporum*, *G. fujikuroi*, and *C. heterostrophus*, respectively.

$3,3\text{-}^2\text{H}_2$, $1,1,1\text{-}^2\text{H}_3$]CJ (CJ-d7, m/z 171) was synthesized according to a reported method³⁴, except that commercially available CJ was used as the starting material. The isotopic purity was established by comparing the data of GC-MS for CJ and CJ-d7. The incorporation of $^2\text{H}_2$ was found to be 94% for CJ-d7. Representative GC-MS/MS chromatograms for authentic CJ are shown in Supplementary Fig. 1, and the results of screening experiment are given in Fig. 2. Peaks corresponding to CJ-d7 (m/z 171), an internal standard, in selected ion monitoring mode were detected in the extracts of all the six species of fungi (Fig. 2A–F, upper panels). However, a peak corresponding to CJ (m/z 164) was only detected in the culture filtrate derived from *L. theobromae* (Fig. 2A, lower panel), and the corresponding MS pattern agreed with that of the authentic standard shown in Supplementary Fig. S1. However, the peak corresponding to CJ was not detected in the filtrates derived from the other five pathogens (Fig. 2B–F, lower panels). A concentration of CJ, calculated based on the peak area ratio between CJ-d7 and fungal-derived CJ, was determined to be 8.7 $\mu\text{g}/\text{mL}$ in the culture filtrate of *L. theobromae*.

Metabolism of deuterium-labeled LA-d5 and *cis*-OPDA-d5 to MeJA-d5 and CJ-d5 in feeding experiments.

CJ in plants is reportedly synthesized using JA and *iso*-OPDA as synthetic intermediates via independent pathways (pathways A and B, respectively, in Fig. 1) that utilize LA and *cis*-OPDA as common intermediates in an early biosynthetic step (Fig. 1)^{13,14}. Based on these reports, we hypothesized that *L. theobromae* also uses these pathways to produce CJ. To determine the biosynthetic pathway affording CJ, *L. theobromae* was administered deuterium-labeled intermediates, including $[18,18,18\text{-}^2\text{H}_3, 17,17\text{-}^2\text{H}_2]$ LA (LA-d5), $[18,18,18\text{-}^2\text{H}_3, 17,17\text{-}^2\text{H}_2]$ *cis*-OPDA (*cis*-OPDA-d5), $[5',5',5'\text{-}^2\text{H}_3, 4',4'\text{-}^2\text{H}_2, 3'\text{-}^2\text{H}_1]$ OPC 8:0 (OPC8-d6), $[5',5',5'\text{-}^2\text{H}_3, 4',4'\text{-}^2\text{H}_2, 3'\text{-}^2\text{H}_1]$ OPC 6:0 (OPC6-d6), $[5',5',5'\text{-}^2\text{H}_3, 4',4'\text{-}^2\text{H}_2, 3'\text{-}^2\text{H}_1]$ OPC 4:0 (OPC4-d6), and $[11,11\text{-}^2\text{H}_2, 10,10\text{-}^2\text{H}_2, 8,8\text{-}^2\text{H}_2, 2,2\text{-}^2\text{H}_2]$ methyl *iso*-12-oxo-phytodienoate (*iso*-MeOPDA-d8) (Fig. 3). A schematic illustration of the experimental procedure is shown in Supplementary Fig. 2. *cis*-OPDA-d5 was synthesized using commercially available LA-d5 according to the method reported by Kajiwara *et al.*³⁵, while OPC8-d6, OPC6-d6, and OPC4-d6 were synthesized according to Matsuura *et al.*³⁶. CJ was detected in the extract without modification, although a portion of the EtOAc extract was treated with a CH_2N_2 solution to convert endogenous JA into MeJA to increase its volatility. Representative GC-MS chromatograms of authentic MeJA are shown in Supplementary Fig. 3. In the feeding experiment using LA-d5 as a substrate, the GC-MS chromatograms obtained in selected ion monitoring mode contained ion peaks at m/z 224 and 229 for MeJA and $[12,12,12\text{-}^2\text{H}_3, 11,11\text{-}^2\text{H}_2]$ MeJA (MeJA-d5), respectively, at Rt. 7.5 min (Fig. 4). The mass fragmentation patterns for the m/z 224 and 229 ion peaks are shown in Supplementary Fig. 4 and substantiated the presence of MeJA and MeJA-d5 in the fungal culture filtrate. The GC-MS chromatograms for the analysis of CJ and $[11,11,11\text{-}^2\text{H}_3, 10,10\text{-}^2\text{H}_2]$ CJ (CJ-d5) when using LA-d5 as a




Figure 3. Chemical structures of deuterium-labeled compounds used in this study. Way of numbering the carbon atoms of OPDA-d5 was according to that of previous report²¹, and those of OPC8-d6, OPC6-d6, and OPC4-d6 were according to the report of Matsuura *et al.*³⁶.

feeding substrate are shown in Fig. 5. These chromatograms contained ion peaks of m/z 164 and 169 for CJ and CJ-d5, respectively, which were detected at Rt. 15.7 min in selected ion monitoring mode. The mass fragmentation patterns of the m/z 164 and 169 ion peaks are shown in Supplementary Fig. 5 and confirmed the presence of CJ and CJ-d5 in the fungal culture filtrate. The results of the feeding experiment using *cis*-OPDA-d5 were identical to those of the LA-d5 experiment (data not shown).

Deuterium-labeled OPC 8:0, OPC 6:0, and OPC 4:0 are metabolized to MeJA but not CJ. In the feeding experiment using OPC8-d6 as a substrate, the GC-MS chromatograms obtained in selected ion monitoring mode contained ion peaks at m/z 224 and 230 for MeJA and $[12,12,12\text{-}^2H_3, 11,11\text{-}^2H_2, 10\text{-}^2H_1]$ MeJA (MeJA-d6), respectively, at Rt. 7.5 min (Fig. 6). The mass fragmentation patterns for the m/z 224 and 230 ion peaks are shown in Supplementary Fig. 6 and confirmed the presence of MeJA and MeJA-d6 in the fungal culture filtrate. The GC-MS chromatograms for the analysis of CJ and $[11,11,11\text{-}^2H_3, 10,10\text{-}^2H_2, 6\text{-}^2H_1]$ CJ (CJ-d6) when using OPC8-d6 as a feeding substrate are shown in Fig. 7. The m/z 164 ion peak was detected at Rt. 15.7 min in the selected ion monitoring chromatogram (Fig. 7B), although the identical ion peak was not detected in the chromatogram obtained from selected ion monitoring at m/z 170 (Fig. 7C). The mass fragmentation pattern of the ion peak at m/z 164 is shown in Supplementary Fig. 7 and confirmed the presence of CJ in the fungal culture filtrate. Because the m/z 170 ion peak could not be detected in a GC-MS chromatogram of acceptable quality (Fig. 7C), it was concluded that OPC8-d6 was not used as a building block to synthesize CJ-d6. The results of the feeding experiments using OPC6-d6 and OPC4-d6 were identical to those obtained using OPC8-d6 (data not shown). The results of the deuterium-labeled OPC 8:0, OPC 6:0, and OPC 4:0 feeding experiments thus suggested that biosynthetic pathway A was not involved in CJ production in *L. theobromae* (Fig. 1).

Feeding deuterium-labeled *iso*-MeOPDA to growing fungal cultures produces labeled CJ as a metabolite but not labeled MeJA. Plants also use *iso*-OPDA as a key intermediate in biosynthetic pathway B to synthesize CJ¹⁴. Furthermore, it has been proposed that the final step in establishing the chemical backbone of CJ is a decarboxylation reaction in which a carboxyl group is eliminated from 3,7-didehydroJA. In a previous report demonstrating biosynthetic pathway B in plants, $[16\text{-}^2H_1, 15\text{-}^2H_1]$ *iso*-OPDA (*iso*-OPDA-d2) was synthesized and administered to the plants to demonstrate CJ metabolism. However, the final decarboxylation step was not fully proven. To examine biosynthetic pathway B in *L. theobromae* as well as the decarboxylation




Figure 4. Representative GC-MS chromatograms for measuring MeJA in feeding experiment using LA-d5. (A) Representative GC-MS chromatogram monitoring total ion. (B) Representative GC-MS chromatogram for measuring fungal derived MeJA using selected ion monitoring at m/z 224. (C) Representative GC-MS chromatogram for measuring fungal derived MeJA-d5 using selected ion monitoring at m/z 229. Representative MS chart of the peak indicated by arrow is given in Supplementary Figure S4.




Figure 5. Representative GC-MS chromatograms for measuring CJ in feeding experiment using LA-d5. (A) Representative GC-MS chromatogram monitoring total ion. (B) Representative GC-MS chromatogram for measuring fungal derived CJ using selected ion monitoring at m/z 164. (C) Representative GC-MS chromatogram for measuring fungal derived CJ-d5 using selected ion monitoring at m/z 169. Representative MS chart of the peak indicated by arrow is given in Supplementary Figure S5.

step, an experiment to administer $[11,11\text{-}^2\text{H}_2, 10,10\text{-}^2\text{H}_2, 8,8\text{-}^2\text{H}_2, 2,2\text{-}^2\text{H}_2]$ methyl *iso*-12-oxo-phytodienoate (*iso*-MeOPDA-d8) was carried out. The carbon backbone of *iso*-MeOPDA-d8 was synthesized according to a reported method³⁷, and deuterium labels were added to the product according to the reported method³⁴ with some modifications, as described in the experimental section. The isotopic purity was established by comparing the data of GC-MS for *iso*-MeOPDA and *iso*-MeOPDA-d8. The incorporation of $^2\text{H}_2$ was found to be 72% for *iso*-MeOPDA-d8. The synthesized compound was fed to *L. theobromae*, and the culture filtrate was analyzed by GC-MS. A representative GC-MS chromatogram is shown in Fig. 8. The m/z 164 and 168 ion peaks were detected at Rt. 15.7 min in selected ion monitoring mode (Fig. 8A and B, respectively). However, the predicted ion peak for m/z 170 was not detected (Fig. 8C). The mass fragmentation patterns of the m/z 164 and 168 ion peaks are shown in Supplementary Fig. 8 and confirmed the presence of CJ and $[4,4\text{-}^2\text{H}_2, 3,3\text{-}^2\text{H}_2]$ CJ (CJ-d4) in the fungal culture filtrate. The labeled CJ-d4 pattern was determined and had specific characteristics. Ion peaks for a $[M\text{-CH}_3]^+$ fragment were observed in the CJ and CJ-d4 MS chromatograms (Supplementary Fig. 8), and a $[M\text{-CD}_3]^+$ fragment was detected in CJ-d7 (m/z 171, Supplementary Fig. 9), indicating that deletion of the methyl group at the C-1 position had occurred during MS fragmentation. This type of deletion was also observed as $[M\text{-CH}_3]^+$ from CJ and $[9\text{-}^2\text{H}_1, 8\text{-}^2\text{H}_1]$ CJ and $[M\text{-CD}_2\text{H}]^+$ from $[4,4\text{-}^2\text{H}_2, 1,1\text{-}^2\text{H}_2]$ CJ in the previous report⁹. These results supported the detection of CJ-d4 as $[4,4\text{-}^2\text{H}_2, 3,3\text{-}^2\text{H}_2]$ CJ. Based on the abovementioned results, it




Figure 6. Representative GC-MS chromatograms for measuring MeJA in feeding experiment using OPC 8:0-d6. (A) Representative GC-MS chromatogram monitoring total ion. (B) Representative GC-MS chromatogram for measuring fungal derived MeJA using selected ion monitoring at m/z 224. (C) Representative GC-MS chromatogram for measuring fungal derived MeJA-d6 using selected ion monitoring at m/z 230. Representative MS chart of the peak indicated by arrow is given in Supplementary Figure S6.




Figure 7. Representative GC-MS chromatograms for measuring CJ in feeding experiment using OPC8-d6. (A) Representative GC-MS chromatogram monitoring total ion. (B) Representative GC-MS chromatogram for measuring fungal derived CJ using selected ion monitoring at m/z 164. (C) Representative GC-MS chromatogram for measuring fungal derived CJ-d5 using selected ion monitoring at m/z 170. Representative MS chart of the peak indicated by arrow is given in Supplementary Figure S7.

appeared reasonable to suppose that CJ was synthesized via biosynthetic pathway B, and the predicted decarboxylation reaction might not be involved in the final biosynthetic step to afford CJ in *L. theobromae*. Interestingly, [5,5- $^2\text{H}_2$, 4,4- $^2\text{H}_2$, 2,2- $^2\text{H}_2$]MeJA (m/z 230) was not detected in the GC-MS data (Supplementary Fig. 10).

In this paper, it was found that *L. theobromae* produced CJ, although *B. cinerea*, *V. longisporum*, *F. oxysporum*, *G. fujikuroi*, and *C. heterostrophus* did not (Fig. 2). It appears that the ability to produce CJ is a unique feature of *L. theobromae*. In addition, it was suggested that the fungus used biosynthetic pathway B (Fig. 1) to produce CJ, and the predicted decarboxylation reaction might not be involved in the final biosynthesis step to produce fungal-derived CJ, suggesting the involvement of an unknown step. However, it has been generally accepted that the deuterium atoms at the α -position tend to be removed by enolization, and the α -methylene group at the C-2 position in 3,7-didehydroJA, which is strongly acidic due to the carboxyl groups, is a typical example of where this type of enolization can occur. However, Koch *et al.*¹³ have proven that deuterium-labeled atoms at the α -positions in [7- $^2\text{H}_1$, 5,5- $^2\text{H}_2$, 2,2- $^2\text{H}_2$] JA remain throughout the biosynthetic steps to give [4,4- $^2\text{H}_2$, 1,1- $^2\text{H}_2$]CJ in lima bean (*Phaseolus lunatus* cv. Sieva). Based on this report, it was presumed that the loss of deuterium atoms detected in the present study might not be due to enolization. To draw a more reliable conclusion, further research, such as




Figure 8. Representative GC MS/MS chromatograms analyzing fungal derived CJ, CJ-d4, and CJ-d6 using *iso*-OPDA-d8 as a substrate for a feeding experiment. **(A)** Representative GC-MS chromatogram for measuring fungal derived CJ using selected ion monitoring at m/z 164. **(B)** Representative GC-MS chromatogram for measuring fungal derived CJ-d4 using selected ion monitoring at m/z 168. **(C)** Representative GC-MS chromatogram for measuring fungal derived CJ-d6 using selected ion monitoring at m/z 170. Representative MS chart of the peak indicated by arrow is given in Figure S8.

feeding experiments applying $[2-^{13}\text{C}]$ and $[1-^{13}\text{C}, 2-^{13}\text{C}]$ sodium acetate to fungal cultures of *L. theobromae*, is needed.

During our research, it was revealed that the fungus, *L. theobromae*, produced CJ. Since CJ has similar chemical structure with that of JA, it was assumed that CJ could mimic the biological activity of JA such as antagonistic effect upon salicylic acid signaling pathway³⁸ that induces plant defense system. Given that CJ had JA like biological activity, *L. theobromae* might produce CJ to inhibit salicylic acid signaling pathway to facilitate invasion into the plants. However, this consideration was just hypothesis, and further research on the production of CJ in the fungus is needed to evaluate why the fungus synthesized CJ.

Experimental Section

General experimental procedures. NMR spectra were recorded in CDCl_3 using a JNM-EX 270 FT-NMR spectrometer (JEOL) for ^1H NMR experiments at 270 MHz and an AMX 500 (Bruker) for ^1H NMR experiments at 500 MHz. FD-MS and FI-MS analyses were performed on a JMS-T100GCV (JEOL) instrument. GC-MS analyses were performed using a Varian instrument. Authentic CJ was purchased from Sigma-Aldrich. LA-d5 (98%) was purchased from Cambridge Isotope Laboratories, Inc.

GC-MS conditions. GC-MS analysis of CJ and MeJA was completed on a Varian CP-3800 gas chromatograph with a Varian 1200 L quadrupole MS/MS in electron ionization mode. For the analysis of CJ, the injection temperature was 150°C , and a fused-silica capillary column (TC-5; $30\text{ m} \times 0.25\text{ mm i.d.}$, $0.25\text{ }\mu\text{m}$ film thickness; GL Sciences) was used. The temperature program started at 40°C for 1 min and subsequently increased at $22^\circ\text{C}/\text{min}$ to 290°C , which was maintained for 20 min. Helium was used as the carrier gas at a linear velocity of $1.2\text{ mL}/\text{min}$, and all spectra were scanned within the range m/z 10–520. For the analysis of MeJA, the injection temperature was 250°C , and a fused-silica capillary column (TC-5) was used. The temperature program started at 80°C for 1 min and subsequently increased at $22^\circ\text{C}/\text{min}$ to 290°C , which was maintained for 20 min.

Identity of fungal sources. *B. cinerea* (MAFF No. 237695), *V. longisporum* (MAFF No. 243295), *G. fujikuroi* (MAFF No. 241712), *L. theobromae* (MAFF No. 306027), and *C. heterostrophus* (MAFF No. 243065) fungal cultures were obtained from The Genetic Resources Center, National Agriculture and Food Research Organization (NARO). The *F. oxysporum* culture (KF192-2) was a culture stock from the Laboratory of Plant Pathology, Research Faculty of Agriculture, Hokkaido University.

Feeding experiments. The fungi were grown in 50 mL flasks containing 20 mL of 2% potato-sucrose stationary medium at 25°C in the dark for 14 days. Deuterium-labeled compounds were dissolved in 1 mM aq. NH_4OH ($500\text{ }\mu\text{L}$) and added to the medium at a final concentration of 1 mM. For extraction, 10 mL of EtOAc was added to the culture together with addition of 1M aq. HCL (0.1 mL), and for the addition of internal standard, CJ-d7 ($1\text{ }\mu\text{g}$) was added to the separated EtOAc extract. The organic solvent in the extract were removed by a stream of N_2 . The resulting residues were dissolved in EtOAc. A portion of the EtOAc solution was subjected to

GC-MS analysis to detect CJ, and another portion was treated with CH_2N_2 and subjected to GC-MS analysis to detect MeJA.

[11,11- $^2\text{H}_2$, 10,10- $^2\text{H}_2$, 8,8- $^2\text{H}_2$, 2,2- $^2\text{H}_2$]methyl 12-oxophytodienoate (iso-MeOPDA-d8). iso-MeOPDA (9 mg), which was synthesized according to a reported method³³, was dissolved in CH_3OD (1.0 mL) in an ampule; CH_3ONa (4 mg) was then added to this solution. The ampule was sealed and heated at 65 °C for 5 days. The reaction mixture was quenched with 10% aq. NH_4Cl and a small volume of water, and the solvent was removed under low pressure. Then, the residue was extracted with diethyl ether (10 mL \times 3), and the solvent was removed under low pressure. The residue was purified by chromatography on silica gel (2 g) using ethyl acetate/*n*-hexane (1:4, v/v) to afford the product in a yield of 3.7 mg (35%). ^1H NMR (500 MHz, CDCl_3): δ 0.97 (t, $J = 7.6$ Hz, 3H), 1.31 (m, 6H), 1.49 (m, 2H), 1.58 (m, 2H), 2.13 (q, $J = 7.4$ Hz, 2H), 2.90 (d, $J = 7.2$ Hz, 2H), 3.64 (s, 3H), 5.19 (m, 1H), and 5.34 (m, 1H). FD-MS (m/z): 314 ($[\text{M}]^+$ (100), 313 (31), 312 (6.3), 311 (2.1), which revealed the labeling ration of iso-MeOPDA-d8 was to be 72%.

[4,4- $^2\text{H}_2$, 3,3- $^2\text{H}_2$, 1,1,1- $^2\text{H}_3$]CJ (CJ-d7). A solution of CJ (51.6 mg) in CH_3OD (1 mL) was added dropwise to a solution of CH_3ONa (40 mg) in 2 mL of CH_3OD . The mixture was refluxed for 5 days. NH_4Cl was added, and the organic solvent was evaporated. The residue was dissolved in water (1 mL). After the product was extracted with Et_2O (50 mL \times 3), the organic layer was dried over Mg_2SO_4 and concentrated *in vacuo* after filtration. The residue was purified by flash column chromatography with silica gel (40 g) using ethyl acetate/*n*-hexane (1:9, v/v) to give [4,4- $^2\text{H}_2$, 3,3- $^2\text{H}_2$, 1,1,1- $^2\text{H}_3$]CJ in a yield of 53.7 mg (99%). GC-MS (m/z): 171 ($[\text{M}]^+$ (100), 170 (4.3), 169 (1.7), which revealed the labeling ration of CJ-d7 was to be 94%. ^1H NMR (270 MHz, CDCl_3): δ 0.96 (t, $J = 7.0$ Hz, 3H), 2.13 (m, 2H), 2.91 (d, $J = 7.1$ Hz, 2H), 5.24 (m, 1H), and 5.34 (m, 1H). FI-MS (m/z): $[\text{M}^+]$ 171.

References

- Thomma, B. *et al.* Separate jasmonate-dependent and salicylate-dependent defense-response pathways in Arabidopsis are essential for resistance to distinct microbial pathogens. *Proc Natl Acad Sci USA* **95**, 15107–15111 (1998).
- Trusov, Y. *et al.* Heterotrimeric G proteins facilitate Arabidopsis resistance to necrotrophic pathogens and are involved in jasmonate signaling. *Plant Physiol* **140**, 210–220, doi:10.1093/pc/140.2.210 (2006).
- van Wees, S., Chang, H., Zhu, T. & Glazebrook, J. Characterization of the early response of Arabidopsis to *Alternaria brassicicola* infection using expression profiling. *Plant Physiol* **132**, 606–617, doi:10.1093/pc/132.3.606 (2003).
- Schaller, A. & Stintzi, A. Enzymes in jasmonate biosynthesis - Structure, function, regulation. *Phytochemistry* **70**, 1532–1538, doi:10.1016/j.phytochem.2009.07.032 (2009).
- Staswick, P. & Tirryaki, I. The oxylipin signal jasmonic acid is activated by an enzyme that conjugates it to isoleucine in Arabidopsis. *Plant Cell* **16**, 2117–2127, doi:10.1105/tpc.104.023549 (2004).
- Wasternack, C. & Kombrink, E. Jasmonates: Structural Requirements for Lipid-Derived Signals Active in Plant Stress Responses and Development. *ACS Chemical Biology* **5**, 63–77, doi:10.1021/cb900269u (2010).
- Xie, D., Feys, B., James, S., Nieto-Rostro, M. & Turner, J. COI1: an Arabidopsis gene required for jasmonate-regulated defense and fertility. *Science* **280**, 1091–1094 (1998).
- Katsir, L., Schillmiller, A., Staswick, P., He, S. & Howe, G. COI1 is a critical component of a receptor for jasmonate and the bacterial virulence factor coronatine. *Proc Natl Acad Sci USA* **105**, 7100–7105, doi:10.1073/pnas.0802332105 (2008).
- Yan, J. B. *et al.* The Arabidopsis CORONATINE INSENSITIVE1 Protein Is a Jasmonate Receptor. *Plant Cell* **21**, 2220–2236, doi:10.1105/tpc.109.065730 (2009).
- Sheard, L. B. *et al.* Jasmonate perception by inositol-phosphate-potentiated COI1-JAZ co-receptor. *Nature* **468**, 400–U301, doi:10.1038/nature09430 (2010).
- Yan, J. B. *et al.* Endogenous Bioactive Jasmonate Is Composed of a Set of (+)-7-iso-JA- Amino Acid Conjugates. *Plant Physiology* **172**, 2154–2164, doi:10.1104/pp.16.00906 (2016).
- Farmer, E., Johnson, R. & Ryan, C. Regulation of Expression of Proteinase Inhibitor Genes by Methyl Jasmonate and Jasmonic Acid. *Plant Physiol* **98**, 995–1002 (1992).
- Koch, T., Bandemer, K. & Boland, W. Biosynthesis of cis-jasmone: A pathway for the inactivation and the disposal of the plant stress hormone jasmonic acid to the gas phase? *Helvetica Chimica Acta* **80**, 838–850 (1997).
- Dabrowska, P. & Boland, W. iso-OPDA: An early precursor of cis-jasmone in plants? *Chembiochem* **8**, 2281–2285, doi:10.1002/cbic.200700464 (2007).
- Taki, N. *et al.* 12-oxo-phytodienoic acid triggers expression of a distinct set of genes and plays a role in wound-induced gene expression in Arabidopsis. *Plant Physiology* **139**, 1268–1283, doi:10.1104/pp.105.067058 (2005).
- Goetz, S. *et al.* Role of cis-12-Oxo-Phytodienoic Acid in Tomato Embryo Development. *Plant Physiology* **158**, 1715–1727, doi:10.1104/pp.111.192658 (2012).
- Flokova, K. *et al.* A previously undescribed jasmonate compound in flowering Arabidopsis thaliana - The identification of cis-(+)-OPDA-11e. *Phytochemistry* **122**, 230–237, doi:10.1016/j.phytochem.2015.11.012 (2016).
- Arnold, M. D. *et al.* The Recently Identified Isoleucine Conjugate of cis-12-Oxo-Phytodienoic Acid Is Partially Active in cis-12-Oxo-Phytodienoic Acid-Specific Gene Expression of Arabidopsis thaliana. *PLoS One* **11**, doi:10.1371/journal.pone.0162829 (2016).
- Schulze, B., Dabrowska, P. & Boland, W. Rapid enzymatic isomerization of 12-oxophytodienoic acid in the gut of lepidopteran larvae. *Chembiochem* **8**, 208–216, doi:10.1002/cbic.200600379 (2007).
- Dabrowska, P., Freitak, D., Vogel, H., Heckel, D. G. & Boland, W. The phytohormone precursor OPDA is isomerized in the insect gut by a single, specific glutathione transferase. *Proc Natl Acad Sci USA* **106**, 16304–16309, doi:10.1073/pnas.0906942106 (2009).
- Dabrowska, P., Shabab, M., Brand, W., Vogel, H. & Boland, W. Isomerization of the Phytohormone Precursor 12-Oxophytodienoic Acid (OPDA) in the Insect Gut A mechanistic and computational study. *J Biol Chem* **286**, 22348–22354, doi:10.1074/jbc.M111.244509 (2011).
- Vadassery, J., Reichelt, M. & Mithofer, A. Direct Proof of Ingested Food Regurgitation by Spodoptera littoralis Caterpillars during Feeding on Arabidopsis. *J Chem Ecol* **38**, 865–872, doi:10.1007/s10886-012-0143-5 (2012).
- Shabab, M., Khan, S. A., Vogel, H., Heckel, D. G. & Boland, W. OPDA isomerase GST16 is involved in phytohormone detoxification and insect development. *FEBS J* **281**, 2769–2783, doi:10.1111/febs.12819 (2014).
- Birkett, M. A. *et al.* New roles for cis-jasmone as an insect semiochemical and in plant defense. *Proc Natl Acad Sci USA* **97**, 9329–9334, doi:10.1073/Pnas.160241697 (2000).
- Bruce, T. J. A. *et al.* cis-Jasmone treatment induces resistance in wheat plants against the grain aphid, Sitobion avenae (Fabricius) (Homoptera: Aphididae). *Pest Manag Sci* **59**, 1031–1036, doi:10.1002/ps.730 (2003).

26. Schulz, S., Boppre, M. & Vanewright, R. I. Specific Mixtures of Secretions from Male Scent Organs of African Milkweed Butterflies (Danainae). *Philos T Roy Soc B* **342**, 161–181, doi:10.1098/Rstb.1993.0144 (1993).
27. Tanaka, K. *et al.* Highly Selective Tuning of a Silkworm Olfactory Receptor to a Key Mulberry Leaf Volatile. *Curr Biol* **19**, 881–890, doi:10.1016/j.cub.2009.04.035 (2009).
28. Powell, W. & Pickett, J. A. Manipulation of parasitoids for aphid pest management: progress and prospects. *Pest Manag Sci* **59**, 149–155, doi:10.1002/ps.550 (2003).
29. Siewers, V., Smedsgaard, J. & Tudzynski, P. The p450 monooxygenase BcABA1 is essential for abscisic acid biosynthesis in *Botrytis cinerea*. *Appl Environ Microbiol* **70**, 3868–3876, doi:10.1128/AEM.70.7.3868-3876.2004 (2004).
30. Miersch, O., Bohlmann, H. & Wasternack, C. Jasmonates and related compounds from *Fusarium oxysporum*. *Phytochemistry* **50**, 517–523, doi:10.1016/s0031-9422(98)00596-2 (1999).
31. Thatcher, L. F., Manners, J. M. & Kazan, K. *Fusarium oxysporum* hijacks COI1-mediated jasmonate signaling to promote disease development in *Arabidopsis*. *Plant J* **58**, 927–939, doi:10.1111/j.1365-313X.2009.03831.x (2009).
32. Tsukada, K., Takahashi, K. & Nabeta, K. Biosynthesis of jasmonic acid in a plant pathogenic fungus, *Lasiodiplodia theobromae*. *Phytochemistry* **71**, 2019–2023, doi:10.1016/j.phytochem.2010.09.013 (2010).
33. Christensen, S. A. *et al.* Maize death acids, 9-lipoxygenase-derived cyclopentane(a)nones, display activity as cytotoxic phytoalexins and transcriptional mediators. *Proc Natl Acad Sci USA* **112**, 11407–11412, doi:10.1073/pnas.1511131112 (2015).
34. Galka, P. W., Ambrose, S. J., Ross, A. R. S. & Abrams, S. R. Syntheses of deuterated jasmonates for mass spectrometry and metabolism studies. *J Label Comp Radiopharm* **48**, 797–809, doi:10.1002/jlcr.994 (2005).
35. Kajiwara, A., Abe, T., Hashimoto, T., Matsuura, H. & Takahashi, K. Efficient Synthesis of (+)-cis-12-Oxo-phytodienoic Acid by an *in Vitro* Enzymatic Reaction. *Biosci Biotechnol Biochem* **76**, 2325–2328, doi:10.1271/bbb.120506 (2012).
36. Matsuura, H., Ohmori, F., Kobayashi, M., Sakurai, A. & Yoshihara, T. Qualitative and quantitative analysis of endogenous jasmonoids in potato plant (*Solanum tuberosum* L.). *Biosci Biotechnol Biochem* **64**, 2380–2387 (2000).
37. Lauchli, R. & Boland, W. Efficient synthesis of H-2(2) -tetrahydrodicranenone B and a 3-oxa-analogue resistant against beta-oxidation. *Tetrahedron* **59**, 149–153 (2003).
38. Niki, T., Mitsuhashi, I., Seo, S., Ohtsubo, N. & Ohashi, Y. Antagonistic effect of salicylic acid and jasmonic acid on the expression of pathogenesis-related (PR) protein genes in wounded mature tobacco leaves. *Plant and Cell Physiol* **39**, 500–507 (1998).

Acknowledgements

The MeJA used as the starting material for the synthesis of deuterium-labeled compounds was kindly provided by the ZEON Corporation (Tokyo, Japan). We acknowledge the assistance of Dr. Eri Fukushi and Mr. Yusuke Takata (Research Faculty of Agriculture, Hokkaido University) in obtaining the spectroscopic data.

Author Contributions

R.M., N.A., K.T., Y.T., and W.S. performed the biochemical experiments. H.M., N.K., and K.M. supported the experiments, and R.M. and H.M. wrote the manuscript.

Additional Information

Supplementary information accompanies this paper at doi:10.1038/s41598-017-05851-7

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017