



Title	Genetic structure of <i>Vaccinium vitis-idaea</i> in lowland cool spot and alpine populations : microrefugia of alpine plants in the midlatitudes
Author(s)	Shimokawabe, Ayuma; Yamaura, Yuichi; Sueyoshi, Masanao; Kudo, Gaku; Nakamura, Futoshi
Citation	Alpine botany, 126(2), 143-151 https://doi.org/10.1007/s00035-016-0169-3
Issue Date	2016-10
Doc URL	http://hdl.handle.net/2115/67222
Rights	The original publication is available at www.springerlink.com .
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Alp. Bot.126-2_143-151.pdf



[Instructions for use](#)

1 Genetic structure of *Vaccinium vitis-idaea* in lowland cool spot and alpine populations:

2 Microrefugia of alpine plants in the mid-latitudes

3

4 Ayuma Shimokawabe¹, Yuichi Yamaura^{1,2}, Masanao Sueyoshi^{1,3}, Gaku Kudo⁴, and Futoshi

5 Nakamura¹

6

7 ¹Graduate School of Agriculture, Hokkaido University, Sapporo, Japan

8 ²Department of Forest Vegetation, Forestry and Forest Products Research Institute, Tsukuba,

9 Ibaraki, Japan

10 ³Aqua Restoration Research Center, Public Works Research Institute, Kakamigahara, Gifu,

11 Japan

12 ⁴Faculty of Environmental Earth Science, Hokkaido University, Sapporo, Hokkaido, Japan

13

14 Corresponding author: yamaura@ffpri.affrc.go.jp

15 Tel: +81-29-829-8224; Fax: +81-29-873-1542

16

17

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

Abstract (246 words: 150–250 words)

Local cool spots (wind-holes) in lowland areas of mid-latitudes may act as microrefugia for cold-adapted species outside of their typical alpine habitats. We examined the genetic structure of *Vaccinium vitis-idaea*, a common alpine species in Japan, in eight lowland wind-hole and five surrounding alpine populations. We collected leaf samples and genotyped seven microsatellite loci. Clonal patches (genets) were common in almost all populations. An analysis of annual shoot growth suggested that individuals in the wind-hole populations were long-lived (>500 years old). Genetic diversity (allelic richness) and differentiation (F_{ST}) of the wind-hole populations were lower and higher than those of the alpine populations, respectively. No significant isolation-by-distance trend in the genetic structure was detected for the wind-hole or alpine populations. All wind-hole populations had negative inbreeding coefficients (F_{IS}), suggesting no tendency toward homozygosity due to inbreeding, regardless of the small populations geographically isolated from the large alpine populations. Therefore, wind-holes may harbor genetically isolated but stable populations due to clonal growth, limited gene flow, and abortion of selfed seeds by early acting inbreeding depression. Analysis of molecular variance demonstrated that genetic variations among and within populations contributed more to regional genetic diversity than those between wind-hole and alpine populations, suggesting that the wind-hole and alpine populations are important for maintaining the genetic diversity of mid-latitude *V. vitis-idaea* populations. On

37 the other hand, Bayesian clustering showed that some wind-hole populations geographically
38 close to the alpine populations had mixed genetic compositions of the alpine and wind-hole
39 populations.

40

41

Introduction

42 Many species inhabiting mountainous areas have upwardly shifted their distributions
43 along with ongoing climate change (Walther et al. 2005; Forister et al. 2010). These
44 elevational shifts are large in alpine plants (Lenoir et al. 2008), and the future distributions of
45 these species are predicted to decline greatly due to the limited area for escape in alpine and
46 geographically isolated habitats (Thuiller et al. 2005; Thuiller 2007). Large-scale range shifts
47 associated with climate change have occurred in past glacial periods, and it has been
48 suggested that small populations persist in local suitable habitats within broadly unsuitable
49 geographical areas (Stewart and Lister 2001; McLachlan et al. 2005; Parducci et al. 2012).
50 These local refugia are called “microrefugia” in contrast to the large, continuous
51 “macrorefugia” (Rull 2009).

52 Microrefugia are characterized by locally specialized microclimates that are only
53 loosely linked to the surrounding macroclimate. As such, they can provide long-lasting small
54 habitats for specialist species (Dobrowski 2011). In the current interglacial period, alpine
55 habitats are macrorefugia, whereas cool spots situated in topographic depressions, heavily
56 incised valley bottoms, or debris-covered glaciers can be microrefugia (Dobrowski 2011;
57 Gentili et al. 2015). As these small cool spots can be distributed over a broad geographical
58 range (Shimokawabe et al. 2015), microrefugia may play important roles in the response of
59 many species to climate change (Mosblech et al. 2011; Hannah et al. 2014).

60 *Vaccinium vitis-idaea* L. (Ericaceae) is a common alpine dwarf shrub inhabiting
61 mid-latitude alpine and subarctic regions. *V. vitis-idaea* also grow in lowland cool spots,
62 including wind-hole micro-topographies (Sato et al. 1993; Sato 1995; Růžička 1999). Wind
63 holes frequently occur at the bottom of talus slopes containing volcanic bedrock (Shimizu
64 2004), where cool conditions are maintained in the vicinity by the preferential flow of cool air
65 generated in interstitial spaces created by rock fragments or colluvia. We call such cool spots
66 “wind-hole sites.” Wind-hole sites harbor various taxa of alpine species that are not found in
67 the surrounding areas (Růžička 1999; Shimokawabe et al. 2015). Shioi (1975) speculated
68 that lowland *V. vitis-idaea* populations have been maintained in local wind-hole sites since the
69 last glacial period in Hokkaido, northern Japan. If this is true, not only alpine habitats
70 (macrorefugia) but also local wind-hole sites (microrefugia) are important habitats for the
71 maintenance of *V. vitis-idaea* population genetic diversity in mid-latitude regions.

72 Populations in small and isolated habitats are subject to heightened extinction risks
73 due to inbreeding depression, genetic drift, and demographic stochasticity (Lande 1988; Reed
74 and Frankham 2003). Several mechanisms contribute to the long-term maintenance of
75 microrefugia populations (Hampe and Jump 2011). One exogenous mechanism is migration
76 from the surrounding populations (Mosblech et al. 2011), and one endogenous mechanism is
77 purging deleterious alleles and self-fertilization/compatibility (Mee and Moore 2014).
78 Wind-hole populations are small and isolated from alpine populations, and the mechanisms of

79 population maintenance may be manifested in their genetic structure.

80 In the present study, we examined the genetic structure of *V. vitis-idaea* populations in
81 northeastern Hokkaido, including alpine and lowland wind-hole populations. We also
82 measured the annual shoot growth to estimate the age of clonal patches. We discuss the
83 mechanism of population maintenance at wind-hole sites and its ecological significance under
84 ongoing climate change.

85

86

87

Methods

88

Species studied and field sampling

89 *V. vitis-idaea* ($2n = 24$) is an evergreen dwarf shrub (5–20 cm in height) with creeping
90 stems under the soil surface (Iwatsuki et al. 1993). Based on genetic analyses, Ikeda et al.
91 (2015) suggested that Japanese *V. vitis-idaea* populations have persisted since before the last
92 glacial period. The major habitats in our study area (Engaru and Kitami: 43.7–43.9° N and
93 143.0–143.4° E, respectively) were alpine zones and lowland wind-hole sites. This species is
94 also known to occur on acid soils in boreal forests and bogs (Garkava-Gustavsson et al. 2005).
95 However, lowland *V. vitis-idaea* populations in Japan other than those at wind-hole sites are
96 limited and fragmentally distributed only in the coastal wetlands of northern/eastern
97 Hokkaido (Umezawa 2007). Flowering occurs in early summer; shrubs are pollinated mainly

98 by bees, and seeds are distributed by animals (Ritchie 1955; Iwatsuki et al. 1993). Upon
99 self-fertilization, partial self-sterility leads to reduced numbers of developed seeds due to
100 early acting inbreeding depression (Guillaume and Jacquemart 1999). *V. vitis-idaea* can form
101 large clonal patches (>30 m) by expanding via underground stems (Persson and Gustavsson
102 2001).

103 The study area was located in northeastern Hokkaido (Fig. 1). Alpine habitats
104 harboring *V. vitis-idaea* cover the western high elevation area (>1,500 m), and wind-hole sites
105 with *V. vitis-idaea* are scattered across the eastern lowland area. This area is topographically
106 complex and includes steep slopes and bedrock dominated by pumice flow deposits
107 (1:200,000 surface geology map by the Ministry of Land, Infrastructure, Transport, and
108 Tourism of Japan). The mean annual temperature during the past 30 years was 5.8°C (mean
109 temperatures of the coldest and warmest months were -8.3 and 19.9°C, respectively) and
110 annual precipitation was 794.5 mm (Japan Meteorological Agency,
111 <http://www.jma.go.jp/jma/indexe.html>). The temperature at ground level remains 7°C lower
112 than the air temperature during the summer (Shimokawabe et al. 2015).

113 We collected leaves in August–September 2013 from eight wind-hole sites and five
114 separate alpine habitats (Fig. 1). The sample size was 23–31 leaves from each site, totaling
115 393 samples (Table 1). No additional wind-hole sites known to harbor *V. vitis-idaea* are
116 present in our study area (Shimokawabe et al. 2015). We uniformly sampled leaves from all

117 areas of the wind-hole sites and from the ridge areas in the alpine habitats (sampling interval:
118 3–10 m for wind-hole sites, 5–10 m for alpine habitats). We recorded the locations of the
119 sampling points using a global positioning system (Garmin GPSMAP 62SJ; Garmin
120 International, Inc., Olathe, KS, USA). Leaf samples were dried in silica gel and preserved at
121 room temperature until analysis.

122

123

Microsatellite analysis

124 We extracted DNA using the cetyltrimethylammonium bromide (CTAB) method. In
125 brief, the sample was crushed in CTAB extraction buffer, chloroform-isoamyl alcohol was
126 added, the aqueous layer containing the DNA was separated, the DNA was purified with
127 isopropanol and ethanol, and the DNA was preserved in Tris-EDTA buffer. Seven
128 microsatellite loci (CA169F, CA236F, NA741, NA800, NA1040, VCC_I2 and VCC_K4:
129 Boches et al. 2005; Appendix A) were amplified and developed for a related species
130 (*Vaccinium corymbosum* L.) using polymerase chain reaction (PCR) and a thermal cycler
131 (Applied Biosystems 2720 Thermal Cycler; Applied Biosystems Inc., Foster City, CA, USA).
132 Following Boches et al. (2005), we used the following PCR program: 3 min at 94°C, followed
133 by 35 cycles of 40 s at 94°C, 40 s at 60°C or 62°C, and 40 s at 72°C with a final rest for 30
134 min at 72°C. Microsatellite fragments were analyzed using the Applied Biosystems 3130
135 Genetic Analyzer, and the genotypes were coded using Peak Scanner ver. 2.0 (Applied

136 Biosystems). GeneScan 500 LIZ Size Standard (Applied Biosystems) was used as the size
137 standard.

138

139 Population genetics analysis

140 A total of 244 multi-locus genotypes in seven loci were identified, and 69 were
141 assigned to more than two different samples (17.6%). Samples with the same genotypes in all
142 loci were treated as the same individual (genet), and the genet was used as the analytical unit.
143 We calculated the probability that two different individuals would have the same genotype
144 (PI: probability of identity) using GenAIEx 6.501 (Peakall and Smouse 2006) to evaluate
145 misassignment of different individuals to genets. We tested departures from Hardy–Weinberg
146 equilibrium based on the chi-square (χ^2) test using GenAIEx 6.501 and obtained the
147 inbreeding coefficient (F_{IS}). We examined the presence of null alleles for each population
148 using GENEPOP 4.3 (Rousset 2008). The Bonferroni correction was used to determine the
149 significance levels of the equilibrium tests.

150 We obtained F_{ST} values as genetic differentiation indices for each pair of populations.

151 The value of this index can be affected by population history, mutation, gene flow, and genetic
152 drift (Marko and Hart 2011). The correlation between the F_{ST} value and geographic distance
153 (isolation by distance, IBD: Wright 1943) was examined based on the Mantel tests for (i) all
154 populations, (ii) wind-hole and wind-hole pairs, and (iii) alpine and alpine pairs. We also

155 examined the differences in genetic differentiation (F_{ST}) and geographic distances among
156 populations for (i) wind-hole and wind-hole pairs, (ii) wind-hole and alpine pairs, and (iii)
157 alpine and alpine pairs with multiple comparison tests (Steel–Dwass method). We obtained
158 the geographic distance as three-dimensional straight-line distances considering elevational
159 differences. The F_{ST} values were calculated, Mantel tests were conducted using GenAlEx
160 6.501, and multiple comparison tests were performed using R ver. 2.15.2 (R Core Team,
161 Vienna, Austria).

162 We calculated allelic richness and private allelic richness (the number of unique alleles
163 for each population, corrected for sample size) as genetic diversity indices (Kalinowski 2004)
164 using HP-Rare v. June-6-2006 (Kalinowski 2005). The diversity indices were compared
165 between wind-hole sites and alpine habitats using *t*-tests. The correlation coefficient between
166 allelic richness and the index of genetic isolation was determined as the mean F_{ST} value for
167 each population. The mean F_{ST} was obtained from the F_{ST} values of all other populations, and
168 the correlations were examined separately for the wind-hole sites and alpine habitats.

169 We conducted analyses of molecular variance (AMOVA) based on these F_{ST} values to
170 divide genetic variation into those occurring within populations, among populations, and
171 among habitats using GenAlEx 6.501. STRUCTURE 2.3.4 (Pritchard et al. 2000) was used to
172 examine individual based Bayesian clustering to estimate the most likely number of
173 genetically differentiated populations (K). An admixture model with the correlated allele

174 frequencies and informative locations was employed. The optimal number of clusters from 1
175 to 13 was inferred by 200,000 interactions following a 100,000 step burn-in with 10 replicate
176 runs for each K -value. STRUCTURE can identify the highest hierarchical level of a
177 population but often fails to detect hierarchical genetic structure at lower levels (Evanno et al.
178 2005). Therefore, subsequent runs were executed with the same method to reveal additional
179 hierarchical genetic clusters when $K > 1$ was identified at the initial run. The STRUCTURE
180 results and the online program STRUCTURE HARVESTER ver. 0.6.94 (Earl and vonHoldt
181 2012) were used to determine the best K number. This program infers the best K value using
182 the delta K based on the rate of change in the log probability of the data (ΔK) between
183 successive K values (the best K has the highest delta K value) (Evanno et al. 2005). Finally,
184 the online program CLUMPAK was used to visualize the spatial genetic structure estimated
185 by STRUCTURE (Kopelman et al. 2015).

186

187 Annual growth measurements

188 We measured the annual growth of shoots and estimated their ages to infer the
189 persistence of individuals at the wind-hole sites. We measured annual growth length of the
190 previous year's shoots from 10 individuals each in the six wind-hole populations during May
191 2015. We thereby obtained a mean value for the 10 individuals in each population. The
192 longest length between ramets was obtained from genetic analyses. We treated this length as

193 the size of the oldest individual in the corresponding population, and estimated its age
194 assuming circular growth: Age = $(0.5 \times \text{distance between ramets})/\text{annual growth}$.

195

196

Results

197 The mean PI across 13 populations, which indicates the probability of misassignment
198 of different individuals to genets, was very low (2×10^{-4}). Therefore, we treated plants with
199 the same genotype as the same individual (genet). Clones were detected at all alpine and
200 wind-hole sites, except at Mt. Mur (Table 1). The longest distances within clones were 14.3–
201 101.8 m (Table 1). Large ramets had between-ramet lengths > 50 m (Appendix B). The mean
202 annual shoot growth of the six wind-hole populations was $3.90 \text{ cm year}^{-1}$. The largest clone
203 was estimated to be 594 years old, and four populations had clones that were > 500 years old
204 (Table 1). Note that our calculations may have underestimated the ages of the clones because
205 we assumed a uniform circular growth pattern. However, this result suggests that *V.*
206 *vitis-idaea* individuals have been maintained at the wind-hole sites for hundreds of years. We
207 observed deviations from Hardy–Weinberg equilibrium at a mean of two populations per
208 locus (adjusted $P < 0.05$). All wind-hole populations had negative F_{IS} values, and six
209 populations differed significantly from zero ($P < 0.05$; Table 1). The existence of null alleles
210 was suggested for multiple loci and populations. However, their frequencies were small, and
211 we used all loci and genets (mean frequency per loci = 0.03).

212 The Mantel tests did not reveal any significant IBD among populations ($r = -0.12$, P
213 $= 0.23$; Fig. 2) or within habitat types (wind-hole–wind-hole pairs: $r = 0.32$, $P = 0.16$; alpine–
214 alpine pairs: $r = 0.05$, $P = 0.54$; Fig. 2). However, the alpine–alpine pairs and the wind-hole–
215 wind-hole pairs had lower and higher F_{ST} values, respectively, than those obtained among
216 habitat pairs (Steel–Dwass method: $P < 0.05$; Fig. 3a). The wind-hole–alpine pairs had larger
217 geographic distances than the other two pairs (Steel–Dwass method: $P < 0.01$; Fig. 3b). These
218 results suggest that the wind-hole–wind-hole pairs had large F_{ST} values regardless of their
219 short geographic distances. Some wind-hole–alpine pairs had F_{ST} values similar to those of
220 alpine–alpine pairs (Fig. 2).

221 The wind-hole populations had lower allelic richness (t -test: $P < 0.01$; Table 1) and
222 larger variations in allelic richness (Fig. 3c) than the alpine populations. No differences in
223 private allelic richness were detected between the wind-hole and alpine populations, and both
224 populations had unique loci (t -test: $P = 0.09$; Fig. 3d, Table 1). Allelic richness and mean F_{ST}
225 values were negatively correlated in the wind-hole populations ($r = -0.87$, $P < 0.01$), but not
226 in the alpine populations ($r = 0.13$, $P = 0.84$; Fig. 4).

227 All three levels (between habitats, among populations, and within populations)
228 significantly contributed to the genetic variation. Variations between habitat types were
229 smaller than those among and within populations, and within-population variations dominated
230 total genetic variation (Table 2).

231 STRUCTURE revealed two main genetic clusters with the highest delta K value (Fig.
232 5a). The alpine populations were assigned to cluster 1, and most genets of the wind-hole
233 populations were assigned to cluster 2. However, some genets of three wind-hole populations
234 (HiH, HiN, and Sin) were assigned to cluster 1. These populations were geographically closer
235 to the alpine sites than the other wind-hole sites (Fig. 1). Therefore, we separated 13
236 populations into two genetic groups with contrasting genetic composition and examined
237 subsequent STRUCTURE runs for each of the two groups (Fig. 5b, c). Subsequent analyses
238 revealed seven genetic clusters within the wind-hole and alpine populations ($K = 4$ for the
239 first group, including five wind-hole populations and $K = 3$ for the second group, including
240 three wind-hole and five alpine populations). Most populations formed unique genetic clusters,
241 indicating high genetic variation among populations, as suggested by AMOVA. However,
242 some populations, which were spatially close to each other, formed similar clusters (e.g., Iga
243 and Mir; HiH, HiN, and Sin; Mt. Hir and Mt. Mur).

244

245

246

Discussion

247

Population maintenance at the wind-hole sites

248

The existence of large clonal patches (>50 m spans) in the wind-hole and alpine

249

populations suggests that individual plants can persist for hundreds of years. Such long clonal

250 lifespans have also been reported in other ericaceous alpine species (Kameyama et al. 2008)
251 and was reported by another genetic study on *V. vitis-idaea* (Persson and Gustavsson 2001). A
252 long lifespan through clonal growth may be important for persistence of small populations,
253 such as wind-hole populations. A low level of heterozygosity is expected for small relict
254 populations if self-fertilization is common (Mee and Moore 2014). However, wind-hole
255 populations had negative F_{IS} values, indicating the existence of mechanisms for avoiding
256 self-fertilization (Wright 1965). This may be because of a very long generation time, which
257 would delay the genetic homogenization caused by inbreeding. On the other hand, early
258 acting inbreeding depression strictly prevents the formation of selfing seeds in *V. vitis-idaea*
259 (Guillaume and Jacquemart 1999). Although restricting seed production by selfing may
260 increase the risk of local extinction by decreasing mating opportunities (Byers and Meagher
261 1992), the existence of long-lived clones would alleviate the extinction risk.

262 Genetic diversity was lower in the wind-hole populations than that in the alpine
263 populations. Although we did not find IBD in the wind-hole populations, a negative
264 correlation between the level of genetic isolation (F_{ST}) and genetic diversity suggests that (1)
265 the wind-hole populations are isolated and subject to genetic drift (Hutchison and Templeton
266 1999) and (2) that gene flow between the wind-hole populations is not dependent on
267 geographic distance. The latter may be partly due to the existence of undiscovered wind-hole
268 populations in the study area. As genetic diversity of a small population can increase by rare

269 immigration of gametes (Ingvarsson 2001), genetic diversity of a wind-hole population would
270 strongly depend on genetic exchange with other populations. In our study, gene flow was
271 lower in the wind-hole–wind-hole pairs than that in the alpine–alpine pairs. Pollinators and
272 seed dispersers would infrequently visit small wind-hole sites surrounded by forest.

273

274 Genetic diversity of the alpine populations

275 We did not find IBD among the alpine populations, and genetic diversity and
276 differentiation were higher and lower than those in in the wind-hole populations, respectively.
277 Although we selected alpine populations from separate mountains, these sites were
278 structurally connected by alpine habitats. Ikeda et al. (2015) also reported the lack of
279 nationwide genetic differentiation in *V. vitis-idaea* populations in Japan using nuclear and
280 chloroplast DNA markers. This finding suggests long-distance and efficient seed dispersal by
281 birds during the post-glacial period. Therefore, pollinators and seed dispersers may move
282 between alpine populations. High genetic diversity would have been maintained in large-sized
283 alpine populations with small effects from genetic drift and metapopulation structure among
284 populations (Billington 1991).

285

286 Ecological significance of wind-hole populations

287 The AMOVA results suggested that genetic variation within and among populations

288 made greater contributions to regional genetic diversity than did differences between the
289 alpine and wind-hole populations. We found unique genetic clustering between most of the
290 wind-hole populations and unique loci in almost all populations, suggesting that the alpine
291 and wind-hole populations are important to regional genetic diversity of *V. vitis-idaea*.
292 Although the wind-hole populations had lower genetic diversity than that of the alpine
293 populations, some wind-hole populations had relatively high genetic diversity and were likely
294 subject to gene flow from surrounding populations. Our results also convincingly demonstrate
295 the long-term persistence and genetic isolation of the wind-hole populations.

296 Microrefugia (wind-hole sites, in our case) are local small habitats, but can be
297 long-lived and span a broad geographical area, at least in our study area (Shimokawabe et al.
298 2015). Růžička et al. (2015) suggested that wind-hole sites in the Czech Republic will
299 maintain cool conditions even in warmer climates. The *V. vitis-idaea* wind-hole populations
300 geographically close to the alpine populations included the genetic composition of the alpine
301 populations, suggesting that wind-hole habitats act as safe sites for wind-hole and alpine
302 populations of cold-adapted species under global warming conditions. Not all cold-adapted
303 species would benefit from microrefugia (Hylander et al. 2015). Nevertheless, because
304 localized unique geographic features, which can be the bases of many microrefugia (Hjort et
305 al. 2015), occur in economically unproductive sites (Lindenmayer and Franklin 2002), we
306 have suggested previously that identifying and protecting potential microrefugia is a robust

307 and cost-effective way to mitigate the impact of climate change (Shimokawabe et al. 2015).

308

309

310 **Acknowledgements**

311 We thank the members of the Department of Forest Science and the Forest Ecosystem

312 Management Group of Hokkaido University for their field assistance and helpful discussions

313 during this study. We are also grateful to A. Hirao for technical comments on this study. The

314 manuscript was greatly improved by the comments from J. Stöcklin and two anonymous

315 reviewers. AS was supported by a Sasakawa Scientific Research Grant from the Japan

316 Science Society (no. 26-542). AS, YY, MS, and FN were supported by the Japan Society for

317 the Promotion of Science (JSPS) KAKENHI Grant no. 23248021, and YY was supported by

318 JSPS KAKENHI Grant no. 20580947.

319

320

321 **References**

- 322 Billington HL (1991) Effect of population size on genetic variation in a dioecious conifer.
323 *Conserv Biol* 5:115–119
- 324 Boches PS, Bassil NV, Rowland LJ (2005) Microsatellite markers for *Vaccinium* from EST
325 and genomic libraries. *Mol Ecol Notes* 5:657–660
- 326 Byers DL, Meagher TR (1992) Mate availability in small populations of plant species with
327 homomorphic sporophytic self-incompatibility. *Heredity* 68:353–359
- 328 Dobrowski SZ (2011) A climatic basis for microrefugia: the influence of terrain on climate.
329 *Global Change Biol* 17:1022–1035
- 330 Earl DA, vonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for
331 visualizing STRUCTURE output and implementing the Evanno method. *Conservation*
332 *Genetics Resources* 4:359–361
- 333 Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using
334 the software structure: a simulation study. *Mol Ecol* 14:2611–2620
- 335 Forister ML et al. (2010) Compounded effects of climate change and habitat alteration shift
336 patterns of butterfly diversity. *Proc Natl Acad Sci USA* 107:2088–2092
- 337 Garkava-Gustavsson L, Persson HA, Nybom H, Rumpunen K, Gustavsson BA, Bartish IV
338 (2005) RAPD-based analysis of genetic diversity and selection of lingonberry
339 (*Vaccinium vitis-idaea* L.) material for *ex situ* conservation. *Genet Resour Crop Evol*
340 52:723–735
- 341 Gentili R, Baroni C, Caccianiga M, Armiraglio S, Ghiani A, Citterio S (2015) Potential
342 warm-stage microrefugia for alpine plants: Feedback between geomorphological and
343 biological processes. *Ecol Complex* 21:87–99
- 344 Guillaume P, Jacquemart A-L (1999) Early-inbreeding depression in *Vaccinium myrtillus* and
345 *V. vitis-idaea*. *Protoplasma* 208:107–114
- 346 Hampe A, Jump AS (2011) Climate relicts: past, present, future. *Annu Rev Ecol Evol Syst*
347 42:313–333
- 348 Hannah L, Flint L, Syphard AD, Moritz MA, Buckley LB, McCullough IM (2014) Fine-grain
349 modeling of species' response to climate change: holdouts, stepping-stones, and
350 microrefugia. *Trends Ecol Evol* 29:390–397
- 351 Hjort J, Gordon JE, Gray M, Hunter ML (2015) Why geodiversity matters in valuing nature's
352 stage. *Conserv Biol* 29:630–639
- 353 Hutchison DW, Templeton AR (1999) Correlation of pairwise genetic and geographic
354 distance measures: inferring the relative influences of gene flow and drift on the
355 distribution of genetic variability. *Evolution* 53:1898–1914
- 356 Hylander K, Ehrlén J, Luoto M, Meineri E (2015) Microrefugia: not for everyone. *Ambio*
357 44:60–68

358 Ikeda H et al. (2015) Persistent history of the bird-dispersed arctic–alpine plant *Vaccinium*
 359 *vitis-idaea* L. (Ericaceae) in Japan. *J Plant Res* 128:437–444
 360 Ingvarsson PK (2001) Restoration of genetic variation lost – the genetic rescue hypothesis.
 361 *Trends Ecol Evol* 16:62–63
 362 Iwatsuki K, Yamazaki T, Boufford DE, Ohba H (1993) Flora of Japan. Volume. IIIa.
 363 Angiospermae Dicotyledoneae Sympetalae(a). Kodansya, Tokyo
 364 Kalinowski ST (2004) Counting alleles with rarefaction: private alleles and hierarchical
 365 sampling designs. *Conserv Genet* 5:539–543
 366 Kalinowski ST (2005) hp-rare 1.0: a computer program for performing rarefaction on
 367 measures of allelic richness. *Mol Ecol Notes* 5:187–189
 368 Kameyama Y, Kasagi T, Kudo G (2008) A hybrid zone dominated by fertile F₁s of two alpine
 369 shrub species, *Phyllodoce caerulea* and *Phyllodoce aleutica*, along a snowmelt
 370 gradient. *J Evol Biol* 21:588–597
 371 Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I (2015) CLUMPAK: a
 372 program for identifying clustering modes and packaging population structure
 373 inferences across *K*. *Mol Ecol Resour* 15:1179–1191
 374 Lande R (1988) Genetics and demography in biological conservation. *Science* 241:1455–
 375 1460
 376 Lenoir J, Gegout JC, Marquet PA, de Ruffray P, Brisse H (2008) A significant upward shift in
 377 plant species optimum elevation during the 20th century. *Science* 320:1768–1771
 378 Lindenmayer DB, Franklin JF (2002) Conserving forest biodiversity: a comprehensive
 379 multiscaled approach. Island Press, Washington, D.C.
 380 Marko PB, Hart MW (2011) The complex analytical landscape of gene flow inference.
 381 *Trends Ecol Evol* 26:448–456
 382 McLachlan JS, Clark JS, Manos PS (2005) Molecular indicators of tree migration capacity
 383 under rapid climate change. *Ecology* 86:2088–2098
 384 Mee JA, Moore J-S (2014) The ecological and evolutionary implications of microrefugia. *J*
 385 *Biogeogr* 41:837–841
 386 Mosblech NAS, Bush MB, van Woesik R (2011) On metapopulations and microrefugia:
 387 palaeoecological insights. *J Biogeogr* 38:419–429
 388 Parducci L et al. (2012) Glacial survival of boreal trees in northern Scandinavia. *Science*
 389 335:1083–1086
 390 Peakall ROD, Smouse PE (2006) genalex 6: genetic analysis in Excel. Population genetic
 391 software for teaching and research. *Mol Ecol Notes* 6:288–295
 392 Persson HA, Gustavsson BA (2001) The extent of clonality and genetic diversity in
 393 lingonberry (*Vaccinium vitis-idaea* L.) revealed by RAPDs and leaf-shape analysis.
 394 *Mol Ecol* 10:1385–1397
 395 Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using

396 multilocus genotype data. *Genetics* 155:945–959
397 Reed DH, Frankham R (2003) Correlation between fitness and genetic diversity. *Conserv Biol*
398 17:230–237
399 Ritchie JC (1955) *Vaccinium vitis-idaea* L. *J Ecol* 43:701–708
400 Rousset F (2008) genepop'007: a complete re-implementation of the genepop software for
401 Windows and Linux. *Mol Ecol Resour* 8:103–106
402 Rull V (2009) Microrefugia. *J Biogeogr* 36:481–484
403 Růžička V (1999) The freezing scree slopes and their arachnofauna. *Decheniana-Beihefte*
404 37:141–147
405 Růžička V, Zacharda M, Šmilauer P, Kučera T (2015) Can paleorefugia of cold-adapted
406 species in talus slopes resist global warming? *Boreal Environ Res* 20:403–412
407 Sato K (1995) An outline of the cool-spots site vegetation of Hokkaido, Japan. *Bull Higashi*
408 *Taisetsu Museum Nat Hist* 17:107–115
409 Sato K, Kudo G, Uemura S (1993) Cool-spots site vegetation in Izariiri-Heide, northern Japan.
410 *Jpn J Ecol* 43:91–98
411 Shibo T (1975) Supplementally report on the periglacial phenomena observed in Kitami
412 District. *Mem Kitami Inst Techn* 7:163–194
413 Shimizu C (2004) An information on the cool air blow holes (wind-holes) including the ice
414 caves in Japan: references to the surrounding landforms and the existence of sporadic
415 perm afrost. *Komazawa Geogr* 40:121–148
416 Shimokawabe A, Yamaura Y, Akasaka T, Sato T, Shida Y, Yamanaka S, Nakamura F (2015)
417 The distribution of cool spots as microrefugia in a mountainous area. *PLoS ONE*
418 10:e0135732
419 Stewart JR, Lister AM (2001) Cryptic northern refugia and the origins of the modern biota.
420 *Trends Ecol Evol* 16:608–613
421 Thuiller W (2007) Climate change and the ecologist. *Nature* 448:550–552
422 Thuiller W, Lavorel S, Araujo MB, Sykes MT, Prentice IC (2005) Climate change threats to
423 plant diversity in Europe. *Proc Natl Acad Sci USA* 102:8245–8250
424 Umezawa S (2007) Wild flowers of Hokkaido. New edn. Hokkaido University Press, Sapporo
425 Walther G-R, Beißner S, Burga CA (2005) Trends in the upward shift of alpine plants. *J Veg*
426 *Sci* 16:541–548
427 Wright S (1943) Isolation by distance. *Genetics* 28:114–138

428

429

430 Table 1. Habitat types and characteristics of *Vaccinium vitis-idaea* populations. Size and perimeter were not measured for the high mountain
 431 population. Asterisks (*) on the F_{IS} value indicates a significant difference from zero (<0.05) based on mean and standard deviation across seven
 432 loci.

Site	Habitat ¹	Ele. ²	Pop. area ³	Peri ⁴	Dist. alp. ⁵	Growth ⁶	Dist. ramets ⁷	Age ⁸	N^9	# genet ¹⁰	AR ¹¹	Pr. AR ¹²	Mean F_{ST}	F_{IS}
Ara	WH	309	2022	278	13.5	6.26 ± 1.66	74.4	594.2	31	10	2.45	0.53	0.19	-0.67*
Goj	WH	519	516	252	18.0	3.46 ± 1.04	32.2	465.3	31	13	2.93	0.05	0.12	-0.45*
Mir	WH	359	555	765	15.2		72.3		31	19	4.07	0.23	0.10	-0.20
Iga	WH	290	866	283	14.5		43.9		30	14	4.35	0.00	0.10	-0.12
HiH	WH	456	248	130	10.8	3.43 ± 1.41	36.2	528.1	31	15	4.81	0.16	0.08	-0.31*
HiN	WH	486	1321	546	10.5	3.62 ± 1.20	37.8	521.4	30	24	4.91	0.25	0.08	-0.17*
Set	WH	635	111	152	19.5	3.16 ± 1.10	14.3	226.3	31	17	4.35	0.38	0.12	-0.33*
Sin	WH	495	671	150	7.2	3.44 ± 0.61	38.8	564.0	31	8	2.43	0.25	0.14	-0.67*

Mt.Ari	HM	1635	-	-	-	55.6	31	27	5.75	0.64	0.08	0.13
Mt.Hir	HM	1771	-	-	-	86.2	31	22	5.39	0.37	0.08	0.04
Mt.Shi	HM	1688	-	-	-	50.7	23	17	5.36	0.42	0.08	-0.03
Mt.Mur	HM	1876	-	-	-	-	31	31	6.64	1.01	0.08	0.21*
Mt.Muk	HM	1759	-	-	-	101.8	31	26	5.57	0.24	0.07	-0.07

433 ¹Habitat: WH (wind-hole); HM (high mountain). ²Elevation (m). ³Population area (m²). ⁴Perimeter of population (m). ⁵Distance from alpine
434 vegetation (km). ⁶Mean growth rate (\pm standard deviation; cm year⁻¹). ⁷Longest distance between ramets (m). We did not detect ramets with the
435 same multi-genotype on Mt. Mur. ⁸Estimated age (years). ⁹Number of individuals analyzed. ¹⁰Number of genets. ¹¹Allelic richness. ¹²Private
436 allelic richness.

437

438 Table 2. Analysis of molecular variance for the *Vaccinium vitis-idaea* samples.

Scale	d.f. ¹	SS ²	Est. var. ³	% variation	<i>P</i>
Between habitat types (wind-holes and high mountains)	1	26.5	0.04	1.4	0.001
Among populations	11	163.0	0.34	12.0	0.001
Within population	475	1129.5	2.48	86.6	0.001


439 ¹Degrees of freedom. ²Sum of squares. ³Estimated variance.

440

441

442

443




444

445 Fig. 1. Wind-hole and high mountain population sampling locations.

446

447




448

449 Fig. 2. Relationships between pairwise F_{ST} values and geographical distance. H-H, high

450 mountain pairs; W-W, wind-hole pairs; W-H, wind-hole–high mountain pairs. None of the

451 relationships were significant correlations based on the Mantel test.

452



453

454 Fig. 3. Genetic distances (a) and geographical distances (b) among habitat types and allelic

455 richness (c) and private allelic richness (d) of *Vaccinium vitis-idaea* populations at the


456 wind-hole and high mountain sites. W, wind-holes; H, high mountains. Different letters within

457 box plots indicate significant differences ($P < 0.05$) based on the Steel–Dwass test for Fig. 3a,

458 b and the t -test for Fig. 3c, d.

459

460




461

462 Fig. 4. Relationships between allelic richness and mean F_{ST} values of the wind-hole and high

463 mountain populations. Only the wind-hole populations were significantly correlated based on

464 Pearson's correlation analysis.

465



466

467 Fig. 5. Genetic clusters of *Vaccinium vitis-idaea* populations in the Engaru and Kitami regions.

468 (a) The initial analysis of all populations showed two genetic clusters ($K = 2$). (b) A

469 subsequent analysis of five wind-hole populations showed four genetic clusters ($K = 4$). (c) A

470 subsequent analysis using three wind-hole populations and five alpine populations showed

471 three genetic clusters ($K = 3$). Colors indicate different genetic clusters within each analysis.

472

473