Title	Analysis of the Relationship Between Enzymatic and Antiviral Activities of the Chicken Oligoadenylate Synthetase-Like
Author(s)	Tag-El-Din-Hassan, Hassan T.; Sasaki, Nobuya; Torigoe, Daisuke; Morimatsu, Masami; Agui, Takashi
Citation	Journal of interferon & cytokine research, 37(2), 71-80 https://doi.org/10.1089/jir.2016.0012
Issue Date	2017-02-01
Doc URL	http://hdl.handle.net/2115/67605
Rights	Final publication is available from Mary Ann Liebert, Inc., publishers http://dx.doi.org/10.1089/jir.2016.0012
Type	article (author version)
File Information	Hassan revised.pdf

[Instructions for use](#)

**Analysis of the Relationship Between Enzymatic and Antiviral
Activities of the Chicken Oligoadenylate Synthetase-Like**

Hassan T. Tag-El-Din-Hassan,^{1,2} Nobuya Sasaki,^{1,3} Daisuke Torigoe,^{1,4} Masami Morimatsu,¹ and
Takashi Agui¹

¹Laboratory of Laboratory Animal Science and Medicine, Department of Disease Control, Graduate
School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

²Poultry Production Department, Faculty of Agriculture, Mansoura University, Mansoura, Egypt.

Present address: ³Laboratory of Laboratory Animal Science and Medicine, Faculty of Veterinary
Medicine, Kitasato University; ⁴Division of Microbiology and Genetics, Institute of Resource
Development and Analysis, Kumamoto University

Corresponding author: T. Agui, Laboratory of Laboratory Animal Science and Medicine,
Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University,
Kita 18, Nishi 9, Kita-ku, Sapporo, Hokkaido 060-0818, Japan
Phone: - +81-11-706-5106. E-mail: agui@vetmed.hokudai.ac.jp

1 ABSTRACT

2

3 The oligoadenylate synthetase (OAS) is well known as an antiviral factor against the flavivirus
4 infection in mammals. It is known that the oligoadenylate synthetase-like (ChOAS-L) protein is
5 only present in the chicken genome. It has been shown in the previous report that the ChOAS-L
6 possesses enzymatic activity to convert ATP into 2'-5'-linked oligoadenylates and antiviral activity
7 against West Nile virus (WNV) replicon. Therefore, this study aimed to investigate the relationship
8 between enzymatic and antiviral activities of ChOAS-L. Eight mutated ChOAS-L proteins were
9 generated using either the site-directed mutagenesis or standard PCR protocol. The wild-typed and
10 mutated proteins were ectopically expressed in HEK293FT to analyze the enzymatic activity, and in
11 BHK-21 and BALB/3T3 cells to analyze the antiviral activity using WNV replicon. The results
12 revealed that all mutated proteins showed no enzymatic activity except for ChOAS-L-AΔUbL2.
13 However, all mutated proteins showed antiviral activity to inhibit the replication of the WNV
14 replicon except for the ChOAS-L-AΔUbL1/UbL2, which showed a partial inhibition compared to
15 the wild-type ChOAS-L-A or other mutated proteins. These results suggest that the ChOAS-L
16 expresses the antiviral activity in a manner independent of enzymatic activity. Our results propose
17 reconsideration of the mechanism of antiviral activity against the [flavivirus replication](#) of the
18 ChOAS-L.

19

1 **Introduction**

2

3 Type I interferon secreted from the virus-infected cells triggers and initiates a cascade of
4 expression of the interferon-stimulated genes, which interact with the infectious pathogens through
5 direct or indirect pathways. The well-recognized proteins as an interferon-stimulated factor are the
6 Mx, 2'-5'oligoadenylate synthetase (OAS), and protein kinase RNA-dependent (PKR) (Der and
7 others 1998; Sadler and Williams 2008).

8 The *OAS* gene families have been studied in human, mouse, rat, cattle, pig, dog, horse, and
9 many other living organisms. The four major types of *OAS* genes (*OAS1*, *OAS2*, *OAS3*, and *OASL*)
10 are conserved, whereas some of the other *OAS* genes have been lost or transformed to pseudo-genes
11 during mammalian evolution (Justesen and others 2000; Perelygin and others 2006; Kjaer and
12 others 2009).

13 Activation of the classical OAS/RNase L pathway results in RNA degradation of both viral
14 and cellular RNA including ribosomal RNA within infected cells. Subsequently it inhibits protein
15 translation in these cells, resulting in effective prevention of viral genome replication. The activity
16 of the OAS/RNase L pathway can be monitored by the degradation of ribosomal RNA in
17 virus-infected cells (Silverman 2007).

18 Recently several lines of evidence suggest that there is alternative RNase L-independent
19 pathway, by which some members of the OAS family can prevent viral replication. Human
20 OAS-L/a (HuOAS-L/a) (Rebouillat and others 1998; Hartmann and others 1998) and murine Oas1b
21 (mOas1b) (Elbahesh and others 2010) are devoid of the enzymatic activity and thus incapable of
22 activating RNase L. Nevertheless, both proteins display antiviral activity against West Nile virus
23 (WNV), suggesting that HuOAS-L/a and mOas1b utilize RNase L-independent pathway (Zhu and
24 others 2014; Marques and others 2008; Kristiansen and others 2011; Moritoh and others 2009;
25 Scherbik and others 2007; Kajaste-Rudnitski and others 2006; Perelygin and others 2002).

26 The OAS proteins belong to the nucleotidyle transferase superfamily involved in diverse

biological functions. The OAS proteins have the motif of the LxxxP, which seems to be important for the synthetize activity, followed by the P-loop, which is involved in the ATP binding activity, the D-D box, which plays an important role in the Mg^{2+} binding, region of 32-36 residues with high content of lysine and arginine named KR rich region (KR-RR), which is important for oligoadenylate binding and involved in the dsRNA binding (Yamamoto and others 2000), and the CFK domain, which is important for the oligomerization of the small OAS proteins (Ghosh and others 1997b). The OAS-L proteins have OAS unit in the N-terminal, which is fused to ubiquitin-like (UbL) domain containing two repetitive domains, UbL1 and UbL2, in the C-terminal (Eskildsen and others 2003).

Up-to-date, the only well characterized avian OAS gene is chicken OAS-like (ChOAS-L) protein, which consists of a single copy gene, locates on chromosome 12, and encodes two alternatively spliced alleles, ChOAS-L-A and ChOAS-L-B. ChOAS-L-B shows 100% identity with its allele ChOAS-L-A except for the partial deletion of 32 amino acids (AA) from the UbL1 (33 AA from 385Ala to 417Cys were substituted with 385Tyr). Interestingly, the ChOAS-L-B possesses the enzymatic activity as well as the ChOAS-L-A (Yamamoto and others 1998; Tatsumi and others 2000, 2003).

Recently, we reported that the ChOAS-L-A protein possesses antiviral activity against WNV replicon replication (Tag-El-Din-Hassan and others 2012). In addition, Cong and others (2013) reported that the ChOAS-L/RNase L pathway might be involved in anti-avian infectious bronchitis virus (IBV) infection. Moreover, Li and others (2007); Lee and others (2014) indicated that the ChOAS-L/RNase L pathway might be involved in anti-avian infectious bursal diseases virus (IBDV) infection. Therefore, the aim of this research is to examine the relationship between enzymatic and antiviral activities in the ChOAS-L against WNV replicon.

MATERIAL AND METHODS

1 ***SITE-DIRECTED MUTAGENESIS***

2 *ChOAS-L-A-FL*, which was *ChOAS-L-A* conjugated with the Flag-Tag DYKDDDDK sequence,
3 was cloned into the pGEM-T easy vector (Promega Corporation, Madison, USA) and used for the
4 generation of eight mutations. The Δ LxxxP domain, Δ P-LooP domain, Δ D-D box domain,
5 KR-RR-K193H (one AA substitution), Δ KR-RR, and ChOAS-L-B (the other allele) were generated
6 by using the QuickChange Lightning Site-Direct Mutagenesis Kit (Stratagene, La Jolla, CA, USA)
7 according to the manufacturer's protocol. The other two mutations, Δ Ubl2 domain and
8 Δ Ubl1/Ubl2 domains, were generated by the standard PCR protocol as follows, denature at 95 °C
9 for 5 min, twenty cycle of denature at 95 °C for 30 sec, annealing at 60 °C for 30 sec, extension at
10 72 °C for 1.5 min, and final extension at 72 °C for 1.5 min. All mutations were confirmed by the
11 DNA sequence analysis. Table 1 shows the primers used to generate these mutations. Figure 1
12 demonstrates the assumed scheme of the mutant proteins.

13 All mutated DNAs conjugated with the Flag-Tag were cut from the pGEM-T easy vector
14 using *NotI* restriction enzyme (Toyobo Co., Ltd. Osaka, Japan) and re-cloned into *NotI* restriction
15 site of the pIRES-EGFP vector (the mammalian expression vector, Clontech Bio. Inc., Shiga, Japan).
16 All cloning reactions were performed using 2x Ligation-Convenience Kit (Wako Nippon Gene Co.,
17 Ltd. Toyama, Japan) according to the manufacturer's protocol. All transformations were done using
18 DH5 α *E. coli* competent cells. The large-scaled plasmid purification was performed using
19 CsCL₂-ethidium bromide gradient purification method as described previously (Sambrook and
20 Russell 2001).

21

22 ***CELL CULTURE***

23

24 HEK293FT, BHK-21, and BALB/3T3 cells were maintained in Dulbecco's Modified Eagle
25 medium (DMEM) (Gibco/Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal bovine
26 serum (FBS) and 1% penicillin/streptomycin/glutamine (Gibco/Invitrogen, Carlsbad, CA, USA).

1 All cell lines were cultured with a passage of every two days in a 5% CO₂ cell culture incubator at
2 37 °C.

4 ***PREPARATION OF CELL LYSATE***

6 Both HEK293FT and BHK-21 cells were cultured in 100-mm dish at the density of 1.5×10^6
7 cells/dish for 24 h, the cells were transfected with 10 µg of pIRES-EGFP (empty vector as a
8 control) or gene-inserted vector, pChOAS-L-A-FL-EGFP, pΔLxxxP-FL-EGFP,
9 pΔP-LooP-FL-EGFP, pΔD-D-Box-FL-EGFP, pKR-RR-K193H-FL-EGFP, pΔKR-RR-FL-EGFP,
10 pChOAS-L-B-FL-EGFP, pΔUbl2-FL-EGFP, or pΔUbl1/Ubl2-FL-EGFP using Lipofectamine
11 2000® (Invitrogen, Carlsbad, CA, USA), according to the manufacturer's protocol. The EGFP
12 expression was analyzed at 24 h post-transfection for all transgenes using a NIKON ECLIPSE
13 TE2000E microscope. In respect of HEK293FT cells, at 48 h post-transfection, the cells washed
14 twice with phosphate-buffered saline (PBS), scratched in 1 ml PBS, spin for 5 min at 780 xg, the
15 supernatant was discarded, then, the cells were suspended in Flag-Tag lyses buffer as described
16 previously (Sarkar and others 2005). The HEK293FT cell lysates were used for the Western blotting
17 and enzymatic activity analysis. Transfected BHK-21 cells were lysed at 72 h post-transfection in
18 the Flag-Tag lyses buffer in the presence or absence of protease inhibitors using the same method
19 (Sarkar and others 2005). The BHK-21 cell lysates were kept at -80°C until used for protein
20 stability assay.

22 ***WESTERN BLOTTING***

24 The protein concentrations were determined using SmartSpec™ (BioRad) according to the
25 Bradford reagent protocol. Twelve µg proteins of each sample were separated in 10%
26 polyacrylamide-SDS gel electrophoresis on 100 V for 1 h, blotted onto PVDF membrane, and then,

1 blocked with PBS containing 3% non-fatty milk and 0.1% Tween-20 for 1 h at room temperature.
2 The membrane was incubated at room temperature with mouse-anti FLAG M2 antibody
3 (Sigma-Aldrich, St. Louis, MO, USA) diluted 1:2,000 with PBS containing 0.1% Tween-20
4 (PBS-Tween-20) for 2 h. The membrane was washed 3 times with PBS-Tween-20 for 10 min, and
5 then, the membrane was incubated at room temperature with HRP-conjugated goat anti-mouse
6 antibody diluted 1:5,000 with PBS-Tween-20 for 1 h. The membrane was washed 3 times with
7 PBS-Tween-20 for 10 min. The immunoreactive bands were detected by ECL™ Western Blotting
8 Detection Reagents (GE Healthcare UK Limited, Little Chalfont, UK) according to the
9 manufacturer's protocol.

11 ***ASSAY OF ENZYMATIC ACTIVITY***

13 The enzymatic activity of ChOAS-L-A protein and all of its mutants was measured as
14 described previously (Sarkar and others 2005), with a slight modification as follows; an aliquot of
15 2.5 µl of HEK293FT whole cell lysates was incubated in incubation buffer (20 mM Tris-HCl at pH
16 7.4, 20 mM magnesium acetate, 2.5 mM dithiothreitol (DTT), 5 mM ATP, 50 µg/ml poly(I):(C), and
17 5 µCi of [α -³²P] ATP (6,000 Ci/mmol) in a final volume of 10 µl for 18 h at 30 °C. The reaction was
18 stopped by boiling for 5 min at 95 °C, followed by adding loading buffer containing 25%
19 formamide, 0.5% bromophenol blue, and 0.5% xylene cyanol.

20 The synthesized 2'-5' (A) products were run in a 20% polyacrylamide-urea (7 M) gel. The
21 electrophoresis was performed for 4 h on 200 V, and then products were visualized by
22 autoradiography with a BAS2500 Bio-Imaging analyzer (FUJIFILM, Tokyo, Japan).

24 ***PROTEIN STABILITY ASSAY***

26 The protein stability assay was performed by incubating the protein samples (12 µg) in the

presence or absence of protease inhibitors at 37°C for 2 h. The incubated samples were subjected to Western blotting as shown above to determine the protein stability.

MEASUREMENT OF INHIBITORY ACTIVITY ON WNV REPLICON REPLICATION

pUC19repWNV/SEAP plasmid containing the WNV genome harboring the secreted alkaline phosphatase reporter gene instead of viral structural genes (Moritoh and others 2011) was propagated in transformed XL10-Gold[®] Ultracompetent *E.coli* Cells (Stratagene, La Jolla, CA, USA) according to the manufacturer's protocol. A PureLink[®] HiPure Plasmid Maxiprep Kit (Invitrogen, Carlsbad, CA, USA) was used for large-scaled purification according to the manufacturer's protocol.

pUC19repWNV/SEAP plasmid was linearized using *FastDigest XbaI* restriction enzyme (Thermo Fisher Scientific, Tokyo, Japan) according to the manufacturer's protocol. The mMMESSAGE mMACHINE[®] SP6 Transcription Kit (Ambion, Austin, TX, USA) was used to synthesize WNV-replicon RNA from the linearized pUC19repWNV/SEAP according to the manufacturer's protocol. Briefly, 1µg of linearized plasmid was incubated in the reaction mixture for 2 h to get the maximum RNA yield as recommended by the manufacture, then the synthesized RNA was recovered using the PureLink[®] RNA Mini Kit (Ambion, Carlsbad, Ca, USA), divided into 10 µg aliquots, and stored at -80°C until used for the measurement of antiviral activity.

One day before the transfection with the wild type *ChOAS-L-A* and mutant genes, the BHK-21 and BALB/3T3 cells were cultured in a 100-mm plate at the density of 1.5×10^6 cells/plate, and then, 10µg of pIRES-EGFP (empty vector as a control), pChOAS-L-A-FL-EGFP, pΔLxxxP-FL-EGFP, pΔP-LooP-FL-EGFP, pΔD-D-Box-FL-EGFP, pKR-RR-K193H-FL-EGFP, pΔKR-RR-FL-EGFP, pChOAS-B-FL-EGFP, pΔUbl2-FL-EGFP, or pΔUbl1/Ubl2-FL-EGFP was transfected using Lipofectamine 2000[®] (Invitrogen, Carlsbad, CA, USA), according to the manufacturer's protocol. The EGFP expression was analyzed at 24 h post-transfection using the NIKON ECLIPSE TE2000E microscope.

1 BHK-21 and BALB/3T3 cells were harvested from non-transfected cells (mock transfection)
2 and from all transfectants. Then, 0.5×10^7 cells were incubated in 500 μ l of PBS on ice with 10 μ g of
3 WNV-replicon RNA in 4 mm cuvette (Nepa Gene Co., Ltd. Chiba, Japan). The electroporation was
4 performed using Gene-Pulser X Cell, Bio-Rad, with pulses of 1.5 kV at 25 μ F and 0.5 kV at 25 μ F
5 of infinite resistance for BHK-21 and BALB/3T3 cells, respectively. The live cells were counted
6 and cultured in triplicate in a 6-well plate (3×10^5 cells/well). Supernatants were collected at 24 and
7 72 h post-electroporation. The inhibitory activity on WNV replicon replication was measured as the
8 secreted alkaline phosphatase (SEAP) activity in the supernatant using Great EscAPe™ SEAP
9 Chemiluminescence Kit 2.0 (Clontech Bio. Inc., Shiga, Japan) and Mithras LB 940 Multimode
10 Microplate Reader according to the manufacturer's protocol.

11

12 ***RNA EXTRACTION AND RT-PCR***

13

14 Total RNA was extracted from HEK293FT cells at 48 h post-transfection and from BHK-21
15 and BALB/3T3 cells before electroporation and at 72 h post-electroporation, using TRIZOL reagent
16 (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol. RNA samples were
17 treated with DNase to remove any DNA contamination using TURBO DNA-free™ Kit (Ambion,
18 Austin, TX, USA) according to manufacturer's protocol. Concentration of purified total RNA
19 samples were determined using SmartSpec™ (BioRad), 1 μ g of total RNA was used to generate
20 cDNAs using *ReverTra Ace*® (Toyobo Co., Ltd. Osaka, Japan) according to manufacturer's protocol.
21 The RT condition was at 42° C for 60 min and at 99 °C for 5 min. The synthesized cDNA was
22 stored at -20°C until used for the PCR, of which conditions were as follow; denature at 95 °C for 5
23 min; 35 cycles of denature at 95 °C for 30 sec, annealing at 60 °C for 30 sec, and extension at 72 °C
24 for 1.5 min; followed by the final extension at 72 °C for 3 min.

25

26 ***LOCALIZATION OF THE CHOAS-L-A AND MUTATED PROTEINS***

1

2 BHK-21 cells were cultured in 96 wells plate (2×10^4 cells/well) for 24, and cells were
3 transfected with 0.1 μ g of *pIRES-EGFP* or *pChOAS-L-A-FL-EGFP* as well as all mutated genes
4 using 0.4 μ l Lipofectamine 2000[®] according to manufacturer's protocol. At 48 h post-transfection,
5 cells were fixed with 4% paraformaldehyde in PBS for 15 min at room temperature and washed
6 twice with 1x washing buffer (PBS containing 0.1% Tween-20). The cells were then permeabilized
7 with 0.1% Triton X-100 in PBS for 5 min at room temperature, washed twice with 1x washing
8 buffer, and followed by blocking in 10% horse serum in PBS for 30 min at room temperature. The
9 mouse-anti FLAG M2 antibody (Sigma-Aldrich, St. Louis, MO, USA) diluted to 1:1000 with the
10 blocking solution was added to wells, incubated for 1 h at room temperature, and followed by
11 washing for three times (5 min each) with 1x washing buffer. The diluted secondary antibodies
12 (Alexa Fluor 568 goat anti-mouse IgG, Invitrogen, Oregon, USA) in 1x PBS were added to wells
13 and incubated for 1 h at room temperature, followed by washing for three times (5 min each) with
14 1x washing buffer. Nuclei counterstaining was performed by incubating cells with DAPI in PBS for
15 5 min at room temperature, followed by washing cells two times (5 min each) with 1x washing
16 buffer, and then the fluorescence images were captured using BZ-X700 All-in-One Florescence
17 Microscope, KEYENCE (Osaka, Japan).

18

19 ***RNASE-L ACTIVATION ASSAY***

20

21 The RNase-L activation assay for ChOAS-L-A protein and all of its mutated proteins was
22 performed using the florescence resonance energy transfer assay (FRET assay) as described
23 previously (Thakur and others 2005) with a slight modification as follows; an aliquot of 2 μ g of
24 BHK-21 whole cell lysates transfected with *pIRES-EGFP* as well as *pChOAS-L-A-FL-EGFP* and
25 all mutated genes was incubated in 50 μ l of 1x incubation buffer (25 mM Tris-HCl at pH 7.4, 0.1 M
26 KCl, 10 mM MgCl₂, 50 μ M ATP, 7 mM β -mercaptoethanol, and 180 nM FRET probe for 0, 5, 10,

1 15, 30, 45, 60, 90, and 120 min at 21°C. As a negative control, lysates of BHK-21 and HEK293FT
2 non-transfected cells were used, whereas as a positive control, BHK-21 and HEK293FT
3 non-transfected cells stimulated with 3,000 IU/ml of IFN- α 2b (INTRON[®] A, MSD K.K., Tokyo,
4 Japan) for 24 h were used. The fluorescence was determined using the Infinite M200 PRO plate
5 reader, Tecan (Mannedorf, Switzerland).

7 **STATISTICAL ANALYSIS**

8
9 All experiments were done three times independently. Collected data were subjected to the
10 one-way analysis of variance (ANOVA) using a SPSS 16.0 software. All values were represented as
11 means $\pm$ SD. Statistical differences were analyzed using Tukey HSD test. Values of $p < 0.05$ and p
12 < 0.01 were considered to be significant.

14 **RESULTS**

16 **ENZYMATIC ACTIVITY OF THE MUTATED CHOAS-L-A**

17
18 Protein and total RNA were extracted at 48 post-transfection from HEK293FT cells
19 transfected with the *pIRES-EGFP* (empty vector as a control), wild-type *ChOAS-L-A* and each
20 mutated *ChOAS-L-A* genes. The expression of all genes were confirmed by RT-PCR and Western
21 blotting.

22 All mutated genes as well as *EGFP* and *Actb* were expressed at 48 h post-transfection (Fig.
23 2A). As shown in Fig. 2B, the cells transfected with pChOAS-L-A-FL-EGFP,
24 pChOAS-L-B-FL-EGFP, and p Δ UbL2-FL-EGFP showed enzymatic activity to convert ATP to 2'-5'
25 (A) with mono-, di-, tri-, and tetramer, whereas the cells transfected with pIRES-EGFP (the empty
26 vector), p Δ LxxxP-FL-EGFP, p Δ P-LooP-FL-EGFP, p Δ D-D-Box-FL-EGFP,

pKR-RR-K193H-FL-EGFP, pΔKR-RR-FL-EGFP, and pΔUbL1/UbL2-FL-EGFP were incapable of synthesizing 2'-5' (A).

ANTIVIRAL ACTIVITY OF THE MUTATED CHOAS-L-A

We did not select chicken-derived cells as host cells for measurement of inhibitory activity on WNV replicon replication to exclude the effect of endogenous ChOAS-L. BHK-21 cells were selected, because this cell line is known to harbor many viruses. BALB/3T3 cells were also selected as cells devoid of mouse Oas1b, which is a flavivirus-specific antiviral factor. Total RNA was extracted from BHK-21 and BALB/3T3 control cells (Mock and pIRES-EGFP transfection), and cells transfected with wild-type *pChOAS-A-FL-EGFP* and mutant genes before the measurement of antiviral activity. The RT-PCR results showed that *ChOAS-L-A* and its mutated genes were all successfully expressed (Figs. 3A and 4A).

The SEAP expression level was measured in the culture supernatant of BHK-21 cells transfected all genes at 24 and 72 h post-electroporation of the WNV-replicon RNA. All mutants significantly ($p<0.01$) showed antiviral activity to reduce the SEAP activity compared to that of the controls, both ChOAS-L-A and ChOAS-L-B showed the highest antiviral activity followed by other mutants. The KR-RR-K193H was significantly ($p<0.05$) different from ChOAS-L-A, ChOAS-L-B or ΔLxxxP (Fig. 3B). Although ΔUbL1/UbL2 showed antiviral activity, its extent was partial when compared to that of ChOAS-L-A, suggesting that UbL domain is necessary for the antiviral activity. However, the result that ChOAS-L-B, partial deletion of UbL1, and ΔUbL2 showed almost full antiviral activity suggest that a single copy of UbL domain is satisfactory to retain the antiviral activity.

The same trend was seen in the BALB/3T3 cells, all mutants significantly ($p<0.01$) showed antiviral activity to reduce the SEAP activity compared with control groups. Moreover, the ΔKR-RR was significantly different from ChOAS-L-A ($p<0.01$), ChOAS-L-B ($p<0.05$) and

1 $\Delta L_{xxx}P$ ($p < 0.05$) (Fig. 4B).

2 Although the WNV-replicon replication level is much lower than those in the BHK-21 cells.
3 The reason for this difference is unknown; however, the BHK-21 cells were reported to have
4 characteristics to allow the efficient replication of WNV (Shi and others 2002; Rossi and others
5 2005; Moritoh and others 2011). Although BALB/3T3 cells might possess resistance to the WNV
6 infection compared to the BHK-21 cells, antiviral activities of ChOAS-L-A and its mutants in
7 BALB/3T3 cells were almost the same as that in BHK-21 cells, suggesting that the results were not
8 dependent on the cell types.

9

10 ***STABILITY OF CHOAS-L-A AND MUTATED PROTEINS***

11

12 Twelve- μ g whole cells lysate of BHK-21 cell transfected with *pChOAS-L-A-FL-EGFP* or
13 other mutated transgenes were incubated on 37°C for 0 and 2 h in the presence or absence of
14 proteases inhibitors in the lysis buffer. Then, the incubated protein samples were used for the
15 western blot analysis. The obtained results revealed that in the presence of the proteases inhibitors in
16 the lysis buffer all protein samples were remained protected. However, in the absence of the
17 proteases inhibitor in the lysis buffer, ChOAS-L-A, $\Delta L_{xxx}P$, ΔP -Loop, ΔD -D, KR-RR-K193H as
18 well as Δ KR-RR proteins showed stability up to 2 h of incubation on 37°C (Fig. 5). On the other
19 hand, ChOAS-L-B (partial deletion in the UbL1), Δ UbL2 and Δ UbL1/UbL2 proteins showed
20 instability after incubation period. This result indicates the importance of the UbL domains for the
21 Ch-OAS-L protein stability.

22

23 ***RNASE-LACTIVITY ASSAY***

24 Two- μ g protein of whole cell lysate of BHK-21 cells transfected with pIRES-EGFP,
25 pChOAS-L-A-FL-EGFP, and other mutated genes, as well as control groups (HEK293FT and
26 BHK-21 non-transfected cells with- and without IFN treatment as positive and negative control,

respectively), were used to perform FRET assay. RNase L activation detection were performed on 0, 5, 10, 15, 30, 45, 60, 90 and 120 minutes post-incubation in the cleavage buffer.

The obtained results showed that, in all time point analysis the HEK293FT/IFN expresses high significant ($p<0.01$) fluorescence level than all protein samples, which indicates that there are high activity of RNase L in HEK293FT/IFN cells (Fig. 6). On the other hand, the ChOAS-L-A protein as well as its mutated proteins showed the same expression as the negative control groups (non-transfected BHK-21 cells). Regarding BHK-21/IFN treated cells, it was statistically insignificant for the 0, 5, 10, 15, and 30 minutes' post incubation. However, starting from 45 minutes' post-incubation, both BHK-21/IFN and HEK293FT/non-transfected showed similar significant ($p<0.05$) increase in the fluorescence activity (Fig. 6). This increasing in the fluorescence activity continued highly significant ($p<0.01$) on 60, 90 and 120 minutes' post-incubation. In contrast, pIRES-EGFP, ChOAS-L-A as well as all mutated proteins expressed almost the same fluorescence activity as for the negative control (BHK-21/non-transfected) with insignificant differences (Fig. 6). These results showed that the ChOAS-L-A protein and all of its mutated protein are incapable of activate the RNase L in BHK-21 cells.

DISCUSSION

ChOAS-L-A protein, a member of the OAS-L protein family characterized by the N-terminal OAS unit (similar to small OAS protein) comprises the motif of the LxxxP followed by the P-Loop, D-D box, region of 32-36 residues with high content of lysine and arginine, and finally the CFK motif. This small OAS unit fused into the C-terminal UbL unit, composed of two repetitive UbL domains, UbL1 and UbL2 (Eskildsen and others 2003). Recently, Tag-El-Din-Hassan and others (2012) reported that the ChOAS-L-A possesses both enzymatic and anti-WNV activities.

Therefore, in the current study, eight *ChOAS-L-A* mutants were generated by using site-directed mutagenesis protocol or ordinary PCR protocol, in which five mutations were introduced in the

domains of the OAS unit and the other three mutations were in the UbL domains, to investigate the relationship between enzymatic and antiviral activities of ChOAS-L-A.

The results revealed that the Δ LxxxP domain mutation resulted in a complete loss of the enzymatic activity (Fig. 2-B). Similarly, Ghosh and others (1997a) found that the deletion of 1 to 9 or 1 to 23 residues from the N-terminal of mouse 9-2 OAS protein resulted in a 100% loss of enzymatic activity. Moreover, the substitution of three conserved amino acids in this motif, L3H, P7Q, A8D, and P7Q/A8D still retain the ability to bind the dsRNA as well as the wild type; however, the enzymatic activity of L3H and A8D showed 30 and 18% of the wild type, respectively, and both P7Q and P7Q/A8D completely lose the enzymatic activity. Further, to confirm the importance of the proline at position 7, Ghosh and others (1997a) generated several substituted proteins, P7Q, P7F, P7Y, P7E, P7K, P7L, and P7T and examined enzymatic activity. All mutations resulted in complete loss of the enzymatic activity compared with the wild type. However, Bandyopadhyay and others (1998) reported that the P7Q mutation showed full enzymatic activity, when the protein expressed in baculovirus/insect cell system. The current study showed that the Δ LxxxP domain mutation did not affect the inhibitory effects on WNV-replicon replication. The SEAP activity in BHK-21 and BALB/3T3 cell expressing Δ LxxxP domain mutant protein was significantly decreased as well as the wild type (ChOAS-L-A) in comparison with the control groups (Fig. 3-B and 4-B). These findings suggest that the LxxxP domain is critically required for the synthetase activity, whereas it is not required for the antiviral activity of the ChOAS-L protein.

The Δ P-LooP domain mutation shows no enzymatic activity (Fig. 2-B). Similarly, Yamamoto and others (2000) examined several mutations in the P-Loop domain of the 42 KDa MuOAS protein, G62A/G63A, K67M, and K67R, all of which mutations resulted in a significant decrease in the enzymatic activity, 0, 2, and 30% of the wild type, respectively. In addition, the K59E mutation in porcine OAS1 resulted in a loss of enzymatic activity (Hartmann and others 2003). With respect to the antiviral activity, the results in the current study shows that the Δ P-LooP mutant protein in both BHK-21 and BALB/3T3 cells resulted in significant inhibition in the WNV-replicon replication

(Figs. 3-B and 4-B). These results show that the P-Loop domain is critically essential for the enzymatic activity, whereas it is not for antiviral properties of ChOAS-L-A protein.

The Δ D-D box domain mutation resulted in a complete loss of enzymatic activity (Fig. 2-B). This result agrees with Yamamoto and others (2000), in which the substitution in D-D box, D76N/D78N, resulted in a 100% loss of enzymatic activity compared with the wild type protein. The same result was obtained by Sarkar and others (1999), in which the enzymatic activity of human p69 OAS was lost, when the D481A and D408A/D410A mutations in D-D box were introduced. In contrast to the enzymatic activity, the Δ D-D box mutation did not affect the antiviral ability of the ChOAS-L-A protein in this report (Figs. 3-B and 4-B). This finding confirms that the D-D box domain is critically important for the enzymatic activity, whereas it is not important for the antiviral activity of ChOAS-L-A protein.

The KR-RR domain mutations in the current study, the mutations either induced by amino acid substitution or complete domain deletion, resulted in a 100% loss of enzymatic activity (Fig. 2-B). These results agreed to the results by Yamamoto and others (2000), in which the K200M and K200R mutations of the 42 KDa MuOAS showed a significant reduction in enzymatic activity to 2 and 50% that of the wild type, respectively. Moreover, the double mutations in both P-loop and KR-RR, K67R/K200R, resulted in a decrease in the enzymatic activity to 10% of the wild type. Similarly, Kon and Suhadolnik (1996) investigated the effects of K199R and K199H mutations of the human 40 KDa OAS. They reported that both mutations resulted in a loss of enzymatic activity; however, both mutants retained the ability to bind poly (I):(C). Similarly, Hartmann and others (2003) induced several mutations in porcine OAS1 protein, R194E, R198E, R198M, R194E/R198E, and K203E, and showed that all of these mutant proteins lost enzymatic activity. Furthermore, Sarkar and others (2002) examined several mutations, Y421P, Y421A, R544A, R544Y, and K547A of the p69 HuOAS. All mutations resulted in a severe reduction in the enzymatic activity to 3% that of the wild type except for K547A, 10% of the control. Furthermore, Ghosh and others (1991) induced several deleting mutations in murine 9-2 Oas protein, one of which deleted the KR-RR.

1 They found that the Δ KR-RR mutated protein retains the dsDNA binding ability almost same as the
2 wild type protein. With respect to the antiviral activity, however, we showed that both mutants
3 remained antiviral activity both in BHK-21 and BALB/3T3 cells as well as the wild-typed
4 ChOAS-L-A protein (Figs. 3-B and 4-B). These results suggest that the KR-RR is critical for the
5 enzymatic activity of the ChOAS-L protein, whereas it is not important for the antiviral activity.

6 Moreover, the Δ LxxxP, Δ P-Loop, Δ D-D Box, KR-RR-K193H or Δ KR-RR mutated proteins was
7 unable to activate the RNase L, using FRET assay, during all studied time points and expresses the
8 same level as the negative control groups (Fig. 6). And all of these mutated protein showed stability
9 up to 2 h post incubation in the absence of the proteases inhibitors and remain detectable as 0.0 h
10 incubation (Fig. 5). As well, Δ LxxxP, Δ P-Loop, Δ D-D Box, KR-RR-K193H or Δ KR-RR mutated
11 proteins was localized in to cytoplasm similar to the wild type ChOAS-L-A protein (Fig. 7). Taken
12 together, these results show that all mutations carried out in the active domains of the ChOAS unit
13 results in deactivation of the enzymatic activity (Fig. 2-B). However, all of these mutated protein
14 still retain the antiviral activity to inhibit the WNV-replicon replication (Fig. 3-B and 4-B), and acts
15 almost same as ChOAS-L-A protein (Fig. 5, 6 and 7), suggesting that the antiviral activity of
16 ChOAS-L is independent of the enzymatic activity.

17 Both ChOAS-L- Δ Ubl2 and ChOAS-L-B, the second allele of ChOAS-L-A possessing 32 AA
18 deletion in Ubl1 generated by the mutagenesis protocol, still possessed the enzymatic activity. This
19 result is consistent with the report by Tatsumi and others (2003), in which the ChOAS-L-B
20 possesses the enzymatic activity. However, the Δ Ubl1/Ubl2 mutation, complete deletion of Ubl
21 domains, did not show the enzymatic activity (Fig. 2-B), which was unexpected result for us. We
22 had supposed that the ChOASL-L-A- Δ Ubl1/Ubl2 should retain the enzymatic activity, because
23 the OAS unit was retained in this mutant. The full function of Ubl domain is not completely
24 uncovered at present. However, in the current study we showed that Ubl domain plays an important
25 role in stabilization of ChOAS-L proteins, where, either ChOAS-L-B (possess partial deletion in
26 Ubl1), ChOAS-L-A- Δ Ubl2 or ChOAS-L-A- Δ Ubl1/Ubl2 proteins, showed instability when

1 incubated in the lysis buffer in the absence of the proteases inhibitors (Fig. 5), similarly Tatsumi and
2 others (2003) found that ChOAS-L-B showed instability if incubated in the absence of the proteases
3 inhibitors, as well, ChOAS-L-B shows age-dependent expression, where its expression in chicken
4 erythrocytes was decreased by aging, if compared with the ChOAS-L-A protein. Although, only one
5 UbL domain is enough for ChOAS-L-A protein to express the enzymatic activity (Fig 2-B).

6 Furthermore, ChOAS-L-B, ChOAS-L-A- Δ UbL2 or ChOAS-L-A- Δ UbL1/UbL2 proteins,
7 showed incapability to activate RNase L in FRET assay, and expresses the same level as the
8 negative control (Fig. 6), and localized into cytoplasm as well as the wild type ChOAS-L-A protein
9 (Fig. 7).

10 As for the antiviral activity, both ChOAS-L-B and ChOAS-L-A- Δ UbL2 mutant proteins still
11 retain and express the antiviral properties to inhibit the WNV-Replicon replication (Fig. 3-B and
12 4-B). However, the ChOAS-L-A- Δ UbL1/UbL2, which supposed to be completely lose the antiviral
13 activity, surprisingly found to be expressing partial antiviral activity against the WNV-replicon
14 replication (Fig. 3-B and 4-B). The possible reason for this results may be as follows, as it was
15 mentioned before, ChOAS-L-A- Δ UbL1/UbL2 still possess the OAS unit, so that, it showed slightly
16 inhibition of the WNV-replicon replication due to the capability of the OAS unit to bind the dsRNA
17 (step of the viral genome replication) inside the cells. Recently, Ibsen and others (2015) showed that
18 the HuOAS-L- Δ UbL_(S) have the ability to bind dsRNA.

19 Taken together, the obtained results indicating that, however the UbL domains are not
20 participated in the enzymatic activity functions of ChOAS-L protein, it is still required at least one
21 UbL domain to express the enzymatic activity. Moreover, at least one UbL domain is critically
22 essential to express the antiviral activity of ChOAS-L.

23 Collectively, in the current study, the FRET Assay stands as the connective edge between the
24 enzymatic and antiviral activities of the OAS proteins and explains why the BHK-21 cell line used
25 in this experiment. Where, the lysates containing the ChOAS-L-A or its mutant proteins showed
26 inability to activate the RNase L same as the negative control group (BHK-21/Shame), even if the

1 protein enzymatically active (such as ChOAS-L-A, ChOAS-L-B and ChOAS-L-A- Δ Ubl2). The
2 incapability to activate RNase L in this experiment is not related to have or haven't the enzymatic
3 activity, however it is related to the cell line itself where, BHK-21 cell express deficiency in the IFN
4 pathway. Which reflects that the antiviral activity of ChOAS-L-A and all of its mutants except for
5 ChOAS-L-A- Δ Ubl1/Ubl2, is for sure due to the OAS/RNase L independent pathway.

6 In conclusion, the obtained results suggest that the ChOAS-L expresses the antiviral activity in a
7 manner independent of the OAS/RNase L classical pathway. However, it may be premature to state
8 ChOAS-L functions without investigating other avian OAS family members for enzymatic and
9 antiviral activities not only against WNV but against specific avian pathogens.

10

11

1 Table 1. Primers used to generate mutations from the wild-typed *ChOAS-L-A* gene and primers for
2 *EGFP* and *Actb* genes.

3

Name		Direction	Primer sequence	No of AA del.
Mutation	ChOAS-L-A-FL Δ LxxxP	Forward	5'-GATTATCGATATGGGGTCGCGCCAGCTGGAGG-3'	5
		Reverse	5'-CCTCCAGCTGGCGCGACCCCATATCGATAATC-3'	
	ChOAS-L-A-FL Δ P-LoopP	Forward	5'-CGTCCTCAAGACCGTCACGGCGCTGCGCAAC-3'	8
		Reverse	5'-GTTGCGCAGCGCCGTGACGGTCTTGAGGACG-3'	
	ChOAS-L-A-FL Δ D-D box	Forward	5'-CGCTGCGCAACAACCTCGGTGGTGCTCTTCATCAACTGC-3'	3
		Reverse	5'-GCAGTTGATGAAGAGCACCACCGAGTTGTTGCGCAGCG-3'	
	ChOAS-L-A-FL-KR RR-K193H	Forward	5'-CAGAAGAACTTTGTGCACTGGCGCCCTGCTAAG-3'	0
		Reverse	5'-CTTAGCAGGGCGCCAGTGCACAAAGTTCTTCTG-3'	
	ChOAS-L-A-FL Δ KR RR	Forward	5'-CCCCAGCTTCACAGAACTGCCCCCAAGTATG-3'	36
		Reverse	5'-CATACTGGGGGGCAGTTCTGTGAAGCTGGGG-3'	
	ChOAS-L-B-FL	Forward	5'-GCTACCAGCAGCGCCTGTACACCGAGCCCCAG-3'	32
		Reverse	5'-CTGGGGCTCGGTGTACAGGCGCTGCTGGTAGC-3'	
	ChOAS-L-A-FL Δ Ubl2	Reverse	5'-TCAGATCTTATCGTCGTCATCCTTGTAAATCCAGCAGCAGCAGCA C-3'	92
	ChOAS-L-A-FL Δ Ubl1/Ubl2	Reverse	5'-TCAGATCTTATCGTCGTCATCCTTGTAAATCCTGCACGTCC-3'	171
ChOAS-L-A-FL		Forward	5'-ATCGATATGGGGTTGGAGAGCGTGAG-3'	
		Reverse	5'-TCAGATCTTATCGTCGTCATCCTTGTAAATCGGAGGGCACGCAGCGTCT GG-3'	
EGFP		Forward	5'-ATGGTGAGCAAGGGCGAGGA-3'	
		Reverse	5'-TACTTGTACAGCTCGTCCA-3'	
ActB		Forward	5'-TGATGGTGGGAATGGGTCAG-3'	
		Reverse	5'-GAAGGCTGGAAAAGAGCCTC-3'	

1 REFERENCES

- 2 Bandyopadhyay S, Ghosh A, Sarkar SN, Sen GC. 1998. Production and purification of recombinant
3 2'-5' oligoadenylate synthetase and its mutants using the baculovirus system.
4 Biochemistry 37(11):3824-3830.
- 5 Cong F, Liu X, Han Z, Shao Y, Kong X, Liu S. 2013. Transcriptome analysis of chicken kidney
6 tissues following coronavirus avian infectious bronchitis virus infection. BMC Genomics
7 14:743-755.
- 8 Der SD, Zhou A, Williams BR, Silverman RH. 1998. Identification of genes differentially regulated
9 by interferon α , β , or γ using oligonucleotide arrays. Proc Natl Acad Sci USA
10 95(26):15623-81562.
- 11 Elbahesh H, Jha BK, Silverman RH, Scherbik SV, Brinton MA. 2010. The *Flv^r*-encoded murine
12 oligoadenylate synthetase 1b (Oas1b) suppresses 2-5A synthesis in intact cells. Virology
13 409(2):262–270.
- 14 Eskildsen S, Justesen J, Schierup MH, Hartmann R. 2003. Characterization of the
15 2'-5'-oligoadenylate synthetase ubiquitin-like family. Nucleic Acids Res
16 31(12):3166-3173.
- 17 Ghosh A, Desai SY, Sarkar SN, Ramaraj P, Ghosh SK, Bandyopadhyay S, Sen GC. 1997a. Effects
18 of mutating specific residues present near the amino terminus of 2'-5'-oligoadenylate
19 synthetase. J Biol Chem 272(24):15452-15458.
- 20 Ghosh A, Sarkar SN, Guo W, Bandyopadhyay S, Sen GC. 1997b. Enzymatic activity of
21 2'-5'-oligoadenylate synthetase is impaired by specific mutations that affect
22 oligomerization of the protein. J Biol Chem 272(52):33220-33226.
- 23 Ghosh, S. K., Kusari, J., Bandyopadhyay, S. K., Samanta, H., Kumar, R., and Sen, G. C., 1991.
24 Cloning, sequencing, and expression of two murine 2'-5'-oligoadenylate synthetases. J
25 Biol Chem. 266(23):15293-15299.
- 26 Hartmann R, Justesen J, Sarkar SN, Sen GC, Yee VC. 2003. Crystal structure of the 2'-specific and

double-stranded RNA-activated interferon-induced antiviral protein 2'-5'-oligoadenylate synthetase. *Mol Cell* 12(5):1173-1185.

Hartmann R, Olsen HS, Widder S, Jorgensen R, Justesen J. 1998. p59OASL, a 2'-5' oligoadenylate synthetase like protein: a novel human gene related to the 2'-5' oligoadenylate synthetase family. *Nucleic Acids Res* 26(18):4121-4128.

Ibsen MS, Gad HH, Andersen LL, Hornung V, Julkunen I, Sarkar SN, Hartmann R. 2015. Structural and functional analysis reveals that human OASL binds dsRNA to enhance RIG-I signaling. *Nucleic Acids Res.* 43(10):5236-48.

Justesen J, Hartmann R, Kjeldgaard NO. 2000. Gene structure and function of the 2'-5'-oligoadenylate synthetase family. *Cell Mol Life Sci* 57(11):1593-1612.

Kajaste-Rudnitski A, Mashimo T, Frenkiel MP, Guénet JL, Lucas M, Desprès P. 2006. The 2'-5'-oligoadenylate synthetase 1b is a potent inhibitor of West Nile virus replication inside infected cells. *J Biol Chem* 281(8):4624-4637.

Kjaer KH, Poulsen JB, Reintamm T, Saby E, Martensen PM, Kelve M, Justesen J. 2009. Evolution of the 2'-5' oligoadenylate synthetase family in eukaryotes and bacteria. *J Mol Evol* 69(6):612-624.

Kon N, Suhadolnik RJ. 1996. Identification of the ATP binding domain of recombinant human 40-kDa 2'-5' oligoadenylate synthetase by photoaffinity labeling with 8-azido-[alpha-³²P] ATP. *J Biol Chem* 271(33):19983-19990.

Kristiansen H, Gad HH, Eskildsen-Larsen S, Despres P, Hartmann R. 2011. The oligoadenylate synthetase family: an ancient protein family with multiple antiviral activities. *J Interferon Cytokine Res* 31(1):41-47.

Lee CC, Wu CC, Lin TL. 2014. Chicken melanoma differentiation-associated gene 5 (MDA5) recognizes infectious bursal disease virus infection and triggers MDA5-related innate immunity. *Arch Virol* 159(7):1671-1686.

Li YP, Handberg KJ, Juul-Madsen HR, Zhang MF, Jørgensen PH. 2007. Transcriptional profiles of

chicken embryo cell cultures following infection with infectious bursal disease virus.
Arch Virol 152(3):463-478.

Marques J, Anwar J, Eskildsen-Larsen S, Rebouillat D, Paludan SR, Sen G, Williams BR, Hartmann R. 2008. The p59 oligoadenylate synthetase-like protein possesses antiviral activity that requires the C-terminal ubiquitin-like domain. J Gen Virol 89(11):2767-2772.

Moritoh K, Maeda A, Nishino T, Sasaki N, Agui T. 2011. Development and application of West Nile virus subgenomic replicon RNA expressing secreted alkaline phosphatase. J Vet Med Sci 73(5): 683-686.

Moritoh K, Yamauchi H, Asano A, Yoshii K, Kariwa H, Takashima I, Isoda N, Sakoda Y, Kida H, Sasaki N, Agui T. 2009. Generation of congenic mouse strains by introducing the virus-resistant genes, Mx1 and Oas1b, of feral mouse-derived inbred strain MSM/Ms into the common strain C57BL/6J. Jpn J Vet Res 57(2):89-99.

Perelygin AA, Scherbik SV, Zhulin IB, Stockman BM, Li Y, Brinton MA. 2002. Positional cloning of the murine Flavivirus resistance gene. Proc Natl Acad Sci USA 99(14):9322-9327.

Perelygin AA, Zharkikh AA, Scherbik SV, Brinton MA. 2006. The mammalian 2'-5' oligoadenylate synthetase gene family: evidence for concerted evolution of paralogous Oas1 genes in Rodentia and Artiodactyla. J Mol Evol 63(4):562-576.

Rebouillat D, Marié I, Hovanessian AG. 1998. Molecular cloning and characterization of two related and interferon-induced 56-kDa and 30-kDa proteins highly similar to 2'-5' oligoadenylate synthetase. Eur J Biochem 257(2):319-30.

Rossi SL, Zhao Q, O'Donnell VK, Mason PW. 2005. Adaptation of West Nile virus replicons to cells in culture and use of replicon-bearing cells to probe antiviral action. Virology 331(2):457-470.

Sadler AJ, Williams BR. 2008. Interferon-inducible antiviral effectors. Nat Rev Immunol 8(7):559-568.

1 Sambrook J, Russell DW. 2001. Molecular cloning: A laboratory manual, 3rd edition, volume
2 1:65-68.

3 Sarkar SN, Ghosh A, Wang HW, Sung SS, Sen GC. 1999. The nature of the catalytic domain of
4 2'-5'oligoadenylate synthetases. J Biol Chem 274(36):25535-25542.

5 Sarkar SN, Miyagi M, Crabb JW, Sen GC. 2002. Identification of the substrate-binding sites of
6 2'-5'-oligoadenylate synthetase. J Biol Chem 277(27):24321-24330.

7 Sarkar SN, Pandey M, Sen GC. 2005. Assays for the interferon-induced enzyme 2'-5'
8 oligoadenylate synthetases. Methods Mol Med 116:81-101.

9 Scherbik SV, Kluetzman K, Pereygin AA, Brinton MA. 2007. Knock-in of the *Oas1b*^(r) allele into a
10 Flavivirus-induced disease susceptible mouse generates the resistant phenotype. Virology
11 368(2):232-237.

12 Shi PY, Tilgner M, Lo MK, Kent KA, Bernard KA. 2002. Infectious cDNA clone of the epidemic
13 west nile virus from New York City. J Virol 76(12):5847-5856.

14 Silverman RH. 2007. Viral encounters with 2'-5'oligoadenylate synthetase and RNase L during the
15 interferon antiviral response. J Virol 81(23):12720-12729.

16 Tag-El-Din-Hassan HT, Sasaki N, Moritoh K, Torigoe D, Maeda A, Agui T. 2012.
17 The chicken 2'-5' oligoadenylate synthetase A inhibits the replication of West Nile virus.
18 Jpn J Vet Res 60(2-3):95-103.

19 Tatsumi R, Hamada K, Sekiya S, Wakamatsu M, Namikawa T, Mizutani M, Sokawa Y. 2002.
20 2'-5'-oligoadenylate synthetase gene in chicken: gene structure, distribution of alleles and
21 their expression. Biochim Biophys Acta 1494(3):263-268.

22 Tatsumi R, Sekiya S, Nakanishi R, Mizutani M, Kojima S, Sokawa Y. 2003. Function of
23 ubiquitin-like domain of chicken 2'-5' oligoadenylate synthetase in conformational
24 stability. J Interferon Cytokine Res 23(11):667-676.

25 Thakur, C.S., Xu, Z., Wang, Z., Novince, Z. and Silverman, R.H., 2005. A convenient and sensitive
26 fluorescence resonance energy transfer assay for RNase L and 2'-5'-oligoadenylates.

1 Methods Mol Med 116:103-113.

2 Yamamoto A, Iwata A, Koh Y, Kawai S, Murayama S, Hamada K, Maekawa S, Ueda S, Sokawa Y.
3 1998. Two types of chicken 2'-5' oligoadenylate synthetase mRNA derived from alleles
4 at a single locus. *Biochim Biophys Acta* 1395(2):181-191.

5 Yamamoto Y, Sono D, Sokawa Y. 2000. Effects of specific mutations in active site motifs of 2'-5'
6 oligoadenylate synthetase on enzymatic activity. *J Interferon Cytokine Res*
7 20(3):337-344.

8 Zhu J, Zhang Y, Ghosh A, Cuevas RA, Forero A, Dhar J, Ibsen MS, Schmid-Burgk JL, Schmidt T,
9 Ganapathiraju MK, Fujita T, Hartmann R, Barik S, Hornung V, Coyne CB, Sarkar SN.
10 2014. Antiviral activity of human OASL protein is mediated by enhancing signaling of the
11 RIG-I RNA sensor. *Immunity* 40(6):936-948.

1 **FIGURE LEGENDS**

2

3 Fig. 1. Schematic diagram of the wild-typed ChOAS-L-A and its mutated proteins as follows;
4 deletion mutation, $\Delta L_{xxx}P$, ΔP -Loop, ΔD -D Box, ΔKR -RR, $\Delta UbL2$, and $\Delta UbL1/UbL2$, point
5 mutation; KR-RR-K193H, and the other allele ChOAS-L-B. Each pattern of column represents each
6 domain and the asterisk represents a point mutation.

7

8 Fig. 2. The measurement of enzymatic activity of the wild-typed ChOAS-L-A and its mutated
9 proteins. A) RT-PCR and Western blot of HEK293FT cells at 48 h post-transfection with
10 pIRES-EGFP (control), pChOAS-L-A-FL-EGFP, and each mutated gene. B) The measurement of
11 enzymatic activity. All mutated proteins showed no enzymatic activity except for ChOAS-L-B and
12 $\Delta UbL2$, both of which showed enzymatic activity with the same extent to that of wild-typed
13 ChOAS-L-A in ChOAS-L-B and slightly less activity in $\Delta UbL2$.

14

15 Fig. 3. Antiviral activity of the ChOAS-L-A and its mutated proteins. BHK-21 cells with mock
16 transfection, pIRES-EGFP, pChOAS-L-A-FL-EGFP, and each of its mutated gene. A) RT-PCR
17 using total RNA extracted from BHK-21 cells before and at 72 h after WNV-replicon RNA
18 electroporation. B) SEAP activity at 24 and 72 h post-WNV-replicon RNA electroporation.

19

20 Fig. 4. Antiviral activity of the ChOAS-L-A and its mutated proteins in BALB/3T3 cells with mock
21 transfection, pIRES-EGFP, pChOAS-L-A-FL-EGFP, and each of its mutated gene. A) RT-PCR
22 using total RNA extracted from BALB/3T3 cells before and at 72 h after WNV-replicon RNA
23 electroporation. B) SEAP activity at 24 and 72 h post-WNV-replicon RNA electroporation.

24

25 Fig. 5. Stability assay of ChOAS-L-A and its mutated protein. Twelve μ g of whole BHK-21 cell
26 lysate transfected with pChOAS-L-A-FL-EGFP and all of its mutants, in the presence or absence of

the proteases inhibitors, were incubated for 0 and 2 h on 37°C, then used for the western blot. All proteins showed stability on 0 and 2 h post-incubation, however, in the absence of the proteases inhibitors, 2 h post-incubation only the protein with mutation in the UbL domains showed degradation.

Fig. 6. RNase L activation assay using the fluorescence resonance energy transfer assay (FRET assay). Two µg protein aliquots of BHK-21 cell lysates transfected with the ChOAS-L-A-FL-EGFP and all of its mutants, were incubated in 50 µl of 1x cleavage buffer with a FRET probe (final concentration 180 ng), for different time points, 0, 5, 10, 15, 30, 45, 60, 90, 120 min at 21°C, then the fluorescence activity was detected. The background fluorescence detected by incubating the FRET probe in the 1x cleavage buffer was subtracted from all reads before statistical analysis.

Fig. 7. Cellular localization of the ChOAS-L-A protein as well as all of its mutant proteins. The BHK-21 cells cultured in 96 well plate, then transfected with the pIRES-EGFP, pChOAS-L-A-FL-EGFP, pΔLxxxP-FL-EGFP, pΔP-LooP-FL-EGFP, pΔD-D-Box-FL-EGFP, pKR-RR-K193H-FL-EGFP, pΔKR-RR-FL-EGFP, pChOAS-L-B-FL-EGFP, pΔUbL2-FL-EGFP, or pΔUbL1/UbL2-FL-EGFP. Forty-eight hours' post-transfection cells were used to perform immunofluorescence protocol. The ChOAS-L-A-FL protein as well as its mutant protein were detected using the mouse-anti FLAG M2 antibody. The Alexa Fluor 568 goat anti-mouse IgG used as secondary antibodies. Nuclei counterstaining was performed by incubating cells with DAPI. The white bar stand for 100 µm.

Fig. 1

Fig. 2-A

Fig. 2-B

Fig 3-A

Fig 3-B

Fig 6A

Fig 6B

1

Fig 5

2
3

Fig. 7

1 **Fig 4**

2
3
4