Title	Selective contribution of the telencephalic arcopallium to the social facilitation of foraging efforts in the domestic chick
Author(s)	Xin, Qiuhong; Ogura, Yukiko; Uno, Leo; Matsushima, Toshiya
Citation	European journal of neuroscience, 45(3), 365-380 https://doi.org/10.1111/ejn.13475
Issue Date	2017-02-05
Doc URL	http://hdl.handle.net/2115/68279
Rights	This is the peer reviewed version of the following article: Xin, Q., Ogura, Y., Uno, L. and Matsushima, T. (2017), Selective contribution of the telencephalic arcopallium to the social facilitation of foraging efforts in the domestic chick. Eur J Neurosci, 45: 365–380, which has been published in final form at 10.1111/ejn.13475. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Type	article (author version)
File Information	EJN45-3 365-380.pdf

[Instructions for use](#)

**Selective contribution of the telencephalic arcopallium to the social facilitation of
foraging efforts in the domestic chick**

Qihong Xin^a, Yukiko Ogura^{b,c}, Leo Uno^a, and Toshiya Matsushima^{d #}

^a Graduate School of Life Science, Hokkaido University, Sapporo, Japan

^b JSPS fellow (PD), Japan Society for Promotion of Sciences, Tokyo, Japan

^c Department of Psychiatry, Graduate School of Medicine, Hokkaido University, Sapporo,
Japan

^d Department of Biology, Faculty of Science, Hokkaido University, Sapporo, Japan

[#] Corresponding author. E-mail: matusima@sci.hokudai.ac.jp

Tel & Fax: +81-11-706-3523

Postal address: N10-W8, Kita-ku, Sapporo, Japan 060-0810, Japan

e-mail addresses

Qihong Xin: qihongxin2013@gmail.com

Yukiko Ogura: y.ogura@med.hokudai.ac.jp

Leo Uno: reouno@eis.hokudai.ac.jp

Toshiya Matsushima: matusima@sci.hokudai.ac.jp

Abstract

To investigate the neural basis of socio-economic behaviors in birds, we examined the effects of bilateral electrolytic lesions of arcopallium (Arco, the major descending pallial area of the avian telencephalon) and the surrounding nuclei in domestic chicks. We tested foraging effort (running distance) in an I-shaped maze with two food patches that delivered food in a biased manner according to a variable interval schedule. Normally, chicks run back and forth between the patches, and the patch use time matches the respective food delivery rate. In the paired phase, even without actual interference of food, chicks showed social facilitation of running effort compared with the single phase. Chicks with lesions in the Arco and lateral Arco showed significant reductions in social facilitation. The lesion effects of the lateral Arco were particularly selective, as it was not accompanied by changes in running distance in the single phase. Lesions of the nidopallium and nucleus taeniae of the amygdala produced no changes in foraging behavior. On the other hand, the Arco lesion did not impair social facilitation of operant peck latency. In accordance with this, anterograde tracing revealed characteristic projections from the lateral Arco to the extended amygdala, hippocampus, and septum, as well as wide areas of limbic nuclei in the hypothalamus and medial areas of the striatum including the nucleus accumbens. Pathways from the lateral Arco could enable chicks to overcome the extra effort investment of social foraging, suggesting functional and anatomical analogies to the anterior cingulate cortex and basolateral amygdala in mammals.

Keywords: social foraging theory, matching law, neuroeconomics, amygdala, anterior cingulate cortex

1. Introduction

To be adaptive, animals must decide how much cost to invest while foraging for food. In mammals, a number of cortical and subcortical structures have been implicated in foraging behavior (Floresco *et al.*, 2008; Wallis & Rushworth, 2014). These include the frontal cortex, specifically the anterior cingulate cortex (ACC) and orbitofrontal cortex (OFC) (Walton *et al.*, 2002, 2003, 2009; Schweimer *et al.*, 2005; Rudebeck *et al.*, 2006), the basolateral amygdala (BLA) (Floresco & Ghods-Sharifi, 2007), and the basal ganglia, specifically the nucleus accumbens (NAc) (Salamone *et al.*, 1994; Cousins *et al.*, 1996; Hauber & Sommer, 2009; but see Walton *et al.*, 2009). Of these, the OFC is critical for processing the time that foragers must wait for food (delay-cost; Rudebeck *et al.*, 2006), whereas the ACC is critical for the amount of effort that foragers must invest to reach and consume food (effort-cost; Walton *et al.*, 2006). Experimental disconnection between the ACC and BLA, as well as inactivation of BLA (Floresco & Ghods-Sharifi, 2007), causes an effort-cost aversion similar to that caused by an ACC lesion. Depending on the type of cost, different subsets of these circuits are recruited during decision-making such that the benefit-cost consequences are appropriate for the ecological and social needs.

Social context is critical when making decisions about effort cost investment. Although classical theories of foraging have assumed single foraging individuals (Charnov, 1976; Stephens & Krebs, 1986), recent behavioral studies clearly support the contribution of inter-individual interactions, such as kleptoparasitism in flocks of birds (Giraldeau & Dubois, 2008), and forms of social cognition such as empathy and perception of fairness in humans (Singer *et al.*, 2006). Additionally, recent imaging studies have revealed a

“mentalizing” network in a region of the ACC that is active when an individual infers the subjective evaluation of rewards generated by another individual (Apps & Ramnani, 2014). However, exactly how social contexts (such as the presence or actions of others) influence foraging decisions remains unclear.

Despite brain structures distinct from mammals, domesticated fowl chicks (*Gallus gallus domesticus*) are useful for studying the social modulation of cost investments and the underlying neural mechanisms. First, chicks make choices depending on the delay-cost (Izawa *et al.*, 2003) and the effort cost (Aoki *et al.*, 2006a), and the relevant brain regions have been localized by lesion experiments. The NAc and surrounding areas in the medial striatum are involved in delay discounting (Izawa *et al.*, 2003), whereas the arcopallium (Arco) is associated with food peck effort (Aoki *et al.*, 2006a) in a doubly dissociated manner (Aoki *et al.*, 2006b). Second, foraging decisions are socially modulated by conspecific individuals. Choice impulsiveness in chicks can be conditionally enhanced by competitive training experiences (Amita *et al.*, 2010), and this enhancement might involve the suppression of striatal neuronal activities, as elicited by the presence of the accompanying forager (Amita & Matsushima, 2014). Furthermore, foraging effort (Ogura & Matsushima, 2011) and operant peck latency (Amita & Matsushima, 2011) are socially facilitated in a reversible and contextual manner. Based on the “drive” theory (Zajonc, 1965), we initially hypothesized that ascending dopaminergic pathway is involved in the social facilitation. Actually, localized lesion to the tegmentum around the substantia nigra (SN, the major ascending dopaminergic nucleus) partially reduced the facilitation. However, selective depletion of dopaminergic neurons failed to reproduce the effect, and the

facilitation remained unchanged (Ogura *et al.*, 2015). Because the lesioned area included fiber tracts descending from the telencephalon (occipito-mesencephalic tract, OM), we subsequently hypothesized that the Arco mediates the social facilitation through its descending projections to the tegmentum.

Functional contributions of the Arco are yet only poorly understood. It is a region of the avian pallium in the telencephalon, and was incorrectly labelled the archistriatum before the nomenclature reform (Reiner *et al.*, 2004). The Arco was thereafter considered to be a homologue of the mammalian amygdala because lesion to Arco reduced escape responses in adult ducks (Phillips, 1964). In accordance with this, electrical stimulation of the Arco caused freely-behaving chickens to emit crow-like vocalizations (Phillips *et al.*, 1972). Another dominant theory held that the Arco was part of the motor/premotor cortex (Veenman *et al.*, 1995), or more specifically, part of the pallial area involved in oro-facial motor control (Wild *et al.*, 1985). In songbirds, localized lesion of the robust nucleus of the Arco impaired song production (Nottebohm *et al.*, 1976). The Arco was also shown to code working memory for sound localization in barn owls (Knudsen & Knudsen, 1996a), suggesting that this region is similar to the lateral prefrontal cortex. Finally, Arco lesions in chicks led to avoidance of a costly option associated with a larger food reward (Aoki *et al.*, 2006a), suggesting a function analogous to the mammalian ACC or BLA.

In the present study, we examined whether and how the Arco functionally contributes to social facilitation. In addition to the foraging effort (running distance), we paid attention to the patch use ratio, which measures how long chick stays at feeders of different food delivery rate. We recently found that a biased food delivery to two feeders caused a biased

stay time, and that the paired chicks showed a significantly better matching to the delivery rate than the chicks in single condition (Q. Xin, Y. Ogura and T. Matsushima, submitted). We assume that ratio of the stay time (patch use ratio) represents the relative value associated with the feeders (as of the cue period neuronal activities in Arco neurons; Aoki *et al.*, 2003). If the patch use ratio changed after the lesion, we may argue that the effects on the foraging efforts could be indirectly caused by the altered value representations. In addition, we measured a behavioral index of motor skill of pecking in some of the chicks tested.

We also examined the lesion effect of nearby structure, the nucleus taeniae of the amygdala (TnA). The TnA is supposed to be homologous to the amygdala (Cheng *et al.*, 1999), particularly to the sub-pallial medial amygdala of mammals (Yamamoto *et al.*, 2005). Furthermore, we examined nidopallium (Nido) located just dorsal to the Arco, because our Arco lesion often extended to Nido. The effect of the Arco lesion could thus be due to the collateral damages to Nido. It is also to be noted that, in pigeons, particularly its caudo-lateral part of Nido (NCL) is supposed to play an analogous function to prefrontal cortex in mammals (Diekamp, Kalt, et al., 2002; Lissek & Güntürkün, 2003).

As a separate experiment, we examined the lesion effect on the operant peck latency. As described above (Amita & Matsushima, 2011), the peck latency is contextually shortened in the social (triplet) condition. If the shortening in peck latency changed after the lesion, we may argue that the effects on the foraging efforts were caused by a more general reason such as incapability to visually perceive conspecifics. Finally, assuming the heterogeneous organization of sub-regions within the Arco, we examined efferent projections by focal

infusions of anterograde tracer (biotinylated dextran amine, BDA), and attempted to relate the efferent projections with the lesion data.

2. Materials and methods

2.1 Subjects and Ethical Note

We used male domestic chicks (*Gallus gallus domesticus*, White Leghorn strain). On post-hatch day 1 (presumed hatching day), we purchased new hatchlings from a local supplier (Iwamura poultry Ltd. Hokkaido establishment, Yubari, Hokkaido, Japan). Chicks were paired and communally housed in transparent plastic cages (15 cm × 28 cm × 12 cm) in a room illuminated according to a 12:12-h light:dark cycle with the light period starting at 08:00 a.m. The temperature of the room was kept constant at ca. 30 °C.

The chicks received two types of food, i.e., grains of millet and chick mash food. The total amount of food per day was adjusted so that (1) the body weight of the chicks gradually increased and (2) the chicks actively consumed food during experiments. Chicks were given mash food from post-hatch day 1. Specifically, they received 2 g (post-hatch days 1–3) and 2.5 g (day 4 and afterwards) of mash food. Grains of millet were added from post-hatch day 2. The amounts of grains were 2 g (days 2–3) and 2.5 g (day 4 and afterwards). Until post-hatch day 2, all chicks were communally fed. From post-hatch day 3, chicks were individually fed in cages that were visually separated by black boards. Water was freely available.

The experiments were conducted under the guidelines and approval of the Committee on Animal Experiments of Hokkaido University. The guidelines are based on the national

regulations for animal welfare in Japan (Law for Humane Treatment and Management of Animals; after a partial amendment No. 68, 2005). After the experiments, the brains of the chicks were dissected under deep anaesthesia. In cases in which surgical operations were not conducted, the chicks were euthanized with carbon dioxide.

2.2 Apparatus

2.2.1 I-shaped maze (Experiments 1 and 2)

We used an I-shaped maze equipped with two lanes (each 12 cm wide, 88 cm long with 40 cm high side walls; Fig. 1A), as reported previously (Ogura & Matsushima, 2011; Ogura *et al.*, 2015). The two lanes were separated by a transparent Plexiglas partition. Except noted otherwise, each lane accommodated one chick at a time. One terminal wall was painted red, and the other yellow (Experiment 1) or blue (Experiment 2). Each wall was equipped with a food patch comprising a pair of food trays, such that there was one tray at each end of a lane. Each tray was a rectangular plastic box (3 cm wide, 4 cm long and 2 cm deep) with a sponge on the bottom of the box. The two trays on each wall were placed side by side, but separated by the Plexiglas wall, so that each chick was individually fed without interference regarding the delivered food. A micro-robot (RCX, LEGO Mindstorms) delivered grains of millet simultaneously to both paired trays in a patch, such that the chicks could be fed concurrently, and thus experienced fictitious interference regarding the food. One micro-robot fed the yellow patch, and another fed the red patch independently. If a lane did not accommodate a chick, food was not delivered to that lane. To prevent the chicks from associating the mechanical sounds made by the micro-robots

with the food reward, we positioned two dummy motors close to the maze. These generated the same motor sounds as the micro-robots at variable intervals (mean = 2.5 s, range = 1.5 to 3.5 s) as distractors. The entire apparatus was placed in a dark room maintained at ca. 25–30 °C and illuminated by four 60 W white light bulbs located above the maze.

We recorded the behavior of the chicks using a video camera/recorder (DCR-SR65, Sony, Japan; 30 frames per sec) placed on the ceiling above the I-maze, providing an aerial view of the subjects. Each chick was marked with a piece of fluorescent colored tape attached to the head. The trajectory of the foraging chicks was tracked offline using Move-Tr/2D 7.0 software (Library Co., Japan) on a PC.

2.2.2 Operant chamber (Experiment 3)

We used a hand-made operant box (20 cm × 20 cm × 19 cm, kept at ca. 27–30 °C) for recording behaviors (Fig. 1C). One of the surrounding walls was equipped with a liquid crystal display (LCD) monitor with the following specifications: size 10.4”, 800 × 600 pixels; Logitech LCM-T102AS, Japan; flash rate: 56–75 Hz; brightness: 230 cd/m²; and pitch size: 0.264 × 0.264 mm. The LCD monitor was covered with a 1-mm thick transparent layer of Plexiglas for protection of the surface, and a frame-type optical touch sensor (size 10.4”, E10D03U-30 without-glass, TouchTEK Co., Yokohama, Japan) for detecting chick pecking. The LCD monitor with the touch sensor was separated from the chick chamber by a 1-mm thick transparent Plexiglas plate. The plate had two open windows (each 2.5 cm × 3 cm, placed side by side at 3 cm, 4 cm above the floor level) through which the chicks pecked at visual stimuli (color cues) displayed on the LCD

monitor at positions corresponding to the windows. Red or green rectangles appeared on the white screen as cue stimuli.

The presentation of visual cues on the monitor and the activity of the solenoids were coordinated, and we recorded the timing of pecking behavior using a programme written using LabView (ver.2010, National Instruments Co., Austin, Texas, USA). The operant box was separated into two chambers by a transparent Plexiglas partition, one accommodating the subject chick and the other accommodating two companion chicks. The chick behavior was monitored using color CCD cameras (250 k pixels, MTV-54B(K)ON, Akizuki-Denshi Co., Tokyo, Japan) placed in the chamber.

2.3 Behavioral procedures

2.3.1 Experiments 1 and 2: running distance, synchrony, and patch use ratio

On post-hatch day 5 and 6, chicks were habituated to the maze in single daily sessions. As part of the habituation process, ca.100 grains of millet were placed at the midway point of the maze, and ca. 40 grains were placed in each of the trays on the sides such that chicks received about ca. 180 grains as starter food. A pair of cagemate chicks was then introduced into the lane with the starter food. Once the chicks had consumed the food, they received a brief intermission of about 1 min before the feeders started to deliver millet grains according to a VI15 schedule, i.e., at variable intervals with mean = 15 s (uniformly in a range of 10–20 s). Two grains of millet were delivered at a time, and the pair of chicks received a total of 120 grains (60 deliveries) from each patch during a ca. 16 min period.

After the delivered food was consumed, the chicks were left in the maze for an additional 2 min.

After habituation, the chicks were tested for 4 days with two test sessions per day. We conducted four tests before and four tests after lesion (pre-lesion and post-lesion conditions, respectively). Each test comprised three consecutive phases: single #1, paired, and single #2 (Fig. 1A). The lesion operation was conducted on the day after the last pre-lesion tests, i.e., post-hatch day 9 or 11 (Experiment 1 or 2, respectively), and the chicks were allowed to recover for 1–2 days before the post-lesion tests.

In Experiment 1, asymmetric food delivery started as soon as the subject chicks were introduced to the maze (Fig. 1B). One feeder supplied food according to a VI10 schedule (mean = 10 s, range = 6.7–13.3 s) and the other had a VI30 schedule (mean = 30 s, range = 20–40 s), so that the former feeder was three times more profitable than the latter. Color assignment was pseudo-randomized among individual chicks. Chicks received two grains of millet at a time. One phase consisted of 45 food deliveries (90 grains) from the more profitable VI10 feeder, and 15 food deliveries (30 grains of millet) from the other VI30 feeder. The duration of each phase was approximately 8 min. In the single #1 phase, chicks were tested individually. After completion of the single #1 phase, a companion chick (cagemate) was introduced into the other lane of the maze, and the paired phase started immediately. During the paired phase, grains were simultaneously delivered to the chicks in both lanes. Finally, in the single #2 phase, the companion was removed from the maze, and the subject chick resumed foraging independently. After the food supply was terminated, the subject was left in the maze for an additional 2 min.

In order to check if the lesion impaired motor function, we examined the pecking food behavior as reported previously (Aoki *et al.*, 2006a). On the first day after the lesion, namely just before the first post-lesion test, single chick placed in the home cage was given a plastic box (4 cm x 6 cm, 2 cm deep) containing 6 grains of millet for four consecutive times with a brief intervals of 5 min. The grains were placed on a sponge on the floor of the box. We counted how many times the chick pecked until the 6 grains were all consumed, averaged the number and divided it by 6 to yield an index to denote the number of pecks per grain (κ ; Matsushima *et al.*, 2008).

As Experiment 2 was actually executed as a pilot experiment before Experiment 1, it had a somewhat different schedule. First, during the acclimatization phase, the chicks consumed 20 grains supplied to each feeder (40 grains in total). Second, both of the terminal patches delivered food in a symmetric manner according to the same VI15 schedule (mean = 15 s, range = 10–20 s), and one grain of millet was delivered at a time. Third, one of the terminal walls was painted blue, rather than yellow. However, we noticed no effect of the blue compared with the yellow wall.

2.3.2 Experiment 3: response peck latency

Chicks were initially habituated to the chamber, and trained to peck the color cue in groups of three individuals. Thereafter, chicks were tested in two phases, i.e., single and triplet (Fig. 1C). In the triplet phase, the subject was separated from two companion chicks by a Plexiglas partition, and received a controlled amount of food that was not scrounged by the companions. Red and green rectangles were displayed as cues. One of the cues was associated with a food reward (S+) and the other with no food (S-). Cue assignment was

randomized among individuals. Fig. 1D shows the procedure of a single trial. First, a color cue was presented for 2 s without any preceding stimuli. When the subject chick pecked at the S+ cue once or more, 6 grains of millet were delivered to the central food tray after a brief mechanical lag (0.29 s on average).

2.3.2.1 Training for measurement of operant peck latency

In order to examine the lesion effects on the response peck latency in those subjects after substantial training, chicks were trained in three phases: habituation, auto-shaping, and differential training. Habituation took place on post-hatch days 2–3, and auto-shaping took place on days 3–4. Both types of training were administered in one block per day. In the habituation phase, triplets of chicks were introduced to the chamber without the Plexiglas partition for 20 min, and given 80 grains of millet without any cues. In the auto-shaping phase, chicks were trained to peck at a white plastic rod (5 mm in outer diameter) that was manually presented to the subjects in 40 trials. Two grains of millet was delivered as soon as one of the three chicks pecked at the rod.

We tested those chicks that successfully associated the color cues with subsequent reward. To do that, on days 5–7, the chicks received two blocks of differential training per day, one block in the single phase and another in the triplet phase (Fig. 1C). The order of the two phases was counter-balanced among individuals. Each block consisted of 50 pseudo-randomly arranged trials including 20 trials with a rewarding color (S+ for 6 grains), 20 trials with a non-rewarding color (S- for no food), and 10 binary choice trials in which the chick had to choose between S+ and S-. In the binary trials, the chicks were not rewarded, regardless of whether they had correctly pecked at the S+. In the triplet phase,

the companion chicks received two grains when the subject successfully pecked and received food in the S+ trials.

2.3.2.2 Pre-lesion and post-lesion tests

On days 8–10, pre-lesion tests were conducted in the same manner as the differential training on days 5–7. On day 11, trained chicks were randomly allocated to two groups, which received either sham or lesion surgery. The chicks were allowed to recover from the operation for 1–2 days, and underwent post-lesion tests on days 13–15, one in the single phase and another in the triplet phase. We measured the response peck latency in single cue trials with the S+ color, and in binary choice trials with the S+ and S- options. Trials in which the chicks successfully pecked at the rewarding S+ option were analysed.

2.4 Surgical procedure for electrolytic lesioning

Chicks were anesthetized via an intramuscular injection of 0.4 ml of a ketamine-xylazine cocktail, which was a 1:1 mixture of 10 mg/ml ketamine (Daichi Sankyo Co., Ltd., Tokyo, Japan) and 2 mg/ml xylazine (Sigma-Aldrich Co., LLC, St. Louis, MO, USA). Supplementary injections (0.1 ml) were given to maintain stable anaesthesia. The chicks were fixed to a stereotaxic apparatus (SR-5N, Narishige, Tokyo, Japan) at the head angle set at ca. 45°. Body temperature was maintained using a heating pad. The skin over the skull surface was incised, areas of the skull removed, and the dura mater was cut to expose the brain. We constructed steel electrodes from electrolytically sharpened insect pins (type #00, max 300µm thick; Shiga Konchu Co., Tokyo, Japan). The pins were coated with enamel paint (Tamiya Inc., Shizuoka, Japan), leaving the ~0.6 mm-long tip unpainted.

Lesions were conducted bilaterally in all experiments. For lesioning the Arco (Experiment 1 and 3), the electrode was vertically inserted toward the coordinates: 2.0–2.3 mm anterior from bregma, 5.5 mm lateral from the midline, and 4.5 mm deep from the brain surface. For lesioning the lateral Arco, the electrode was vertically inserted toward the coordinates: 2.0–2.3 mm anterior from bregma, 6.5–6.6 mm lateral from the midline, and 3.7–3.9 mm deep from the brain surface. For lesioning the Nido, the coordinates were: 2.0–2.2 mm anterior from bregma, 5.5 mm lateral from the midline, and 2.0–2.5 mm deep from the brain surface. For lesioning the TnA, the coordinates were: 0.9–1.0 mm anterior from bregma, 4.1 mm lateral from the midline, and 6.0–6.2 mm deep from the brain surface. For the sham control group, we anaesthetized the subjects, placed them into the stereotaxic apparatus, incised the brain surface, and inserted an electrode, but did not apply current.

Chicks received one lesion per hemisphere. We applied pulses with constant current and alternating polarity (amplitude: ± 1.5 mA, pulse duration: 50 ms, repetition rate: 10 Hz, 1 min in duration) to the inserted pin electrode using an electric stimulator (SEN-3301 and isolating unit SS-403J, Nihon Kohden, Tokyo, Japan). A silver wire was placed on the caudal skull as a reference electrode. After applying current, the electrode was left in place for 5 min before withdrawing. We covered the incised skull with the skin flap using superglue (Aron Alpha®, Toa Gosei Co., Ltd., Tokyo, Japan). After the surgery, chicks recovered individually overnight, and were then housed with their former cagemates.

2.5 Histological examination of the lesion site

On the day after the final test, the chicks were deeply anesthetized via an

intramuscular injection of approximately 0.8 ml of ketamine-xylazine cocktail, and transcardially perfused with a fixative (4% paraformaldehyde in 0.1 M PB, pH 7.4). The dissected brain samples were post-fixed in the same fixative for one day at 4 °C, and then cryoprotected in 20% sucrose-PBS for one day at 4 °C. The brain tissue was frozen at –80°C for storage, and cut into 50 µm frontal sections using a sliding microtome with a freezing unit (Yamato Kohki Industrial Co., Ltd., Saitama, Japan). In some cases, in order to prevent brain regions (such as telencephalon and diencephalon) to fragment, the samples were embedded in yolk and post-fixed in the same fixative for >2 days at 4 °C. The tissue was then cut into 100 µm frontal sections using a vibrating microslicer (DTK-1000, Dosaka EM, Kyoto, Japan). Sections were mounted, air-dried, stained with cresyl violet, dehydrated, cleared using xylene, and cover-slipped.

The coordinates were based on the stereotaxic atlas of the chick brain by Kuenzel and Masson (1988) (Kuenzel & Masson, 1988), and terminology adhered to the nomenclature reform (Reiner *et al.*, 2004).

2.6 Experiment 4: anterograde tract tracing

2.6.1 Injection of BDA

To examine the efferent terminals projecting from different sites within the Arco, we used biotinylated dextran amine (BDA, 10% in distilled water, 10 kDa; D22910, Molecular Probes®, Thermo Fisher Scientific Inc., Waltham, MA, USA) as an anterograde tracer. We used 15 chicks: 13 for different sites within the Arco, and 2 for the Nido. On approximately post-hatch day 7, we injected BDA using a Nanoject II (Drummond Scientific Co.,

Broomall, PA, USA) under ketamine-xylazine anaesthesia, as described in the methods for the lesion experiment. The tracer was deposited via a slow pressure injection lasting for 10–15 min (13.8 nl per injection × 11 injections, 150 nl per site). After the injection, the glass capillary was left in place for an additional 10 min to minimize leakage of the dye along the injection track.

2.6.2 Perfusion and sectioning

Seven days after the BDA injection, chicks were deeply anaesthetized and perfused with 4% paraformaldehyde in 0.1 M phosphate buffer (PB, pH 7.4), as in the post-lesion histological examination. Dissected brains were post-fixed, cryoprotected, and sectioned into 50- μ m sections on a frontal plane using a freezing microtome. Sections were stored in 4°C phosphate-buffered saline (PBS) for histochemical processing.

2.6.3 Histochemical processing of BDA-labelled terminal fibers

Sections were pretreated in 3% H₂O₂ in PBS (pH 7.4) for 30 min, rinsed in PBS (4×), incubated in avidin-biotinylated horseradish peroxidase complex (ABC, Vectastain® Elite ABC Kit, Vector Laboratories, Inc., Burlingame, CA, USA) for 1 hr at room temperature, and then rinsed in PBS (4×) and in distilled water (1×). BDA labelling was developed using nickel-enhanced diaminobenzidine (3,3-diaminobenzidine tetrahydrochloride, DAB, DAB Peroxidase Substrate kit, SK-4100, Vector Laboratories). The sections were incubated in distilled water containing Tris buffer, DAB, H₂O₂, and nickel-ammoniumsulphate (1:2:1:1) for 5–10 min. The reaction was visually controlled and terminated by rinsing the sections in PBS (3×). After the staining was developed, the sections were mounted onto amino silane -

coated glass slides (Matsunami glass Ind., Ltd., Japan), air-dried, dehydrated, cleared, and coverslipped.

2.7 Data analysis

2.7.1 Measurements of behaviors

In Experiments 1 and 2, we assessed the following behavioral indices. (1) Running distance was given by the sum of the distance that the subject ran in each trial in the single #1, paired, and single #2 phases. (2) The synchrony index was given as the ratio of time in which both chicks stayed in the same side of the maze, divided by the total trial time. When the two chicks were perfectly synchronized, the synchrony index was 1.0 (in-phase synchrony). When two chicks moved independently, the index was at the chance level, i.e., 0.5 (asynchrony). (3) The patch use ratio was given as the proportion of time in which the subject stayed close (<10 cm) to the VII0 feeder, divided by the total stay time at both feeders. The running distance and the patch use ratio in the two single phase sessions were averaged giving rise to $\overline{\text{single}} = \frac{\text{single \#1} + \text{single \#2}}{2}$. In Experiment 3, we assessed the response peck latency of rewarding trials in the single and triplet phases.

2.7.2 Statistical analyses of behavior data

We focused on the post-lesion running distance, synchrony index, patch use ratio (Experiments 1 and 2), and response peck latency (Experiment 3) as response variables. *Group:* Arco, lateral Arco, Nido lesions and sham control in Experiment 1; the TnA lesion and sham control in Experiment 2; and the Arco lesion and sham in Experiment 3. *Phase:* single and paired in Experiments 1 and 2; single and triplets in Experiment 3.

For Experiments 1 and 2, we made inter-group comparisons of running distance and patch use ratio by a two-way analysis of variance (ANOVA) with repeated measures based on type III sums of squares. The synchrony index was compared using a one-way ANOVA (Experiment 1) or Student *t*-test (Experiment 2). For Experiment 3, we made inter-group comparisons of response peck latency using a two-way ANOVA with repeated measures based on type I sums of squares.

Post-hoc pairwise comparisons were made after a Holm correction. The significance level was set at $P = 0.05$. All statistical calculations were performed using R (version 3.1.3, Windows version) and the “ANOVAKUN 4.8.0” add-on (programmed by Dr. Ryuta Iseki, <http://riseki.php.xdomain.jp/index.php?ANOVA%E5%90%9B>).

To compare the number of pecks per grain (κ index), we adopted a non-parametric permuted Brunner-Munzel test according to the R script developed by Dr. Haruhiko Okumura (Mie University, Tsu-city, Mie, Japan; <http://oku.edu.mie-u.ac.jp/~okumura/stat/brunner-munzel.html>). Pairwise comparisons were made among groups. P value was shown after Bonferroni correction.

2.7.3 *Quantitative analysis of BDA projection patterns*

In Experiment 4, labelled fibers and terminal arborizations were localized according to the brain atlas (Kuenzel & Masson, 1988; Puelles *et al.*, 2007). Further abbreviations of neural nuclei and nuclei boundaries followed the nomenclature reform (Reiner *et al.*, 2004) and also the anatomical data reported by Montagnese *et al.* (2003) and Hanics *et al.* (2016). Based on visual inspections of the intensity of labels, blind to the injection site, we determined five grades of fiber density: abundant, moderate, low, sparse (only 2–3 fibers in

the entire region), and absent. We used hierarchical clustering analysis (Everitt *et al.*, 2011) with the Euclidean metric and the average agglomeration method to create clustering of the efferent regions. We generated a heatmap with dendrogram classification using the “pheatmap” package (version 1.0.8) of R statistics.

3. Results

Sixty-seven chicks underwent surgical operations. Of these, four chicks were discarded in Experiment 1, as three chicks died during or after the surgery, and one stopped foraging. One and two chicks were similarly discarded in Experiments 2 and 3, respectively. The present results are based on data obtained from the remaining 60 chicks. All of these chicks actively foraged in their homecage and the apparatus, maintaining normal postures and locomotor activities throughout the experiments, including post-lesion. Of the additional 16 chicks that received BDA micro-injections (Experiment 4), 15 showed successful staining.

3.1 Behavioral effects of lesions (Experiments 1, 2, and 3)

3.1.1 Histology

In Fig. 2A, ablated areas are superimposed for 5 groups of chicks examined in Experiments 1, 2, and 3. For the Arco, the tissue damage included the intermediate Arco, dorsal Arco, and part of the lateral Arco, but the TnA appeared undamaged (Fig. 2Bb). The Nido was also partially lesioned in some of the chicks. For the lateral Arco, the lateral Arco was fully ablated in all chicks, while the dorsal Arco and intermediate Arco were partially damaged in some chicks. For the Nido, damage was localized to the areas dorsal to those

lesioned in the Arco chicks. For the TnA, the bilateral lesion was successful, sparing the Arco (Fig. 2Bc).

3.1.2 Behavioral effects of Arco lesion (Experiment 1)

Chicks actively ran back and forth between the two patches. Representative trajectories of chicks with sham (n=11), Arco (n=7), lateral Arco (n=6), and Nido (n=7) lesions are shown in Fig. 3. Examples from the first session on post-lesion day 1 are included. In the single phase, chicks in all groups tended to stay longer at the yellow patch (bottom) compared with the red patch (top), suggesting biased patch use. Note that the food delivery timing (arrowheads) did not strictly entrain the chick behaviors. In the paired phase, both the sham chicks and chicks with Nido lesions exhibited facilitated running and a high degree of synchrony. However, the chicks with Arco lesions were slower in the single phase, and often desynchronized in the paired phase. The chicks with lateral Arco lesions were also desynchronized in the paired phase, but reduced running was not apparent in the single phase.

In Fig. 4Aa,b, running distance and patch use ratio are plotted along eight successive sessions, four pre-lesion and four post-lesion. In the pre-lesion sessions, the running distance increased in the paired phase compared with the single #1 and #2 phases. The patch use ratio was slightly higher in the paired vs. the single phases. In the post-lesion tests, chicks with Arco lesions (green) exhibited a reduced running distance in both phases. Chicks with lateral Arco lesions (orange) showed a selectively suppressed running distance in the paired phase. On the other hand, chicks with Nido lesions (blue) behaved similarly to

the sham controls. We found no significant differences in patch use ratio among the groups of chicks in the post-lesion tests.

We further compared the running distance data in the post-lesion tests among the four groups of chicks in the two phases (Fig. 4Ac); data in the single #1 and #2 phases were merged to give an average (*single*). A two-way ANOVA revealed significant main effects of group ($F_{3,27} = 9.450, P < 0.001$), phase ($F_{1,27} = 264.604, P < 0.001$), and the group $\times$ phase interaction ($F_{3,27} = 17.221, P < 0.001$). Multiple comparisons failed to reveal significant differences in the *single* phase data, but a suggestive level of difference between the Arco vs. sham groups ($t = 2.744, P = 0.064$) should be noted. In the paired phase, we found significant differences between the Arco vs. sham ($t = 6.159, P < 0.001$), Arco vs. Nido ($t = 5.347, P < 0.001$), lateral Arco vs. sham ($t = 3.465, P = 0.007$), and lateral Arco vs. Nido ($t = 2.945, P = 0.020$) groups, but not between the sham vs. Nido ($t = 0.248, P = 0.806$) or Arco vs. lateral Arco ($t = 2.192, P = 0.074$) groups.

The synchrony index data are compared in Fig. 4Ad. A one-way ANOVA detected a significant effect of group ($F_{3,27} = 11.076, P < 0.001$). Multiple comparisons by Holm's tests revealed significant differences between the Arco vs. sham ($t = 4.132, P = 0.002$), Arco vs. Nido ($t = 4.218, P = 0.002$), lateral Arco vs. sham ($t = 3.901, P = 0.002$) and lateral Arco vs. Nido ($t = 4.019, P = 0.002$) groups, but not between the sham vs. Nido ($t = 0.033, P > 0.999$) or Arco vs. lateral Arco ($t = 0.531, P > 0.999$) groups.

The patch use ratios are compared in Fig. 4Ae. A two-way ANOVA detected significant main effects of phase ($F_{1,27} = 4.759$, $P = 0.038$), but not group ($F_{3,27} = 0.226$, $P = 0.878$) or group $\times$ phase interaction ($F_{3,27} = 0.786$, $P = 0.512$).

The κ index (number of pecks per grain) was compared in four groups of chicks in Experiment 1; 1.12 ± 0.04 (mean $\pm$ SEM, sham, $n = 7$; median = 1.10), 29.01 ± 20.05 (Arco, $n = 5$; median = 6.50), 2.70 ± 0.48 (lateral Arco, $n = 6$; median = 2.59), 1.07 ± 0.02 (Nido, $n = 7$; median = 1.08). We did not measure the index in some chicks in sham and Arco groups. It is also to be noted that the large SEM value in the Arco group was due to 2 chicks with exceptionally high κ index (107.50 and 25.11). Permuted Brunner-Munzel test revealed significant difference between Arco vs. sham ($P = 0.015$), and marginally significant difference between lateral Arco vs. sham ($P = 0.049$), but not between Nido vs. sham ($P > 0.999$).

3.1.3 Behavioral effects of TnA lesion (Experiment 2)

The effects of the TnA lesion ($n=6$) on running distance are shown in Fig. 4B. Food delivery was not biased in Experiment 2. In terms of running distance (4Ba), a two-way ANOVA revealed significant main effects of phase ($F_{1,15} = 966.044$, $P < 0.001$), but not group ($F_{1,15} = 1.916$, $P = 0.187$) or interaction ($F_{1,15} = 0.196$, $P = 0.664$). Student t -tests did not reveal a significant difference in synchrony indices (4Bb) ($t_{15} = 1.525$, $P = 0.148$). In terms of the patch use ratio (4Bc), we found no significant effects of group ($F_{1,15} = 1.923$, $P = 0.186$), phase ($F_{1,15} = 1.717$, $P = 0.210$), or interaction ($F_{1,15} = 0.007$, $P = 0.935$).

3.1.4 Effects of Arco lesion on peck latency (Experiment 3)

The peck latency did not differ between the group of chicks with the Arco lesion (n=6) and the sham controls (n=6; Fig. 4C). A two-way ANOVA detected a significant main effect of phase ($F_{1,10} = 38.377, P < 0.001$), but not group ($F_{1,10} = 0.525, P = 0.485$) or interaction ($F_{1,10} = 1.494, P = 0.250$).

3.2 Anterograde tracing of efferent projections from the Arco

Fig. 5A shows representative injection sites in three chicks, and Fig. 5B–D show anterogradely labelled fibers with dense varicosities. Retrogradely labelled cell bodies (Fig. 5Cg) were found in some cases (see below). Intensive labels appeared almost exclusively in the hemisphere ipsilateral to the injection site. In a few cases, however, sparse labels occurred in the contralateral midbrain. Furthermore, we found projections to the contralateral telencephalon (including the contralateral Arco) in one chick. In the following analyses, we disregarded these contralateral projections. After the BDA injection into the lateral Arco, we found dense terminal labels in the hippocampus (Hp, Fig. 5Ba), septum (Sept, 5Bb), and nucleus accumbens (NAc, 5Bc), as well as other regions. Injection into the dorsal Arco (5C) yielded terminal labels in a wide range of areas such as the intermediate medial mesopallium (IMM, 5Ca), caudal extended amygdala (CEA, 5Cb), and ventral tegmental area (VTA, 5Cc), which was also labelled after the lateral Arco injection. On the other hand, we found labelling in the following areas after injection into the dorsal and the intermediate Arco: terminals in the medial spiriformis nucleus (SPM, 5Cd), the

intercollicular nucleus (ICo, 5Ce), and the optic tectum (TeO, 5Cf). In addition, injection into the dorsal Arco yielded retrogradely labelled cell bodies in the Nido (5Cg), whereas no such labels appeared after the lateral Arco injection. When BDA was injected into the Nido, anterograde labels were found in limited brain regions. These included the Arco, lateral striatum (LSt, 5D), MSt, rostral and caudal part of extended amygdala (rEA and cEA).

The results of hierarchical cluster analysis of the efferent projection patterns are shown in Fig. 6. Chicks were categorized into three main clusters (arbitrarily labelled as α , β , and γ), each of which proved to correspond to the injection site. Chicks in cluster α had received a Nido injection ($n = 2$), and their projection patterns were distinct from those in the other groups. Chicks in cluster β corresponded to the lateral Arco ($n = 5$), and chicks in cluster γ included three sub-clusters; γ_1 for lesions in the PoA and ventral Arco, γ_2 for the dorsal Arco, and γ_3 for the intermediate Arco (see the injection sites below).

The regions of the efferent fibers and terminals were categorized into 4 clusters, δ , ϵ , ζ , and η (η_1 , η_2 , and η_3). Many telencephalic nuclei were allocated to clusters δ and ϵ . Some “limbic” structures such as the hippocampus (Hp) and septum (Sept) were included in cluster ϵ , which mainly received efferents from the lateral Arco. Chicks with BDA injection to the ventral Arco were grouped in cluster δ , but the terminals were sporadic and sparse if present. Brain areas categorized in the ζ cluster (rEA and cEA) received projections from both lateral and other Arco regions. Nuclei in the thalamus, hyper thalamus, tegmental areas, and optic tectum (TeO) were categorized in cluster η . The η_1 cluster included the LSt and MSt, the former of which selectively received efferents from the dorsal and the ventral Arco (or collectively, the medial Arco). It should be noted that the Nido strongly projected

to the LSt (Fig. 5D). The LSt labels after the medial Arco injection could therefore be due to a diffusion of BDA from the injection site. The η_2 cluster included the midbrain central gray (GCT), thalamic anterior dorsomedial nucleus (DMA), thalamic posterior dorsomedial nucleus (DMP), thalamic anterior dorsolateral nucleus (DLA), thalamic posterior dorsolateral nucleus (DLP), and thalamic posterior dorsointermediate nucleus (DIP). The η_2 cluster received projections from both the lateral and more medial Arco regions. Midbrain areas such as the intercollicular nucleus (ICo) and optic tectum (TeO) were included in cluster η_3 , which selectively received projections from the medial Arco. In the Nido, retrogradely labelled cell bodies appeared only in those chicks that had received an injection in the medial regions of the Arco.

4. Discussion

Bilateral electrolytic lesions of the Arco and lateral Arco significantly suppressed social facilitation of the foraging effort (Fig. 4Aa,c), but had no effect on matching behavior or peck latency (Fig. 4Ab,C). The effect of the lateral Arco lesion was selective, as it was not accompanied by changes in the running distance in the single phase. In accordance with this, anterograde tracing revealed distinct patterns of efferents from the lateral Arco (Figs. 5, 6 and 7). On the other hand, TnA lesion had no effects (Fig. 4B), though it is noted that the task of Experiment 2 is slightly different from Experiment 1. In the following, we will discuss the neural substrates responsible for social facilitation, and the possible correspondence between the avian and mammalian social neuro-economic systems.

The high κ index found in the Arco and lateral Arco groups warrants a careful consideration on the lesion effects on running distance. When consuming food, chicks usually peck 2-4 times per sec in average (Matsushima et al., 2008). In Experiment 1 of the present study, chicks were given two grains of millet at a time. Therefore, in these chicks with a high κ value, the consumption time (involved in the patch stay time) could be inevitably longer when chick encountered food, thus reducing the time for running. The suppressed running time in the single condition in the Arco group of chicks could be, at least partially, due to a side effect of the pecking impairment (Fig. 4Ac). The lateral Arco chicks showed a suppressed social facilitation without effects on the running distance, even though the κ index was significantly lower. These results are in concert with the idea that Arco plays a functional role in oro-facial motor control (Wild et al., 1985; for further discussions, see below in section 4.1.1). On the other hand, the observed suppression in the social facilitation by the lesion cannot be accounted for by the pecking impairment that accompanies.

4.1 Neural substrates of social facilitation, effort investment, matching behavior, and operant peck latency

4.1.1 Social facilitation

A straightforward interpretation of the present lesion study is that the Arco, particularly the lateral region, enables chicks to overcome the extra effort that must be paid in a social foraging context. Most probably, the lateral Arco corresponds to the amygdalopiriform area of the Arco (abbreviated as APir; Puelles *et al.*, 2007, Hanics *et al.*, 2016), which is characterized by its dense projections to the NAc and BSTl. In a previous study, we

revealed that Arco lesions resulted in handling-cost aversion in a binary choice task, in which the larger option was accompanied by more pecks (Aoki *et al.*, 2006a). In present study, we focused on the foraging effort involved in approaching a food patch (Ogura & Matsushima, 2011; Ogura *et al.*, 2015), and found that chicks paid extra effort, even when it was not accompanied by extra food gain.

However, it would be premature to argue that the lateral Arco is specifically involved in social facilitation. First, no distinctive cytoarchitectonic differences have been found among the presumed regions of the Arco, despite the results of the present tracing study. Second, electrolytic lesions cannot unequivocally distinguish the contribution of the lateral Arco. Most probably, the lesions placed in the dorsal and intermediate Arco also damaged the efferent fibers issued from the lateral Arco, as has been reported previously (Atoji *et al.* 2006, Hanics *et al.* 2016). Additionally, lesions in the lateral Arco appeared to also affect the medial Arco regions to some degree. Future research should include localized pharmacological and/or molecular manipulations of the efferent Arco pathways.

An alternative interpretation of the present lesion study is that the Arco contributes to the visual perception of conspecifics, rather than foraging effort. This idea is compatible with the recent finding that visual exposure to alive conspecific activated amygdaloid nuclei in visually naïve chicks (Mayer, Rosa-Salva *et al.* 2017). If it is the case, we may assume that the observed suppression in facilitation is due to the incapability of the lesioned chick to discriminate companion chicks. However, this idea is not compatible with the finding that Arco lesions failed to suppress the shortened peck latency (Fig. 4C). Although

the Arco might be involved in social perception, it is not critically required in all aspects of inter-individual interactions.

The Arco lesion might have directly suppressed locomotor activity. The Arco is a heterogeneous structure comprising a somato-motor region and a limbic region (Wild et al., 1985; Veenman et al., 1995). The somato-motor region, including the intermediate and dorsal Arco, is thought to be analogous to the mammalian premotor/motor cortex (Shanahan et al., 2013). This region is also involved in sensori-motor control (Zeier, 1971), specifically oro-facial control (Wild et al., 1985). In pigeons, lesioning the intermediate Arco caused deficient feeding, and lesioned subjects could not hold grains with their beaks (Zeier, 1971). The increase in κ index found in the present study is in a good concert with these report.

More specifically, we may argue that the Arco lesion reduced the maximum limit of the running speed, making social facilitation less distinct. The reduced running in the single phase (Fig. 3) might thus be linked to the diverse projections of the medial Arco regions to midbrain areas including the tegmentum and TeO (Figs. 6 and 7). Indeed, reduced action in the descending reticulo-spinal pathway (see review by Grillner, 2006) may have slowed the running speed of the chicks after the Arco lesion. However, the lateral Arco did not project to the LSt (category η_1) or the tectum/tegmentum (η_3).

The Nido is not considered to be involved in social facilitation, despite its dense projections to the MSt and LSt (category η_1). Previous lesion (Diekamp, Gagliardo, *et al.*, 2002) and single-unit recording studies (Diekamp, Kalt, *et al.*, 2002; Veit *et al.*, 2014) in birds suggest that the caudo-lateral part of the Nido (NCL) is critical for working memory,

similar to the prefrontal cortex (PFC) in mammals. However, our behavioral paradigm in the present study does not require working memory. Furthermore, the lesioned Nido area in chicks may be dissimilar from the NCL region studied in pigeons. It may be necessary to further characterize more lateral and caudal Nido areas.

The TnA lesion was also not effective, although previous studies have suggested that the TnA has amygdaloid features in birds (Cheng *et al.*, 1999; Absil *et al.*, 2002). Starlings with TnA lesions showed reduced “social facilitation” of foraging behavior in terms of behavioral synchronization (Cheng *et al.*, 1999), although these findings were not replicated in the present study. We should also notice that it remains controversial about the location and identification of nucleus taeniae in the avian brain (Puelles *et al.*, 2007, Hanics *et al.*, 2016).

4.1.2 Investment of foraging effort

Ogura *et al.* (2015) reported a dissociation of the neural substrates involved in foraging effort and social facilitation of foraging. Specifically, lesioning the MSt suppressed foraging effort in the single phase, but did not lead to impaired social facilitation. The present tracing results show that the lateral Arco projects to the NAc and MSt, whereas the intermediate and dorsal Arco project mainly to the MSt (Figs. 6 and 7). A recent neuroanatomical study also revealed amygdalo-fugal terminals in the MSt/NAc with the morphological features of excitatory synapses (Hanics *et al.*, 2012). Taken together, these data indicate that Arco-MSt/NAc pathways are involved in determining foraging effort.

4.1.3 Matching behavior

The patch use ratio matched the biased food delivery rate between the two patches (Figs. 3 and 4Ab) (Q. Xin, Y. Ogura and T. Matsushima, unpublished), and was unchanged by the lesions. Thus, the mechanisms that evaluate food patches, and proportionately allocate stay time, appear to be distinct from those involved in social facilitation. A similar dissociation was found by Aoki *et al.* (2006a), who found that handling-cost aversion occurred without impairments to amount-based choices. To our knowledge, however, the neural mechanisms of matching have not been addressed in birds. In mammals, neurons in the parietal cortex represent the relative value of behaviors in a dynamic foraging environment (Sugrue *et al.*, 2004). However, the avian counterpart to the mammalian parietal cortex has not been established (see below for further comparative arguments). In primates, activity of phasically active neurons (PANs) in the primate striatum co-vary with the action-value, suggesting that PANs participate in the encoding of action values (Lau & Glimcher, 2008). The striatal pathways described above may also be involved in matching in chicks.

4.1.4 Operant peck latency

The Arco lesion did not change the peck latency (Fig. 4C). The response latency is often supposed to represent the subjective value of a predicted food reward (Lauwereyns & Wisniewski, 2006). In chicks, however, experimental manipulation of handling-cost did not affect the peck latency, indicating that the latency does not represent the accompanying cost (Aoki *et al.*, 2006a). Most probably, the peck latency is socially facilitated by distinct mechanisms that are responsible for the foraging effort. In rats, dopamine depletion in the caudate nucleus has been associated with an increased response latency (Amalric & Koob,

1987). In future research, similar depletion effects in the striatum (LSt, MSt, and NAc) could be examined with respect to peck latency in chicks.

4.2 Candidate Arco efferents involved in social facilitation

Assuming that the lateral Arco is selectively involved in social facilitation, the extended amygdala (rEA and cEA) may contribute to this behavior, as they both receive dense projections from the Arco. Currently, these regions are both recognized as part of the amygdaloid complex in the avian brain (Reiner *et al.*, 2004). Further research involving localized lesions to these amygdaloid nuclei may help to clarify this issue.

A projection from the lateral Arco to the NAc might be another candidate efferent. In rats, pharmacological manipulation of dopaminergic activity in the NAc biases behavior away from costly actions without disrupting discrimination between rewards of different magnitudes (Salamone *et al.*, 1994). In chicks, electrolytic lesioning of the MSt/NAc suppressed foraging efforts, but dopamine depletion in these regions had no effect (Ogura *et al.*, 2015). A projection to the Sept may also be a candidate. In primates, a class of Sept neurons code reward uncertainty; these may process risk-cost and related emotional responses (Monosov & Hikosaka, 2013). Distinct projections from the lateral Arco to the Sept (Figs. 5Bb, 6 and 7) might play a critical role in the social foraging.

The involvement of a descending projection to midbrain tegmental areas is also an important consideration. As lesioning the tegmentum around the SN suppresses social facilitation (Ogura *et al.*, 2015), damage to the major descending fiber bundle (OM; Zeier & Karten, 1971; Davies *et al.*, 1997) may be a critical factor. However, the lateral Arco

only sparsely projects to the tegmental nuclei. The specific involvement of the lateral Arco-tegmental pathway is thus questionable. As we observed only a partial loss of social facilitation after a considerable part of the Arco was lesioned, the Arco could be one of a number of involved substrates, each of which partially contributes to social facilitation.

4.3 Is the avian Arco analogous to the prefrontal cortex or amygdala in mammals?

Based on the present lesion data, we may argue that the Arco plays multiple functions in a manner analogous to the mammalian prefrontal cortex, particularly the ACC. Lesioning the mammalian ACC caused an effort-cost aversion, biasing behavior away from a costly action (Walton *et al.*, 2003, 2009; Schweimer *et al.*, 2005; Rudebeck *et al.*, 2006). As with the chicken Arco (Phillips & Youngren, 1986), a taming effect appeared after an ACC lesion in humans, specifically personality changes, such as lack of distress (Tow & Whitty, 1953; Cohen *et al.*, 2001). The avian Arco is involved in sensori-motor control (Zeier, 1971; Knudsen & Knudsen, 1996b), and this may hold true for the mammalian ACC. For instance, bilateral lesions of the rostral ACC is reported to cause oculomotor deficits, i.e., deficits in central gaze fixation (Paus *et al.*, 1991).

This issue should also be discussed in anatomical terms. As with the lateral Arco, the ACC/mPFC has connections with the hippocampal-parahippocampal region (Carmichael & Price, 1995; Chiba *et al.*, 2001; Kondo *et al.*, 2005; Mohedano-Moriano *et al.*, 2007; Passingham & Wise, 2012). In addition, the projections of lateral Arco to other limbic regions, such as the hypothalamus, dorsal thalamus, extended amygdala (rEA and cEA), MSt, and NAc should not be ignored (Figs. 6 and 7). In the ACC, similarly strong

connections appear with limbic regions such as the hypothalamus (Ongür *et al.*, 1998), amygdala (Morecraft *et al.*, 2007), and striatum (Haber *et al.*, 1995).

Such functional and anatomical considerations enable the alternative idea of comparing the lateral Arco to the mammalian amygdala, particularly the BLA. Inactivation of the BLA also causes effort-cost aversion (Floresco & Ghods-Sharifi, 2007). Lesions of the chicken Arco (Phillips & Youngren, 1986; Lowndes & Davies, 1995; Saint-Dizier *et al.*, 2009) and its electrical stimulation (Phillips & Youngren, 1971) have suggested that Arco is critical for fear- or anxiety-related behavior, similarly to the mammalian BLA in terms of emotion processing (Vazdarjanova *et al.*, 2001; Tye *et al.*, 2011).

Anatomically, as pointed out by Hanics *et al.* (Hanics *et al.*, 2016), the efferent projections from the lateral Arco to the NAc and BSTl suggest a correspondence to the mammalian BLA. Similarly, the BLA has projections to the hippocampal formation (Saunders *et al.*, 1988; Pitkänen *et al.*, 2000) and Sept (Calderazzo *et al.*, 1996) in mammals. Based on common neurochemical characterization, Martinez-Garcia *et al.* (Martínez-García *et al.*, 2009) hypothesized that the Arco complex is homologous with the mammalian BLA.

Convergent evolution of “prefrontal” and “amygdala” functions between birds and mammals, together with their functional multiplicity and anatomicalies, should be carefully examined in future.

Acknowledgements

We are grateful to colleagues in the sub-department of Behavioral Neurobiology in the Department of Biological Sciences, Hokkaido University, particularly for Dr. Masayo Soma and Dr. Kazuhiro Wada. It is also acknowledged that the apparatus used in this study was manufactured with the technical aid of the skilled engineer, Mr. Yasutaka Sasaki in the machine laboratory of Hokkaido University. T.M. was funded by the Japan Society for the Promotion of Science (JSPS, KAKENHI; Grant-in-Aid for Scientific Research (B) #25291071 and Grant-in-Aid for Challenging Exploratory Research #26650114). Y.O. was also funded by JSPS (Grant-in-Aid for JSPS Fellows, #26-8054). Q.X. was financially supported by a Japanese Ministry of Education, Culture, Sports, Science, and Technology (MEXT) scholarship for postgraduate study. Q.X., Y.O. and T.M. designed the experiments. Q.X., Y.O. and L.U. conducted the experiments and analysed data. Q.X., Y.O. and T.M. wrote the manuscript. All authors have approved the manuscript for publication. We declare no competing interests.

Abbreviations

Abbreviation	English name
ACC	anterior cingulate cortex
AMH	anterior medial hypothalamic nucleus
Apir	amygdalopiriform area
Arco	arcopallium
BLA	basolateral amygdala
CA	anterior commissure
DIP	thalamic posterior dorsointermediate nucleus
DLA	thalamic anterior dorsolateral nucleus
DLP	thalamic posterior dorsolateral nucleus
DMA	thalamic anterior dorsomedial nucleus
DMP	thalamic posterior dorsomedial nucleus
EA	extended amygdala
GCt	midbrain central gray
GP	globus pallidus
HbL	lateral habenula
HbM	medial habenula
Hp	hippocampus
ICo	intercollicular nucleus
IH	inferioris hypothalamic nucleus

IMM	intermediate medial mesopallium
LSt	lateral striatum
Lhy	lateral hypothalamus
LoC	locus coeruleus
MLd	dorsal lateral mesencephalic nucleus
MSt	medial striatum
NAc	nucleus accumbens
NCL	nidopallium caudolaterale
NIII	oculomotor (third) nerve
Nido	nidopallium
OFC	orbitofrontal cortex
OM	occipito-mesencephalic tract
OTu	olfactory tubercle
PHN	hypothalamic periventricular nucleus
PMI	thalamic nucleus paramedianus internus
POM	medial preoptic nucleus
PVN	paraventricular nucleus
Pap	papillioformis nucleus
PoA	posterior pallial amygdala
Ru	nucleus ruber
SC	nucleus subceruleus

SCE	stratum cellulare externum
SN	substantia nigra
SPL	lateral spiriformis nucleus
SPM	medial spiriformis nucleus
Sept	septum
TeO	optic tectum
TnA	nucleus taeniae of the amygdala
VL	lateral ventricle
VMN	hypothalamic ventromedial nucleus
VP	ventral pallidum
VT	third ventricle
VTA	ventral tegmental area
cBSTl	caudal bed nucleus of the stria terminalis
rBSTl	rostral bed nucleus of the stria terminalis

- Absil, P., Braquenier, J.B., Balthazart, J., & Ball, G.F. (2002) Effects of lesions of nucleus taeniae on appetitive and consummatory aspects of male sexual behavior in Japanese quail. *Brain Behav Evol*, **60**, 13–35.
- Amalric, M. & Koob, G.F. (1987) Depletion of dopamine in the caudate nucleus but not in nucleus accumbens impairs reaction-time performance in rats. *J. Neurosci.*, **7**, 2129–2134.
- Amita, H., Kawamori, A., & Matsushima, T. (2010) Social influences of competition on impulsive choices in domestic chicks. *Biol. Lett.*, **6**, 183–186.
- Amita, H. & Matsushima, T. (2011) Instantaneous and cumulative influences of competition on impulsive choices in domestic chicks. *Front. Neurosci.*, **5**, 1–8.
- Amita, H. & Matsushima, T. (2014) Competitor suppresses neuronal representation of food reward in the nucleus accumbens/medial striatum of domestic chicks. *Behav. Brain Res.*, **268**, 139–149.
- Aoki, N., Izawa, E.-I., Yanagihara, S., & Matsushima, T. (2003) Neural correlates of memorized associations and cued movements in archistriatum of the domestic chick. *Eur. J. Neurosci.*, **17**, 1935–1946.
- Aoki, N., Csillag, A., & Matsushima, T. (2006a) Localized lesions of arcopallium intermedium of the lateral forebrain caused a handling-cost aversion in the domestic chick performing a binary choice task. *Eur. J. Neurosci.*, **24**, 2314–2326.

- Aoki, N., Suzuki, R., Izawa, E.-I.I., Csillag, A., & Matsushima, T. (2006b) Localized lesions of ventral striatum, but not arcopallium, enhanced impulsiveness in choices based on anticipated spatial proximity of food rewards in domestic chicks. *Behav. Brain Res.*, **168**, 1–12.
- Apps, M.A.J. & Ramnani, N. (2014) The Anterior Cingulate Gyrus Signals the Net Value of Others' Rewards. *J. Neurosci.*, **34**, 6190–6200.
- Atoji, Y., Saito, S., & Wild, M. (2006) Fiber connections of the compact division of the posterior pallial amygdala and lateral part of the bed nucleus of the stria terminalis in the pigeon (*Columba livia*). *J. Comp. Neurol.*, **499**, 161–182.
- Calderazzo, L., Cavalheiro, E.A., Macchi, G., Molinari, M., & Bentivoglio, M. (1996) Branched connections to the septum and to the entorhinal cortex from the hippocampus, amygdala, and diencephalon in the rat. *Brain Res. Bull.*, **40**, 245–251.
- Carmichael, S.T. & Price, J.L. (1995) Limbic connections of the orbital and medial prefrontal cortex in macaque monkeys. *J. Comp. Neurol.*, **363**, 615–641.
- Charnov, E.L. (1976) Optimal foraging, the marginal value theorem. *Theor. Popul. Biol.*, **9**, 129–136.
- Cheng, M.F., Chaiken, M., Zuo, M., & Miller, H. (1999) Nucleus taenia of the amygdala of birds: Anatomical and functional studies in ring doves (*Streptopelia risoria*) and European starlings (*Sturnus vulgaris*). *Brain. Behav. Evol.*, **53**, 243–270.

- Chiba, T., Kayahara, T., & Nakano, K. (2001) Efferent projections of infralimbic and prelimbic areas of the medial prefrontal cortex in the Japanese monkey , *Macaca fuscata*. *Brain Res*, **888**, 83–101.
- Cohen, R.A., Paul, R., Zawacki, T.M., Moser, D.J., Sweet, L., & Wilkinson, H. (2001) Emotional and Personality Changes Following Cingulotomy **I**, 38–50.
- Cousins, M.S., Atherton, a., Turner, L., & Salamone, J.D. (1996) Nucleus accumbens dopamine depletions alter relative response allocation in a T-maze cost/benefit task. *Behav. Brain Res.*, **74**, 189–197.
- Davies, D.C., Csillag, A., Székely, A.D., & Kabai, P. (1997) Efferent connections of the domestic chick archistriatum: A phaseolus lectin anterograde tracing study. *J. Comp. Neurol.*, **389**, 679–693.
- Diekamp, B., Gagliardo, A., & Güntürkün, O. (2002) Nonspatial and subdivision-specific working memory deficits after selective lesions of the avian prefrontal cortex. *J. Neurosci.*, **22**, 9573–9580.
- Diekamp, B., Kalt, T., & Güntürkün, O. (2002) Working memory neurons in pigeons. *J. Neurosci.*, **22**, RC210.
- Everitt, B.S., Landau, S., Leese, M., & Stahl, D. (2011) *Cluster Analysis*, 5th edn. John Wiley & Sons, Ltd., New York.
- Floresco, S.B. & Ghods-Sharifi, S. (2007) Amygdala-prefrontal cortical circuitry regulates effort-based decision making. *Cereb. Cortex*, **17**, 251–260.

- Floresco, S.B., St Onge, J.R., Ghods-Sharifi, S., & Winstanley, C. (2008) Cortico-limbic-striatal circuits subserving different forms of cost-benefit decision making. *Cogn. Affect. Behav. Neurosci.*, **8**, 375–389.
- Giraldeau, L.A. & Dubois, F. (2008) Social foraging and the study of exploitative behavior. *Advances in the Study of Behavior.*, **38**, pp. 59–104.
- Grillner, S. (2006) Biological Pattern Generation: The Cellular and Computational Logic of Networks in Motion. *Neuron*, **52**, 751–766.
- Haber, S.N., Kunishio, K., Mizobuchi, M., & Lynd-Balta, E. (1995) The orbital and medial prefrontal circuit through the primate basal ganglia. *J. Neurosci.*, **15**, 4851–4867.
- Hanics, J., Bálint, E., Milanovich, D., Zachar, G. Ádám, Á., & Csillag, A. (2012) Amygdalofugal axon terminals immunoreactive for L-aspartate or L-glutamate in the nucleus accumbens of rats and domestic chickens: a comparative electron microscopic immunocytochemical study combined with anterograde pathway tracing. *Cell Tissue Res.*, **350**, 409–423.
- Hanics, J., Teleki, G., Alpár, A., Székely, A.D., & Csillag, A. (2016) Multiple amygdaloid divisions of arcopallium send convergent projections to the nucleus accumbens and neighboring subpallial amygdala regions in the domestic chicken: a selective pathway tracing and reconstruction study. *Brain Struct. Funct.*, **221**, 1–15.
- Hauber, W. & Sommer, S. (2009) Prefrontostriatal circuitry regulates effort-related decision making. *Cereb. Cortex*, **19**, 2240–2247.

- Izawa, E.I., Zachar, G., Yanagihara, S., & Matsushima, T. (2003) Localized lesion of caudal part of lobus parolfactorius caused impulsive choice in the domestic chick: Evolutionarily conserved function of ventral striatum. *J. Neurosci.*, **23**, 1894–1902.
- Knudsen, E.I. & Knudsen, P.F. (1996a) Disruption of auditory spatial working memory by inactivation of the forebrain archistriatum in barn owls. *Nature*, **383**, 428–431.
- Knudsen, E.I. & Knudsen, P.F. (1996b) Contribution of the forebrain archistriatal gaze fields to auditory orienting behavior in the barn owl. *Exp. brain Res.*, **108**, 23–32.
- Kondo, H., Saleem, K.S., & Price, J.L. (2005) Differential connections of the perirhinal and parahippocampal cortex with the orbital and medial prefrontal networks in macaque monkeys. *J. Comp. Neurol.*, **493**, 479–509.
- Kuenzel, W.J. & Masson, M. (1988) *A Stereotaxic Atlas of the Brain of the Chick (Gallus Domesticus)*. Johns Hopkins University Press, Baltimore.
- Lau, B. & Glimcher, P.W. (2008) Value representations in the primate striatum during matching behavior. *Neuron*, **58**, 451–463.
- Lauwereyns, J. & Wisniewski, R.G. (2006) A reaction-time paradigm to measure reward-oriented bias in rats. *J. Exp. Psychol. Anim. Behav. Process.*, **32**, 467–473.
- Lissek, S. & Güntürkün, O. (2003) Dissociation of extinction and behavioral disinhibition: The role of NMDA receptors in the pigeon associative forebrain during extinction. *J. Neurosci.*, **23**, 8119–8124.

- Lowndes, M. & Davies, D.C. (1995) The effect of archistriatal lesions on “open field” and fear/avoidance behaviour in the domestic chick. *Behav. Brain Res.*, **72**, 25–32.
- Martínez-García, F., Novejarque, A., & Lanuza, E. (2009) The evolution of the amygdala in vertebrates. In Kaas J. H. (Eds), *Evolutionary Neuroscience*. Academic Press, Oxford, pp. 313-392.
- Matsushima, T., Kawamori, A., & Bem-Sojka T. (2008) Neuro-economics in chicks: foraging choices based on delay, cost and risk. *Brain Res. Bull.*, **76**, 245-252.
- Mayer, U., Rosa-Salva, O., and Vallortigara, G. (2017) First exposure to an alive conspecific activates septal and amygdaloid nuclei in visually-naïve domestic chicks (*Gallus gallus*). *Behav. Brain Res.*, **317**, 71-81.
- Mohedano-Moriano, A., Pro-Sistiaga, P., Arroyo-Jimenez, M.M., Artacho-Pérula, E., Insausti, A.M., Marcos, P., Cebada-Sánchez, S., Martínez-Ruiz, J., Muñoz, M., Blaizot, X., Martinez-Marcos, A., Amaral, D.G., & Insausti, R. (2007) Topographical and laminar distribution of cortical input to the monkey entorhinal cortex. *J. Anat.*, **211**, 250–260.
- Monosov, I.E. & Hikosaka, O. (2013) Selective and graded coding of reward uncertainty by neurons in the primate anterodorsal septal region. *Nat. Neurosci.*, **16**, 756–762.
- Montagnese, C.M., Mezey, S.E., & Csillag, A. (2003) Efferent connections of the dorsomedial thalamic nuclei of the domestic chick (*Gallus domesticus*). *J. Comp. Neurol.*, **459**, 301–326.

- Morecraft, R.J., Mcneal, D.W., Stilwell-Morecraft, K.S., Gedney, M., Ge, J., Schroeder, C.M., & Van Hoesen, G.W. (2007) Amygdala interconnections with the cingulate motor cortex in the rhesus monkey. *J. Comp. Neurol.*, **500**, 134–165.
- Nottebohm, F., Stokes, T.M., & Leonard, C.M. (1976) Central control of song in the canary, *Serinus canarius*. *J. Comp. Neurol.*, **165**, 457–486.
- Ogura, Y., Izumi, T., Yoshioka, M., & Matsushima, T. (2015) Dissociation of the neural substrates of foraging effort and its social facilitation in the domestic chick. *Behav. Brain Res.*, **294**, 162–176.
- Ogura, Y. & Matsushima, T. (2011) Social facilitation revisited: Increase in foraging efforts and synchronization of running in domestic chicks. *Front. Neurosci.*, **5**, 1–12.
- Ongür, D., An, X., & Price, J.L. (1998) Prefrontal cortical projections to the hypothalamus in macaque monkeys. *J. Comp. Neurol.*, **401**, 480–505.
- Passingham, R.E. & Wise, S.P. (2012) *The Neurobiology of the Prefrontal Cortex: Anatomy, Evolution, and the Origin of Insight*. Oxford University Press. Oxford.
- Paus, T., Kalina, M., Patocková, L., Angerová, Y., Cerný, R., Mecir, P., Bauer, J., & Krabec, P. (1991) Medial vs lateral frontal lobe lesions and differential impairment of central-gaze fixation maintenance in man. *Brain*, **114**, 2051–2067.
- Phillips, R.E. (1964) “Wildness” in the Mallard duck: Effects of brain lesions and stimulation on “escape behavior” and reproduction. *J. Comp. Neurol.*, **122**, 139–155.

Phillips, R.E. & Youngren, O.M. (1971) Brain stimulation and species-typical behaviour:

Activities evoked by electrical stimulation of the brains of chickens (*Gallus gallus*).

Anim. Behav., **19**, 757–779.

Phillips, R.E. & Youngren, O.M. (1986) Unilateral kainic acid lesions reveal dominance of

right archistriatum in avian fear behavior. *Brain Res.*, **377**, 216–220.

Phillips, R.E., Youngren, O.M., & Peek, F.W. (1972) Repetitive vocalizations evoked by

local electrical stimulation of avian brains I. Awake chickens (*Gallus gallus*). *Anim.*

Behav., **20**, 689–705.

Pitkänen, A., Pikkarainen, M., Nurminen, N., & Ylinen, A. (2000) Reciprocal connections

between the amygdala and the hippocampal formation, perirhinal cortex, and

postrhinal cortex in rat. A review. *Ann. N. Y. Acad. Sci.*, **911**, 369–391.

Puelles, L., Martinez-de-la-Torre, S., Paxinos, G., Watson, C.R.R., & Martinez, M. (2007)

The Chick Brain in Stereotaxic Coordinates: An Atlas Featuring Neuromeric

Subdivisions and Mammalian Homologies. Academic Press, London.

Reiner, A., Perkel, D.J., Bruce, L.L., Butler, A.B., Csillag, A., Kuenzel, W., Medina, L.,

Paxinos, G., Shimizu, T., Striedter, G., Wild, M., Ball, G.F., Durand, S., Gütürkün, O.,

Lee, D.W., Mello, C. V., Powers, A., White, S.A., Hough, G., Kubikova, L., Smulders,

T. V., Wada, K., Dugas-Ford, J., Husband, S., Yamamoto, K., Yu, J., Siang, C., &

Jarvis, E.D. (2004) Revised nomenclature for avian telencephalon and some related

brainstem nuclei. *J. Comp. Neurol.*, **473**, 377–414.

Rudebeck, P.H., Walton, M.E., Smyth, A.N., Bannerman, D.M., & Rushworth, M.F.S.

(2006) Separate neural pathways process different decision costs. *Nat. Neurosci.*, **9**, 1161–1168.

Saint-Dizier, H., Constantin, P., Davies, D.C., Leterrier, C., Lévy, F., & Richard, S. (2009)

Subdivisions of the arcopallium/posterior pallial amygdala complex are differentially involved in the control of fear behaviour in the Japanese quail. *Brain Res. Bull.*, **79**, 288–295.

Salamone, J.D., Cousins, M.S., & Bucher, S. (1994) Anhedonia or anergia? Effects of

haloperidol and nucleus accumbens dopamine depletion on instrumental response selection in a T-maze cost/benefit procedure. *Behav. Brain Res.*, **65**, 221–229.

Saunders, R.C., Rosene, D.L., & Van Hoesen, G.W. (1988) Comparison of the efferents of

the amygdala and the hippocampal formation in the rhesus monkey: II. Reciprocal and non-reciprocal connections. *J. Comp. Neurol.*, **271**, 185–207.

Schweimer, J., Saft, S., & Hauber, W. (2005) Involvement of catecholamine

neurotransmission in the rat anterior cingulate in effort-related decision making. *Behav. Neurosci.*, **119**, 1687–1692.

Shanahan, M., Bingman, V.P., Shimizu, T., Wild, M., & Güntürkün, O. (2013) Large-scale

network organization in the avian forebrain: a connectivity matrix and theoretical analysis. *Front. Comput. Neurosci.*, **7**, 89.

- Singer, T., Seymour, B., Doherty, J.P.O., Stephan, K.E., Dolan, R.J., & Frith, C.D. (2006) Empathic neural responses are modulated by the perceived fairness of others. *Nature*, **439**, 466–469.
- Stephens, D.W. & Krebs, J.R. (1986) *Foraging Theory*. Princeton University Press, Princeton.
- Sugrue, L.P., Corrado, G.S., & Newsome, W.T. (2004) Matching behavior and the representation of value in the parietal cortex. *Science*, **304**, 1782–1787.
- Tow, P.M. & Whitty, C.W. (1953) Personality changes after operations on the cingulate gyrus in man. *J. Neurol. Neurosurg. Psychiatry*, **16**, 186–193.
- Tye, K.M., Prakash, R., Kim, S.-Y., Fenno, L.E., Grosenick, L., Zarabi, H., Thompson, K.R., Gradinaru, V., Ramakrishnan, C., & Deisseroth, K. (2011) Amygdala circuitry mediating reversible and bidirectional control of anxiety. *Nature*, **471**, 358–362.
- Vazdarjanova, A., Cahill, L., & McGaugh, J.L. (2001) Disrupting basolateral amygdala function impairs unconditioned freezing and avoidance in rats. *Eur. J. Neurosci.*, **14**, 709–718.
- Veenman, C.L.E.O., Wild, J.M., & Reiner, A. (1995) Organization of the avian “corticostriatal” projection system: A retrograde and anterograde pathway tracing study in pigeons. *J. Comp. Neurol.*, **354**, 87–126.
- Veit, L., Hartmann, K., & Nieder, A. (2014) Neuronal Correlates of Visual Working Memory in the Corvid Endbrain. *J. Neurosci.*, **34**, 7778–7786.

- Wallis, J.D. & Rushworth, M.F.S. (2014) Integrating benefits and costs in decision making. In Glimcher, P.W. & Fehr, E., *Neuroeconomics*, 2nd edn. Academic Press, London, pp. 411–433.
- Walton, M.E., Bannerman, D.M., Alterescu, K., & Rushworth, M.F.S. (2003) Functional specialization within medial frontal cortex of the anterior cingulate for evaluating effort-related decisions. *J. Neurosci.*, **23**, 6475–6479.
- Walton, M.E., Bannerman, D.M., & Rushworth, M.F.S. (2002) The role of rat medial frontal cortex in effort-based decision making. *J. Neurosci.*, **22**, 10996–11003.
- Walton, M.E., Groves, J., Jennings, K. a., Croxson, P.L., Sharp, T., Rushworth, M.F.S., & Bannerman, D.M. (2009) Comparing the role of the anterior cingulate cortex and 6-hydroxydopamine nucleus accumbens lesions on operant effort-based decision making. *Eur. J. Neurosci.*, **29**, 1678–1691.
- Walton, M.E., Kennerley, S.W., Bannerman, D.M., Phillips, P.E.M., & Rushworth, M.F.S. (2006) Weighing up the benefits of work: Behavioral and neural analyses of effort-related decision making. *Neural Networks*, **19**, 1302–1314.
- Wild, J.M., Arends, J.J.A., & Zeigler, H.P. (1985) Telencephalic connections of the trigeminal system in the pigeon (*Columba livia*): A trigeminal sensorimotor circuit. *J. Comp. Neurol.*, **234**, 441–464.

- Yamamoto, K., Sun, Z., Wang, H.B., & Reiner, A. (2005) Subpallial amygdala and nucleus taeniae in birds resemble extended amygdala and medial amygdala in mammals in their expression of markers of regional identity. *Brain Res. Bull.*, **66**, 341-347.
- Zajonc, R.B. (1965) Social facilitation. *Science*, **149**, 269-274.
- Zeier, H. (1971) Archistriatal lesions and response inhibition in the pigeon. *Brain Res.*, **31**, 327–339.
- Zeier, H. & Karten, H.J. (1971) The archistriatum of the pigeon: Organization of afferent and efferent connections. *Brain Res.*, **31**, 313–326.

Figure captions

Fig. 1. Experimental apparatus and behavioral procedures for foraging behavior (A, B) and operant responses (C, D).

(A) Apparatus and procedure of Experiments 1 and 2. An I-shaped maze was separated by a transparent Plexiglas partition to two lanes, each equipped with two food trays on both sides. One test block comprised three consecutive phases: single #1, paired, and single #2. (B) Examples of the food delivery schedules with arrowheads indicating the timing of delivery. In Experiment 1, the delivery was asymmetric as one patch followed the VI10 schedule (variable interval with the mean = 10 sec), and the other the VI30 (with the mean = 30 sec). Colors were randomly assigned among individuals. In Experiment 2, the delivery was symmetric and both patches followed the VI15 schedule. (C) An operant box used in Experiment 3. Visual stimuli (red or green rectangles) were displayed on a LCD monitor, and an infra-red (IR) touch sensor detected chick pecking behavior. Chicks were tested either individually or in groups of three. In the triplet phase, the subject was separated from the two companions by transparent Plexiglas. (D) Trial procedure in Experiment 3. Color cues were presented for 2 sec. When the subject pecked at the rewarding cue, 6 grains of millet were delivered to the central food tray. No food followed when the subject pecked the non-rewarding cue.

Fig. 2. Histological reconstruction of lesions. (A) Areas with damaged tissue are superimposed onto frontal sections. The levels of the sections (A6.4 to A8.2) follow the chick atlas by Kuenzel & Masson (1988). n denotes the number of chicks in each group.

(B) Examples of Nissl-stained sections from the sham (a), Arco lesion (b), and TnA lesion (c) groups. See Abbreviations.

Fig. 3. Representative running trajectories of chicks with sham (dark grey), Arco (green), lateral Arco (orange), and Nido (blue) lesions. Trajectories from single subjects in the single (left) and paired (right) phases are shown. The trajectory of the naïve companion is superimposed in light grey. The y-axis indicates the position along the I-shaped maze (red: top, yellow: bottom), and the x-axis represents time. Horizontal dashed lines indicate the midpoint of the maze. Arrowheads denote the timing of food delivery. The delivery was biased, as the yellow patch followed the VI10 schedule (bottom) while the red patch followed the VI30 schedule (top).

Fig. 4. Behavioral effects of lesions. (A) Effect of the Arco (green), lateral Arco (orange), and Nido (blue) lesions compared with the sham control (grey) in Experiment 1. (a) Averaged running distance plotted against the sessions, which were each composed of three consecutive phases: single #1, paired, and single #2. Mean $\pm$ SEM are shown in this and the following graphs. (b) Patch use ratio at the VI10 feeder plotted against the sessions. Dashed horizontal lines indicate the level of 0.75. (c-e) Post-lesion data (4 sessions) compared among the four groups of chicks. (c) Average running distance in $\overline{\text{single}}$ (mean of single #1 and single #2) and paired trials. (d) Averaged synchrony index during the paired phase. (e) Patch use ratio in $\overline{\text{single}}$ (mean of single #1 and single #2) and paired trials. (B) Effect of the TnA lesion (pink) compared with the sham control (grey) in Experiment 2. (a) Running

distance, (b) synchrony index, and (c) patch use ratio. Dashed horizontal lines in (c) indicate the level of 0.5. (C) We compared the effect of the Arco lesion (green) with the sham control (grey) in Experiment 3. Operant peck latency (ms) in rewarding trials. See text for statistical tests.

Fig. 5. Anterograde tract tracing using biotinylated dextran amine (BDA). (A)

Examples of BDA injection sites in the lateral Arco (a), dorsal Arco (b), and Nido (c).

Levels of the frontal sections follow the atlas by Kuenzel & Masson (1988). (B)

Anterogradely labelled terminals in the hippocampus (Hp, a), septum (Sept, b), and nucleus accumbens (NAc, c) after injection in the lateral Arco. (C) Anterogradely labelled terminals in the intermediate medial mesopallium (IMM, a), caudal extended amygdala (cEA, b), ventral tegmental area (VTA, c), medial spiriformis nucleus (SPM, d), intercollicular nucleus (ICo, e), and optic tectum (TeO, f) after injection in the dorsal Arco. We also found retrogradely labelled cell bodies in the Nido (g). Dashed lines indicate the boundaries of the regions. (D) Anterogradely labelled terminals in the lateral striatum (LSt) after injection in the Nido. See Abbreviations.

Fig. 6. Hierarchical clustering of efferent projections. Rows indicate the recipient brain regions, and columns represent the subjects. We semi-quantitatively determined five grades of terminal density: abundant, moderate, low, sparse, and absent. Subjects were grouped into three main clusters; α , β , and γ , and cluster γ was further divided into three sub-clusters: γ_1 , γ_2 , and γ_3 . The recipient brain regions were also grouped into four clusters δ , ϵ , ζ , and η ,

and cluster η was further divided into three sub-clusters, η_1 , η_2 , and η_3 ; where n denotes the number of chicks. At the bottom, retrogradely labelled cell density in the Nido is indicated as +++, ++, and $\pm$. See the text for details and see Abbreviations. The BDA injection sites are illustrated below. Different colors indicate different clusters of subject chicks and numbers in parenthesis (1~15) denote the injection site in individual chicks.

Fig. 7. Major efferent projections from sub-regions in the Arco. Recipient areas in the telencephalon (A8.8), diencephalon (A7.0), and midbrain (A3.4) are schematically shown. Projections from the lateral Arco (orange), medial regions of the Arco (green), and the Nido (blue) are labelled using the same colors in Fig. 6. Areas with abundant and moderate efferent terminals are shown on the right columns. Refer to the abbreviation table.

A Experiment 1 & 2

B Experiment 1 & 2

a Experiment 1 Arco / Nido lesion

or

b Experiment 2 TnA lesion

C Experiment 3

D Experiment 3

A Injection site

Lateral Arco

Dorsal Arco

Nido

B Lateral Arco

C Dorsal Arco

D Nido

XIN Figure 6