



Title	THE HYDROSTATIC STOKES SEMIGROUP AND WELL-POSEDNESS OF THE PRIMITIVE EQUATIONS ON SPACES OF BOUNDED FUNCTIONS
Author(s)	Giga, Yoshikazu; Gries, Mathis; Hieber, Matthias; Hussein, Amru; Kashiwabara, Takahito
Citation	Hokkaido University Preprint Series in Mathematics, 1109, 1-30
Issue Date	2018-02-08
DOI	10.14943/81805
Doc URL	<a href="http://hdl.handle.net/2115/68294">http://hdl.handle.net/2115/68294</a>
Type	bulletin (article)
File Information	HydrostaticStokesSemigroup.pdf



[Instructions for use](#)

# THE HYDROSTATIC STOKES SEMIGROUP AND WELL-POSEDNESS OF THE PRIMITIVE EQUATIONS ON SPACES OF BOUNDED FUNCTIONS

YOSHIKAZU GIGA, MATHIS GRIES, MATTHIAS HIEBER, AMRU HUSSEIN, AND TAKAHITO KASHIWABARA

ABSTRACT. Consider the 3-d primitive equations in a layer domain  $\Omega = G \times (-h, 0)$ ,  $G = (0, 1)^2$ , subject to mixed Dirichlet and Neumann boundary conditions at  $z = -h$  and  $z = 0$ , respectively, and the periodic lateral boundary condition. It is shown that this equation is globally, strongly well-posed for arbitrary large data of the form  $a = a_1 + a_2$ , where  $a_1 \in C(\overline{G}; L^p(-h, 0))$ ,  $a_2 \in L^\infty(G; L^p(-h, 0))$  for  $p > 3$ , and where  $a_1$  is periodic in the horizontal variables and  $a_2$  is sufficiently small. In particular, no differentiability condition on the data is assumed. The approach relies on  $L_H^\infty L_z^p(\Omega)$ -estimates for terms of the form  $t^{1/2} \|\partial_z e^{tA_\sigma} \mathbb{P}f\|_{L_H^\infty L_z^p(\Omega)} \leq C e^{t\beta} \|f\|_{L_H^\infty L_z^p(\Omega)}$  for  $t > 0$ , where  $e^{tA_\sigma}$  denotes the hydrostatic Stokes semigroup. The difficulty in proving estimates of this form is that the hydrostatic Helmholtz projection  $\mathbb{P}$  fails to be bounded with respect to the  $L^\infty$ -norm. The global strong well-posedness result is then obtained by an iteration scheme, splitting the data into a smooth and a rough part and by combining a reference solution for smooth data with an evolution equation for the rough part.

## 1. INTRODUCTION

The primitive equations are a model for oceanic and atmospheric dynamics and are derived from the Navier-Stokes equations by assuming a hydrostatic balance for the pressure term, see [17–19]. These equations are known to be globally and strongly well-posed in the three dimensional setting for arbitrarily large data belonging to  $H^1$  by the celebrated result of Cao and Titi [5]. The latter considers the case of Neumann boundary conditions and this result also holds true for the case mixed Dirichlet and Neumann boundary conditions, again for data in  $H^1$ , as shown by Kukavika and Ziane [14].

Several approaches have been developed in the last years aiming for extending the above two results to the case of rough initial data. One approach is based on the theory of weak solutions, see e.g. [13, 16, 23, 24]. Although the existence of weak solutions to the primitive equations for initial data in  $L^2$  is known since the pioneering work by Lions, Temam and Wang [17], its uniqueness remains an open problem until today. Li and Titi [16] proved uniqueness of weak solutions assuming that the initial data are small  $L^\infty$ -perturbations of continuous data or data belonging to  $\{v \in L^6 : \partial_z v \in L^2\}$ , where  $z$  denotes the vertical variable. By a weak-strong uniqueness argument, these unique weak solutions regularize and even become strong solutions. For a survey of known results, see also [15].

A different approach to the primitive equations is based on a semilinear evolution equation for the hydrostatic Stokes operator within the  $L^p$ -setting, see [11]. There, the existence of a unique, global, strong solution to the primitive equations for initial data belonging to  $H^{2/p, p}$  was proved for the case of mixed Dirichlet-Neumann boundary conditions. This approach was transferred in [8, 9] to the case of pure Neumann boundary conditions and global, strong well-posedness of the primitive equations was obtained for data  $a$  of the form  $a = a_1 + a_2$ , where  $a_1 \in C(\overline{G}; L^1(-h, 0))$  and  $a_2 \in L^\infty(G; L^1(-h, 0))$  with  $a_2$  being small. These spaces are scaling invariant and represent the anisotropic character of the primitive equations.

---

2010 *Mathematics Subject Classification.* Primary: 35Q35; Secondary: 47D06, 76D03, 86A05.

*Key words and phrases.* hydrostatic Stokes semigroup,  $L^\infty$ -estimates, primitive equations, global strong well-posedness  
This work was partly supported by the DFG International Research Training Group IRTG 1529 and the JSPS Japanese-German Graduate Externship on Mathematical Fluid Dynamics. The first author is partly supported by JSPS through grant Kiban S (No. 26220702), Kiban A (No. 17H01091), Kiban B (No. 16H03948), the second and fourth author are supported by IRTG 1529 at TU Darmstadt, the fifth author is supported by JSPS Grant-in-Aid for Young Scientists B (No. 17K14230).

Note that the choice of boundary conditions has a severe impact on the linearized primitive equations. In the setting of layer domains, i.e.,  $\Omega = G \times (-h, 0) \subset \mathbb{R}^3$  with  $G = (0, 1)^2$  and  $h > 0$ , this is illustrated best by the hydrostatic Stokes operator  $A_{\bar{\sigma}}$ . The latter can be represented formally by the differential expression

$$(1.1) \quad \mathcal{A}v = \Delta v + \frac{1}{h} \nabla_H (-\Delta_H)^{-1} \operatorname{div}_H \left( \partial_z v \Big|_{z=-h} \right),$$

restricted to hydrostatically solenoidal vector fields, where for  $z = -h$  Dirichlet and for  $z = 0$  Neumann boundary conditions are imposed and periodicity is assumed horizontally, see [7] for details. In particular, in the case of pure Neumann boundary conditions, the hydrostatic Stokes operator reduces to the Laplacian, i.e.  $A_{\bar{\sigma}}v = \Delta v$ .

It is the aim of this article to study properties of the hydrostatic Stokes semigroup and terms of the form  $\nabla e^{tA_{\bar{\sigma}}}\mathbb{P}$  on spaces of bounded functions. These properties yield then the global, strong well-posedness result of the primitive equations in the case of mixed Dirichlet-Neumann boundary conditions. More precisely, we prove global, strong well-posedness of the primitive equations for initial data of the form

$$a = a_1 + a_2, \quad a_1 \in C(\bar{G}; L^p(-h, 0)), \quad \text{and} \quad a_2 \in L^\infty(G; L^p(-h, 0)) \quad \text{for } p > 3,$$

where  $a_1$  is periodic in the horizontal variables and  $a_2$  is sufficiently small. Our strategy is to introduce a reference solution for the smoothed part of the initial data and to combine this with an evolution equation approach for the remaining rough part.

The main difficulty when dealing with the primitive equations on spaces of bounded functions is that the hydrostatic Helmholtz projection  $\mathbb{P}$  *fails to be bounded* with respect to the  $L^\infty$ -norm. This is similar to the case of the classical Stokes semigroup, for which  $L^\infty$ -theory was developed in [1] and [2].

In Sections 6 and 7 we prove that the combination of the three main players,  $\nabla$ ,  $\mathbb{P}$ ,  $e^{tA_{\bar{\sigma}}}$ , nevertheless give rise to bounded operators on  $L_H^\infty L_z^p(\Omega)$ , which in addition satisfy typical global, second order parabolic decay estimates of the form

$$\begin{aligned} t^{1/2} \|\partial_i e^{tA_{\bar{\sigma}}}\mathbb{P}f\|_{L_H^\infty L_z^p(\Omega)} &\leq C e^{t\beta} \|f\|_{L_H^\infty L_z^p(\Omega)}, \\ t^{1/2} \|e^{tA_{\bar{\sigma}}}\mathbb{P}\partial_j f\|_{L_H^\infty L_z^p(\Omega)} &\leq C e^{t\beta} \|f\|_{L_H^\infty L_z^p(\Omega)}, \\ t \|\partial_i e^{tA_{\bar{\sigma}}}\mathbb{P}\partial_j f\|_{L_H^\infty L_z^p(\Omega)} &\leq C e^{\beta t} \|f\|_{L_H^\infty L_z^p(\Omega)}, \end{aligned}$$

for  $t > 0$ , where  $\partial_i, \partial_j \in \{\partial_x, \partial_y, \partial_z\}$ .

Note that the choice of the boundary conditions involved affects to a very great extent the difficulty in proving these estimates. For the case of mixed Dirichlet-Neumann boundary conditions, our approach relies on the representation (1.1) of the linearized problem. The constraint  $p > 3$  arises from embedding properties for the reference solution and estimates for the linearized problem in  $L^\infty(G; L^p(-h, 0))$ .

Our approach is based on an iteration scheme, which is inspired by the classical schemes to the Navier-Stokes equations. Here, the iterative construction of a unique, local solution relies on  $L_H^\infty L_z^p(\Omega)$ -estimates for the crucial terms of the form  $e^{tA_{\bar{\sigma}}}\mathbb{P}\operatorname{div}(u \otimes v)$ , where  $u = (v, w)$  is the full velocity and  $v$  its horizontal component. Let us note that the above linear estimates are of independent interest for further considerations.

The use of a reference solution allows us to obtain the smallness condition on the  $L_H^\infty L_z^p$ -perturbation  $a_2$  of  $a_1$  by means of an absolute constant, while for Neumann boundary conditions it is needed that  $a_2$  is small compared to  $a_1$ , cf. [8]. Also, Li and Titi assume in [16] that  $a_2$  is small compared to the  $L^4$ -norm of  $a_1$ .

Comparing our result with the one by Li and Titi in [16], which has been obtained for Neumann boundary conditions, we observe that the initial data allowed in our approach are of anisotropic nature and require no conditions on the derivatives of the initial data, such as e.g.  $\partial_z v \in L^2$  as in [16].

This article is structured as follows: In Section 2 we collect preliminary facts and fix the notation. In Section 3 we state our main results concerning the global strong well-posedness of the primitive equations for rough data and the crucial estimates for the linearized problem. The proof of our main results starts with a discussion of anisotropic  $L^p$ -spaces in Section 4, which is followed in Section 5 by estimates for the Laplacian in anisotropic spaces. The subsequent Sections 6 and 7 are devoted to the development

of an  $L^\infty(G; L^p(-h, 0))$ -theory for the hydrostatic Stokes equations and its associated resolvent problem. Finally, in Section 8 we present our iteration scheme yielding the global, strong well-posedness of the primitive equations for rough initial data.

## 2. PRELIMINARIES

Let  $\Omega = G \times (-h, 0)$  where  $G = (0, 1)^2$ . We consider the primitive equations on  $\Omega$  given by

$$(2.1) \quad \begin{aligned} \partial_t v - \Delta v + (u \cdot \nabla)v + \nabla_H \pi &= 0 & \text{on } \Omega \times (0, \infty), \\ \partial_z \pi &= 0 & \text{on } \Omega \times (0, \infty), \\ \operatorname{div}_H \bar{v} &= 0 & \text{on } G \times (0, \infty), \\ v(0) &= a & \text{on } \Omega, \end{aligned}$$

using the notations  $\operatorname{div}_H v = \partial_x v_1 + \partial_y v_2$  and  $\nabla_H \pi = (\partial_x \pi, \partial_y \pi)^T$ , while  $\bar{v} = \frac{1}{h} \int_{-h}^0 v(\cdot, z) dz$  is the vertical average,  $\pi: G \rightarrow \mathbb{R}$  denotes the surface pressure,  $u = (v, w)$  is the velocity field with horizontal and vertical components  $v: \Omega \rightarrow \mathbb{R}^2$  and  $w: \Omega \rightarrow \mathbb{R}$  respectively, where  $w = w(v)$  is given by the relation

$$(2.2) \quad w(x, y, z) = - \int_h^z \operatorname{div}_H v(x, y, r) dr.$$

This is supplemented by mixed Dirichlet and Neumann boundary conditions

$$(2.3) \quad \partial_z v = 0 \text{ on } \Gamma_u \times (0, \infty), \quad \pi, v \text{ periodic on } \Gamma_l \times (0, \infty), \quad v = 0 \text{ on } \Gamma_b \times (0, \infty),$$

where the boundary is divided into  $\Gamma_u = G \times \{0\}$ ,  $\Gamma_l = \partial G \times [-h, 0]$  and  $\Gamma_b = G \times \{0\}$ .

In the following we will be dealing with anisotropic  $L^p$ -spaces on cylindrical sets of the type  $U = \Omega$  or  $U = \mathbb{R}^2 \times \mathbb{R}$ . More precisely, if  $U = U' \times U_3 \subset \mathbb{R}^2 \times \mathbb{R}$  is a product of measurable sets and  $q, p \in [1, \infty]$  we define

$$L_H^q L_z^p(U) := L^q(U'; L^p(U_3)) := \{f: U \rightarrow \mathbb{K} \text{ measurable, } \|f\|_{L_H^q L_z^p(U)} < \infty\},$$

for  $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$  with norm

$$\|f\|_{L_H^q L_z^p(U)} := \begin{cases} \left( \int_{U'} \|f(x', \cdot)\|_{L^p(U_3)}^q dx' \right)^{1/q}, & q \in [1, \infty), \\ \operatorname{ess\,sup}_{x' \in U'} \|f(x', \cdot)\|_{L^p(U_3)}, & q = \infty. \end{cases}$$

Endowed with this norm,  $L_H^q L_z^p(U)$  is a Banach space for all  $p, q \in [1, \infty]$ .

We will denote the  $W^{k,p}$ -closure of  $C_{\text{per}}^\infty(\bar{\Omega})$  by  $W_{\text{per}}^{k,p}(\Omega)$ , where  $C_{\text{per}}^\infty(\bar{\Omega})$  denotes the space of smooth functions  $v$  on  $\bar{\Omega}$  that such that  $\partial_x^\alpha v$  and  $\partial_y^\alpha v$  are periodic on  $\Gamma_l$  with period 1 in the variables  $x$  and  $y$  for all  $\alpha \in \mathbb{N}$ , but not necessarily periodic with respect to the vertical direction  $z$ . Moreover, by  $C^{m,\alpha}(\bar{\Omega})$ ,  $C^{m,\alpha}(\bar{G})$  we denote the spaces of  $m$ -times differentiable functions with Hölder-continuous derivatives of exponents  $\alpha \in (0, 1)$  and the subspaces of functions periodic on  $\Gamma_l$  and  $\partial G$  will be denoted by  $C_{\text{per}}^{m,\alpha}(\bar{\Omega})$  and  $C_{\text{per}}^{m,\alpha}(\bar{G})$ , respectively. For a Banach space  $E$  we denote by  $C_{\text{per}}([0, 1]^2; E)$  the set of continuous functions  $f: [0, 1]^2 \rightarrow E$  such that  $f(0, y) = f(1, y)$  and  $f(y, 0) = f(y, 1)$  for all  $x, y \in [0, 1]$ .

In order to include the condition  $\operatorname{div}_H \bar{v} = 0$  one defines the *hydrostatic Helmholtz projection*  $\mathbb{P}$  as in [7, 11] using the two-dimensional Helmholtz projection  $Q$  with periodic boundary conditions given by  $Qg = g - \nabla_H \pi$  for  $g: G \rightarrow \mathbb{R}^2$  solving  $\Delta_H \pi = \operatorname{div}_H g$  for  $\pi$  periodic on  $\partial G$ , where  $\Delta_H g = \partial_x^2 g + \partial_y^2 g$ . The hydrostatic Helmholtz projection is then defined as

$$\mathbb{P}f = f - (1 - Q)\bar{f} = f + \frac{1}{h} \nabla_H (-\Delta_H)^{-1} \operatorname{div}_H \bar{f} = f - \nabla_H \pi.$$

The range of  $\mathbb{P}: L^p(\Omega)^2 \rightarrow L^p(\Omega)^2$ ,  $p \in (1, \infty)$ , is denoted by  $L_\sigma^p(\Omega)$  and is given by

$$\overline{\{v \in C_{\text{per}}^\infty(\bar{\Omega})^2 : \operatorname{div}_H \bar{v} = 0\}}^{\|\cdot\|_{L^p(\Omega)}}.$$

Further characterizations of  $L_\sigma^p(\Omega)$  are given in [11, Proposition 4.3].

Since  $\mathbb{P}$  fails to be bounded on  $L^\infty(\Omega)^2$  it is not evident which space is a suitable substitute for  $L_\sigma^p(\Omega)$  in the case  $p = \infty$ . In this article, we will be considering the spaces

$$(2.4) \quad X := C_{\text{per}}([0, 1]^2; L^p(-h, 0))^2 \quad \text{and} \quad X_{\bar{\sigma}} := X \cap L_\sigma^p(\Omega), \quad p \in (1, \infty).$$

The linearization of equation (2.1), called the *hydrostatic Stokes equation*, is given by

$$(2.5) \quad \partial_t v - \Delta v + \nabla_H \pi = f, \quad \text{div}_H \bar{v} = 0, \quad v(0) = a$$

and subject to boundary conditions (2.3). The dynamics of this evolution equation is governed by the hydrostatic Stokes operator, and its  $X_{\bar{\sigma}}$ -realization  $A_{\bar{\sigma}}$  is given by

$$A_{\bar{\sigma}} v := \mathcal{A}v, \quad D(A_{\bar{\sigma}}) = \{v \in W_{\text{per}}^{2,p}(\Omega)^2 \cap X_{\bar{\sigma}} : \partial_z v|_{\Gamma_u} = 0, v|_{\Gamma_b} = 0, \mathcal{A}v \in X_{\bar{\sigma}}\},$$

where  $\mathcal{A}v$  is defined by (1.1). It will be proved that  $A_{\bar{\sigma}}$  generates a strongly continuous, analytic semigroup  $e^{tA_{\bar{\sigma}}}$  on  $X_{\bar{\sigma}}$ . Information on the linear theory in  $L_\sigma^p(\Omega)$  for  $p \in (1, \infty)$  can be found in [7].

### 3. MAIN RESULTS

Our first main result concerns the global well-posedness of the primitive equations for *arbitrarily large* initial data in  $X_{\bar{\sigma}}$ , while the second result extends this situation to the case of small perturbations in  $L_H^\infty L_z^p(\Omega)$ . Here, a *strong solution* means – as in [11] – a solution  $v$  to the primitive equations satisfying

$$(3.1) \quad v \in C^1((0, \infty); L^p(\Omega))^2 \cap C((0, \infty); W^{2,p}(\Omega))^2.$$

Our third main result concerns  $L_H^\infty L_z^p$ -estimates for the hydrostatic Stokes semigroup. These estimates are essential for proving the above two results on the non-linear problem. They are also of independent interest.

**Theorem 3.1.** *Let  $p \in (3, \infty)$ . Then for all  $a \in X_{\bar{\sigma}}$  there exists a unique, global, strong solution  $v$  to the primitive equations (2.1) with  $v(0) = a$  satisfying*

$$v \in C([0, \infty); X_{\bar{\sigma}}), \quad t^{1/2} \nabla v \in L^\infty((0, \infty); X), \quad \limsup_{t \rightarrow 0^+} t^{1/2} \|\nabla v(t)\|_{L_H^\infty L_z^p(\Omega)} = 0.$$

*The corresponding pressure satisfies*

$$\pi \in C((0, \infty); C^{1,\alpha}([0, 1]^2)), \quad \alpha \in (0, 1 - 3/p)$$

*and is unique up to an additive constant.*

**Theorem 3.2.** *Let  $p \in (3, \infty)$ . Then there exists a constant  $C_0 > 0$  such that if  $a = a_1 + a_2$  with  $a_1 \in X_{\bar{\sigma}}$  and  $a_2 \in L_H^\infty L_z^p(\Omega)^2 \cap L_\sigma^p(\Omega)$  with*

$$\|a_2\|_{L_H^\infty L_z^p(\Omega)} \leq C_0,$$

*then there exists a unique, global, strong solution  $v$  to the primitive equations (2.1) with  $v(0) = a$  satisfying*

$$v \in C([0, \infty); L_\sigma^p(\Omega)) \cap L^\infty((0, T); L_H^\infty L_z^p(\Omega))^2$$

*as well as*

$$t^{1/2} \nabla v \in L^\infty((0, \infty); X), \quad \limsup_{t \rightarrow 0^+} t^{1/2} \|\nabla v\|_{L_H^\infty L_z^p(\Omega)} \leq C \|a_2\|_{L_H^\infty L_z^p},$$

*where  $C > 0$  does not depend on the data, and the pressure has the same regularity as in Theorem 3.1.*

Taking advantage of the regularization of solutions for  $t > 0$  one passes into the setting discussed in [11] and [9], and thus we obtain the following corollary.

**Corollary 3.3.** *For  $t > 0$  the solution  $v, \pi$  in Theorem 3.1 and Theorem 3.2 are real analytic in time and space, and the velocity  $v$  decays exponentially as  $t \rightarrow \infty$ .*

Our main result on the hydrostatic semigroup acting on  $X_{\bar{\sigma}}$  reads as follows.

**Theorem 3.4.** *Let  $p \in (3, \infty)$ . Then the following assertions hold true:*

a)  $A_{\bar{\sigma}}$  is the generator of a strongly continuous, analytic and exponentially stable semigroup  $e^{tA_{\bar{\sigma}}}$  on  $X_{\bar{\sigma}}$  of angle  $\pi/2$ .

b) There exist constants  $C > 0$ ,  $\beta > 0$  such that for  $\partial_i, \partial_j \in \{\partial_x, \partial_y, \partial_z\}$

- (i)  $t^{1/2} \|\partial_j e^{tA_{\bar{\sigma}}} f\|_{L_H^\infty L_z^p(\Omega)} \leq C e^{\beta t} \|f\|_{L_H^\infty L_z^p(\Omega)}, \quad t > 0, f \in X_{\bar{\sigma}},$
- (ii)  $t^{1/2} \|e^{tA_{\bar{\sigma}}} \mathbb{P} \partial_j f\|_{L_H^\infty L_z^p(\Omega)} \leq C e^{\beta t} \|f\|_{L_H^\infty L_z^p(\Omega)}, \quad t > 0, f \in X_{\bar{\sigma}},$
- (iii)  $t \|\partial_i e^{tA_{\bar{\sigma}}} \mathbb{P} \partial_j f\|_{L_H^\infty L_z^p(\Omega)} \leq C e^{\beta t} \|f\|_{L_H^\infty L_z^p(\Omega)}, \quad t > 0, f \in X_{\bar{\sigma}};$

c) For all  $f \in X_{\bar{\sigma}}$

$$\lim_{t \rightarrow 0^+} t^{1/2} \|\nabla e^{tA_{\bar{\sigma}}} f\|_{L_H^\infty L_z^p(\Omega)} = 0.$$

**Remarks 3.5.** a) We note that when in the situation of Theorem 3.2 the initial data do not belong to  $X$ , i.e. when  $a_2 \neq 0$ , the solution fails to be continuous at  $t = 0$  with respect to the  $L_H^\infty L_z^p$ -norm.

b) The condition  $p > 3$  is due to the embeddings

$$v_{\text{ref}}(t_0) \in B_{pq}^{2-2/q}(\Omega)^2 \hookrightarrow C^1(\bar{\Omega})^2 \quad \text{and} \quad W^{2,p}(\Omega) \hookrightarrow C^{1,\alpha}(\bar{\Omega}) \quad \text{for } p \in (3, \infty),$$

cf. [25, Section 3.3.1]. Since the two-dimensional Helmholtz projection  $Q$  fails to be bounded with respect to the  $L^\infty$ -norm, we instead estimate it in spaces of Hölder continuous functions  $C_{\text{per}}^{0,\alpha}([0, 1]^2) = C^{0,\alpha}(\mathbb{T}^2)$  for  $\alpha \in (0, 1)$  where  $\mathbb{T}^2$  denotes the two-dimensional torus. In fact  $Q$  is bounded with respect to the  $C^{0,\alpha}$ -norm. This follows by the theory of Fourier multipliers on Besov spaces, compare e.g. [3, Theorem 6.2] for the whole space, and the periodic case follows using periodic extension.

c) In Theorem 3.4 one can even consider  $f \in L_H^\infty L_z^p(\Omega)^2$  for  $p \in (3, \infty)$ . Then the corresponding semigroup is still analytic, but it fails to be strongly continuous. The estimates (i) – (iii) still hold, whereas property c) in Theorem 3.4 has to be replaced by

$$\limsup_{t \rightarrow 0^+} t^{1/2} \|\nabla e^{tA_{\bar{\sigma}}} v\|_{L_H^\infty L_z^p(\Omega)} \leq C \|v\|_{L_H^\infty L_z^p(\Omega)}$$

for some  $C > 0$ , where with a slight abuse of notation  $e^{tA_{\bar{\sigma}}}$  denotes also the hydrostatic Stokes semigroup on  $L_H^\infty L_z^p(\Omega)$ .

d) Some words about our strategy for proving the global well-posedness results are in order:

- (i) We will first construct a local, mild solution to the problem (2.1), i.e. a function satisfying the relation

$$(3.2) \quad v(t) = e^{tA_{\bar{\sigma}}} a + \int_0^t e^{(t-s)A_{\bar{\sigma}}} \mathbb{P} F(v(s)) ds, \quad t \in (0, T)$$

for some  $T > 0$ , where  $F(v) = -(u \cdot \nabla)v$ . We will then show that  $v$  regularizes for  $t_0 > 0$  and using the result of [11, Theorem 6.1] or [9, Theorem 3.1], we may take  $v(t_0)$  as a new initial value to extend the mild solution to a global, strong solution on  $(t_0, \infty)$  and then on  $(0, \infty)$  by uniqueness. The additional regularity for  $t \rightarrow 0^+$  results from the construction of the mild solutions.

- (ii) In order to construct a mild solution we decompose  $a = a_{\text{ref}} + a_0$  such that  $a_{\text{ref}}$  is sufficiently smooth and  $a_0$  can be taken to be arbitrarily small.
- (iii) Using previously established results concerning the existence of solutions to the primitive equations for *smooth* data, we obtain a reference solution  $v_{\text{ref}}$  and construct then  $V := v - v_{\text{ref}}$  via an iteration scheme using  $L^\infty$ -type estimates for terms of the form  $\nabla e^{tA_{\bar{\sigma}}} \mathbb{P}$  given in Theorem 3.4.

#### 4. PROPERTIES OF ANISOTROPIC SPACES

In this section, we will discuss properties of anisotropic  $L^q$ - $L^p$ -spaces. We will write  $C(U'; L^p(U_3))$  for the set of continuous  $L^p(U_3)$ -valued functions on  $U'$  and likewise

$$L^q(U'; C(U_3)) := \{f \in L_H^q L_z^\infty(U) : f(x', \cdot) \in C(U_3) \text{ for almost all } x' \in U'\},$$

and  $C_c(U'; L^p(U_3))$  and  $L^q(U'; C_c(U_3))$  for the subsets of functions with compact support in horizontal and vertical variables, respectively. For  $p, q \in [1, \infty)$  the space  $C_c^\infty(U)$  is dense in these spaces as well as in  $L_H^q L_z^p(U)$ , and furthermore we have

$$\overline{C_c^\infty(\mathbb{R}^3)}^{\|\cdot\|_{L_H^\infty L_z^p}} = C_0(\mathbb{R}^2; L^p(\mathbb{R})), \quad \overline{C_c^\infty(\mathbb{R}^3)}^{\|\cdot\|_{L_H^q L_z^\infty}} = L^q(\mathbb{R}^2; C_0(\mathbb{R})),$$

as well as

$$\overline{C_{\text{per}}^\infty(\overline{\Omega})}^{\|\cdot\|_{L_H^\infty L_z^p}} = X, \quad \overline{C_{\text{per}}^\infty(\overline{\Omega})}^{\|\cdot\|_{L_H^q L_z^\infty}} = L^q(G; C[-h, 0]).$$

Observe that even  $C_{\text{per}}^\infty([0, 1]^2; C_c^\infty(-h, 0))^2$  is dense in  $X$  and  $L_H^q L_z^p(\Omega)^2$ . If  $p = q = \infty$ , then

$$\overline{C_c^\infty(\mathbb{R}^3)}^{\|\cdot\|_{L_H^\infty L_z^\infty}} = C_0(\mathbb{R}^3), \quad \overline{C_{\text{per}}^\infty(\overline{\Omega})}^{\|\cdot\|_{L_H^\infty L_z^\infty}} = C_{\text{per}}([0, 1]^2; C[-h, 0]).$$

Here  $C_0(\mathbb{R}^d)$  denotes the set of functions vanishing at infinity. These density results follow from the fact that if  $E$  is a Banach space over  $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ , then the linear space generated by elementary tensor functions  $f \otimes e$  for measurable  $f : U' \rightarrow \mathbb{K}$  and  $e \in E$  is dense in  $L^q(U'; E)$  for  $q \in [1, \infty)$ , since it contains the simple  $E$ -valued functions. It is also dense in  $C_0(U'; E)$ , if one only considers continuous functions  $f$ , due to a generalization of the Stone-Weierstrass theorem, see e.g. [12].

In the case that  $U \subset \mathbb{R}^3$  is bounded, we also have

$$L_H^{q_1} L_z^{p_1}(U) \hookrightarrow L_H^{q_2} L_z^{p_2}(U)$$

whenever  $q_1 \geq q_2$  and  $p_1 \geq p_2$ . See [11, Section 5] for more details.

Another important property of the  $L_H^q L_z^p$ -norm is its behaviour under operations like multiplication and convolution. For the former one, we obviously obtain

$$\|fg\|_{L_H^q L_z^p(U)} \leq \|f\|_{L_H^{q_1} L_z^{p_1}(U)} \|g\|_{L_H^{q_2} L_z^{p_2}(U)}$$

whenever  $1/p_1 + 1/p_2 = 1/p$  and  $1/q_1 + 1/q_2 = 1/q$ . For the latter one, the following variant of Young's inequality holds true.

**Lemma 4.1.** [10, Theorem 3.1]. *Let  $f \in L_H^q L_z^p(\mathbb{R}^3)$  for  $p, q \in [1, \infty]$  and  $g \in L^1(\mathbb{R}^3)$ . Then*

$$\|g * f\|_{L_H^q L_z^p(\mathbb{R}^3)} \leq \|g\|_{L^1(\mathbb{R}^3)} \|f\|_{L_H^q L_z^p(\mathbb{R}^3)}.$$

## 5. LINEAR ESTIMATES FOR THE LAPLACE OPERATOR

In this section we establish resolvent and semigroup estimates for Laplace operators with a focus on anisotropic  $L_H^q L_z^p$ -spaces, where  $p, q \in [1, \infty]$ .

First, we consider the resolvent problem for the Laplacian on the full space for

$$\lambda \in \Sigma_\theta = \{\lambda \in \mathbb{C} \setminus \{0\} : |\arg(\lambda)| < \theta\}, \quad \theta \in (0, \pi),$$

i.e.

$$(5.1) \quad \Delta v - \lambda v = f \text{ on } \mathbb{R}^3, \quad f \in C_c^\infty(\mathbb{R}^3),$$

and for  $\partial_j \in \{\partial_x, \partial_y, \partial_z\}$

$$(5.2) \quad \Delta w - \lambda w = \partial_j f \text{ on } \mathbb{R}^3, \quad f \in C_c^\infty(\mathbb{R}^3).$$

It is well known that the solution to problem (5.1) is given by the convolution  $v = K_\lambda * f$  and the one to problem (5.2) by  $v = \partial_j K_\lambda * f$ , where  $K_\lambda$  is explicitly given by

$$K_\lambda(x) = \frac{1}{4\pi} \frac{e^{-\lambda^{1/2}|x|}}{|x|}, \quad x \in \mathbb{R}^3 \setminus \{0\}.$$

Using this representation one easily obtains the following uniform  $L^1(\mathbb{R}^3)$ -estimates.

**Lemma 5.1.** *For all  $\theta \in (0, \pi)$  there exists  $C_\theta > 0$  such that for all  $\lambda \in \Sigma_\theta$  one has*

$$|\lambda| \cdot \|K_\lambda\|_{L^1(\mathbb{R}^3)} + |\lambda|^{1/2} \|\nabla K_\lambda\|_{L^1(\mathbb{R}^3)} \leq C_\theta.$$

*Proof.* Set  $\psi := \arg(\lambda) \in (-\theta, \theta)$ . Since  $K_\lambda$  is radially symmetric we use spherical coordinates to obtain

$$\int_{\mathbb{R}^3} |K_\lambda(x)| dx = \int_0^\infty r e^{-|\lambda|^{1/2} \cos(\psi/2)r} dr$$

as well as

$$\int_{\mathbb{R}^3} |\nabla K_\lambda(x)| dx \leq \int_0^\infty (1 + |\lambda|^{1/2}r) e^{-|\lambda|^{1/2} \cos(\psi/2)r} dr.$$

So,  $|\lambda| \cdot \|K_\lambda\|_{L^1(\mathbb{R}^3)} = \sec(\psi/2)^2$  and  $|\lambda|^{1/2} \|\nabla K_\lambda\|_{L^1(\mathbb{R}^3)} \leq \sec(\psi/2) + \sec(\psi/2)^2$ , and thus we obtain the desired result.  $\square$

From this and Young's inequality for convolutions in anisotropic spaces, cf. Lemma 4.1, one immediately obtains suitable  $L_H^q L_z^p$ -estimates for the resolvent problems (5.1) and (5.2) for  $q, p \in [1, \infty]$ .

**Corollary 5.2.** *Let  $\lambda \in \Sigma_\theta$  for some  $\theta \in (0, \pi)$ . Assume one of the following cases:*

- (i)  $p, q \in [1, \infty)$  and  $f \in L_H^q L_z^p(\mathbb{R}^3)$ , or
- (ii)  $p \in [1, \infty)$ ,  $q = \infty$ , and  $f \in L_H^q L_z^p(\mathbb{R}^3)$  with compact support in horizontal direction, or
- (iii)  $p = \infty$ ,  $q \in [1, \infty)$ , and  $f \in L_H^q L_z^p(\mathbb{R}^3)$  with compact support in vertical direction.

Then the functions

$$v = K_\lambda * f \quad \text{and} \quad w = \partial_j K_\lambda * f$$

are the unique solutions to the problems (5.1) and (5.2) in  $L_H^q L_z^p(\mathbb{R}^3)$ , respectively, and there exists a constant  $C_\theta > 0$  such that

$$(5.3) \quad |\lambda| \cdot \|v\|_{L_H^q L_z^p(\mathbb{R}^3)} + |\lambda|^{1/2} \|\nabla v\|_{L_H^q L_z^p(\mathbb{R}^3)} + \|\Delta v\|_{L_H^q L_z^p(\mathbb{R}^3)} \leq C_\theta \|f\|_{L_H^q L_z^p(\mathbb{R}^3)},$$

$$(5.4) \quad |\lambda|^{1/2} \|w\|_{L_H^q L_z^p(\mathbb{R}^3)} \leq C_\theta \|f\|_{L_H^q L_z^p(\mathbb{R}^3)}.$$

**Remark 5.3.** In the case  $q, p \in [1, \infty)$  we have that  $C_c^\infty(\mathbb{R}^3)$  is dense in  $L_H^q L_z^p(\mathbb{R}^3)$ , so we may assume that  $f$  is essentially bounded and has compact support, yielding  $\partial_i(K_\lambda * f) = (\partial_i K_\lambda) * f$ . In the cases where  $q$  and/or  $p$  is infinite we add this as an assumption.

We now investigate for the Laplacian on  $\Omega$  with boundary conditions (2.3) the resolvent problems

$$(5.5) \quad \lambda v - \Delta v = f \text{ on } \Omega,$$

and for  $\partial_i \in \{\partial_x, \partial_y, \partial_z\}$

$$(5.6) \quad \lambda w - \Delta w = \partial_i f \text{ on } \Omega.$$

**Lemma 5.4.** *Let  $\theta \in (0, \pi)$  and  $f \in L_H^q L_z^p(\Omega)$  for  $q \in [1, \infty], p \in [1, \infty)$ . Then there exists  $\lambda_0 > 0$  such that for  $\lambda \in \Sigma_\theta$  with  $|\lambda| \geq \lambda_0$  the problems (5.5) and (5.6) have unique solutions  $v \in L_H^q L_z^p(\Omega)$  and  $w \in L_H^q L_z^p(\Omega)$ , respectively, and there exists a constant  $C_\theta > 0$  such that*

$$(5.7) \quad |\lambda| \cdot \|v\|_{L_H^q L_z^p(\Omega)} + |\lambda|^{1/2} \|\nabla v\|_{L_H^q L_z^p(\Omega)} + \|\Delta v\|_{L_H^q L_z^p(\Omega)} \leq C_\theta \|f\|_{L_H^q L_z^p(\Omega)},$$

$$(5.8) \quad |\lambda|^{1/2} \|w\|_{L_H^q L_z^p(\Omega)} \leq C_\theta \|f\|_{L_H^q L_z^p(\Omega)}.$$

In particular for  $q = \infty$  and  $p \in (2, \infty)$  one can chose  $\lambda_0 = 0$ .

To prove this lemma, we will need some facts concerning isotropic  $L^p$ -spaces. So, for  $p \in (1, \infty)$  denote by  $\Delta_p$  the Laplace operator on  $L^p(\Omega)$  defined by

$$\Delta_p v = \Delta v, \quad D(\Delta_p) = \{v \in W_{\text{per}}^{2,p}(\Omega) : \partial_z v|_{\Gamma_u} = 0, v|_{\Gamma_b} = 0\}.$$

One has  $\rho(-\Delta_p) \subset \mathbb{C} \setminus [\delta, \infty)$ , for some  $\delta > 0$ , i.e.  $0 \in \rho(-\Delta_p)$ , cf. [21, Remark 8.23], and the resolvent satisfies for some  $C_{\theta,p} > 0$  the estimate

$$(5.9) \quad |\lambda| \cdot \|(\Delta_p - \lambda)^{-1} f\|_{L^p(\Omega)} + \|\Delta_p (\Delta_p - \lambda)^{-1} f\|_{L^p(\Omega)} \leq C_{\theta,p} \|f\|_{L^p(\Omega)}, \quad f \in L^p(\Omega),$$



where  $\lambda \in \Sigma_\theta$ ,  $\theta \in (0, \pi)$ . Furthermore,  $-\Delta_p$  possesses a bounded  $\mathcal{H}^\infty$ -calculus of angle 0, see e.g. [21], and therefore

$$(5.10) \quad D((-\Delta_p)^\vartheta) = [L^p(\Omega), D(\Delta_p)]_\vartheta \subset W^{2\vartheta, p}(\Omega), \quad \vartheta \in [0, 1],$$

where  $[\cdot, \cdot]$  denotes the complex interpolation functor. In particular  $\partial_j(-\Delta_p)^{-1/2}$  is bounded on  $L^p(\Omega)$  for  $\partial_j \in \{\partial_x, \partial_y, \partial_z\}$  and by taking adjoints the same holds true for the closure of  $(-\Delta_p)^{-1/2}\partial_j$ . This yields the estimates

$$(5.11) \quad \begin{aligned} |\lambda|^{1/2} \|\partial_j(\Delta_p - \lambda)^{-1} f\|_{L^p(\Omega)} + |\lambda|^{1/2} \|(\Delta_p - \lambda)^{-1} \partial_j f\|_{L^p(\Omega)} &\leq C_{\theta, p} \|f\|_{L^p(\Omega)}, \quad f \in L^p(\Omega), \\ \|\partial_j(\Delta_p - \lambda)^{-1} \partial_i f\|_{L^p(\Omega)} &\leq C_{\theta, p} \|f\|_{L^p(\Omega)}, \quad f \in L^p(\Omega) \end{aligned}$$

for  $\lambda \in \Sigma_\theta$ ,  $\theta \in (0, \pi)$ , and some  $C_{\theta, p} > 0$ .

*Proof of Lemma 5.4.* First, we apply the following density arguments:

- (i) For  $q, p \in [1, \infty)$  and  $f \in L_H^q L_z^p(\Omega)$  we assume that  $f \in C_{\text{per}}^\infty([0, 1]^2; C_c^\infty(-h, 0))$  since  $C_{\text{per}}^\infty([0, 1]^2; C_c^\infty(-h, 0))$  is a dense subspace of  $L_H^q L_z^p(\Omega)$ .
- (ii) For  $q = \infty$  and  $f \in L^\infty(G; L^p(-h, 0))$ , we assume that  $f \in L^\infty(G; C_c^\infty(-h, 0))$ , as the latter space is dense in  $L_H^\infty L_z^p(\Omega)$ .

In particular, in either case we may assume that  $f = 0$  on  $\Gamma_u \cup \Gamma_b$  and  $f \in L^\infty(\Omega)$ . The existence of a unique solution to the problems (5.5) and (5.6) in  $L_H^q L_z^p(\Omega)$  for such smooth  $f$  follows from the properties of the mappings  $(\lambda - \Delta)^{-1}$  and  $(\lambda - \Delta)^{-1} \partial_i$  in  $L^r(\Omega)$  for  $\lambda \in \Sigma_\theta$  since

$$v \in W^{2, r}(\Omega) \hookrightarrow L^\infty(\Omega) \hookrightarrow L_H^q L_z^p(\Omega) \quad \text{and} \quad w \in W^{1, r}(\Omega) \hookrightarrow L^\infty(\Omega) \hookrightarrow L_H^q L_z^p(\Omega), \quad r > 3.$$

It therefore suffices to prove the estimates (5.7) and (5.8). This is done in the following by localizing the results of Lemma 5.2.

For this purpose we first make use of the extension operator

$$E = E_z^{\text{even, odd}} \circ E_H^{\text{per}}$$

where  $E_H^{\text{per}}$  is the periodic extension operator from  $G$  to  $\mathbb{R}^2$  in horizontal direction and  $E_z^{\text{even, odd}}$  extends from  $(-h, 0)$  to  $(-2h, h)$  in vertical direction via even and odd reflexion at the top and bottom part of the boundary respectively.

Second, we utilize a family of cut-off-functions  $\chi_r \in C_c^\infty(\mathbb{R}^3)$  for  $r \in (0, \infty)$  of the form  $\chi_r(x, y, z) = \varphi_r(x, y) \psi_r(z)$  where  $\varphi_r \in C_c^\infty(\mathbb{R}^2)$  and  $\psi_r \in C_c^\infty(\mathbb{R})$  satisfy

$$\begin{aligned} \varphi_r &\equiv 1 \quad \text{on } [-1/4, 5/4]^2, & \varphi_r &\equiv 0 \quad \text{on } ((-\infty, -r - 1/4] \cup [5/4 + r, \infty))^2, \\ \psi_r &\equiv 1 \quad \text{on } [-5h/4, h/4], & \psi_r &\equiv 0 \quad \text{on } (-\infty, -r - 5h/4] \cup [h/4 + r, \infty), \end{aligned}$$

and there is a constant  $M > 0$  independent of  $r$  such that

$$\|\varphi_r\|_\infty + \|\psi_r\|_\infty + r (\|\nabla_H \varphi_r\|_\infty + \|\partial_z \psi_r\|_\infty) + r^2 (\|\Delta_H \varphi_r\|_\infty + \|\partial_z^2 \psi_r\|_\infty) \leq M.$$

Here, we consider  $0 < 4r < 3 \min\{1, h\}$  which implies that  $\varphi_r$  and  $\psi_r$  are supported on  $(-1, 2)$  and  $(-2h, h)$  respectively. We now define an extension of  $v$  from  $\Omega$  onto the whole space  $\mathbb{R}^3$  via

$$u(x, y, z) = \chi_r(x, y, z)(Ev)(x, y, z)$$

for a suitable value of  $r$  which we will specify later on. If  $v$  solves problem (5.5) then  $u$  solves the problem

$$\lambda u - \Delta u = F \text{ on } \mathbb{R}^3, \quad F := \chi_r E f - 2(\nabla \chi_r) \cdot E(\nabla v) - (\Delta \chi_r) E v.$$

Here we made use of the fact that  $E$  commutes with derivatives of  $v$ .

Note that not only does  $F$  have compact support, but we also have  $F \in L^\infty(\mathbb{R}^3)$  since we may assume that  $f \in L^\infty(\Omega)$  and  $v \in W^{1, \infty}(\Omega)$  by the above approximation argument. Thus we may now apply Lemma 5.2, and estimate (5.3) yields

$$|\lambda| \cdot \|u\|_{L_H^q L_z^p(\mathbb{R}^3)} + |\lambda|^{1/2} \|\nabla u\|_{L_H^q L_z^p(\mathbb{R}^3)} \leq C_\theta \|F\|_{L_H^q L_z^p(\mathbb{R}^3)}.$$

To estimate  $F$  we use that  $\chi_r$  is supported on  $(-1, 2)^2 \times (-2h, h)$ , and therefore

$$\begin{aligned} \|\chi_r E f\|_{L_H^q L_z^p(\mathbb{R}^3)} &\leq 27M^2 \|f\|_{L_H^q L_z^p(\Omega)}, \\ \|(\nabla \chi_r) \cdot E(\nabla v)\|_{L_H^q L_z^p(\mathbb{R}^3)} &\leq 27M^2 r^{-1} \|\nabla v\|_{L_H^q L_z^p(\Omega)}, \\ \|(\Delta \chi_r) E v\|_{L_H^q L_z^p(\mathbb{R}^3)} &\leq 27M^2 r^{-2} \|v\|_{L_H^q L_z^p(\Omega)}. \end{aligned}$$

Next, we set  $r = \eta|\lambda|^{-1/2}$  to obtain

$$\|F\|_{L_H^q L_z^p(\mathbb{R}^3)} \leq 27M^2 \left( \|f\|_{L_H^q L_z^p(\Omega)} + 2\eta^{-1} |\lambda|^{1/2} \|\nabla v\|_{L_H^q L_z^p(\Omega)} + \eta^{-2} |\lambda| \cdot \|v\|_{L_H^q L_z^p(\Omega)} \right).$$

Now assume that  $\eta > 0$  is sufficiently large enough such that  $54C_\theta M^2 \eta^{-1} < 1/2$ ,  $27C_\theta M^2 \eta^{-2} < 1/2$  and then assume that  $\lambda_0 > 0$  is large enough such that  $4\eta\lambda_0^{-1/2} < 3 \min\{1, h\}$ . This and the fact that  $u$  is an extension of  $v$  then yields

$$|\lambda| \cdot \|v\|_{L_H^q L_z^p(\Omega)} + |\lambda|^{1/2} \|\nabla v\|_{L_H^q L_z^p(\Omega)} \leq 54C_\theta M^2 \|f\|_{L_H^q L_z^p(\Omega)} \quad \text{for } |\lambda| \geq \lambda_0.$$

In the case  $q = \infty$ ,  $p \in (2, \infty)$  we obtain the estimate for the full range of  $\lambda \in \Sigma_\theta$  by setting  $\lambda_1 := \frac{\lambda_0}{|\lambda|}$  for  $0 < |\lambda| < \lambda_0$ . Then  $f \in L_H^\infty L_z^p(\Omega) \hookrightarrow L^p(\Omega)$  yields

$$|\lambda| \cdot \|v\|_{L^p(\Omega)} + |\lambda|^{1/2} \|\nabla v\|_{L^p(\Omega)} + \|\Delta v\|_{L^p(\Omega)} \leq C_{\theta,p} \|f\|_{L^p(\Omega)}$$

by (5.9) and since  $\lambda_1 v - \Delta v = f + (\lambda_1 - \lambda)v$  we obtain

$$|\lambda_1| \cdot \|v\|_{L_H^\infty L_z^p(\Omega)} + |\lambda_1|^{1/2} \|\nabla v\|_{L_H^\infty L_z^p(\Omega)} + \|\Delta v\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p} \left( \|f\|_{L_H^\infty L_z^p(\Omega)} + |\lambda_1 - \lambda| \cdot \|v\|_{L_H^\infty L_z^p(\Omega)} \right)$$

where we can further estimate  $|\lambda_1 - \lambda| < \lambda_0$ , and  $p \in (1, \infty)$  yields

$$\|v\|_{L_H^\infty L_z^p(\Omega)} \leq C_p \|v\|_{W_H^{2,p} L_z^p(\Omega)} \leq C_p \|v\|_{W^{2,p}(\Omega)} \leq C_p \|\Delta v\|_{L^p(\Omega)} \leq C_p \|f\|_{L^p(\Omega)} \leq C_p \|f\|_{L_H^\infty L_z^p(\Omega)}$$

where we used  $W^{2,p}(G) \hookrightarrow L^\infty(G)$  and that  $\Delta_p$  is invertible on  $L^p(\Omega)$ . Since  $|\lambda_1| = \lambda_0 > |\lambda|$ , this yields the desired result for the full range of  $\lambda \in \Sigma_\theta$ ,  $\theta \in (0, \pi)$ .

If  $v$  instead solves problem (5.6) with  $\partial_i = \partial_z$  then  $u$  solves the problem

$$\lambda u - \Delta u = G \text{ on } \mathbb{R}^3, \quad G := \chi_r E(\partial_z f) - 2(\nabla \chi_r) \cdot E(\nabla v) - (\Delta \chi_r) E v.$$

We rewrite

$$-2(\nabla \chi_r) \cdot E(\nabla v) - (\Delta \chi_r) E v = -2\operatorname{div}(\nabla \chi_r E v) + (\Delta \chi_r) E v, \quad \chi_r E(\partial_z f) = \partial_z(\chi_r s E f) - (\partial_z \chi_r) s E f$$

where

$$s(z) = \begin{cases} 1, & z \in (-2h, 0), \\ -1, & x \in (0, h). \end{cases}$$

Here, by the density argument above, we may assume  $f = 0$  on  $\Gamma_u \cup \Gamma_b$ . This yields  $u = u_1 + u_2$  where

$$\begin{aligned} \lambda u_1 - \Delta u_1 &= \partial_z G_1 + \operatorname{div}_H G_2 && \text{on } \mathbb{R}^3, \quad G_1 := \chi_r s E f, \quad G_2 := -2(\nabla \chi_r) E v, \\ \lambda u_2 - \Delta u_2 &= G_3 && \text{on } \mathbb{R}^3, \quad G_3 := -(\partial_z \chi_r) s E f + (\Delta \chi_r) E v. \end{aligned}$$

Since  $G_i$ ,  $i \in \{1, 2, 3\}$ , are bounded and have compact support, we may apply Lemma 5.2 to obtain the estimate

$$|\lambda|^{1/2} \|u\|_{L_H^q L_z^p(\mathbb{R}^3)} \leq C_\theta \left( \|G_1\|_{L_H^q L_z^p(\mathbb{R}^3)} + \|G_2\|_{L_H^q L_z^p(\mathbb{R}^3)} + |\lambda|^{-1/2} \|G_3\|_{L_H^q L_z^p(\mathbb{R}^3)} \right).$$

Proceeding as above we obtain

$$\begin{aligned} \|G_1\|_{L_H^q L_z^p(\mathbb{R}^3)} &\leq 27M^2 \|f\|_{L_H^q L_z^p(\Omega)}, \\ \|G_2\|_{L_H^q L_z^p(\mathbb{R}^3)} &\leq 54M^2 \eta^{-1} |\lambda|^{1/2} \|v\|_{L_H^\infty L_z^p(\Omega)}, \\ \|G_3\|_{L_H^\infty L_z^p(\mathbb{R}^3)} &\leq 27M^2 \eta^{-1} |\lambda|^{1/2} \|f\|_{L_H^\infty L_z^p(\Omega)} + 27M^2 \eta^{-2} |\lambda| \cdot \|v\|_{L_H^\infty L_z^p(\Omega)}. \end{aligned}$$

The above assumptions on  $\eta$  and  $\lambda_0$  then yield the desired result for  $|\lambda| > \lambda_0$ . The case  $\partial_i \in \{\partial_x, \partial_y\}$  is analogous where for  $f \in L^\infty(G; C_c^\infty(-h, 0))$  horizontal derivatives are understood in the sense of distributions, and otherwise derivatives can be treated using smooth approximations as above.

For the case  $q = \infty$  and  $p \in (2, \infty)$ , to extend this estimate to the full range of  $\lambda \in \Sigma_\theta$  one proceeds as above to obtain

$$|\lambda|^{1/2} \cdot \|v\|_{L^p(\Omega)} + \|\nabla v\|_{L^p(\Omega)} \leq C_{\theta,p} \|f\|_{L^p(\Omega)}$$

from (5.11), as well as

$$|\lambda_1|^{1/2} \cdot \|v\|_{L_H^\infty L_z^p(\Omega)} + \|\nabla v\|_{L_H^\infty L_z^p(\Omega)} \leq C_\theta \left( \|f\|_{L_H^\infty L_z^p(\Omega)} + |\lambda_1|^{-1/2} |\lambda_1 - \lambda| \cdot \|v\|_{L_H^\infty L_z^p(\Omega)} \right).$$

Since we have  $|\lambda_1|^{-1/2} |\lambda_1 - \lambda| \leq \lambda_0^{1/2}$  and  $p \in (2, \infty)$  this yields

$$\|v\|_{L_H^\infty L_z^p(\Omega)} \leq C_p \|v\|_{W_H^{1,p} L_z^p(\Omega)} \leq C_p \|v\|_{W^{1,p}(\Omega)} \leq C_p \|\nabla v\|_{L^p(\Omega)} \leq C_p \|f\|_{L^p(\Omega)} \leq C_p \|f\|_{L_H^\infty L_z^p(\Omega)},$$

where we used the embedding  $W^{1,p}(G) \hookrightarrow L^\infty(G)$  and the Poincaré inequality  $\|v\|_{L^p(\Omega)} \leq C_p \|\nabla v\|_{L^p(\Omega)}$  for  $v$  with  $v|_{\Gamma_b} = 0$ .  $\square$

**Remark 5.5.** The results of Lemma 5.4 also hold true if the condition  $\partial_z v|_{\Gamma_u} = 0$  is replaced by  $v|_{\Gamma_u} = 0$  or if  $L_H^q L_z^p(\Omega)$  is replaced by  $C_{\text{per}}([0, 1]^2; L^p(-h, 0))$ . For pure Dirichlet boundary conditions one extends by an odd reflexion at both  $z = 0$  and  $z = -h$  replacing  $E_z^{\text{even, odd}}$  by  $E_z^{\text{odd, odd}}$  and setting  $s(z) \equiv 1$  in the proof.

Since  $\Omega = G \times (-h, 0)$  is a cylindrical domain the semigroup generated by the Laplacian with the above boundary conditions satisfies

$$e^{t\Delta}(f \otimes g) = e^{t\Delta_H} f \otimes e^{t\Delta_z} g, \quad f : G \rightarrow \mathbb{R}^2, \quad g : (-h, 0) \rightarrow \mathbb{R},$$

where  $(f \otimes g)(x, y, z) := f(x, y)g(z)$  is an elementary tensor,  $\Delta_H := \partial_x^2 + \partial_y^2$  is the Laplacian on  $G$  with periodic boundary conditions and  $\Delta_z$  is defined by

$$\Delta_z v := \partial_z^2 v, \quad D(\Delta_z) = \{f \in W^{2,p}(-h, 0) : f(-h) = \partial_z f(0) = 0\}.$$

We now investigate these operators separately, starting with the vertical one, cf. [6, 21].

**Lemma 5.6.** *Let  $p \in (1, \infty)$ . Then the operator  $\Delta_z$  generates a strongly continuous, exponentially stable, analytic semigroup on  $L^p(-h, 0)$ .*

**Lemma 5.7.** *Let  $\theta \in (0, \pi/2)$ . Then there exists a constant  $C_\theta > 0$  such that for all  $\tau \in \Sigma_\theta$  we have*

$$|\tau|^{1/2} \|\nabla_H e^{\tau\Delta_H} Qf\|_{L^\infty(G)} \leq C_\theta \|f\|_{L^\infty(G)}, \quad f \in L^\infty(G).$$

**Remark 5.8.** Note that although the two-dimensional Helmholtz projector with periodic boundary conditions  $Q$  is unbounded on  $L^\infty(G)$ , the composition  $\nabla_H e^{\tau\Delta_H} Q$  defines a bounded operator for  $\tau \in \Sigma_\theta$ .

*Proof of Lemma 5.7.* Let  $Q_{\mathbb{R}^2}$  and  $Q$  be the Helmholtz projection on  $\mathbb{R}^2$  and  $\mathbb{T}^2$ , respectively, and  $E_H^{\text{per}}$  be the periodic extension operator from  $G$  onto  $\mathbb{R}^2$ . Then  $E_H^{\text{per}} Qf = Q_{\mathbb{R}^2} E_H^{\text{per}} f$  for all  $f : G \rightarrow \mathbb{R}^2$  and

$$E_H^{\text{per}} |\tau|^{1/2} \nabla_H e^{\tau\Delta_H} Qf = |\tau|^{1/2} \nabla_H e^{\tau\Delta_H} E_H^{\text{per}} Qf = |\tau|^{1/2} \nabla_H e^{\tau\Delta_H} Q_{\mathbb{R}^2} E_H^{\text{per}} f.$$

Since  $\|E_H^{\text{per}} f\|_{L^\infty(\mathbb{R}^2)} = \|f\|_{L^\infty(G)}$  it therefore suffices to consider the operator  $\Delta_H$  on the full space  $\mathbb{R}^2$ . Recall that  $1 - Q_{\mathbb{R}^2}$  is given by  $(R_j R_k)_{1 \leq j, k \leq 2}$  where  $R_j$  is the Riesz transform in the  $j$ -th direction. We therefore investigate the family of Fourier multipliers

$$m_{\tau, j, k, l}(\xi) = \begin{cases} |\tau|^{1/2} \xi_l \left( \delta_{j, k} - \frac{\xi_j \xi_k}{|\xi|^2} \right) e^{-\tau|\xi|^2}, & \xi \in \mathbb{R}^2 \setminus \{0\}, \\ 0, & \xi = 0, \end{cases} \quad \text{for } 1 \leq j, k, l \leq 2.$$

Using the invariance under rescaling and replacing  $\xi$  with  $|\tau|^{-1/2} \xi$ , we may assume that  $\tau = e^{i\psi}$  where  $|\psi| < \theta$ . We show that for each of these symbols we have  $m = \hat{g}$  for some  $g \in L^1(\mathbb{R}^2)$  such that  $\|g\|_{L^1(\mathbb{R}^2)} \leq C_\theta$ . The desired estimate then follows from Young's inequality. Since this family of symbols belongs to  $C(\mathbb{R}^2) \cap C^\infty(\mathbb{R}^2 \setminus \{0\})$  we verify the Mihlin condition

$$(5.12) \quad \max_{|\alpha| \leq 2} \sup_{\xi \in \mathbb{R}^2 \setminus \{0\}} |\xi|^{|\alpha| + \delta} |D^\alpha m(\xi)| < M < \infty,$$

for some  $\delta > 0$ . Elementary calculations using the homogeneity of the first factor show that for an arbitrary multi-index  $\alpha \in \mathbb{N}^2$  we have

$$\sup_{\xi \in \mathbb{R}^2 \setminus \{0\}} |\xi|^\alpha \left| D^\alpha \frac{\xi_j \xi_k}{|\xi|^2} \right| < M_\alpha < \infty, \quad \sup_{\xi \in \mathbb{R}^2 \setminus \{0\}} |\xi|^{\alpha+\delta} \left| D^\alpha \xi_l e^{-e^{i\psi} |\xi|^2} \right| < M_{\alpha,\delta,\psi} \leq M_{\alpha,\delta,\theta} < \infty$$

for  $\delta \in (0, 1)$  which together with the product rule yield that (5.12) is satisfied. Analogously we verify the condition

$$(5.13) \quad |\xi|^{|\alpha|} |D^\alpha m(\xi)| \leq C_\alpha |\xi|, \quad |\xi| \leq 1, \xi \neq 0$$

for  $0 < |\alpha| \leq 2$  by noting that

$$|\xi|^{|\alpha|} \left| D^\alpha \frac{\xi_j \xi_k \xi_l}{|\xi|^2} \right| \leq C_\alpha |\xi|, \quad |\xi| \leq 1, \xi \neq 0$$

and

$$|\xi|^{|\alpha|} \left| D^\alpha e^{-e^{i\psi} |\xi|^2} \right| + |\xi|^{|\alpha|} \left| D^\alpha \xi_l e^{-e^{i\psi} |\xi|^2} \right| \leq C_{\alpha,\delta,\psi} \leq C_{\alpha,\delta,\theta} \quad |\xi| \leq 1, \xi \neq 0.$$

We now split the symbol into  $m = \varphi m + (1 - \varphi)m$  where  $\varphi \in C_c^\infty(\mathbb{R}^2)$  is a cut-off function satisfying  $\varphi(\xi) = 1$  for  $|\xi| \leq 2$ . Applying [4, Lemma 8.2.3 and 8.2.4] to the terms  $(1 - \varphi)m$  and  $\varphi m$  respectively then yields the desired results.  $\square$

## 6. LINEAR ESTIMATES FOR THE HYDROSTATIC STOKES OPERATOR: PART 1

A key element in the proof of our global existence results are the estimates for the hydrostatic Stokes semigroup in  $X_{\bar{\sigma}}$ . To this end, we prove first estimates in the larger space  $X$ , where we make use of representation (1.1). We thus define the operator  $A$  by

$$Av := \mathcal{A}v, \quad D(A) = \{v \in W_{\text{per}}^{2,p}(\Omega)^2 \cap X : \partial_z v|_{\Gamma_u} = 0, v|_{\Gamma_b} = 0, \mathcal{A}v \in X\}.$$

It is the aim of this section to prove the following claim.

**Claim 6.1.** *Let  $p \in (3, \infty)$ . Then*

- a)  *$A$  is the generator of a strongly continuous, analytic semigroup on  $X$ .*
- b) *There exist constants  $C > 0$ ,  $\beta \in \mathbb{R}$  such that for  $\partial_i \in \{\partial_x, \partial_y, \partial_z\}$ ,  $t > 0$  and  $f \in X$  one has that*

$$(i) \quad t^{1/2} \|\partial_i e^{tA} f\|_{L_H^\infty L_x^p(\Omega)} \leq C e^{\beta t} \|f\|_{L_H^\infty L_x^p(\Omega)}, \quad t > 0, f \in X,$$

for  $\partial_j \in \{\partial_x, \partial_y\}$

$$(ii) \quad t^{1/2} \|\partial_j e^{tA} \mathbb{P}f\|_{L_H^\infty L_x^p(\Omega)} \leq C e^{\beta t} \|f\|_{L_H^\infty L_x^p(\Omega)}, \quad t > 0, f \in X,$$

$$(iii) \quad t^{1/2} \|e^{tA} \mathbb{P}\partial_j f\|_{L_H^\infty L_x^p(\Omega)} \leq C e^{\beta t} \|f\|_{L_H^\infty L_x^p(\Omega)}, \quad t > 0, f \in X,$$

$$(iv) \quad t \|\partial_i e^{tA} \mathbb{P}\partial_j f\|_{L_H^\infty L_x^p(\Omega)} \leq C e^{\beta t} \|f\|_{L_H^\infty L_x^p(\Omega)}, \quad t > 0, f \in X.$$

c)  *$X_{\bar{\sigma}}$  is an invariant subspace of  $A$ , and its restriction is  $A_{\bar{\sigma}}$ . The semigroup  $e^{tA}$  restricts to an exponentially stable, strongly continuous, analytic semigroup of angle  $\pi/2$  on  $X_{\bar{\sigma}}$ .*

d) *Furthermore, for all  $v \in X_{\bar{\sigma}}$*

$$\lim_{t \rightarrow 0^+} t^{1/2} \|\nabla e^{tA} v\|_{L_H^\infty L_x^p(\Omega)} = 0.$$

In order to solve equation (2.5) in  $X_{\bar{\sigma}}$ , we collect first several facts concerning the corresponding theory in  $L_{\bar{\sigma}}^p(\Omega)$ . To this end, let  $p \in (1, \infty)$ , and define  $A_{p,\bar{\sigma}}: D(A_{p,\bar{\sigma}}) \rightarrow L_{\bar{\sigma}}^p(\Omega)$  by

$$A_{p,\bar{\sigma}}v := \mathbb{P}\Delta v, \quad D(A_{p,\bar{\sigma}}) = \{v \in W_{\text{per}}^{2,p}(\Omega)^2 : \text{div}_H \bar{v} = 0, \partial_z v|_{\Gamma_u} = 0, v|_{\Gamma_b} = 0.\}.$$

Consider furthermore  $A_p: D(A_p) \rightarrow L^p(\Omega)^2$  defined by

$$A_p v := \Delta_p v + Bv, \quad D(A_p) := D(\Delta_p)^2, \quad Bv := \frac{1}{h}(1 - Q)\partial_z v|_{\Gamma_b},$$

where  $\Delta_p$  denotes the Laplacian in  $L^p(\Omega)^2$  as in the last section. By [7], the operator  $A_p$  is an extension of  $A_{p,\bar{\sigma}}$ . The idea is that the pressure term may be recovered by applying the vertical average and horizontal divergence to (2.5), yielding

$$(6.1) \quad \Delta_H \pi = \operatorname{div}_H \bar{f} - \operatorname{div}_H \frac{1}{h} \partial_z v \Big|_{\Gamma_b},$$

or equivalently since  $1 - Q$  agrees with  $\nabla_H(-\Delta_H)^{-1} \operatorname{div}_H$  one has  $\nabla_H \pi = (1 - Q)\bar{f} - Bv$ .

Note that the following inclusions hold

$$(6.2) \quad A \subset A_p \quad \text{and} \quad A_{\bar{\sigma}} \subset A_{p,\bar{\sigma}},$$

and that  $e^{tA_{p,\bar{\sigma}}}$ ,  $e^{tA_p}$ ,  $e^{tA}$  and  $e^{tA_{\bar{\sigma}}}$  are consistent semigroups.

*Proof of Claim 6.1.* Let  $\lambda_0 > 0$  with  $\lambda_0 \in \rho(A_p)$ ,  $\theta \in (0, \pi/2)$ , and

$$\lambda \in \Sigma_{\theta+\pi/2} \cap B_{\lambda_0}(0)^c \subset \rho(A_p).$$

By (6.2) it follows that  $\lambda - A$  is injective for  $\lambda \in \rho(A_p)$  and likewise  $\lambda - A_{\bar{\sigma}}$  is injective for  $\lambda \in \rho(A_{\bar{\sigma}})$ . Since  $X \hookrightarrow L^p(\Omega)^2$  the existence of a unique  $v \in D(A_p)$  for  $p \in (1, \infty)$  follows from the  $L^p$ -theory for  $A_p$ , cf. [7], and since  $W_{\text{per}}^{2,p}(\Omega)^2 \hookrightarrow X$  for  $p \in (3/2, \infty)$  it follows that  $v \in D(A)$ . Since  $(A_p - \lambda)^{-1}$  further leaves  $L_{\bar{\sigma}}^p(\Omega)$  invariant,  $f \in X_{\bar{\sigma}}$  implies  $v \in D(A_{\bar{\sigma}})$ . Hence,

$$(6.3) \quad \rho(A_p) \subset \rho(A) \quad \text{and} \quad \rho(A_{p,\bar{\sigma}}) \subset \rho(A_{\bar{\sigma}}).$$

In particular the resolvent sets are non-empty and thus the operators are closed.

Since the semigroup estimates follow from resolvent estimates by arguments involving the inverse Laplace transform, it now remains to prove suitable resolvent estimates in  $X$ . To this end we observe first that  $v = (\lambda - A)^{-1}f$  is equivalent to

$$(6.4) \quad v = (\lambda - \Delta_p)^{-1}(f + Bv),$$

and second, using the fact that  $Q$  is continuous on  $C_{\text{per}}^{0,\alpha}([0, 1]^2)$  for  $\alpha \in (0, 1)$ , that

$$\|Bv\|_{L_{\bar{H}}^\infty L_z^p(\Omega)} \leq h^{1/p} \|Bv\|_{L^\infty(\Omega)} \leq h^{1/p} \|Bv\|_{C^{0,\alpha}([0,1]^2)} \leq C \|\partial_z v\|_{\Gamma_b} \|C^{0,\alpha}([0,1]^2)\| \leq C \|v\|_{C^{1,\alpha}(\bar{\Omega})}.$$

Assuming  $p \in (3, \infty)$  we have  $W^{2,p}(\Omega) \hookrightarrow C^{1,\alpha}(\bar{\Omega})$  for some  $\alpha = \alpha_p \in (0, 1 - 3/p)$ . Using the resolvent estimate for  $A_p$  in  $L^p(\Omega)^2$  we obtain

$$\|v\|_{C^{1,\alpha}(\bar{\Omega})} \leq C_p \|v\|_{W^{2,p}(\Omega)} \leq C_p (\|v\|_{L^p(\Omega)} + \|Av\|_{L^p(\Omega)}) \leq C_p (1 + |\lambda|^{-1}) \|f\|_{L^p(\Omega)}.$$

This and  $|\lambda| > \lambda_0$  yield  $\|Bv\|_{L_{\bar{H}}^\infty L_z^p(\Omega)} \leq C_p (1 + \lambda_0^{-1}) \|f\|_{L_{\bar{H}}^\infty L_z^p(\Omega)}$ . So, using Lemma 5.4 we obtain

$$(6.5) \quad |\lambda| \cdot \|v\|_{L_{\bar{H}}^\infty L_z^p(\Omega)} + |\lambda|^{1/2} \|\nabla v\|_{L_{\bar{H}}^\infty L_z^p(\Omega)} + \|Av\|_{L_{\bar{H}}^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0} \|f\|_{L_{\bar{H}}^\infty L_z^p(\Omega)},$$

where we used that for  $\lambda$  as above and  $p \in (3, \infty)$  one has

$$\|Av\|_{L_{\bar{H}}^\infty L_z^p(\Omega)} \leq \|\Delta v\|_{L_{\bar{H}}^\infty L_z^p(\Omega)} + \|Bv\|_{L_{\bar{H}}^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0} \|f\|_{L_{\bar{H}}^\infty L_z^p(\Omega)}.$$

Note that if one instead considers  $f \in X_{\bar{\sigma}}$ , then  $\lambda_0 > 0$  can be taken to be arbitrarily small and  $\theta$  arbitrarily close to  $\pi/2$  by [7, Theorem 3.1]. Since  $0 \in \rho(A_{p,\bar{\sigma}}) \subset \rho(A_{\bar{\sigma}})$ , compare [11, Theorem 3.1] and (6.3) it follows that the spectral bound

$$\beta := \sup\{\operatorname{Re}(\lambda) : \lambda \in \sigma(A_{\bar{\sigma}})\}$$

is negative implying exponential decay, and estimate (6.5) is valid for all  $\lambda \in \Sigma_\theta$ ,  $\theta \in (0, \pi)$  and  $f \in X_{\bar{\sigma}}$ .

To verify that  $D(A)$  and  $D(A_{\bar{\sigma}})$  are dense in  $X$  and  $X_{\bar{\sigma}}$  respectively, observe that the space

$$C_{\text{per}}^\infty([0, 1]^2; C_c^\infty((-h, 0)))^2$$

is contained in  $D(A)$  and dense in  $X$ , so the semigroup generated by  $A$  is strongly continuous on  $X$ . Since it leaves  $L_{\bar{\sigma}}^p(\Omega)$  invariant, the restriction of the semigroup on  $X \cap L_{\bar{\sigma}}^p(\Omega) = X_{\bar{\sigma}}$  is strongly continuous as well and generated by the restriction of  $A$  onto  $D(A) \cap L_{\bar{\sigma}}^p(\Omega) = D(A_{\bar{\sigma}})$ , i.e.  $A_{\bar{\sigma}}$ , which is therefore densely defined on  $X_{\bar{\sigma}}$ . Thus we have proven a), c) and estimate (i) in b).

To prove the remaining semigroup estimates in *b*) we consider the corresponding resolvent estimates. Since  $X \hookrightarrow L^p(\Omega)^2$  and  $\mathbb{P}$  is bounded on  $L^p(\Omega)^2$  the existence of

$$v := (\lambda - A_{p,\bar{\sigma}})^{-1}\mathbb{P}f \in D(A_{p,\bar{\sigma}}) \hookrightarrow W_{\text{per}}^{2,p}(\Omega)^2 \hookrightarrow X$$

for  $f \in X$  follows from the  $L^p$ -theory for  $A_{p,\bar{\sigma}}$ , and it suffices to extend the  $L^p$ -estimate

$$(6.6) \quad |\lambda|^{1/2} \|\partial_i(\lambda - A_{p,\bar{\sigma}})^{-1}f\|_{L^p(\Omega)} + |\lambda|^{1/2} \|(\lambda - A_{p,\bar{\sigma}})^{-1}\partial_i f\|_{L^p(\Omega)} \leq C_{\theta,p} \|f\|_{L^p(\Omega)}, \quad f \in L_{\bar{\sigma}}^p(\Omega),$$

where  $\partial_i \in \{\partial_x, \partial_y\}$ ,  $\theta \in (0, \pi)$ ,  $C_{\theta,p} > 0$ , to  $X$ , i.e. to prove the estimate

$$(6.7) \quad |\lambda|^{1/2} \|\nabla_H v\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p} \|f\|_{L_H^\infty L_z^p(\Omega)}, \quad \lambda \in \Sigma_\theta.$$

Recall that  $\mathbb{P}f = f - (1 - Q)\bar{f} = \tilde{f} + Q\bar{f}$ , and that if  $f \in X$  then  $\bar{f} \in C_{\text{per}}([0, 1]^2)^2$  satisfies  $\|\bar{f}\|_\infty \leq C\|f\|_{L_H^\infty L_z^p(\Omega)}$  for any  $p \in [1, \infty]$ . Using (6.4) we rewrite

$$v = (\lambda - A_{\bar{\sigma}})^{-1}\mathbb{P}f = (\lambda - \Delta)^{-1}(\tilde{f} + Bv + Q\bar{f}),$$

and since the term  $\tilde{f} + Bv$  can be dealt with as before, it suffices to show the estimate

$$(6.8) \quad |\lambda|^{1/2} \|\nabla_H(\lambda - \Delta)^{-1}Q\bar{f}\|_{L_H^\infty L_z^p(\Omega)} \leq C_\theta \|\bar{f}\|_{L^\infty(G)}.$$

Since  $Q\bar{f}$  does not depend on  $z$  we can write  $Q\bar{f} = Q\bar{f} \otimes 1$ , and so for  $\lambda = |\lambda|e^{i\psi}$  with  $\psi \in (-\pi/2 + \varepsilon, \pi/2 - \varepsilon)$  for small  $\varepsilon > 0$  we have

$$|\lambda|^{1/2} \nabla_H(\lambda - \Delta)^{-1}(Q\bar{f} \otimes 1) = |\lambda|^{1/2} \int_0^\infty e^{-\lambda t} (\nabla_H e^{t\Delta_H} Q\bar{f} \otimes e^{t\Delta_z} 1) dt,$$

where  $e^{t\Delta_z}$  denotes the semigroup from Lemma 5.6. Applying the estimates in Lemma 5.7 and 5.6 yields

$$\begin{aligned} |\lambda|^{1/2} \|\nabla_H(\lambda - \Delta)^{-1}(Q\bar{f} \otimes 1)\|_{L_H^\infty L_z^p(\Omega)} &\leq |\lambda|^{1/2} \int_0^\infty e^{-\lambda t} \|\nabla_H e^{t\Delta_H} Q\bar{f}\|_{L^\infty(G)} \|e^{t\Delta_z} 1\|_{L^p(-h,0)} dt \\ &\leq C|\lambda|^{1/2} \left( \int_0^\infty e^{-|\lambda| \cos(\psi)t} t^{-1/2} dt \right) \|\bar{f}\|_{L^\infty(G)} \\ &\leq C \frac{\sqrt{\pi}}{\sqrt{\cos(\pi/2 - \varepsilon)}} \|\bar{f}\|_{L^\infty(G)}. \end{aligned}$$

To include the full range of angles  $\psi$  one simply replaces  $\Delta_H$  and  $\Delta_z$  with  $e^{i\theta}\Delta_H$  and  $e^{i\theta}\Delta_z$  respectively where  $\theta \in (-\pi/2, \pi/2)$  is a suitable angle.

Since an elementary calculation shows that  $\nabla_H$  commutes with  $A$  and  $\mathbb{P}$  we obtain

$$\partial_i(\lambda - A)^{-1}f = (\lambda - A)^{-1}\partial_i f, \quad \partial_i(\lambda - A)^{-1}\mathbb{P}f = (\lambda - A)^{-1}\mathbb{P}\partial_i f$$

for horizontal derivatives  $\partial_i \in \{\partial_x, \partial_y\}$  and  $f \in C_{\text{per}}^\infty([0, 1]^2; C_c^\infty[-h, 0])^2$ . Note that for any  $v \in W_{\text{per}}^{2,p}(\Omega)$  the horizontal derivatives  $\partial_x v$  and  $\partial_y v$  are periodic on  $\Gamma_l$  as well. This yields suitable estimates for the right-hand sides.

To verify *d*), we first make use of the density of the domains of the generators. So, let  $\varepsilon > 0$  and  $v' \in D(A_{\bar{\sigma}})$  such that  $\|v - v'\|_{L_H^\infty L_z^p(\Omega)} < \varepsilon/2C_0$ . By *b*) (*i*) we have

$$t^{1/2} \|\nabla e^{tA} v\|_{L_H^\infty L_z^p(\Omega)} \leq C_0 \|v\|_{L_H^\infty L_z^p(\Omega)}$$

for all  $v \in X$  and  $t > 0$ . Then

$$t^{1/2} \|\nabla e^{tA} v\|_{L_H^\infty L_z^p(\Omega)} \leq \frac{\varepsilon}{2} + t^{1/2} \|\nabla e^{tA} v'\|_{L_H^\infty L_z^p(\Omega)}$$

and we can further estimate

$$\|\nabla e^{tA} v'\|_{L_H^\infty L_z^p(\Omega)} \leq h^{1/p} \|e^{tA} v'\|_{C^1(\bar{\Omega})} \leq C_p \|e^{tA} v'\|_{D(A_{p,\bar{\sigma}})}.$$

This and the invertibility of  $A_{p,\bar{\sigma}}$  on  $L_{\bar{\sigma}}^p(\Omega)$  yield

$$t^{1/2} \|\nabla e^{tA_{p,\bar{\sigma}}} v'\|_{L_{\bar{\sigma}}^p(\Omega)} \leq C_p t^{1/2} \|A_{p,\bar{\sigma}} e^{tA_{p,\bar{\sigma}}} v'\|_{L_{\bar{\sigma}}^p(\Omega)} = C_p t^{1/2} \|e^{tA_{p,\bar{\sigma}}} A_{p,\bar{\sigma}} v'\|_{L_{\bar{\sigma}}^p(\Omega)} \leq C_p t^{1/2} \|A_{p,\bar{\sigma}} v'\|_{L_{\bar{\sigma}}^p(\Omega)}$$

and since  $A_{p,\bar{\sigma}} v' \in L_{\bar{\sigma}}^p(\Omega)$  the claim follows.  $\square$

## 7. LINEAR ESTIMATES FOR THE HYDROSTATIC STOKES OPERATOR: PART 2

This section is devoted to prove that the estimates of Claim 6.1 in the case of vertical derivatives, i.e. that the estimates (ii), (iii) and (iv) in Claim 6.1 are valid even for  $\partial_j = \partial_z$ .

**Claim 7.1.** *Under the assumptions of Claim 6.1 there exist constants  $C > 0$  and  $\beta \in \mathbb{R}$  such that*

$$(7.1) \quad t^{1/2} \|\partial_z e^{tA} \mathbb{P} f\|_{L_H^\infty L_z^p(\Omega)} \leq C e^{t\beta} \|f\|_{L_H^\infty L_z^p(\Omega)},$$

$$(7.2) \quad t^{1/2} \|e^{tA} \mathbb{P} \partial_z f\|_{L_H^\infty L_z^p(\Omega)} \leq C e^{t\beta} \|f\|_{L_H^\infty L_z^p(\Omega)},$$

$$(7.3) \quad t \|\partial_i e^{tA} \mathbb{P} \partial_j f\|_{L_H^\infty L_z^p(\Omega)} \leq C e^{t\beta} \|f\|_{L_H^\infty L_z^p(\Omega)},$$

where  $\partial_i, \partial_j \in \{\partial_x, \partial_y, \partial_z\}$ , for all  $t > 0$  and  $f \in X$ .

As in the last section, these semigroup estimates follow from suitable resolvent estimates and standard arguments involving the inverse Laplace transform.

Before investigating the estimate for  $\partial_z(\lambda - A)\mathbb{P}$  we present an anisotropic version of an interpolation inequality. We use the notation  $(x, y, z) =: (x', z)$  and let  $B(x'_0; r) = \{x' \in \mathbb{R}^2 : |x' - x'_0| < r\}$  denote a disk in  $\mathbb{R}^2$ .

**Lemma 7.2.** *Let  $p \in (2, \infty)$ ,  $q \in [1, \infty]$ ,  $r > 0$ , and  $x'_0 \in \mathbb{R}^2$ . Then, for  $v \in W^{1,p}(B(x'_0; r); L_z^q)$ ,  $L_z^q = L^q(-h, 0)$  we have*

$$\|v\|_{L^\infty(B(x'_0; r); L_z^q)} \leq C r^{-2/p} (\|v\|_{L^p(B(x'_0; r); L_z^q)} + r \|\nabla_H v\|_{L^p(B(x'_0; r); L_z^q)}),$$

where the constant  $C = C_{\Omega, p, q} > 0$  is independent of  $r$  and  $x'_0$ .

*Proof.* We put  $w(x') := (\int_{-h}^0 |v(x', z)|^q dz)^{1/q}$  and apply a two-dimensional interpolation inequality, compare [20, Lemma 3.1.4] to have

$$(7.4) \quad \|w\|_{L^\infty(B(x'_0; r))} \leq C r^{-2/p} (\|w\|_{L^p(B(x'_0; r))} + r \|\nabla_H w\|_{L^p(B(x'_0; r))}).$$

One sees that  $\|w\|_{L^p(B(x'_0; r))} = \|v\|_{L^p(B(x'_0; r); L_z^q)}$ . To estimate the second term we compute  $\partial_i w$  for  $\partial_i \in \{\partial_x, \partial_y\}$  as follows:

$$\partial_i w(x') = \left( \int_{-h}^0 |v(x', z)|^q dz \right)^{1/q-1} \int_{-h}^0 |v(x', z)|^{q-2} (\partial_i v(x', z) \cdot v(x', z)) dz.$$

Using Hölder's inequality we obtain

$$|\partial_i w(x')| \leq \left( \int_{-h}^0 |v(x', z)|^q dz \right)^{1/q-1} \int_{-h}^0 |v(x', z)|^{q-1} |\partial_i v(x', z)| dz \leq \left( \int_{-h}^0 |\partial_i v(x', z)|^q dz \right)^{1/q}$$

and substituting this into (7.4) proves the estimate for  $q < \infty$ . The case  $q = \infty$  is a straightforward result of (7.4).  $\square$

It is well known that  $1 - Q = -\nabla_H(-\Delta_H)^{-1} \operatorname{div}_H = \nabla_H \Delta_H^{-1} \operatorname{div}_H$  with periodic boundary conditions is a singular integral operator which fails to be bounded in  $L^\infty(G)^2$ . However, if one allows for a logarithmic (and therefore divergent) factor, some  $L^\infty$ -type estimate are still available. In this spirit we give a local  $L^p$ -estimate for the operator  $\nabla_H(-\Delta_H)^{-1} \operatorname{div}_H$  corresponding to the scale of the  $L^\infty$ -norm.

**Proposition 7.3.** *Let  $p \in (1, \infty)$ ,  $x'_0 \in G$ . Then there exists  $r_0 > 0$  such that for all  $r \in (0, r_0)$  the weak solution of*

$$(7.5) \quad \Delta_H \pi = \operatorname{div}_H F \quad \text{in } G, \quad \pi|_{\partial G} : \text{periodic}, \quad \int_G \pi dx' = 0,$$

for  $F \in L^\infty(G)^2$  satisfies

$$\|\nabla_H \pi\|_{L^p(B(x'_0; r))} \leq C r^{2/p} (1 + |\log r|) \|F\|_{L^\infty(G)}.$$

Here the constant  $C = C_{G, p} > 0$  is independent of  $x'_0$  and  $r$ .

*Proof.* By applying a periodic extension we may assume that (7.5) holds in a larger square  $G' := (-2, 3)^2$ . We choose  $r_0 < 1/8$  to obtain  $B(x'_0; 4r_0) \subset (-1/2, 3/2)^2$  and utilize two cut-off functions  $\omega, \theta \in C_c^\infty(\mathbb{R}^2)$ ,  $\theta = \theta_r$ , satisfying the following properties:

$$\begin{aligned} \omega &\equiv 1 \text{ on } [-1, 2]^2, & \text{supp}(\omega) &\subset G', & \|\nabla_H^k \omega\|_{L^\infty(\mathbb{R}^2)} &\leq C, \\ \theta &\equiv 1 \text{ on } B(x'_0; 2r), & \text{supp}(\theta) &\subset B(x'_0; 4r), & \|\nabla_H^k \theta\|_{L^\infty(\mathbb{R}^2)} &\leq Cr^{-k} \end{aligned}$$

for  $k = 0, 1, 2$ ; compare the proof of Lemma 5.4. From (7.5) we see that  $\omega\pi$  satisfies

$$\Delta_H(\omega\pi) = \text{div}_H(\omega F) - \nabla_H \omega \cdot F + 2\text{div}_H((\nabla_H \omega)\pi) - (\Delta_H \omega)\pi \quad \text{in } \mathbb{R}^2.$$

Then, letting  $\Psi(x', y') := \frac{1}{2\pi} \log|x' - y'|$  be the Green's function for the Laplacian in  $\mathbb{R}^2$ , we obtain

$$(\omega\pi)(x') = - \int_{\mathbb{R}^2} (\nabla_{y'} \Psi)(x', y') \cdot [\omega F + 2(\nabla_{y'} \omega)\pi](y') dy' - \int_{\mathbb{R}^2} \Psi(x', y') [(\nabla_H \omega) \cdot F + (\Delta_H \omega)\pi](y') dy'.$$

Therefore, for  $x' \in B(x'_0; r)$  we have the representation

$$\begin{aligned} \nabla_H \pi(x') &= - \int_{\mathbb{R}^2} (\nabla_{x'} \nabla_{y'} \Psi)(x', y') [\omega F + 2(\nabla_{y'} \omega)\pi](y') dy' - \int_{\mathbb{R}^2} (\nabla_{x'} \Psi)(x', y') [(\nabla_H \omega) \cdot F + (\Delta_H \omega)\pi](y') dy' \\ &= - \int_{\mathbb{R}^2} (\nabla_{x'} \nabla_{y'} \Psi)(x', y') [\theta F + \omega(1 - \theta)F + 2(\nabla_{y'} \omega)\pi](y') dy' \\ &\quad - \int_{\mathbb{R}^2} (\nabla_{x'} \Psi)(x', y') [(\nabla_H \omega) \cdot F + (\Delta_H \omega)\pi](y') dy' \\ &=: \Pi_1(x') + \Pi_2(x') + \Pi_3(x') + \Pi_4(x') + \Pi_5(x') \end{aligned}$$

where in the second step we used  $\omega\theta = \theta$ . We derive  $L^p(B(x'_0; r))$ -estimates for each of the above terms as follows: By the Calderón–Zygmund inequality we have

$$\|\Pi_1\|_{L^p(\mathbb{R}^2)} \leq C \|\theta F\|_{L^p(\mathbb{R}^2)} \leq C \|\theta\|_{L^p(\mathbb{R}^2)} \|F\|_{L^\infty(G')} \leq Cr^{2/p} \|F\|_{L^\infty(G')}.$$

For the second term note that we have  $|\nabla_{x'} \nabla_{y'} \Psi(x', y')| \leq C|x' - y'|^{-2}$  and

$$\text{supp}(\omega(1 - \theta)) = \text{supp}(\omega - \theta) \subset \text{supp}(\omega) \setminus B(x'_0; 2r)$$

yields  $\text{supp}(\omega(1 - \theta)) \subset \{r \leq |x' - y'| \leq 4\}$  and therefore

$$\begin{aligned} \|\Pi_2\|_{L^p(B(x'_0; r))} &\leq \|1\|_{L^p(B(x'_0; r))} \left( \sup_{x' \in B(x'_0; r)} \int_{r \leq |x' - y'| \leq 4} C|x' - y'|^{-2} dy' \right) \|\omega(1 - \theta)F\|_{L^\infty(G')} \\ &\leq Cr^{2/p} (1 + |\log r|) \|F\|_{L^\infty(G)}. \end{aligned}$$

The condition  $\text{supp}(\nabla_H \omega) \subset G' \setminus [-1, 2]$  yields

$$\|\Pi_3\|_{L^p(B(x'_0; r))} \leq \|1\|_{L^p(B(x'_0; r))} \left( \sup_{1/2 \leq |x' - y'| \leq 3} C|x' - y'|^{-2} \right) \|2(\nabla_H \omega)\pi\|_{L^1(G')} \leq Cr^{2/p} \|\pi\|_{L^1(G')}.$$

It follows from Poincaré's inequality and the  $L^2$ -theory for (7.5) that

$$\|\pi\|_{L^1(G')} \leq C \|\pi\|_{L^2(G)} \leq C \|\nabla_H \pi\|_{L^2(G)} \leq C \|F\|_{L^2(G)} \leq C \|F\|_{L^\infty(G)}$$

and therefore  $\|\Pi_3\|_{L^p(B(x'_0; r))} \leq Cr^{2/p} \|F\|_{L^\infty(G)}$ . Similarly to  $\Pi_3$ , we have

$$\begin{aligned} \|\Pi_4 + \Pi_5\|_{L^p(B(x'_0; r))} &\leq \|1\|_{L^p(B(x'_0; r))} \left( \sup_{1/2 \leq |x' - y'| \leq 3} C|x' - y'|^{-1} \right) (\|F\|_{L^1(G)} + \|\pi\|_{L^1(G)}) \\ &\leq Cr^{2/p} \|F\|_{L^\infty(G)}. \end{aligned}$$

Combining these estimates yields the desired estimate.  $\square$

**Remark 7.4.** Note that the Calderón–Zygmund inequality we have used to estimate  $\Pi_1$  does not hold for  $p \in \{1, \infty\}$  while the arguments of Section 6 can be adapted to cover the case  $p = \infty$ .



We now turn to prove the estimate  $|\lambda|^{1/2}\|\partial_z(\lambda - A)^{-1}\mathbb{P}f\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0}\|f\|_{L_H^\infty L_z^p(\Omega)}$  for  $\lambda \in \Sigma_\theta$ ,  $|\lambda| > \lambda_0$  and for  $\theta \in (0, \pi)$ ,  $p > 3$ . For this purpose we observe that the solution  $v$  to the resolvent problem

$$\lambda v - Av = \mathbb{P}f \quad \text{on } \Omega$$

with boundary conditions (2.3) is decomposed as  $v = v_1 + v_2$ , where  $(v_1, \pi_1)$  and  $(v_2, \pi_2)$  solve

$$(7.6) \quad \lambda v_1 - \Delta v_1 + \nabla_H \pi_1 = f \text{ on } \Omega, \quad \Delta_H \pi_1 = -h^{-1} \operatorname{div}_H(\partial_z v|_{\Gamma_b}) \text{ on } G,$$

and

$$(7.7) \quad \lambda v_2 - \Delta v_2 + \nabla_H \pi_2 = 0 \text{ on } \Omega, \quad \Delta_H \pi_2 = \operatorname{div}_H \bar{f} \text{ on } G,$$

respectively, both equipped with the boundary conditions (2.3) and periodic boundary conditions for  $\pi_i$  on  $\partial G$ , as  $\pi := \pi_1 + \pi_2$  satisfies (6.1). Since (7.6) is equivalent to  $v_1 = (\lambda - \Delta)^{-1}(f + Bv)$  we obtain

$$(7.8) \quad |\lambda|^{1/2}\|\partial_z v_1\|_{L_H^\infty L_z^p(\Omega)} \leq |\lambda|^{1/2}\|\nabla v_1\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0}\|f\|_{L_H^\infty L_z^p(\Omega)}$$

for  $|\lambda| > \lambda_0$  by the same argument used to derive (6.5). This,  $\nabla_H v_2 = \nabla_H v - \nabla_H v_1$ , and estimate (6.7) yield

$$(7.9) \quad |\lambda|^{1/2}\|\nabla_H v_2\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0}\|f\|_{L_H^\infty L_z^p(\Omega)}, \quad \lambda \in \Sigma_\theta.$$

In order to prove estimate (7.1) it thus remains to establish the following.

**Proposition 7.5.** *Let  $p \in (3, \infty)$  and  $\theta \in (0, \pi)$ . Then there exists constants  $\lambda_0 > 0$  and  $C_{\theta,p,\lambda_0} > 0$  such that for all  $\lambda \in \Sigma_\theta$  with  $|\lambda| > \lambda_0$  and  $f \in X$  the solution  $v_2$  of (7.7) satisfies*

$$|\lambda|^{1/2}\|\partial_z v_2\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0}\|f\|_{L_H^\infty L_z^p(\Omega)}.$$

**Remark 7.6.** The estimate

$$|\lambda|^{1/2}\|\partial_z(\lambda - A)^{-1}\mathbb{P}f\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p}\|f\|_{L_H^\infty L_z^p(\Omega)}, \quad f \in X$$

actually holds for the full range of  $\lambda \in \Sigma_\theta$ ,  $\theta \in (0, \pi)$ , i.e. one can take  $\lambda_0 = 0$ . This is obtained by using that  $\mathbb{P}f \in L_{\bar{\sigma}}^p(\Omega)$  yields  $v := (\lambda - A)^{-1}\mathbb{P}f \in D(A_{p,\bar{\sigma}})$  and therefore

$$\|v\|_{L_H^\infty L_z^p(\Omega)} \leq C_p\|v\|_{W^{2,p}(\Omega)} \leq C_p\|Av\|_{L_{\bar{\sigma}}^p(\Omega)} \leq C_p\|\mathbb{P}f\|_{L_{\bar{\sigma}}^p(\Omega)} \leq C_p\|f\|_{L_{\bar{\sigma}}^p(\Omega)} \leq C_p\|f\|_{L_H^\infty L_z^p(\Omega)},$$

so the same argument as in the proof of Lemma 5.4 applies.

*Proof of Proposition 7.5.* We will simply write  $(v, \pi)$  instead of  $(v_2, \pi_2)$  for the solution of (7.7). By applying a periodic extension in the horizontal variables we may assume that (7.7) holds in a larger domain allowing us to replace  $\Omega$  and  $G$  by  $\Omega' := G' \times (-h, 0)$  and  $G' := (-2, 3)^2$  respectively. We decompose the boundary of  $\Omega'$  into  $\Gamma'_u = G' \times \{0\}$ ,  $\Gamma'_l := \partial G' \times [-h, 0]$  and  $\Gamma'_b = G \times \{-h\}$ . For simplicity we continue to denote the periodic extensions of  $v$ ,  $\pi$  and  $f$  in the same manner.

Let  $\eta > 1$  be a parameter to be fixed later, and let  $\lambda_0$  be a positive number such that

$$(7.10) \quad r_0 := \eta \lambda_0^{-1/2} < \min\{1/8, h/4\}.$$

We fix arbitrary  $\lambda \in \Sigma_\theta$ ,  $|\lambda| > \lambda_0$ , put  $r := \eta|\lambda|^{-1/2} < r_0$ , and introduce two cut-off functions  $\alpha = \alpha_r$ ,  $\beta = \beta_r$ , satisfying

$$\begin{aligned} \alpha &\in C^\infty([-h, 0]), & \alpha &\equiv 0 \text{ on } [-h, -h+r], & \alpha &\equiv 1 \text{ on } [-h+2r, 0], & |\partial_z^k \alpha(z)| &\leq Cr^{-k}, \\ \beta &\in C^\infty([-h, 0]), & \beta &\equiv 1 \text{ on } [-h, -h+2r], & \beta &\equiv 0 \text{ on } [-h+3r, 0], & |\partial_z^k \beta(z)| &\leq Cr^{-k} \end{aligned}$$

for  $k = 0, 1, 2$ , compare the proof of Lemma 5.4. We then split the estimate for  $\partial_z v$  into the ‘‘upper’’ and ‘‘lower’’ parts in  $\Omega$  as

$$(7.11) \quad \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)} \leq \|\partial_z(\alpha v)\|_{L_H^\infty L_z^p(\Omega)} + \|\partial_z(\beta v)\|_{L_H^\infty L_z^p(\Omega)}.$$

**Step 1.** Let us first focus on  $\partial_z(\alpha v)$ . By Lemma 7.2 with radius  $|\lambda|^{-1/2}$  and  $p = q$  we have (7.12)

$$|\lambda|^{1/2} \|\partial_z(\alpha v)\|_{L_H^\infty L_z^p(\Omega)} \leq C_p |\lambda|^{1/p} \sup_{x'_0 \in G} \left( |\lambda|^{1/2} \|\partial_z(\alpha v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))} + \|\nabla_H \partial_z(\alpha v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))} \right),$$

where  $C(x'_0; |\lambda|^{-1/2})$  denotes the cylinder  $B(x'_0; |\lambda|^{-1/2}) \times (-h, 0)$  and we used that

$$\|f\|_{L_H^\infty L_z^p(\Omega)} = \sup_{x'_0 \in G} \|f\|_{L^\infty(B(x'_0, R); L_z^p)}, \quad R > 0.$$

In the following we fix arbitrary  $x'_0 \in G$  and introduce a cut-off function  $\theta = \theta_r \in C_c^\infty(\mathbb{R}^2)$  such that

$$\theta \equiv 1 \text{ in } \overline{B(x'_0; |\lambda|^{-1/2})}, \quad \text{supp } \theta \subset B(x'_0; r), \quad \|\nabla_H^k \theta\|_{L^\infty(\mathbb{R}^2)} \leq Cr^{-k}$$

for  $k = 0, 1, 2$ . Then  $\theta \alpha v$  solves

$$\begin{aligned} \lambda(\theta \alpha v) - \Delta(\theta \alpha v) &= -\theta \alpha \nabla_H \pi - 2\nabla(\theta \alpha) \cdot \nabla v - (\Delta(\theta \alpha))v \quad \text{on } \Omega', \\ \partial_z(\theta \alpha v)|_{\Gamma'_a \cup \Gamma'_b} &= 0, \quad \theta \alpha v \text{ periodic on } \Gamma'_l. \end{aligned}$$

We further differentiate this equation with respect to  $z$  to obtain

$$\lambda(\theta \partial_z(\alpha v)) - \Delta(\theta \partial_z(\alpha v)) = F_1 + \partial_z F_2 \quad \text{on } \Omega', \quad \theta \partial_z(\alpha v)|_{\Gamma'_a \cup \Gamma'_b} = 0, \quad \theta \partial_z(\alpha v) \text{ periodic on } \Gamma'_l,$$

where

$$\begin{aligned} F_1 &:= -\theta(\partial_z \alpha)(\nabla_H \pi) - (\Delta_H \theta)(\partial_z \alpha)v - (\Delta_H \theta)\alpha(\partial_z v), \\ F_2 &:= -2(\nabla_H \theta)\alpha \cdot (\nabla_H v) - 2\theta(\partial_z \alpha)(\partial_z v) - \theta(\partial_z^2 \alpha)v. \end{aligned}$$

By (5.7) and (5.8) for  $\Omega'$  in the case  $q = p$ , we obtain the estimate

$$(7.13) \quad |\lambda|^{1/2} \|\partial_z(\theta \alpha v)\|_{L^p(\Omega')} + \|\nabla \partial_z(\theta \alpha v)\|_{L^p(\Omega')} \leq C_\theta \left( |\lambda|^{-1/2} \|F_1\|_{L^p(\Omega')} + \|F_2\|_{L^p(\Omega')} \right).$$

and since  $\theta \equiv 1$  on  $C(x'_0; |\lambda|^{-1/2}) \subset \Omega'$  by (7.10), we further have

$$(7.14) \quad \begin{aligned} \|\partial_z(\alpha v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))} &\leq \|\partial_z(\theta \alpha v)\|_{L^p(\Omega')}, \\ \|\nabla \partial_z(\alpha v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))} &\leq \|\nabla \partial_z(\alpha v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))}. \end{aligned}$$

Let us estimate each term on this right-hand side of (7.13) as follows: Denoting  $\|\cdot\|_{L_H^p} := \|\cdot\|_{L^p(B(x'_0; r))}$  and  $\|\cdot\|_{L_z^p} := \|\cdot\|_{L^p(-h, 0)}$ , we first observe that the cut-off functions satisfy

$$\|\theta\|_{L_H^p} \leq Cr^{2/p}, \quad \|\nabla_H \theta\|_{L_H^p} \leq Cr^{2/p-1}, \quad \|\Delta_H \theta\|_{L_H^p} \leq Cr^{2/p-2}$$

as well as

$$\|\partial_z \alpha\|_{L_z^p} \leq Cr^{1/p-1}, \quad \|\partial_z^2 \alpha\|_{L_z^p} \leq Cr^{1/p-2}.$$

By Proposition 7.3 we then have

$$\|\theta(\partial_z \alpha)(\nabla_H \pi)\|_{L^p(\Omega')} \leq \|\theta\|_\infty \|\partial_z \alpha\|_{L_z^p} \|\nabla_H \pi\|_{L_H^p} \leq C_p r^{3/p-1} (1 + |\log r|) \|f\|_{L_H^\infty L_z^p(\Omega)}.$$

We further have the Poincaré inequality

$$(7.15) \quad \|f\|_{L^\infty(G'; L^p(-h, -h+d))} \leq d \|\partial_z f\|_{L_H^\infty L_z^p}, \quad 0 \leq d \leq h, \quad f|_{\Gamma'_b} = 0$$

and hence using Hölder's inequality yields

$$\|(\Delta_H \theta)(\partial_z \alpha)v\|_{L^p(\Omega')} \leq \|\Delta_H \theta\|_{L_H^p} \|\partial_z \alpha\|_\infty \|v\|_{L^\infty(G'; L^p(-h, -h+2r))} \leq Cr^{2/p-2} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}.$$

For the third term in  $F_1$  we simply have

$$\|(\Delta_H \theta)\alpha(\partial_z v)\|_{L^p(\Omega')} \leq \|\Delta_H \theta\|_{L_H^p} \|\alpha\|_\infty \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)} \leq Cr^{2/p-2} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}.$$

The first term in  $F_2$  is estimated via (7.9), yielding

$$\|(\nabla_H \theta)\alpha(\nabla_H v)\|_{L^p(\Omega')} \leq \|\nabla_H \theta\|_{L_H^p} \|\alpha\|_\infty \|\nabla_H v\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta, p, \lambda_0} r^{2/p-1} |\lambda|^{-1/2} \|f\|_{L_H^\infty L_z^p(\Omega)},$$

whereas for the second term in  $F_2$  we simply have

$$\|\theta(\partial_z \alpha)(\partial_z v)\|_{L^p(\Omega')} \leq \|\theta\|_\infty \|\partial_z \alpha\|_\infty \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)} \leq Cr^{2/p-1} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)},$$

and by the Poincaré inequality (7.15) we estimate the last term by

$$\|\theta(\partial_z^2 \alpha)v\|_{L^p(\Omega')} \leq \|\theta\|_{L_H^p} \|\partial_z^2 \alpha\|_\infty \|v\|_{L_H^\infty L_z^p(\Omega)} \leq Cr^{2/p-1} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}.$$

Collecting the above estimates, using (7.12), (7.13) and (7.14), as well as  $r = \eta|\lambda|^{-1/2}$ , we obtain that

$$(7.16) \quad \begin{aligned} |\lambda|^{1/2} \|\partial_z(\alpha v)\|_{L_H^\infty L_z^p(\Omega)} &\leq C_{\theta,p,\lambda_0} \left( \eta^{2/p-2} + \eta^{3/p-2} |\lambda|^{-1/2p} + \eta^{2/p-1} r^{1/p} |\log(r)| \right) \|f\|_{L_H^\infty L_z^p(\Omega)} \\ &\quad + C_{\theta,p} (\eta^{2/p-1} + \eta^{2/p-2}) |\lambda|^{1/2} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}. \\ &\leq C_{\theta,p,\lambda_0} \eta^{2/p-1} (1 + r^{1/p} |\log r|) \|f\|_{L_H^\infty L_z^p(\Omega)} \\ &\quad + C_{\theta,p} (\eta^{2/p-1} + \eta^{2/p-2}) |\lambda|^{1/2} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}. \end{aligned}$$

**Step 2:** Now we shall estimate  $\partial_z(\beta v)$ . We apply Lemma 7.2 as in the previous step to obtain

$$(7.17) \quad |\lambda|^{1/2} \|\partial_z(\beta v)\|_{L_H^\infty L_z^p(\Omega)} \leq C_p |\lambda|^{1/p} \sup_{x'_0 \in G} \left( |\lambda|^{1/2} \|\partial_z(\beta v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))} + \|\nabla_H \partial_z(\beta v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))} \right).$$

In the following we fix an arbitrary point  $x'_0 \in G$ . With the same cut-off function  $\theta \in C_c^\infty(\mathbb{R}^2)$  as in Step 1, we find that  $\theta\beta v$  solves

$$\lambda(\theta\beta v) - \Delta(\theta\beta v) = F_3 \quad \text{in } \Omega', \quad \partial_z(\theta\beta v)|_{\Gamma'_u} = 0, \quad \theta\beta v|_{\Gamma'_l} = 0, \quad \theta\beta v \text{ periodic on } \Gamma'_l$$

where

$$F_3 := -\theta\beta(\nabla_H \pi) - 2(\nabla_H \theta)\beta \cdot (\nabla_H v) - 2\theta(\partial_z \beta)(\partial_z v) - (\Delta_H \theta)\beta v - 2\theta(\partial_z^2 \beta)v.$$

We apply estimate (5.7) on  $\Omega'$  with  $q = p$  to obtain

$$(7.18) \quad |\lambda|^{1/2} \|\nabla(\theta\beta v)\|_{L^p(\Omega')} + \|\Delta(\theta\beta v)\|_{L^p(\Omega')} \leq C_\theta \|F_3\|_{L^p(\Omega')}$$

where we further have, compare (7.14), that

$$(7.19) \quad \begin{aligned} \|\partial_z(\beta v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))} &\leq \|\nabla(\theta\beta v)\|_{L^p(\Omega')}, \\ \|\nabla_H \partial_z(\beta v)\|_{L^p(C(x'_0; |\lambda|^{-1/2}))} &\leq \|\nabla_H \partial_z(\theta\beta v)\|_{L^p(\Omega')} \leq \|\theta\beta v\|_{W^{2,p}(\Omega')} \leq C_p \|\Delta(\theta\beta v)\|_{L^p(\Omega')} \end{aligned}$$

by the invertibility of the Laplace operator with mixed Neumann and Dirichlet boundary conditions, compare Section 5.

We now estimate the right-hand side of (7.18) as follows: Note that  $\beta$  satisfies the estimates

$$\|\beta\|_{L_z^p} \leq Cr^{1/p}, \quad \|\partial_z \beta\|_{L_z^p} \leq Cr^{1/p-1}, \quad \|\partial_z^2 \beta\|_{L_z^p} \leq Cr^{1/p-2}$$

since  $\text{supp}(\beta) \subset [-h, -h+3r]$ . It follows from Proposition 7.3 that

$$\|\theta\beta(\nabla_H \pi)\|_{L^p(\Omega')} \leq \|\theta\|_\infty \|\beta\|_{L_z^p} \|\nabla_H \pi\|_{L_H^p} \leq C_p r^{3/p} (1 + |\log r|) \|f\|_{L_H^\infty L_z^p(\Omega)}.$$

The estimate (7.9) implies that

$$\|(\nabla_H \theta)\beta \cdot (\nabla_H v)\|_{L_H^p} \leq \|\nabla_H \theta\|_{L_H^p} \|\beta\|_\infty \|\nabla_H v\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0} r^{2/p-1} |\lambda|^{-1/2} \|f\|_{L_H^\infty L_z^p(\Omega)},$$

and for the term containing vertical derivatives we have

$$\|\theta(\partial_z \beta)(\partial_z v)\|_{L^p(\Omega')} \leq \|\theta\|_{L_H^p} \|\partial_z \beta\|_\infty \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)} \leq Cr^{2/p-1} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}.$$

By the Poincaré inequality (7.15) we have

$$\|(\Delta_H \theta)\beta v\|_{L^p(\Omega')} \leq \|\Delta_H \theta\|_{L_H^p} \|\beta\|_\infty \|v\|_{L^\infty(G; L^p(-h, -h+3r))} \leq Cr^{2/p-1} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}$$

as well as

$$\|\theta(\partial_z^2 \beta)v\|_{L^p(\Omega')} \leq \|\theta\|_{L_H^p} \|\partial_z^2 \beta\|_\infty \|v\|_{L^\infty(G; L^p(-h, -h+3r))} \leq Cr^{2/p-1} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}.$$

Combining the above estimates with (7.17), (7.18) and (7.19) as well as  $r = \eta|\lambda|^{-1/2}$  then yields

$$(7.20) \quad \begin{aligned} |\lambda|^{1/2} \|\partial_z(\beta v)\|_{L_H^\infty L_z^p(\Omega)} &\leq C_{\theta,p,\lambda_0} \left( \eta^{2/p-1} + \eta^{3/p} |\lambda|^{-1/2p} (1 + |\log(\eta|\lambda|^{-1/2})|) \right) \|f\|_{L_H^\infty L_z^p(\Omega)} \\ &\quad + C_{\theta,p} \eta^{2/p-1} |\lambda|^{1/2} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)}. \end{aligned}$$

We now substitute (7.16) and (7.20) into (7.11). Since all constants  $C > 0$  do not depend on the parameter  $\eta > 0$ , we can take it to be sufficiently large and so similarly to the proof of Lemma 5.4 we obtain

$$|\lambda|^{1/2} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0} \left( \eta^{2/p-1} (1 + r^{1/p} |\log(r)|) + \eta^{3/p} |\lambda|^{-1/2p} (1 + |\log(|\lambda|)|) \right) \|f\|_{L_H^\infty L_z^p(\Omega)}.$$

Since

$$\sup_{0 < r < r_0} r^{1/p} |\log(r)| < \infty, \quad \sup_{|\lambda| > \lambda_0} |\lambda|^{-1/2p} (1 + |\log(|\lambda|)|) < \infty,$$

for any  $r_0, \lambda_0 > 0$  and  $p \in (1, \infty)$ , this implies the desired estimate  $|\lambda|^{1/2} \|\partial_z v\|_{L_H^\infty L_z^p(\Omega)} \leq C \|f\|_{L_H^\infty L_z^p(\Omega)}$  for  $|\lambda| \geq \lambda_0$ .  $\square$

We now turn to the problem

$$(7.21) \quad \lambda v - Av = \mathbb{P} \partial_z f \text{ on } \Omega$$

with boundary conditions (2.3) for  $f \in X$ . Since

$$(7.22) \quad \mathbb{P} \partial_z f = \partial_z f - (1 - Q) \overline{\partial_z f} = \partial_z f,$$

whenever  $f = 0$  on  $\Gamma_u \cup \Gamma_b$  and  $C_{\text{per}}^\infty([0, 1]^2; C_c^\infty(-h, 0))^2$  is dense in  $X$  we may assume without loss of generality that (7.22) holds. Moreover, in view of periodic extension we may assume that (7.21) holds in a larger domain  $\Omega' := G' \times (-h, 0)$ ,  $G' := (-2, 3)^2$ . Since the problem is well-posed in  $L_\sigma^p(\Omega)$  by (6.6), estimate (7.2) then follows from the following:

**Proposition 7.7.** *Let  $p \in (2, \infty)$  and  $\theta \in (0, \pi)$ . Then there exists constants  $\lambda_0 > 0$  and  $C_{\theta,p,\lambda_0} > 0$  such that for all  $\lambda \in \Sigma_\theta$  with  $|\lambda| > \lambda_0$  and  $f \in X$  the solution to the problem (7.21) satisfies*

$$|\lambda|^{1/2} \|v\|_{L_H^\infty L_z^p(\Omega)} \leq C_{\theta,p,\lambda_0} \|f\|_{L_H^\infty L_z^p(\Omega)}.$$

To prove this estimate, we adopt a duality argument combined with the use of a regularized delta function, which is based on the methodology known in  $L^\infty$ -type error analysis of the finite element method, cf. [22].

In order to prove this estimate we first introduce some notation. Using periodicity, one sees that for any  $\varepsilon \in (0, 1)$  we have  $B(x'_0, \varepsilon) \subset G'$  for  $x'_0 \in G$  and

$$\|v\|_{L_H^\infty L_z^p(\Omega)}^p = \sup_{x'_0 \in G} \sup_{x' \in B(x'_0; \varepsilon)} \int_{-h}^0 |v(x', z)|^p dz,$$

where by  $B(x'_0; \varepsilon)$  we continue to denote a disk in  $\mathbb{R}^2$ , compare Lemma 7.2. In the following we fix arbitrary  $x'_0 \in G$ ,  $x' \in B(x'_0; \varepsilon)$  and choose  $\varepsilon = |\lambda|^{-\frac{p}{2(p-2)}}$  for  $\lambda$  as above.

Letting  $\delta \geq 0$  be a smooth nonnegative function in the variables  $(x, y) =: x'$  such that  $\text{supp } \delta \subset B(0; 1)$  and  $\int_{\mathbb{R}^2} \delta dx' = 1$ , we introduce a rescaled function as

$$(7.23) \quad \delta_\varepsilon(x') := \frac{1}{\varepsilon^2} \delta\left(\frac{x'}{\varepsilon}\right), \quad \delta_{\varepsilon, x'_0}(x') := \delta_\varepsilon(x' - x'_0).$$

We then obtain

$$(7.24) \quad \int_{-h}^0 |v(x', z)|^p dz = \int_{-h}^0 \int_{G'} (|v(x', z)|^p - |v(y', z)|^p) \delta_{\varepsilon, x'_0}(y') dy' dz + (v, \delta_{\varepsilon, x'_0} |v|^{p-2} v^*)_{\Omega'} =: I_1(x') + I_2,$$

where  $v^*$  means the complex conjugate of  $v$  and  $(\cdot, \cdot)_{\Omega'}$  denotes the inner product on  $L^2(\Omega')^2$ . In the following we estimate the two terms on the right-hand side separately, beginning with  $I_1$ .

**Lemma 7.8.** *Under the assumptions of Proposition 7.7 we have for all for all  $x'_0 \in G$  and  $x' \in B(x'_0; \varepsilon)$ ,  $\varepsilon = |\lambda|^{-\frac{p}{2(p-2)}}$ , that*

$$|I_1(x')| = \left| \int_{\Omega'} (|v(x', z)|^p - |v(y', z)|^p) \delta_{\varepsilon, x'_0}(y') dy' dz \right| \leq C_{\theta, p} |\lambda|^{-1/2} \|f\|_{L_H^\infty L_z^p(\Omega)} \|v\|_{L_H^\infty L_z^p(\Omega)}^{p-1}.$$

*Proof.* Since  $\int_{\mathbb{R}^2} \delta_{\varepsilon, x'_0}(y') dy' = 1$  and  $\text{supp } \delta_{\varepsilon, x'_0} \subset B(x'_0; \varepsilon)$  we obtain

$$\begin{aligned} |I_1(x')| &\leq \sup_{y' \in B(x'_0; \varepsilon)} \int_{-h}^0 \left| |v(x', z)|^p - |v(y', z)|^p \right| dz \\ &\leq C \sup_{y' \in B(x'_0; \varepsilon)} \int_{-h}^0 (|v(x', z)|^{p-1} + |v(y', z)|^{p-1}) |v(x', z) - v(y', z)| dz, \end{aligned}$$

where we have used the elementary inequality

$$|a^p - b^p| \leq p \max\{a, b\}^{p-1} |a - b| \leq p(a+b)^{p-1} |a - b| \leq p2^{p-2} (a^{p-1} + b^{p-1}) |a - b|$$

for all  $a, b \geq 0$ , where we used that  $p \in [2, \infty)$  implies that  $x \mapsto x^{p-1}$  is a convex function. Hölder's inequality then implies that

$$\int_{-h}^0 (|v(x', z)|^{p-1} + |v(y', z)|^{p-1}) |v(x', z) - v(y', z)| dz \leq (\|v(x')\|_{L_z^p}^{p-1} + \|v(y')\|_{L_z^p}^{p-1}) \|v(x') - v(y')\|_{L_z^p}.$$

Hence we have

$$\sup_{x' \in B(x'_0; \varepsilon)} |I_1(x')| \leq C \sup_{y' \in B(x'_0; \varepsilon)} \|v(y')\|_{L_z^p}^{p-1} \sup_{y' \in B(x'_0; \varepsilon)} \|v(x') - v(y')\|_{L_z^p} \leq C \|v\|_{L_H^\infty L_z^p}^{p-1} \varepsilon^\alpha \|v\|_{C_H^\alpha L_z^p(\Omega)},$$

where  $\alpha := 1 - 2/p > 0$  and  $\|v\|_{C_H^\alpha L_z^p(\Omega)}$  denotes the space of  $L^p(-h, 0)$ -valued Hölder continuous functions of exponent  $\alpha$  on  $\overline{G}$ .

The assumption  $\varepsilon = |\lambda|^{-\frac{p}{2(p-2)}}$  then yields  $\varepsilon^\alpha = |\lambda|^{-1/2}$ . We now use the Sobolev embedding  $W^{1,p}(G) \hookrightarrow C^\alpha(\overline{G})$  to obtain the estimate  $\|v\|_{C_H^\alpha L_z^p} \leq C \|v\|_{W^{1,p}(\Omega)}$ . In addition, the Poincaré inequality yields

$$\|v\|_{W^{1,p}(\Omega)} \leq C_p \|\nabla v\|_{L^p(\Omega)} = C_p \|\nabla(\lambda - A_{p, \bar{\sigma}})^{-1} \partial_z f\|_{L^p(\Omega)} \leq C_{\theta, p} \|f\|_{L^p(\Omega)} \leq C_{\theta, p} \|f\|_{L_H^\infty L_z^p(\Omega)},$$

where we used that  $\nabla(-A_{p, \bar{\sigma}})^{-1/2}$ ,  $A_{p, \bar{\sigma}}(\lambda - A_{p, \bar{\sigma}})^{-1}$  and  $(-A_{p, \bar{\sigma}})^{-1/2} \partial_z$  are (uniformly) bounded on  $L_{\bar{\sigma}}^p(\Omega)$  for  $\lambda \in \Sigma_\theta$  by [7]. Combining these results then gives the desired estimate.  $\square$

In order to estimate  $I_2$  we perform a duality argument. For this purpose we introduce an auxiliary problem corresponding to (7.21) as follows:

$$(7.25) \quad \begin{aligned} \lambda^* w - \Delta w + \nabla_H \Pi &= \delta_{\varepsilon, x'_0} |v|^{p-2} v^* \quad \text{in } \Omega', \\ \partial_z \Pi &= 0 \quad \text{in } \Omega', \\ \text{div}_H \bar{w} &= 0 \quad \text{in } G', \\ \partial_z w|_{\Gamma'_u} &= 0, \quad w|_{\Gamma'_l} = 0, \quad w, \Pi \text{ periodic on } \Gamma'_l, \end{aligned}$$

where the upper script  $*$  means complex conjugate as before. We establish an  $L_H^1 L_z^q$ -estimate to this problem, where  $q := p/(p-1)$  is the dual index of  $p$ .

**Proposition 7.9.** *Let  $p \in (2, \infty)$ ,  $1/p + 1/q = 1$  and  $\theta \in (0, \pi)$ . Then there exists a sufficiently large  $\lambda_0 > 0$  and a constant  $C_{p, \lambda_0, \theta} > 0$  such that the solution of (7.25) satisfies*

$$|\lambda|^{1/2} \|\partial_z w\|_{L_H^1 L_z^q(\Omega')} \leq C_{\theta, p} \left(1 + |\lambda|^{-1/2q} \varepsilon^{2/s-2}\right) \|v\|_{L_H^\infty L_z^p(\Omega)}^{p-1},$$

for all  $\varepsilon \in (0, 1)$ ,  $s \in (1, q]$ ,  $x'_0 \in G$ ,  $\lambda \in \Sigma_\theta$ ,  $|\lambda| > \lambda_0$ , and  $v \in X$ .

**Remark 7.10.** If one even has  $p \in (3, \infty)$  then this result can be extended to the full range of  $\lambda \in \Sigma_\theta$  by a similar argument as in the proof of Lemma 5.4, compare Remark 7.6.

For simplicity, we write  $L_H^p L_z^q$  to refer to  $L_H^p L_z^q(\Omega') = L^p(G'; L^q(-h, 0))$  when there is no ambiguity. First we introduce the following result.

**Lemma 7.11.** *Let  $\varepsilon \in (0, 1)$ ,  $x'_0 \in G$ ,  $p \in (1, \infty)$ ,  $1/p + 1/q = 1$  and  $v \in X$  be arbitrary. Then, for  $\delta_{\varepsilon, x'_0}$  defined as in (7.23) and  $s \in [1, q]$  we have*

$$\|\delta_{\varepsilon, x'_0} |v|^{p-2} v^*\|_{L^s(\Omega')} \leq C \|\delta_{\varepsilon, x'_0} |v|^{p-2} v^*\|_{L_H^s L_z^q} \leq C \varepsilon^{2/s-2} \|v\|_{L_H^\infty L_z^p(\Omega)}^{p-1}$$

for a constant  $C > 0$  not depending on  $\varepsilon$ ,  $x'_0$  and  $v$ .

*Proof.* We set  $F := \delta_{\varepsilon, x'_0} |v|^{p-2} v^*$ . Noting that  $|F|^q = \delta_{\varepsilon, x'_0}^q |v|^p$  and that  $\delta_{\varepsilon, x'_0}$  is independent of  $z$ , we obtain

$$\begin{aligned} \|F\|_{L_H^s L_z^q} &= \left[ \int_{G'} \left( \int_{-h}^0 \delta_\varepsilon(x' - x'_0)^q |v(x', z)|^p dz \right)^{s/q} dx' \right]^{1/s} \\ &\leq \left( \int_{G'} \delta_\varepsilon(x' - x'_0)^s dx' \right)^{1/s} \left[ \sup_{x' \in G'} \left( \int_{-h}^0 |v(x', z)|^p dz \right)^{1/p} \right]^{p/q} \\ &\leq C \varepsilon^{2/s-2} \|v\|_{L_H^\infty L_z^p(\Omega)}^{p-1}, \end{aligned}$$

where we used the periodicity of  $v$  in the last step. This completes the proof.  $\square$

*Proof of Proposition 7.9.* We set  $r := \eta|\lambda|^{-1/2}$ , where  $\eta > 0$  is a large number to be fixed later and  $|\lambda| > \lambda_0$ , where  $\lambda_0 > 0$  is sufficiently large such that  $\eta\lambda_0^{-1/2} < 1$ . We introduce two cut-off functions  $\alpha = \alpha_r$ ,  $\beta = \beta_r$  in the vertical direction as follows:

$$\begin{aligned} \alpha &\in C^\infty([-h, 0]), \quad \alpha \equiv 0 \text{ in } [-h, -h+r], \quad \alpha \equiv 1 \text{ in } [-h+2r, 0], \quad |\partial_z^k \alpha(z)| \leq Cr^{-k}, \\ \beta &\in C^\infty([-h, 0]), \quad \beta \equiv 1 \text{ in } [-h, -h+2r], \quad \beta \equiv 0 \text{ in } [-h+3r, 0], \quad |\partial_z^k \beta(z)| \leq Cr^{-k} \end{aligned}$$

for  $k = 0, 1, 2$ . Then we may split the estimate for  $\partial_z w$  into the ‘‘upper’’ and ‘‘lower’’ parts in  $\Omega'$  as

$$(7.26) \quad \|\partial_z w\|_{L_H^1 L_z^q} \leq \|\partial_z(\alpha w)\|_{L_H^1 L_z^q} + \|\partial_z(\beta w)\|_{L_H^1 L_z^q}.$$

**Step 1.** We consider  $\alpha w$ , which satisfies

$$\begin{aligned} \lambda^* \alpha w - \Delta(\alpha w) &= \alpha F - \alpha(\nabla_H \Pi) - 2(\partial_z \alpha)(\partial_z w) - (\partial_z^2 \alpha)w, \\ \partial_z(\alpha w) &= 0 \quad \text{on } \Gamma'_u \cup \Gamma'_b, \quad \alpha w \text{ periodic on } \Gamma'_l \end{aligned}$$

where  $F := \delta_{\varepsilon, x'_0} |v|^{p-2} v^*$  as in the proof of Lemma 7.11. Differentiating this with respect to  $z$  yields

$$\begin{aligned} \lambda^* \partial_z(\alpha w) - \Delta(\partial_z(\alpha w)) &= \partial_z [\alpha F - 2(\partial_z \alpha)(\partial_z w) - (\partial_z^2 \alpha)w] - (\partial_z \alpha)(\nabla_H \Pi) \quad \text{in } \Omega', \\ \partial_z(\alpha w) &= 0 \quad \text{on } \Gamma'_u \cup \Gamma'_b, \quad \partial_z(\alpha w) \text{ periodic on } \Gamma'_l. \end{aligned}$$

Applying Lemma 5.4 in  $L_H^1 L_z^q(\Omega')$  we obtain

$$\begin{aligned} |\lambda|^{1/2} \|\partial_z(\alpha w)\|_{L_H^1 L_z^q} &\leq C \left( \|\alpha F\|_{L_H^1 L_z^q} + \|(\partial_z \alpha)(\partial_z w)\|_{L_H^1 L_z^q} + \|(\partial_z^2 \alpha)w\|_{L_H^1 L_z^q} \right) \\ &\quad + C|\lambda|^{-1/2} \|(\partial_z \alpha)(\nabla_H \Pi)\|_{L_H^1 L_z^q}. \end{aligned}$$

We now estimate each term on the right-hand side. By Lemma 7.11 with  $s = 1$  we have

$$\|\alpha F\|_{L_H^1 L_z^q} \leq \|\alpha\|_\infty \|F\|_{L_H^1 L_z^q} \leq C \|v\|_{L_H^\infty L_z^p(\Omega)}^{p-1}.$$

Using the estimate on derivatives of  $\alpha$  we obtain

$$\|(\partial_z \alpha)(\partial_z w)\|_{L_H^1 L_z^q} \leq Cr^{-1} \|\partial_z w\|_{L_H^1 L_z^q},$$

and by the Poincaré inequality we have

$$\|(\partial_z^2 \alpha)w\|_{L_H^1 L_z^q} \leq Cr^{-2} \|w\|_{L_H^1 L_z^q(G' \times (-h, -h+2r))} \leq Cr^{-1} \|\partial_z w\|_{L_H^1 L_z^q}.$$

Using  $L^s(G') \hookrightarrow L^1(G')$  as well as the estimate on the pressure term, cf. [11, Theorem 3.1.], in  $L^s(\Omega)$  for  $s \in (1, q]$ , we obtain

$$\|(\partial_z \alpha)(\nabla_H \Pi)\|_{L_H^1 L_z^q} \leq C \|\partial_z \alpha\|_{L_z^q} \|\nabla_H \Pi\|_{L^s(G')} \leq C r^{1/q-1} \|F\|_{L^s} \leq C r^{1/q-1} \varepsilon^{2/s-2} \|v\|_{L_H^\infty L_z^p}^{p-1},$$

Collecting the above estimates and plugging in  $r = \eta|\lambda|^{-1/2}$  yields

$$(7.27) \quad |\lambda|^{1/2} \|\partial_z(\alpha w)\|_{L_H^1 L_z^q} \leq C(1 + \eta^{1/q-1} |\lambda|^{-1/2q} \varepsilon^{2/s-2}) \|v\|_{L_H^\infty L_z^p}^{p-1} + C\eta^{-1} |\lambda|^{1/2} \|\partial_z w\|_{L_H^1 L_z^q}.$$

**Step 2.** We consider  $\beta w$ , which satisfies

$$\begin{aligned} \lambda^* \beta w - \Delta(\beta w) &= \beta F - \beta \nabla_H \Pi - 2(\partial_z \beta)(\partial_z w) - (\partial_z^2 \beta)w \quad \text{in } \Omega', \\ \partial_z(\beta w) &= 0 \text{ on } \Gamma'_u, \quad \beta w = 0 \text{ on } \Gamma'_b, \quad \partial_z(\beta w) \text{ periodic on } \Gamma'_l. \end{aligned}$$

Applying Lemma 5.4 in  $L_H^1 L_z^q$  we obtain

$$|\lambda|^{1/2} \|\partial_z(\beta w)\|_{L_H^1 L_z^q} \leq C(\|\beta F\|_{L_H^1 L_z^q} + \|(\partial_z \beta)(\partial_z w)\|_{L_H^1 L_z^q} + \|(\partial_z^2 \beta)w\|_{L_H^1 L_z^q} + \|\beta(\nabla_H \Pi)\|_{L_H^1 L_z^q}).$$

A calculation similar to Step 1 then gives

$$(7.28) \quad |\lambda|^{1/2} \|\partial_z(\beta w)\|_{L_H^1 L_z^q} \leq C(1 + \eta^{1/q} |\lambda|^{-1/2q} \varepsilon^{2/s-2}) \|v\|_{L_H^\infty L_z^p}^{p-1} + C\eta^{-1} |\lambda|^{1/2} \|\partial_z w\|_{L_H^1 L_z^q}.$$

Substituting (7.27) and (7.28) into (7.26) and choosing sufficiently large  $\eta$  enable us to absorb the term  $|\lambda|^{1/2} \|\partial_z w\|_{L_H^1 L_z^q}$  from the right-hand side, which leads to

$$|\lambda|^{1/2} \|\partial_z w\|_{L_H^1 L_z^q} \leq C(1 + |\lambda|^{-1/2q} \varepsilon^{2/s-2}) \|v\|_{L_H^\infty L_z^p}^{p-1}$$

This completes the proof.  $\square$

With the preparations above, we are now in the position to prove Proposition 7.7.

*Proof of Proposition 7.7.* By (7.24) and Lemma 7.8 we have

$$(7.29) \quad \|v\|_{L_H^\infty L_z^p}^p \leq C|\lambda|^{-1/2} \|f\|_{L_H^\infty L_z^p} \|v\|_{L_H^\infty L_z^p}^{p-1} + I_2,$$

with  $I_2$  as defined in (7.24). Substituting (7.25) and integrating by parts, we find that

$$\begin{aligned} I_2 &= (v, \delta_{\varepsilon, x'_0} |v|^{p-2} v^*)_{\Omega'} = (v, \lambda^* w - \Delta w + \nabla_H \Pi)_{\Omega'} = (\lambda v - \Delta v + \nabla_H \pi, w)_{\Omega'} = (\partial_z f, w)_{\Omega'} \\ &= -(f, \partial_z w)_{\Omega'}, \end{aligned}$$

where we have used that  $(v, \nabla_H \Pi)_{\Omega'} = 0 = (\nabla_H \pi, w)_{\Omega'}$  since  $\operatorname{div}_H \bar{v} = 0 = \operatorname{div}_H \bar{w}$  for the third and  $f|_{\Gamma_u \cup \Gamma_b} = 0$  for the last equality. Using  $1/p + 1/q = 1$  and applying Proposition 7.9 we obtain

$$|I_2| \leq \|f\|_{L_H^\infty L_z^p} \|\partial_z w\|_{L_H^1 L_z^q} \leq C|\lambda|^{-1/2} \|f\|_{L_H^\infty L_z^p} \|v\|_{L_H^\infty L_z^p}^{p-1} \left(1 + |\lambda|^{-1/2q} \varepsilon^{2/s-2}\right).$$

We set  $\varepsilon = |\lambda|^{-\frac{p}{2(p-2)}}$  for  $|\lambda| > 1$  and  $s = \min\{\frac{4p}{3p+2}, \frac{p}{p-1}\} \in (1, q]$ . This yields

$$-\frac{1}{2q} + \left(1 - \frac{1}{s}\right) \frac{p}{p-2} = -\frac{1}{2} + \frac{1}{2p} + \left(1 - \frac{1}{s}\right) \frac{p}{p-2} \leq -\frac{1}{4} + \frac{1}{2p} < 0$$

which implies that  $1 + |\lambda|^{-1/2q} \varepsilon^{2/s-2} \leq 2$  for  $|\lambda| > 1$  and therefore

$$(7.30) \quad |I_2| \leq C|\lambda|^{-1/2} \|f\|_{L_H^\infty L_z^p} \|v\|_{L_H^\infty L_z^p}^{p-1}, \quad |\lambda| > 1.$$

The desired estimate then follows from (7.29) and (7.30) after dividing by  $\|v\|_{L_H^\infty L_z^p}^{p-1}$ .  $\square$

*Proof of Claim 7.1.* Estimate (7.1) now follows from (7.8) and Proposition 7.5, whereas estimate (7.2) follows from Proposition 7.7. Estimate 7.3 follows from (7.1), (7.2) and Claim 6.1.  $\square$

## 8. PROOF OF THE MAIN RESULTS

Theorem 3.4 is a direct consequence of Claims 6.1 and 7.1.

For the non-linear problem in the space  $X$  we will make use of the following estimates.

**Lemma 8.1.** *Let  $p > 3$ . Then exists a constant  $C > 0$  such that for all  $t > 0$  and  $v_i \in X_{\overline{\sigma}}$  satisfying  $\nabla v_i \in X$  and  $v_i|_{\Gamma_b} = 0$  with  $u_i = (v_i, w_i)$  as in (2.2) for  $i = 1, 2$  we have*

- (i)  $\|e^{tA}\mathbb{P}(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} \leq Ct^{-1/2}\|\nabla v_1\|_{L_H^\infty L_z^p}\|v_2\|_{L_H^\infty L_z^p},$
- (ii)  $\|\nabla e^{tA}\mathbb{P}(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} \leq Ct^{-1/2}\|\nabla v_1\|_{L_H^\infty L_z^p}\|\nabla v_2\|_{L_H^\infty L_z^p},$
- (iii)  $\|\nabla e^{tA}\mathbb{P}(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} \leq Ct^{-1}\|\nabla v_1\|_{L_H^\infty L_z^p}\|v_2\|_{L_H^\infty L_z^p},$

as well as

$$(iv) \quad \|e^{tA}\mathbb{P}(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} \leq C \left( t^{-1/2}\|\nabla v_i\|_{L_H^\infty L_z^p}\|v_j\|_{L_H^\infty L_z^p} + \|\nabla v_1\|_{L_H^\infty L_z^p}\|\nabla v_2\|_{L_H^\infty L_z^p} \right)$$

where  $\{i, j\} = \{1, 2\}$ .

*Proof.* We begin by noting that

$$\|(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} \leq (\|v_1\|_{L^\infty(\Omega)} + \|w_1\|_{L^\infty(\Omega)}) \|\nabla v_2\|_{L_H^\infty L_z^p}.$$

So, using Sobolev embeddings, the Poincaré inequality and  $X \hookrightarrow L^p(\Omega)^2$  we obtain

$$\|v_i\|_{L^\infty(\Omega)} \leq C\|v_i\|_{W^{1,p}(\Omega)} \leq C\|\nabla v_i\|_{L^p(\Omega)} \leq C\|\nabla v_i\|_{L_H^\infty L_z^p}.$$

Similarly one has

$$\|w_i\|_{L^\infty(\Omega)} \leq C\|\operatorname{div}_H v_i\|_{L_H^\infty L_z^p} \leq C\|\nabla v_i\|_{L_H^\infty L_z^p}.$$

This allows us to obtain (ii) via Claim 6.1 and 7.1 as well as

$$\|\nabla e^{tA}\mathbb{P}(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} \leq Ct^{-1/2}\|(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} \leq Ct^{-1/2}\|\nabla v_1\|_{L_H^\infty L_z^p}\|\nabla v_2\|_{L_H^\infty L_z^p}.$$

To prove (i) we proceed analogously as above to obtain

$$\|v_1 \otimes v_2\|_{L_H^\infty L_z^p} \leq C\|\nabla v_i\|_{L_H^\infty L_z^p}\|v_j\|_{L_H^\infty L_z^p}, \quad \|w_1 v_2\|_{L_H^\infty L_z^p} \leq C\|\nabla v_1\|_{L_H^\infty L_z^p}\|v_2\|_{L_H^\infty L_z^p}$$

where  $\{i, j\} = \{1, 2\}$  and since  $\operatorname{div} u_i = 0$  we can write

$$(u_1 \cdot \nabla)v_2 = \nabla \cdot (u_1 \otimes v_2) = \nabla_H \cdot (v_1 \otimes v_2) + \partial_z(w_1 v_2)$$

which allows us to apply Claim 6.1 and 7.1 yielding

$$\begin{aligned} \|e^{tA}\mathbb{P}(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} &= \|e^{tA}\mathbb{P}\nabla \cdot (u_1 \otimes v_2)\|_{L_H^\infty L_z^p} \\ &\leq \|e^{tA}\mathbb{P}\nabla_H \cdot (v_1 \otimes v_2)\|_{L_H^\infty L_z^p} + \|e^{tA}\mathbb{P}\partial_z(w_1 v_2)\|_{L_H^\infty L_z^p} \\ &\leq Ct^{-1/2} \left( \|v_1 \otimes v_2\|_{L_H^\infty L_z^p} + \|w_1 v_2\|_{L_H^\infty L_z^p} \right) \\ &\leq Ct^{-1/2}\|\nabla v_1\|_{L_H^\infty L_z^p}\|v_2\|_{L_H^\infty L_z^p}, \end{aligned}$$

and estimate (iii) is obtained analogously via

$$\|\nabla e^{tA}\mathbb{P}(u_1 \cdot \nabla)v_2\|_{L_H^\infty L_z^p} \leq Ct^{-1} \left( \|v_1 \otimes v_2\|_{L_H^\infty L_z^p} + \|w_1 v_2\|_{L_H^\infty L_z^p} \right) \leq Ct^{-1}\|\nabla v_1\|_{L_H^\infty L_z^p}\|v_2\|_{L_H^\infty L_z^p}.$$

To prove (iv) we observe that  $w_i = 0$  on  $\Gamma_u \cup \Gamma_b$  implies that

$$\mathbb{P}\partial_z(w_1 v_2) = \partial_z(w_1 v_2) = -(\operatorname{div}_H v_1)v_2 + w_1 \partial_z v_2$$

and the right-hand side is further estimated via

$$\|(\operatorname{div}_H v_1)v_2\|_{L_H^\infty L_z^p} \leq C\|\nabla v_1\|_{L_H^\infty L_z^p}\|v_2\|_{L^\infty(\Omega)} \leq C\|\nabla v_1\|_{L_H^\infty L_z^p}\|\nabla v_2\|_{L_H^\infty L_z^p},$$

and

$$\|w_1 \partial_z v_2\|_{L_H^\infty L_z^p} \leq \|w_1\|_{L^\infty(\Omega)}\|\partial_z v_2\|_{L_H^\infty L_z^p} \leq C\|\nabla v_1\|_{L_H^\infty L_z^p}\|\nabla v_2\|_{L_H^\infty L_z^p}.$$



Applying Claim 6.1 then yields that for  $\{i, j\} = \{1, 2\}$  we have

$$\|e^{tA} \mathbb{P} \nabla_H \cdot (v_1 \otimes v_2)\|_{L_H^\infty L_z^p} \leq Ct^{-1/2} \|v_1 \otimes v_2\|_{L_H^\infty L_z^p} \leq Ct^{-1/2} \|\nabla v_i\|_{L_H^\infty L_z^p} \|v_j\|_{L_H^\infty L_z^p},$$

as well as

$$\|e^{tA} \mathbb{P} \partial_z (w_1 v_2)\|_{L_H^\infty L_z^p} \leq C \|\nabla v_1\|_{L_H^\infty L_z^p} \|\nabla v_2\|_{L_H^\infty L_z^p}$$

which implies (iv) and completes the proof.  $\square$

It has been proven in [7] that the operator  $A_{p, \bar{\sigma}}$  possesses maximal  $L^q$ -regularity. In [9] the authors applied this to develop a solution theory for initial data

$$a \in X_\gamma := (L_{\bar{\sigma}}^p(\Omega), D(A_p))_{1-1/q, q} \subset B_{pq}^{2-2/q}(\Omega)^2 \cap L_{\bar{\sigma}}^p(\Omega)$$

where  $p, q \in (1, \infty)$  satisfy  $1/p + 1/q \leq 1$ . In particular, one has the following result.

**Lemma 8.2.** *Let  $a \in X_\gamma$ . Then there exists a unique strong solution to the primitive equations (2.1) with boundary conditions (2.3) satisfying*

$$v \in C([0, \infty); X_\gamma).$$

This enables a key step in the proof of our main result as it guarantees the existence of smooth reference solutions  $v_{\text{ref}}$  to the primitive equations given sufficiently smooth reference data  $a_{\text{ref}}$ . In order to construct  $v$  as a solution to problem (2.1) with initial data  $a$  we construct  $V := v - v_{\text{ref}}$  by an iterative method using initial data  $a_0 := a - a_{\text{ref}}$ . Before we do so, we establish an auxiliary lemma.

**Lemma 8.3.** *Let  $(a_n)_{n \in \mathbb{N}}$  be a sequence of positive real numbers such that*

$$a_{m+1} \leq a_0 + c_1 a_m^2 + c_2 a_m \quad \text{for all } m \in \mathbb{N}$$

*and constants  $c_1 > 0$  and  $c_2 \in (0, 1)$  such that  $4c_1 a_0 < (1 - c_2)^2$ . Then  $a_m < \frac{2}{1-c_2} a_0$  for all  $m \in \mathbb{N}$ .*

*Proof.* Let  $x_0$  be the smallest solution to the equation  $x = a_0 + c_1 x^2 + c_2 x$ . Then

$$0 < x_0 = \frac{(1 - c_2) - \sqrt{(1 - c_2)^2 - 4c_1 a_0}}{2c_1} = \frac{1}{2c_1} \frac{4c_1 a_0}{(1 - c_2) + \sqrt{(1 - c_2)^2 - 4c_1 a_0}} < \frac{2}{1 - c_2} a_0,$$

and since  $p(x) = a_0 + c_1 x^2 + c_2 x$  is an increasing function on  $[0, \infty)$  it follows that  $p(x) \leq x_0$  for  $x \in [0, x_0]$ . The condition  $c_2 \in (0, 1)$  further yields

$$(1 - c_2) + \sqrt{(1 - c_2)^2 - 4c_1 a_0} < 2$$

from which it follows that  $a_0 < x_0$  and thus the claim is easily derived by induction.  $\square$

We now prove our main result.

*Proof of Theorem 3.1. Step 1: Decomposition of data.*

Given an initial value  $a \in X_{\bar{\sigma}}$  we will split it into a smooth part  $a_{\text{ref}}$  and a small rough part  $a_0$ , where  $a = a_{\text{ref}} + a_0$ , as follows: Since  $A_{\bar{\sigma}}$  is densely defined on  $X_{\bar{\sigma}}$  we take  $a_{\text{ref}} \in D(A_{\bar{\sigma}})$  such that  $a_0 := a - a_{\text{ref}}$  can be assumed to be arbitrarily small in  $X_{\bar{\sigma}}$ . Now let  $q \in (1, \infty)$  be such that  $1/q + 1/p \leq 1$  and  $2/q + 3/p < 1$ . The latter condition on  $q$  then yields the embedding  $X_\gamma \hookrightarrow C^1(\bar{\Omega})^2$ . Due to  $D(A_{\bar{\sigma}}) \subset D(A_{p, \bar{\sigma}}) \subset X_\gamma$  it follows from Lemma 8.2 that taking  $a_{\text{ref}}$  as initial data of the primitive equations, there exists a function  $v_{\text{ref}} \in C([0, \infty); X_\gamma)$  solving the primitive equations with initial data  $v_{\text{ref}}(0) = a_{\text{ref}}$ .

**Step 2: Estimates for the construction of a local solution.**

We will show that there exists a constant  $C_0 > 0$  such that if  $a_0 \in X_{\bar{\sigma}}$  satisfies  $\|a_0\|_{L_H^\infty L_z^p} < C_0$  then there exists a time  $T > 0$  and a unique function

$$V \in \mathcal{S}(T) := \{V \in C([0, T]; X_{\bar{\sigma}}) : \|\nabla V(t)\|_{L_H^\infty L_z^p} = o(t^{-1/2})\},$$

where

$$\|V\|_{\mathcal{S}(T)} = \max \left\{ \sup_{0 < t < T} \|V(t)\|_{L_H^\infty L_z^p}, \sup_{0 < t < T} t^{1/2} \|\nabla V(t)\|_{L_H^\infty L_z^p} \right\}$$

such that  $v = v_{\text{ref}} + V$  solves problem (2.1) on  $(0, T)$  with initial value  $v(0) = a$ . In order to construct  $V$  we define the iterative sequence of functions  $(V_m)_{m \in \mathbb{N}}$  via

$$(8.1) \quad V_0(t) = e^{tA} a_0, \quad V_{m+1}(t) = e^{tA} a_0 + \int_0^t e^{(t-s)A} F_m(s) ds$$

where

$$F_m := -\mathbb{P}((U_m \cdot \nabla)V_m + (U_m \cdot \nabla)v_{\text{ref}} + (u_{\text{ref}} \cdot \nabla)V_m)$$

and  $U_m = (V_m, W_m)$ ,  $u_{\text{ref}} = (v_{\text{ref}}, w_{\text{ref}})$  with the vertical component  $w$  given by the horizontal component  $v$  via the relation (2.2). We will now estimate this sequence in  $\mathfrak{S}(T)$  for some value  $T > 0$  to be fixed later on. Since  $\mathbb{P}a_0 = a_0$  we have

$$\|V_0\|_{\mathfrak{S}(T)} \leq C \|a_0\|_{L_H^\infty L_z^p}, \quad T \in (0, \infty)$$

by Lemma 6.1. For  $m \geq 1$  we will first consider the gradient estimates. We have already estimated the term  $\nabla e^{tA} a_0$ , whereas for the convolution integrals we have

$$\begin{aligned} \left\| \int_0^{t/2} \nabla e^{(t-s)A} \mathbb{P}((U_m(s) \cdot \nabla)V_m(s)) ds \right\|_{L_H^\infty L_z^p} &\leq C \left( \int_0^{t/2} (t-s)^{-1} s^{-1/2} ds \right) K_m(t) H_m(t) \\ &= Ct^{-1/2} K_m(t) H_m(t) \end{aligned}$$

by Lemma 8.1 (iii) where

$$K_m(t) := \sup_{0 < s < t} s^{1/2} \|\nabla V_m(s)\|_{L_H^\infty L_z^p}, \quad H_m(t) := \sup_{0 < s < t} \|V_m(s)\|_{L_H^\infty L_z^p}$$

and via Lemma 8.1 (ii) we obtain

$$\begin{aligned} \left\| \int_{t/2}^t \nabla e^{(t-s)A} \mathbb{P}(U_m(s) \cdot \nabla V_m(s)) ds \right\|_{L_H^\infty L_z^p} &\leq C \left( \int_{t/2}^t (t-s)^{-1/2} s^{-1} ds \right) K_m(t)^2 \\ &\leq Ct^{-1/2} K_m(t)^2. \end{aligned}$$

Finally applying Lemma 8.1 (ii) to the two remaining mixed terms yields

$$\begin{aligned} \left\| \int_0^t \nabla e^{(t-s)A} \mathbb{P}(U_m(s) \cdot \nabla)v_{\text{ref}}(s) ds \right\|_{L_H^\infty L_z^p} &\leq C \left( \int_0^t (t-s)^{-1/2} s^{-1/2} ds \right) \sup_{0 < s < t} \|\nabla v_{\text{ref}}(s)\|_{L_H^\infty L_z^p} K_m(t) \\ &= C \sup_{0 < s < t} \|\nabla v_{\text{ref}}(s)\|_{L_H^\infty L_z^p} K_m(t), \\ \left\| \int_0^t \nabla e^{(t-s)A} \mathbb{P}(u_{\text{ref}}(s) \cdot \nabla)V_m(s) ds \right\|_{L_H^\infty L_z^p} &\leq C \sup_{0 < s < t} \|\nabla v_{\text{ref}}(s)\|_{L_H^\infty L_z^p} K_m(t). \end{aligned}$$

We set  $R := \sup_{0 \leq t \leq T_0} \|\nabla v_{\text{ref}}(t)\|_{L_H^\infty L_z^p}$  and note that  $0 < R < \infty$  by Lemma 8.2, since  $v_{\text{ref}} \in C([0, \infty); X_\gamma)$  and  $2/q + 3/p < 1$  implies that  $X_\gamma \subset B_{pq}^{2-2/q}(\Omega)^2 \hookrightarrow C^1(\overline{\Omega})^2$  via embedding theory, cf. [25, Section 3.3.1]. Taking these estimates together yields

$$(8.2) \quad t^{1/2} \|\nabla V_{m+1}(t)\|_{L_H^\infty L_z^p} \leq C_1 \left( \|a_0\|_{L_H^\infty L_z^p} + K_m(t) H_m(t) + K_m(t)^2 + Rt^{1/2} K_m(t) \right).$$

To estimate  $\|V_{m+1}(t)\|_{L_H^\infty L_z^p}$  we apply Lemma 8.1 (i) to obtain

$$\begin{aligned} \left\| \int_0^t e^{(t-s)A} \mathbb{P}((U_m(s) \cdot \nabla)V_m(s)) ds \right\|_{L_H^\infty L_z^p} &\leq C \left( \int_0^t (t-s)^{-1/2} s^{-1/2} ds \right) K_m(t) H_m(t) \\ &= CK_m(t) H_m(t) \end{aligned}$$

whereas for the mixed terms Lemma 8.1 (iv) yields

$$\begin{aligned} \|e^{(t-s)A}\mathbb{P}(U_m(s) \cdot \nabla)v_{\text{ref}}(s)\|_{L_H^\infty L_z^p} &\leq C \left( (t-s)^{-1/2} \|\nabla v_{\text{ref}}(s)\|_{L_H^\infty L_z^p} \|V_m(s)\|_{L_H^\infty L_z^p} \right. \\ &\quad \left. + \|\nabla V_m(s)\|_{L_H^\infty L_z^p} \|\nabla v_{\text{ref}}(s)\|_{L_H^\infty L_z^p} \right) \end{aligned}$$

and therefore

$$\begin{aligned} \left\| \int_0^t e^{(t-s)A}\mathbb{P}((U_m(s) \cdot \nabla)v_{\text{ref}}(s)) ds \right\|_{L_H^\infty L_z^p} &\leq C \left( \int_0^t (t-s)^{-1/2} ds \right) RH_m(t) + C \left( \int_0^t s^{-1/2} ds \right) RK_m(t) \\ &= CRt^{1/2}(H_m(t) + K_m(t)), \end{aligned}$$

and the other mixed term can be treated analogously due to the symmetry of the right-hand side in (iv). Taking these estimates together yields

$$(8.3) \quad \|V_{m+1}(t)\|_{L_H^\infty L_z^p} \leq C_1 \left( \|a_0\|_{L_H^\infty L_z^p} + K_m(t)H_m(t) + t^{1/2}H_m(t) + t^{1/2}K_m(t) \right).$$

Since the right-hand sides of (8.2) and (8.3) are increasing functions we obtain for  $t > 0$  that

$$(8.4) \quad \begin{aligned} K_{m+1}(t) &\leq C_1 \left( \|a_0\|_{L_H^\infty L_z^p} + K_m(t)H_m(t) + K_m(t)^2 + Rt^{1/2}K_m(t) \right), \\ H_{m+1}(t) &\leq C_1 \left( \|a_0\|_{L_H^\infty L_z^p} + K_m(t)H_m(t) + Rt^{1/2}H_m(t) + Rt^{1/2}K_m(t) \right). \end{aligned}$$

Now let  $T \in (0, T_0)$  where  $T_0 > 0$  is chosen in such a way that

$$8C_1RT_0^{1/2} < 1.$$

Then for all  $0 < t \leq T < T_0$  we have

$$\|V_{m+1}\|_{\mathcal{S}(t)} \leq C_1 \|a_0\|_{L_H^\infty L_z^p} + 2C_1 \|V_m\|_{\mathcal{S}(t)}^2 + \frac{1}{4} \|V_m\|_{\mathcal{S}(t)}.$$

By Lemma 8.3 it follows that if  $8C_1^2 \|a_0\|_{L_H^\infty L_z^p} < (1 - 1/4)^2$ , then for all  $m \in \mathbb{N}$  we have

$$(8.5) \quad \|V_m\|_{\mathcal{S}(t)} \leq \frac{8}{3} C_1 \|a_0\|_{L_H^\infty L_z^p}, \quad t \in (0, T].$$

The property  $\lim_{t \rightarrow 0^+} t^{1/2} \|\nabla V_m(t)\|_{L_H^\infty L_z^p} = 0$  is then easily obtained via induction and Claim 6.1 (e).

**Step 3: Convergence.**

We now show that  $(V_m)_{m \in \mathbb{N}}$  is a Cauchy sequence in  $\mathcal{S}(T)$  if  $\|a_0\|_{L_H^\infty L_z^p}$  is sufficiently small. For this purpose we consider the new sequence

$$\tilde{V}_m := V_{m+1} - V_m, \quad m \geq 0.$$

Using the previous estimates we already know that  $\|\tilde{V}_0\|_{\mathcal{S}(T)} < \infty$ . To estimate this sequence further we use

$$F_m - F_{m-1} = \left( \tilde{U}_{m-1} \cdot \nabla \right) V_m + (U_{m-1} \cdot \nabla) \tilde{V}_{m-1} + \left( \tilde{U}_{m-1} \cdot \nabla \right) v_{\text{ref}} + (U_{\text{ref}} \cdot \nabla) \tilde{V}_{m-1}$$

and proceed as above to obtain

$$(8.6) \quad \begin{aligned} t^{1/2} \|\nabla \tilde{V}_m(t)\|_{L_H^\infty L_z^p} &\leq C_2 \left( 2H_m(t)\tilde{K}_{m-1}(t) + 2\tilde{H}_{m-1}(t)K_{m-1}(t) + K_m(t)\tilde{K}_{m-1}(t) \right. \\ &\quad \left. + K_{m-1}(t)\tilde{K}_{m-1}(t) + 2Rt^{1/2}\tilde{K}_{m-1} \right) \end{aligned}$$

as well as

$$(8.7) \quad \|\tilde{V}_m(t)\|_{L_H^\infty L_z^p} \leq C_2 \left( \tilde{K}_{m-1}(t) [H_m(t) + H_{m-1}(t)] + Rt^{1/2} [\tilde{K}_{m-1}(t) + \tilde{H}_{m-1}(t)] \right),$$

where

$$\tilde{K}_m(t) := \sup_{0 < s < t} s^{1/2} \|\nabla \tilde{V}_m(t)\|_{L_{\tilde{H}}^\infty L_z^p}, \quad \tilde{H}_m(t) := \sup_{0 < s < t} \|\tilde{V}_m(t)\|_{L_{\tilde{H}}^\infty L_z^p}.$$

By (8.5) it follows that if

$$\max\{2RT_0^{1/2}, 16C_1\|a_0\|_{L_{\tilde{H}}^\infty L_z^p}\} < 1/4C_2,$$

then for  $m \geq 1$  and  $0 < t \leq T < T_0$  we have

$$\|\tilde{V}_m(t)\|_{\mathfrak{S}(t)} \leq C_2 \left( 16C_1\|a_0\|_{L_{\tilde{H}}^\infty L_z^p} + 2Rt^{1/2} \right) \|\tilde{V}_{m-1}(t)\|_{\mathfrak{S}(t)} < \frac{1}{2} \|\tilde{V}_{m-1}(t)\|_{\mathfrak{S}(t)}.$$

Therefore, since  $\mathfrak{S}(T)$  is a Banach space,  $(V_m)_{m \in \mathbb{N}}$  converges in  $\mathfrak{S}(T)$ . We denote the limit by  $V$  and see that it satisfies

$$(8.8) \quad V(t) = e^{tA}a_0 - \int_0^t e^{(t-s)A} \mathbb{P} \left( (U(s) \cdot \nabla)V(s) + (U(s) \cdot \nabla)v_{\text{ref}}(s) + (u_{\text{ref}}(s) \cdot \nabla)V(s) \right) ds$$

for  $t \in (0, T)$  and thus  $v := V + v_{\text{ref}}$  is a solution to the primitive equations (2.1).

**Step 4: Extending to a global solution.**

Using  $V \in \mathfrak{S}(T)$ , the embedding  $L_{\tilde{H}}^\infty L_z^p(\Omega) \hookrightarrow L^p(\Omega)$ , as well as the semigroup estimates

$$t^\vartheta \|e^{tA} \mathbb{P} f\|_{D((-A_{p,\bar{\sigma}})^\vartheta)} \leq C \|f\|_{L^p(\Omega)}, \quad t^{1/2} \|e^{tA} \mathbb{P} \nabla \cdot f\|_{L^p(\Omega)} \leq C \|f\|_{L^p(\Omega)}, \quad t > 0, \quad \vartheta \in [0, 1]$$

compare [11, Lemma 4.6] and [7, Theorem 3.7], one easily obtains that  $V(t_0) \in D((-A_{p,\bar{\sigma}})^\vartheta)$  for  $t_0 > 0$ , and thus  $v(t_0) \in D((-A_{p,\bar{\sigma}})^{1/p})$  as well, so  $v$  can be extended to a global solution that is strong on  $(t_0, \infty)$ .

**Step 5: Uniqueness.**

To see that  $v$  is a unique solution and thus strong on  $(0, t_0)$  as well, we consider  $v^{(1)}$  and  $v^{(2)}$  both to be solutions in the sense of Theorem 3.1 with initial value  $a$  and set

$$t^* := \inf\{t \in [0, \infty) : v^{(1)}(t) \neq v^{(2)}(t)\}.$$

Assume that  $t^* \in (0, \infty)$ . Then using continuity of the solutions

$$a^* := v^{(1)}(t^*) = v^{(2)}(t^*) = a_{\text{ref}}^* + a_0^*$$

where  $a_0^* \in X_{\bar{\sigma}}$  is sufficiently small and  $a_{\text{ref}}^* \in D(A_{\bar{\sigma}})$ . Let  $v_{\text{ref}}^*$  be the reference solution to the initial data  $a_{\text{ref}}^*$  and

$$V^{(i)}(t) := v^{(i)}(t^* + t) - v_{\text{ref}}^*(t^*), \quad i = 1, 2.$$

Then  $V^{(1)}, V^{(2)} \in \mathfrak{S}(T^*)$  both satisfy the condition (8.8) for arbitrary  $t \in (0, T^*)$ ,  $T^* \in (0, \infty)$ . We set  $\tilde{V} := V^{(1)} - V^{(2)}$  and observe that proceeding analogously as before one obtains

$$\begin{aligned} \tilde{H}(t) &\leq C_3 \left( t^{1/2} (\tilde{H}(t) + \tilde{K}(t)) + H^{(1)}(t) \tilde{K}(t) + K^{(2)}(t) \tilde{H}(t) \right), \\ \tilde{K}(t) &\leq C_3 \left( 2t^{1/2} \tilde{K}(t) + H^{(1)}(t) \tilde{K}(t) + K^{(2)}(t) \tilde{H}(t) + K^{(1)} \tilde{K}(t) + K^{(2)} \tilde{K}(t) \right), \end{aligned}$$

where  $\tilde{H}, H^{(i)}, \tilde{K}, K^{(i)}$  are defined analogously to above. This yields

$$(8.9) \quad \|\tilde{V}\|_{\mathfrak{S}(t)} \leq C_3 \left( t^{1/2} + H^{(1)}(t) + H^{(2)}(t) + K^{(1)}(t) + K^{(2)}(t) \right) \|\tilde{V}\|_{\mathfrak{S}(t)}, \quad t \in (0, T^*).$$

By taking  $T^* > 0$  to be small the terms  $(T^*)^{1/2}$  and  $K^{(1)}(T^*), K^{(2)}(T^*)$  can be taken to be arbitrarily small due to  $\|\nabla V^{(i)}(t)\| = o(t^{-1/2})$ , which in the case  $t^* = 0$  follows from the regularity of  $v$  and in the case  $t^* > 0$  this follows from  $\|\nabla v(t^*)\| \in L_{\tilde{H}}^\infty L_z^p(\Omega)^2$ .

As for  $H^{(1)}$  and  $H^{(2)}$ , using the same arguments that derived (8.3) one obtains for  $t \in (0, T^*)$  that

$$(8.10) \quad H^{(i)}(t) \leq C_1 \left( \|a_0^*\|_{L_{\tilde{H}}^\infty L_z^p} + K^{(i)}(t) H^{(i)}(t) + R^* t^{1/2} H^{(i)}(t) + R^* t^{1/2} K^{(i)}(t) \right),$$

where  $R^* := \sup_{0 \leq t \leq T^*} \|\nabla v_{\text{ref}}^*(t)\|_{L_{\tilde{H}}^\infty L_z^p}$ . Now, we choose  $T \in (0, T^*)$  so small that

$$K^{(i)}(T) H^{(i)}(T^*) + R^* T^{1/2} H^{(i)}(T^*) + R^* T^{1/2} K^{(i)}(T^*) \leq \|a_0^*\|_{L_{\tilde{H}}^\infty L_z^p}.$$

Now, taking  $\|a_0^*\|_{L_H^\infty L_z^p}$  to be sufficiently small, using that the constants  $C_i > 0$ ,  $i = 1, 2, 3$ , are independent of  $\|a_0^*\|_{L_H^\infty L_z^p}$ , we obtain that the pre-factor in (8.9) is smaller 1. Hence, it follows that  $\|\tilde{V}\|_{S(t)} = 0$  for  $t \in (0, T)$  and thus  $v^{(1)} = v^{(2)}$  on  $[0, t^* + T)$  which is a contradiction.

**Step 6:** Additional regularity.

By [11, Theorem 6.1] we thus have

$$v \in C^1((0, \infty); L_{\bar{\sigma}}^p(\Omega)) \cap C((0, \infty); W^{2,p}(\Omega))^2, \quad \pi \in C((0, \infty); W^{1,p}(G)).$$

The additional regularity  $v \in C([0, \infty]; X_{\bar{\sigma}})$  follows from the strong continuity of the semigroup on  $X_{\bar{\sigma}}$ .

For the pressure we have  $\pi(t) \in W^{1,p}(G) \hookrightarrow C^{0,\alpha}([0, 1]^2)$  for  $\alpha \in (0, 1 - 2/p)$ . To obtain the regularity of  $\nabla_H \pi$ , observe that

$$\nabla_H \pi = -Bv - (1 - \mathbb{P})(u \cdot \nabla)v = -Bv - (1 - Q)\overline{(u \cdot \nabla)v}$$

where we used that  $(1 - \mathbb{P})f = (1 - Q)\bar{f}$ . In the proof of Claim 6.1 we have already proven that  $Bv(t) \in C^{0,\alpha}([0, 1]^2)$  for  $\alpha \in (0, 1 - 3/p)$  if  $v(t) \in W^{2,p}(\Omega)^2$ . Likewise, since  $1 - Q$  is continuous on  $C_{\text{per}}^{0,\alpha}([0, 1]^2)^2$  and  $v \in C((0, \infty); W^{2,p}(\Omega))^2$ , we obtain that the remaining terms belong to  $C((0, \infty); C^{0,\alpha}([0, 1]^2))^2$ .  $\square$

*Proof of Theorem 3.2.* Here, we make use of the fact that the relevant estimates in Claim 6.1 and Claim 7.1 can also be applied in  $L_H^\infty L_z^p(\Omega)^2$ , compare Remark 3.5 (c).

Let  $a = a_1 + a_2$  be as in Theorem 3.2. Next, we introduce a decomposition setting  $a_0 := a_2 + (a_1 - a_{\text{ref}})$  where

$$a_{\text{ref}} \in D(A_{\bar{\sigma}}), \quad a_1 \in X_{\bar{\sigma}} \quad \text{and} \quad a_2 \in L_H^\infty L_z^p(\Omega)^2 \cap L_{\bar{\sigma}}^p(\Omega),$$

where  $a_{\text{ref}}$  is such that  $a_0$  satisfies the smallness condition of Theorem 3.1.

Then the same iteration scheme as in the previous proof can be used to construct  $V$  for the initial value  $a_0$  and, in turn,  $v$  to the initial value  $a$ .

The property

$$v \in C([0, \infty); L_{\bar{\sigma}}^p(\Omega)) \cap L^\infty((0, T); L_H^\infty L_z^p(\Omega))^2$$

follows from the boundedness and exponential stability of the semigroup on  $L_{\bar{\sigma}}^p(\Omega)$  and  $L_H^\infty L_z^p(\Omega)^2 \cap L_{\bar{\sigma}}^p(\Omega)$ , as well as the strong continuity on  $L_{\bar{\sigma}}^p(\Omega)$ . Since the solution regularizes at  $t_0 > 0$ , compare Step 4 in the previous proof, we further obtain  $v \in C((0, \infty); X_{\bar{\sigma}})$  from the strong continuity on  $X_{\bar{\sigma}}$ .

The condition

$$(8.11) \quad \limsup_{t \rightarrow 0^+} t^{1/2} \|\nabla v\|_{L_H^\infty L_z^p(\Omega)} \leq C \|a_2\|_{L_H^\infty L_z^p},$$

is verified as follows. Since Claim 6.1 yields

$$\limsup_{t \rightarrow 0^+} t^{1/2} \|\nabla e^{tA}(a_1 - a_{\text{ref}})\|_{L_H^\infty L_z^p} = 0, \quad t^{1/2} \|\nabla e^{tA} a_2\|_{L_H^\infty L_z^p} \leq C_4 \|a_2\|_{L_H^\infty L_z^p}, \quad t > 0,$$

one obtains

$$\limsup_{t \rightarrow 0^+} t^{1/2} \|\nabla V_0(t)\|_{L_H^\infty L_z^p} \leq C_4 \|a_2\|_{L_H^\infty L_z^p}.$$

We now prove  $\limsup_{t \rightarrow 0^+} t^{1/2} \|\nabla V_m(t)\|_{L_H^\infty L_z^p} \leq 2C_4 \|a_2\|_{L_H^\infty L_z^p}$  by induction. Assuming the claim holds for  $m \in \mathbb{N}$  we obtain

$$\limsup_{t \searrow 0} t^{1/2} \|\nabla V_{m+1}(t)\|_{L_H^\infty L_z^p} \leq \left(1 + 2C_1 \|a_0\|_{L_H^\infty L_z^p} + 4C_4 \|a_2\|_{L_H^\infty L_z^p}\right) C_4 \|a_2\|_{L_H^\infty L_z^p}$$

in the same manner as (8.2). Assuming that  $\|a_0\|_{L_H^\infty L_z^p} < 1/4C_1$  and  $\|a_2\|_{L_H^\infty L_z^p} < 1/8C_4$  it follows that the claim holds for all  $m \in \mathbb{N}$  and by taking the limit the same estimate holds for  $V$ . Using  $v_{\text{ref}} \in C([0, \infty); C^1(\bar{\Omega})^2)$ , we obtain that  $v = V + v_{\text{ref}}$  satisfies (8.11).

To prove uniqueness we make the following modifications. If  $v^{(1)}$  and  $v^{(2)}$  are both solutions in the sense of Theorem 3.2, we again define  $t^* := \inf\{t \in [0, \infty) : v^{(1)}(t) \neq v^{(2)}(t)\}$ .

In the case  $t^* > 0$  we have  $a^* = v^{(1)}(t^*) = v^{(2)}(t^*) \in D((-A_{p,\bar{\sigma}})^\vartheta)$  for any  $\vartheta \in [0, 1]$ , compare Step 4 of the proof of Theorem 3.1. Choosing  $2\vartheta - 3/p > 0$  we have that  $D((-A_{p,\bar{\sigma}})^\vartheta) \hookrightarrow X_{\bar{\sigma}}$  and thus we can decompose  $a^* = a_{\text{ref}}^* + a_0^*$  as before and the same argument applies.

If we instead have  $t^* = 0$  we continue to use the decomposition  $a = a_{\text{ref}} + a_0$  where  $a_0 = a_2 + (a_1 - a_{\text{ref}})$ . In this case we have  $\lim_{t \rightarrow 0^+} K^{(i)}(t) \leq C \|a_2\|_{L_H^\infty L_z^p}$  for an absolute constant  $C > 0$  and thus the quantities on the right-hand side of (8.9) can again be taken to be sufficiently small, where on the right-hand side of (8.10) one has  $\|a_0\|_{L_H^\infty L_z^p}$  instead of  $\|a_0^*\|_{L_H^\infty L_z^p}$ , which again yields uniqueness.

This completes the proof.  $\square$

## REFERENCES

- [1] K. Abe, Y. Giga. Analyticity of the Stokes semigroup on spaces of bounded functions. *Acta Math.*, 211, 1-46, 2013.
- [2] K. Abe, Y. Giga, M. Hieber. Stokes resolvent estimates in spaces of bounded functions. *Ann. Sci. Ec. Norm. Super.*, 48, 537-559, 2015.
- [3] H. Amann. Operator-valued Fourier multipliers, vector-valued Besov spaces, and applications. *Math. Nachr.*, 186, 5–56, 1997.
- [4] W. Arendt, C.J.K. Batty, M. Hieber and F. Neubrander. *Vector-valued Laplace transforms and Cauchy problems*. Birkhäuser Verlag, Basel, 2011.
- [5] Ch. Cao and E. Titi. Global well-posedness of the three-dimensional viscous primitive equations of large scale ocean and atmosphere dynamics. *Annals of Mathematics*, 166:245–267, 2007.
- [6] R. Denk, M. Hieber, and J. Prüss.  $\mathcal{R}$ -boundedness, Fourier multipliers and problems of elliptic and parabolic type. *Mem. Amer. Math. Soc.*, 166, 2003.
- [7] Y. Giga, M. Gries, M. Hieber, A. Hussein and T. Kashiwabara. Bounded  $H^\infty$ -Calculus for the Hydrostatic Stokes Operator on  $L^p$ -Spaces and Applications. *Proc. Amer. Math. Soc.*, 145(9):3865–3876, 2017.
- [8] Y. Giga, M. Gries, M. Hieber, A. Hussein and T. Kashiwabara. The primitive equations in the scaling invariant space  $L^\infty(L^1)$ . Submitted, Preprint arXiv:1710.04434, 2017.
- [9] Y. Giga, M. Gries, M. Hieber, A. Hussein and T. Kashiwabara. Analyticity of solutions to the primitive equations. Submitted, Preprint arXiv:1710.04860, 2017.
- [10] W. Grey and G. Sinnamon. Product operators on mixed norm spaces. *Linear and Nonlinear Analysis*, 2(2): 189–197, 2016.
- [11] M. Hieber and T. Kashiwabara. Global Strong Well-Posedness of the Three Dimensional Primitive Equations in  $L^p$ -Spaces. *Arch. Rational Mech. Anal.*, 221(3): 1077–1115, 2016.
- [12] L. Khan. On a Stone-Weierstrass theorem for vector-valued functions. *Punjab Univ. J. Math. (Lahore)*, 12/13: 11–14, 1979–1980.
- [13] I. Kukavica, Y. Pei, W. Rusin and M. Ziane. Primitive equations with continuous initial data. *Nonlinearity*, 27:1135–1155, 2014.
- [14] I. Kukavica and M. Ziane. On the regularity of the primitive equations of the ocean. *Nonlinearity*, 20(12):2739–2753, 2007.
- [15] J. Li and E. Titi. Recent Advances Concerning Certain Class of Geophysical Flows. In *Handbook of Mathematical Analysis in Mechanics of Viscous Fluids*, Springer, 2016.
- [16] J. Li and E. Titi. Existence and uniqueness of weak solutions to viscous primitive equations for certain class of discontinuous initial data. *SIAM J. Math. Anal.*, 49(1):1–28, 2017.
- [17] J. L. Lions, R. Temam and Sh. H. Wang. New formulations of the primitive equations of atmosphere and applications. *Nonlinearity*, 5(2):237–288, 1992.
- [18] J. L. Lions, R. Temam and Sh. H. Wang. On the equations of the large-scale ocean. *Nonlinearity*, 5(5):1007–1053, 1992.
- [19] J. L. Lions, R. Temam and Sh. H. Wang. Models for the coupled atmosphere and ocean. (CAO I,II). *Comput. Mech. Adv.*, 1:3–119, 1993.
- [20] A. Lunardi. *Analytic Semigroups and Optimal Regularity in Parabolic Problems* Birkhäuser Verlag, Basel, 1995
- [21] T. Nau  $L^p$ -theory of cylindrical boundary value problems. *An Operator-Valued Fourier Multiplier and Functional Calculus Approach*. Springer Spektrum, Wiesbaden, 2012.
- [22] R. Rannacher and R.Scott. Some optimal error estimates for piecewise linear finite element approximations. *Math. Comp.*, 38(158):437–445, 1982.
- [23] M. Petcu, R. Temam and M. Ziane. Some mathematical problems in geophysical fluid dynamics. In *Handbook of numerical analysis. Vol. XIV. Special volume: computational methods for the atmosphere and the oceans*, 14:577–750, 2009.
- [24] T. Tachim Medjo. On the uniqueness of  $z$ -weak solutions of the three-dimensional primitive equations of the ocean. *Nonlinear Anal. Real World Appl.*, 11(3):1413–1421, 2010.
- [25] H. Triebel. *Theory of Function Spaces*. (Reprint of 1983 edition) Springer AG, Basel, 2010.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA 3-8-1, MEGURO-KU, TOKYO, 153-8914, JAPAN

*E-mail address:* `labgiga@ms.u-tokyo.ac.jp`

DEPARTEMENT OF MATHEMATICS, TU DARMSTADT, SCHLOSSGARTENSTR. 7, 64289 DARMSTADT, GERMANY

*E-mail address:* `gries@mathematik.tu-darmstadt.de`

DEPARTEMENT OF MATHEMATICS, TU DARMSTADT, SCHLOSSGARTENSTR. 7, 64289 DARMSTADT, GERMANY

*E-mail address:* `hieber@mathematik.tu-darmstadt.de`

DEPARTEMENT OF MATHEMATICS, TU DARMSTADT, SCHLOSSGARTENSTR. 7, 64289 DARMSTADT, GERMANY

*E-mail address:* `hussein@mathematik.tu-darmstadt.de`

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO, TOKYO 153-8914, JAPAN

*E-mail address:* `tkashiwa@ms.u-tokyo.ac.jp`