Title	Effects of seawater transfer and fasting on the endocrine and biochemical growth indices in juvenile chum salmon (<i>Oncorhynchus keta</i>)
Author(s)	Taniyama, Natsumi; Kaneko, Nobuto; Inatani, Yu; Miyakoshi, Yasuyuki; Shimizu, Munetaka
Citation	General and Comparative Endocrinology, 236, 146-156 https://doi.org/10.1016/j.ygcen.2016.07.020
Issue Date	2016-09-15
Doc URL	http://hdl.handle.net/2115/68544
Rights	© 2016. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Taniyama16HUSCAP.pdf

[Instructions for use](#)

Title

Effects of seawater transfer and fasting on the endocrine and biochemical growth indices in juvenile chum salmon (*Oncorhynchus keta*)

Authors

Natsumi Taniyama¹, Nobuto Kaneko¹, Yu Inatani¹, Yasuyuki Miyakoshi², and Munetaka Shimizu^{1*}

Affiliations

¹Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato, Hakodate, Hokkaido 041-8611, Japan

²Salmon and Freshwater Fisheries Research Institute, Hokkaido Research Organization, 3-373 Kitakashiwagi, Eniwa, Hokkaido 061-1433, Japan

*Corresponding author: e-mail: mune@fish.hokudai.ac.jp; Office/Fax: +81-138-40-8897

Abstract

Insulin-like growth factor (IGF)-I, IGF-binding protein (IGFBP)-1 and RNA/DNA ratio are endocrine and biochemical parameters used as growth indices in fish, however, they are subjected to environmental modulation. Chum salmon (*Oncorhynchus keta*) migrate from freshwater (FW) to seawater (SW) at fry/juvenile stage weighing around 1 g and suffer growth-dependent mortality during the early phase of their marine life. In order to reveal environmental modulation of the IGF/IGFBP system and establish a reliable growth index for juvenile chum salmon, we examined effects of SW transfer and fasting on IGF-I, IGFBP-1 and RNA/DNA ratio, and correlated them to individual growth rate. Among serum IGF-I, liver and muscle *igf-1*, *igfbp-1a*, *igfbp-1b* and RNA/DNA ratio examined, muscle RNA/DNA ratio and muscle *igfbp-1a* responded to SW transfer. Serum IGF-I, liver *igf-1* and liver RNA/DNA ratio were sensitive to nutritional change by being reduced in 1 week in both FW and SW while muscle *igf-1* was reduced 2 weeks after fasting. In contrast, *igfbp-1a* in both tissues was increased by 2 weeks of fasting and *igfbp-1b* in the liver of SW fish was increased in 1 week. These results suggest that the sensitivity of *igf-1* and *igfbp-1s* to fasting differs between tissues and subtypes, respectively. When chum salmon juveniles in SW were marked and subjected to feeding or fasting, serum IGF-I showed the highest correlation with individual growth rate. Liver *igfbp-1a* and *-1b*, and muscle *igf-1* exhibited moderate correlation coefficients with growth rate. These results show that serum IGF-I is superior to the other parameters as a growth index in juvenile chum salmon in term of its stability to salinity change, high sensitivity to fasting and strong relationship with growth rate. On the one hand, when collecting blood from chum salmon fry/juveniles is not practical, measuring liver *igfbp-1a* and *-1b*, or/and muscle *igf-1* is an alternative.

Keywords

insulin-like growth factor-I, insulin-like growth factor binding protein, salmon, salinity, fasting, growth index

1. Introduction

Insulin-like growth factor (IGF)-I is a 7.5-kDa polypeptide structurally similar to proinsulin and promotes growth of animals (Daughaday and Rotwein, 1989). IGF-I circulates in the blood at a relatively high level (~200 ng/ml) and acts as a classical hormone. Liver is the major site of circulating IGF-I producing 75% of endocrine IGF-I under stimulation by growth hormone (GH). IGF-I is also expressed in many tissues and exerts its actions through paracrine and autocrine manners (Daughaday and Rotwein, 1989; Le Roith et al., 2001). IGF-I is essential for normal postnatal growth, where local IGF-I is more important than endocrine IGF-I in mice (Le Roith et al., 2001; Ohlsson et al., 2009). Liver-derived endocrine IGF-I contributes to 30% of postnatal growth and plays a critical role in regulating GH secretion by the pituitary gland via a negative feedback loop (Le Roith et al., 2001; Ohlsson et al., 2009).

IGF-I is distinct from insulin in terms of the presence of high-affinity binding proteins. IGF-binding proteins (IGFBPs) regulate availability of IGF-I to target tissues and modulate its activity (Jones and Clemmons, 1995; Rajaram et al., 1997; Firth and Baxter, 2002). There are six IGFBPs in mammals and different types have different functions. IGFBP-1 is one of the major circulating IGFBPs and generally an inhibitor of IGF-I actions by preventing it from interacting with the receptor (Lee et al., 1993; Kajimura and Duan, 2007). IGFBP-1 is mainly produced by the liver but other tissues also express it. IGFBP-1 shows dynamic fluctuations in response to fasting and stress, and its increased level is often a sign of catabolic states.

In teleosts, the importance of IGF-I in growth, development, osmoregulation and maturation has been widely recognized (Wood et al., 2005; Reinecke 2010). Although the relative importance of endocrine and local IGF-I in fish growth has not been elucidated, circulating IGF-I generally reflects growth status (Picha et al., 2008a; Beckman, 2011). IGFBPs are also present in teleosts and their structure is well conserved (Wood et al., 2005; Kelley et al., 2006). However, due to an extra round of whole genome duplication in the teleost lineage, fish often have two copies of the each member of IGFBPs (Daza et al., 2011). Two *igfbp-1* subtypes were first identified in zebrafish (*Danio rerio*) (Kamei et al., 2008). Zebrafish IGFBP-1 co-orthologs were expressed predominantly in the liver and exhibited overlapping inhibitory actions on cell proliferation, although temporal expression patterns during embryonic development, response to hypoxia and affinity for IGF were different (Kamei et al., 2008). We identified two of three major IGFBPs of Chinook salmon (*Oncorhynchus tshawytscha*) as

IGFBP-1a and -1b and showed that both subtypes were induced under osmotic stress, suggesting that they have inhibitory actions on IGF-I (Shimizu et al., 2011). However, sites of their expression were different; *igfbp-1b* was exclusively expressed in the liver while *igfbp-1a* was detected in a variety of tissues (Shimizu et al., 2011). Recent detailed genomic analyses by Macqueen et al. (2013) revealed that salmonid species have 19 IGFBPs due to the tetraploidy origin of this group and the two IGFBP-1 subtypes of Chinook salmon corresponded to *igfbp-1a1* and *-1b1*, respectively. Tissue distributions of Atlantic salmon (*Salmo salar*) *igfbp-1a1* and *-1b1* were also different as is the case for Chinook salmon (Macqueen et al., 2013). These findings suggest that salmon IGFBP-1 subtypes partition their IGF-inhibitory actions, where the modes of IGFBP-1a and -1b actions are mainly local and systemic, respectively.

Analyses of the environmental modulation of IGF-I and IGFBPs in fish are important to understand how growth is adjusted to maximize performance and survival. Food availability, temperature, stress, salinity and water quality are major environmental factors affecting fish growth. Among these factors, the effect of feeding ration or/and fasting on circulating IGF-I in fish has been most intensively studied (Picha et al., 2008a; Beckman, 2011). In post-smolt coho salmon (*O. kisutch*), plasma IGF-I well reflected feeding status and growth rate (Beckman et al., 2004). In contrast, plasma IGFBP-1a and -1b levels were inversely related to feeding and growth status (Kawaguchi et al., 2013). Acute stress such as hypoxia, direct seawater transfer and handling increased tissue or circulating *igfbp-1*/IGFBP-1 in several fish species (Kelley et al., 2006; Kamei et al., 2008; Shimizu et al., 2011) and chronic stress reduces growth most likely through suppressing circulating IGF-I levels. Although the liver is the major site of production of circulating IGF-I and IGFBP-1, other tissues also express them and locally regulate growth. However, it is not well understood how local *igf-1* and *igfbp-1* respond to a change in food availability.

Adjusting growth under different salinities is crucial for euryhaline fishes and such modulation should be achieved by changes in IGF-I and IGFBPs. Salinity could affect IGF-I and IGFBPs directly at local tissues or indirectly through changes in GH and cortisol (Reinecke, 2010). Gill *igf-1* and in some case liver *igf-1* were shown to change when rainbow trout (*O. mykiss*) were transferred from freshwater (FW) to seawater (SW) (Sakamoto and Hirano, 1993; Poppinga et al., 2007). Plasma IGF-I in trout was also increased in a higher salinity (66‰ SW; Shepherd et al., 2005). Despite relatively a large number studies examined response of local or circulating IGF-I to salinity change in fish, there is no consensus on the direction of response; It

appears to vary depending on species, stages, salinity and duration (Reinecke et al., 2010; Beckman, 2011). On one hand, only a few studies using salmonids examined effect of salinity on plasma IGFBPs (Shepherd et al., 2006; Shimizu et al., 2007; 2011) and responses of *igfbp* mRNAs to salinity change are scarce.

In addition to the mechanistic importance of IGF-I and IGFBPs in growth regulation, they have drawn much attention from fish endocrinologists/biologists because of their potential utility as growth indices (Picha et al., 2008a; Beckman, 2011). Estimating growth rate of fish is important in aquaculture, and IGF-I and IGFBP-1 have been proposed as positive and negative growth indices, respectively. Evaluating growth status of larvae, fry and juvenile fish is also relevant to stock assessment since growth-dependent mortality is a key determinant of stock recruitment (Beamish and Mahnken, 2001). Circulating IGF-I is so far the most reliable growth index in several fishes (Picha et al., 2008a; Beckman, 2011). However, it is not clear whether *igf-1* mRNA can be used as an alternative of plasma IGF-I. Moreover, utility of *igfbp-1a* and *-1b* as negative growth indices are not fully validated.

Chum salmon (*O. keta*) is one of the eight Pacific salmon and an obligatory anadromous species; They hatch in freshwater and all juveniles weighing around 1 g migrate to SW in the first spring. This species is an important commercial fish in Japan and target of intensive hatchery release. However, their returns fluctuate between years and local regions (Miyakoshi et al., 2013). They are also believed to suffer growth-dependent mortality during the early phase of their marine life (Bax, 1983; Fukuwaka and Suzuki, 2002; Wertheimer and Thrower, 2007). Therefore, an accurate assessment of growth status of juvenile chum salmon at the estuary and nearshore is important to assess their survival. We have recently shown that plasma IGF-I is a good growth index in juvenile chum salmon in SW (Kaneko et al., 2015). However, it is not known if salinity change affects basal IGF-I levels and its sensitivity to food deprivation. Moreover, since chum salmon juveniles are sometimes so small that collecting blood at field survey is not practical, availability of other growth indices that can be used as alternative to serum IGF-I is valuable. The aims of the present study are twofold: 1) to analyze responses of IGF-I/*igf-1* and *igfbp-1s* to salinity change and fasting, and 2) to examine their utility as growth indices by comparing with a biochemical growth index, RNA/DNA ratio.

2. Materials and methods

2.1. Fish and rearing experiments

In May 2014, underyearling chum salmon (fork length (FL): 4.58 ± 0.08 cm; body weight (BW): 0.64 ± 0.03 g) were obtained from a local hatchery (Kamisato Hatchery) in Abashiri area, northeastern Hokkaido, Japan, transferred to the rearing facility at Faculty of Fisheries Sciences, Hokkaido University and reared in freshwater glass aquariums ($60 \times 29.5 \times 36$ cm³; water volume: 60L) in a temperature-controlled room (10°C). Each aquarium was a closed-circulation system installed with a portable upper filter system. Fish were fed daily on a commercial diet (Marubeni Nisshin Feed Co. Ltd., Tokyo, Japan) to satiety. Three days after transfer, fish were divided into two groups: FW and SW groups. SW group was acclimated to full-strength (33 ppt) artificial SW (Marine Salt Pro; Spectrum Brands Inc., Tokyo, Japan) by increasing the salinity stepwisely over 5 days (6.6 ppt/day). Salinity was monitored by using a salinity meter (MotherTool, Nagano, Japan). They were transferred to four SW tanks and acclimated for 2 additional days. No mortality occurred during SW transfer. FW group was transferred to four FW tanks and acclimated for 2 days. Fish were fed during the acclimation/transfer period. Each group was further divided into Fed and Fasted groups to make four treatments (20 fish/treatment): FW-Fed, FW-Fasted, SW-Fed and SW-Fasted. Two tanks were used for each treatment (i.e. duplicated tanks/treatment). Fed groups were fed twice daily to satiety for 2 weeks while Fasted groups received no feed for the same period. Salinity was kept at full-strength seawater (31-34 ppt) and water temperature was maintained between 11.0-11.5°C during the experiment. Water was replaced one time at 1 week after fasting. Water quality (pH, NO₃⁻, NO₂⁻, HCO₃⁻, Cl₂, and general and carbonate hardness) was monitored by using a test kit (Tetra test; Spectrum Brands, Tokyo, Japan). The experiment was carried out in accordance with the guideline of the Hokkaido University Animal Care and Use Committee. Eight fish from each treatment were sampled at 0, 1, 2 and 3 weeks after the beginning of SW acclimation. Fish were anesthetized using 3.3% 2-phenoxyethanol (Kanto Chemical, Tokyo, Japan). After FL and BW were measured, the tail was cut and blood was withdrawn using 10 or 20 µL plain glass tubes (Microcap; Drummond Scientific Company, Broomall, PA, USA). Blood was allowed to clot overnight at 4°C and centrifuged at 10,000 rpm for 15 min. Serum was collected and stored at -80°C until use. Gills were collected and immediately frozen on dry ice and stored at -80°C. Two pieces of the liver were collected; One piece was immersed in RNAlater (Thermo Fisher Scientific, Waltham, MA), sit at 4°C overnight and stored at -30°C, and the other piece was frozen on dry ice and stored at -80°C. Muscle samples were collected similar to the liver.

In June 2014, juveniles (FL: 5.80 ± 0.17 cm; BW: 1.37 ± 0.12 g) that had been

acclimated to artificial SW were lightly anesthetized using 2-phenoxyethanol (approximately 1%) and individually marked with PIT-tags (size $\phi 1.4 \text{ mm} \times 8.4 \text{ mm}$, Biomark, Boise, ID). They were randomly placed into two 60 L SW tanks (25 fish per tank) and one group was fed twice daily on the commercial diet with a ration at 3.0% body weight/day for 10 days. The other group was fasted throughout the experimental period. SW was not replaced during the experiment. The experiment was carried out in accordance with the guideline of the Hokkaido University Animal Care and Use Committee. FL and BW of all fish were measured at the beginning of the experiment and 10 days after treatment. Condition factor (K) was calculated as follows: $(\text{BW (g)}) \times 1000 / (\text{FL (cm)})^3$. Specific growth rate (SGR) was calculated as follows: $\text{SGR (\%/day)} = \ln(s_2 - s_1) \times (d_2 - d_1)^{-1} \times 100$, where s_2 is length or weight on day₁ and $d_2 - d_1$ is the number of days among measurements. On day 10, fish from each treatment (Fed: $n = 25$, Fasted $n = 24$) were sampled for blood, liver and muscle as described above.

2.3. Na^+, K^+ -ATPase activity assay

Gill NKA activity was measured according to Quabius et al. (1997) with minor modifications. This method is based on the ability of NKA to hydrolyze ATP to give ADP and inorganic phosphorus with or without presence of ouabain at 37°C for 10 min. Liberated inorganic phosphorus reacted with ammonium molybdate was quantified by measuring absorbance at 630 nm using a spectrophotometer (Corona Electronic, Ibaraki, Japan). Protein concentration was measured by using BCA (bicinchoninic acid) Protein Assay Kit (Thermo Scientific, IL). The activity was expressed as Pi (μmol) per protein (mg) per period (h).

2.4. Time-resolved fluoroimmunoassay (TR-FIA) for IGF-I

Prior to the assay, serum IGF-I was extracted with an acid-ethanol as described in Shimizu et al. (2000). IGF-I was quantified by TR-FIA based on the method described in Small and Peterson (2005) using recombinant salmon/trout IGF-I (GroPep, Adelaide, SA, Australia) for standard and labeling with europium, and anti-barramundi IGF-I (GroPep) as a primary antiserum.

2.5. Measurement of RNA/DNA ratio

RNA/DNA ratio was measured by a spectrofluorimetric method modified from Grémare and

Vétion (1994) as described in Kawaguchi et al. (2013). Briefly, amount of total nucleic acids (DNA + RNA) was measured using 4 µg/mL Thiazole orange (Sigma-Aldrich, St. Louis, MO) and that of DNA using 0.02 mg/mL Hoechst 33258 (Dojindo, Kumamoto, Japan). RNA/DNA ratio was calculated from these values.

2.6. RNA extraction and cDNA synthesis

Total RNA was extracted from livers and muscles using ISOGEN (Nippon gene; Tokyo, Japan) according to the manufacturer's instruction. One and half µg RNA was reverse-transcribed using SuperScript VILO cDNA Synthesis kit (Invitrogen, Carlsbad, CA, USA) in a 10-µl reaction according to the manufacturer's instruction. cDNA was stored at -30°C until use.

2.7. Real-time quantitative PCR (qPCR)

Sequences of primers for qPCR of *igf-1*, *igfbp-1a*, *igfbp-1b* and *ef-1a* were the same as described in Kawaguchi et al. (2013). Reverse transcribed-PCRs using these primers were performed to prepare assay standards for chum salmon. PCR products run on 1.5% agarose gel were excised and purified using QIAEX II Gel Extraction Kit (Qiagen, Valencia, CA, USA). Copy numbers of the purified amplicon were calculated from the molecular weight of the amplicon and concentration. The standard cDNA were serially diluted from 1×10^7 to 3×10^2 copies.

qPCR was set up using Power SYBR Green PCR Master Mix (Applied Biosystems, Carlsbad, CA, USA) in a reaction volume of 20 µl with primer concentration of 100 nM. qPCR was run on a 7300 Sequence Detector (Applied Biosystems) using the manufacturer's recommended cycling conditions: 50°C for 2 min, 95°C for 10 min followed by 40 cycles at 95°C for 15 se and 60°C for 1 min. Measured values were normalized to those of *ef-1a*. Performance of qPCR was evaluated by confirming a single peak of the dissociation curve in each assay and calculating the amplification efficiencies of the standard curves, which were within the range of 97-100%. Coefficients of determination of the standard curves were also between 0.99-1.00.

2.8. Statistical analysis

Results of the experiments were analyzed first by three-way ANOVA (salinity x time x nutrition) using the JMP program (SAS Institute Inc., Cary, NC, USA). When significant effects were found, differences were further identified by one-way ANOVA followed by the Fisher's protected least significant difference (PLSD) test. Differences among groups were considered to be significant at $P < 0.05$. SGRs from the second experiment were biased to high and low values. In order to avoid anchor effects, the SGR data were first categorized into high, med and low, and eight samples were randomly selected from each category to analyze relationships between growth rate and the endocrine/biochemical parameters by simple linear regression.

3. Results

3.1. Effect of SW transfer

Average FL, BW and K of each treatment are shown in Table 1. There were no effects of SW transfer for 1 week on FL, BW and K. Gill NKA activity was lower in fish in FW (Fig. 1). There was no effect of SW transfer on serum IGF-I (Fig. 2). SW transfer had no effect on liver RNA/DNA ratio whereas muscle RNA/DNA was higher in fish in SW (Fig. 3a,b). Both liver and muscle *igf-1* were not affected by SW transfer (Fig. 3c,d). Liver *igfbp-1a* levels were not influenced by SW transfer but those in the muscle significantly increased 1 week after SW transfer (Fig. 4a,b). There was no effect of SW transfer on liver *igfbp-1b* (Fig. 4c).

3.2. Combined effects of salinity and fasting

FL, BW and K were reduced by fasting for 2 weeks in both salinities (Table 1). There were overall effects of salinity and fasting on gill NKA activity, and fasting for 2 weeks significantly reduced the activity in both salinities (Fig. 5). Serum IGF-I levels were reduced by fasting in both salinities and there was no interaction with salinity (Fig. 6). In addition, serum IGF-I levels in SW-Fed fish was higher than those in FW-Fed fish (Fig. 6). Liver RNA/DNA was reduced by fasting and prolonged fasting further reduced it in SW fish (Fig. 7a). In contrast, the effect of fasting on muscle RNA/DNA was found only in fish in SW after 2 weeks (Fig. 7b). There were overall effects of fasting to reduce liver and muscle *igf-1* (Fig. 7c,d). However, fasting effect on liver *igf-1* was seen in 1 week while it took 2 weeks of fasting to see the reduction of muscle *igf-1* (Fig. 7c,d). In addition, an overall effect of salinity on muscle *igf-1* was seen (Fig. 7d).

There were overall effects of fasting to increase both liver and muscle *igfbp-1a*, while no significant difference between Fed and Fasted groups 1 week after fasting was detected in both tissues (Fig. 8a,b). In muscle, effect of fasting on *igfbp-1a* was seen only in SW group at week 2 (Fig. 8b). Fasting also increased liver *igfbp-1b* in both salinities and the effect was first detected in 1 week in SW group (Fig. 8c).

3.3. Relationships of endocrine and biochemical parameters with growth

Data on FL, BW, K, SGR-FL and SGR-BW of juvenile chum salmon fed or fasted for 10 days in SW are provided in Supplemental Table 1. The relationships between the endocrine/biochemical parameters and growth parameters were analyzed (Table 2). No endocrine/biochemical parameters correlated with FL. Serum IGF-I showed a moderate positive relationship with BW and weak negative relationships were seen for liver *igfbp-1a* and *-1b*. Liver RNA/DNA and serum IGF-I showed moderate positive relationships with K while liver and muscle *igfbp-1a* and liver *igfbp-1b* were negatively correlated with K. Serum IGF-I showed the highest positive relationship with SGR-FL. Negative, weak relationships with SGR-FL were seen for liver *igfbp-1b* and muscle *igf-1*. Serum IGF-I also had the highest regression coefficient with SGR-BW. In addition, muscle *igf-1* had a moderate positive relationship with SGR-BW whereas negative relationships were seen for liver *igfbp-1a* and *-1b*. Scattered plots between the parameters and SGR-BW are shown in Figure 9.

4. Discussion

Chum salmon juveniles experience changes in salinity and food availability during their seaward migration. The general belief is that growth soon after sea entry is key for their survival (Bax, 1983; Fukuwaka and Suzuki, 2002; Wertheimer and Thrower, 2007). The present study examined effects of SW transfer and fasting on IGF-I/*igf-1* and *igfbp-1s* in order to analyze environmental modulation of the IGF/IGFBP system in chum salmon as well as assess their utility as growth indices.

Chum salmon is an obligatory anadromous species migrating to the sea in their first spring at fry/juvenile stage often weighing less than 1 g. This species is unique in terms of acquiring high hypo-osmoregulatory ability soon after yolk absorption and maintaining it for relatively long period of time in freshwater (Iwata et al., 1982; Hasegawa et al., 1986). Kojima

et al. (1993) reported a temporal decrease in hypo-osmoregulatory ability in chum salmon juveniles weighing 3-4 g but those fish were still able to restore increased plasma sodium ion concentrations in 24 hr after direct SW transfer. In the present study, chum salmon fry/juveniles (0.6 g) were acclimated stepwisely to full-strength SW (33 ppt) over 5 days. In addition to their high hypo-osmoregulatory ability before entering SW, the gradual acclimation employed in the present study should completely acclimated fish to SW without stress. Although we did not measure plasma osmolarity or ion concentrations, or plasma cortisol due to the limitation of serum volume, the high gill NKA activity before SW transfer and no further activation of the activity suggest their high hypo-osmoregulatory ability. In addition, fish were fed during acclimation. Thus, we considered changes in the parameters during the first week of SW transfer were responses to salinity change rather than osmotic stress or fasting.

There are a number of studies examined effect of salinity change on tissue *igf-1* or circulating IGF-I in fish. Sakamoto and Hirano (1993) reported transfer of rainbow trout to 26 ppt (80%) SW resulted in transient increases in gill and body kidney *igf-1* at 1 day and 8 days, respectively, but not in liver *igf-1*. In contrast, liver *igf-1* in the same species increased 12 hr after transfer to 20 ppt (61%) SW (Poppinga et al., 2007). Exposure of Caspian trout (*S. trutta caspius*) to brackish water (10.5 ppt) caused an increase in liver *igf-1* in 5 days (Sharif et al., 2015). Shepherd et al. (2005) reported up-regulation of plasma IGF-I in rainbow trout 5 days after gradual transfer to 22 ppt (66%) SW. In the present study, serum IGF-I, liver *igf-1* and muscle *igf-1* were not altered by transfer to full SW (33 ppt) for up to 3 weeks. It is hard to compare these responses to other studies since species, salinity, feeding status and experimental settings are different. However, one study by Iwata et al. (2012) used chum salmon fry/juveniles similar to the present study. When chum salmon juveniles (fry) were transferred to SW, hepatic *igf-1* levels were similar, or even lower, to those in FW during first 2 weeks and became higher 4 weeks after transfer (Iwata et al., 2012). These results suggest that effect of SW transfer on hepatic *igf-1* in juvenile chum salmon is not acute and our finding supports the notion. The present study also reports no acute effect of SW transfer on muscle *igf-1* in chum salmon.

Effects of salinity on IGFBPs have been investigated in a few studies using trout/salmon. Shepherd et al. (2005) acclimated rainbow trout to 22 ppt SW and found increases in plasma levels of 42-50-kDa (most likely IGFBP-2b) and 21-kDa IGFBPs (most likely IGFBP-1b). A direct exposure to full-strength SW of underyearling Chinook salmon fall smolts had no effect on serum IGFBP-2b but induced both serum IGFBP-1a and -1b (Shimizu et al., 2011), suggesting the involvement of IGFBP-1s in osmoregulation or/and stress response. On

one hand, no studies examined effect of salinity on tissue *igfbp-1*. In the present study using juvenile chum salmon, both liver *igfbp-1a* and *-1b* did not respond to SW transfer, while muscle *igfbp-1a* levels increased by SW transfer but returned to the basal levels 2 weeks after transfer. These responses may be adjustment by the muscle to different salinity rather than stress response. Specific role of IGFBP-1a in the muscle during the early phase of SW adaptation is a subject of future research.

High activity of branchial NKA is essential for salmon to maintain the internal osmolarity and ion balance in SW. As in its name, NKA consumes ATP to extrude sodium ions in exchange with potassium ions. Under the shortage of nutritional input, decreased gill NKA activity and increased Cl⁻ ions were observed in rainbow trout and Atlantic salmon (Jürss et al., 1987; Stefansson et al., 2009). Consistent with the previous studies, a negative effect of fasting on gill NKA activity was observed in juvenile chum salmon. A decrease in gill NKA activity during fasting may be adaptive in terms of saving energy, but long-term disturbance of ion balance under such condition should increase risk of mortality of juvenile salmon in wild.

Circulating IGF-I has been shown to be sensitive to changes in nutritional input. Beckman et al. (2004) reported plasma IGF-I levels in post-smolt coho salmon responded well to varying feeding ration, being higher in fish with high feeding ration. Fasting reduced plasma IGF-I levels in Chinook salmon as fast as 4 days (Pierce et al., 2005). In the present study, fasting for 1 week reduced serum IGF-I levels in fish in both FW and SW and prolonged fasting for another week tended to further reduce it. Thus, the response of serum IGF-I levels in juvenile chum salmon to fasting is consistent with the previous studies. It is worth noting that IGF-I levels in SW-Fed fish were higher than those in FW-Fed fish 3 weeks after transfer, suggesting that activation of the somatotrophic axis in SW.

The liver is the major site of production of circulating IGF-I while virtually all tissues express *igf-1*. Responses of *igf-1* in the liver and muscle to fasting have been compared in several fish species. In rainbow trout and channel catfish (*Ictalurus punctatus*), fasting for 30 days reduced both liver and muscle *igf-1* and refeeding for 15 days was enough to restore its levels (Gabillard et al., 2006; Peterson and Waldbieser, 2009). On the other hand, a higher sensitivity of muscle *igf-1* to fasting than liver *igf-1* has been reported in Mozambique tilapia (*Oreochromis mossambicus*), yellowtail (*Seriola quinqueradiata*) and a cichlid fish (*Cichlasoma dimerus*) (Fox et al., 2010; Fukada et al., 2012; Delgadin et al., 2015), suggesting local regulation of *igf-1* is important to adjust tissue growth under fasting condition. It should be emphasized that the results of these studies do not necessarily deny the importance of hepatic

igf-1 in growth regulation since the mRNA level is a reflection of the balance between translation and degradation. In the present study, the basal mRNA levels of *igf-1* were higher in the liver than in the muscle, and hepatic *igf-1* responded to fasting for 1 week while muscle *igf-1* levels were not reduced by fasting for 1 week but 2 weeks. These results suggest that in juvenile chum salmon, liver *igf-1* is more sensitive to fasting. Thus, the sensitivity of tissue *igf-1* to fasting appears to vary depending on species, stage and conditions. Salinity doesn't seem to have a strong impact on tissue (liver and muscle) *igf-1* although there was a small but significant difference in muscle *igf-1* levels between FW-Fed and SW-Fed groups.

The present study compared the responses of liver and muscle *igfbp-1a* to fasting as well as those between *igfbp-1a* and *-1b* in the liver. Fasting fish usually causes an increase in hepatic *igfbp-1a/b* (Kamei et al., 2008; Pedroso et al., 2009; Peterson and Waldbieser, 2009; Kawaguchi et al., 2013; Breves et al., 2014), which is adaptive to save energy under nutritional deficiency (Kajimura and Duan, 2007). Selection for large size resulted in lower muscle expression of *igfbp-1a* and *-1b* in zebrafish (Amaral and Johnston, 2012), suggesting IGFBP-1 negatively regulate growth. However, there is no study comparing their responses to fasting between different tissues. In the present study, *igfbp-1a* was expressed both in the liver and muscle at the similar levels while *igfbp-1b* was detectable only in the liver. The exclusive hepatic expression of *igfbp-1b* has been also reported in tilapia (Breves et al., 2014). Hepatic and muscle *igfbp-1a* were increased by fasting in FW as well as in SW, suggesting the importance of muscle IGFBP-1a under fasting conditions. Our finding is in good agreement with those in zebrafish (Amaral and Johnston, 2011, 2012) where muscle *igfbp-1a* responded to fasting and refeeding. In zebrafish, *igfbp-1b* was also expressed in the muscle and showed the similar responses to fasting and refeeding although the levels were much lower than *igfbp-1a* (Amaral and Johnston, 2011). Liver *igfbp-1b* was increased by fasting in both FW and SW as is the case for liver *igfbp-1a*, but a significant increase was first seen in SW fish in 1 week. In addition, a larger response of muscle *igfbp-1a* to fasting was found in SW fish. These results suggest that the sensitivity of *igfbp-1s* is higher in SW although how salinity influences the fasting response of *igfbp-1s* is not known at present. IGFBP-1s produced in the liver under fasting condition should be released into the bloodstream and act on circulating IGF-I to adjust growth rate.

One of the objectives of the present study was to validate the endocrine and biochemical parameters as growth indices for juvenile chum salmon. There are considerable interests in assessing growth status of fish using growth indices in aquaculture industry and

stock assessment. The latter is based on the hypothesis that high growth-dependent mortality soon after SW entry occurs in juvenile chum salmon (Bax, 1980; Fukwaka and Suzuki, 2002; Wertheimer and Thrower, 2007). However, since chum salmon juveniles experience large changes in environment such as salinity and food availability during downstream and coastal migration, good candidates of growth indices for this species should be stable during salinity change, sensitive to nutritional change and well correlated with growth rate. We have recently reported that serum IGF-I in juvenile chum salmon acclimated in SW was strongly, positively correlated with growth rate (Kaneko et al., 2015). The present study revealed that IGF-I was stable after SW transfer and there was no interaction between fasting response and salinity. The results of the present study strengthen the utility of serum IGF-I as a growth index in juvenile chum salmon. In addition, the present study compared IGF-I with the other candidates of growth index in terms of correlation with growth rate. As a result, serum IGF-I still showed the strongest relationship with growth rate. This result confirms that serum IGF-I is a good growth index for juvenile chum salmon.

The reliability of circulating IGF-I as growth index has been examined and discussed (Picha et al., 2008a; Beckman, 2011). Although the basal levels of circulating IGF-I fluctuates in response to season, sudden water temperature change, developmental stage and maturation status, it is so far the most reliable growth index in fish (Picha et al., 2008a; Beckman, 2011). However, collecting blood from small fish is sometimes not possible/practical. Chum salmon start migrating to the ocean even they are below 1 g. In such case, measuring mRNA levels of growth indices is an alternative. The present study assessed the utility of *igf-1* and *igfbp-1s* as alternative growth indices for juvenile chum salmon. Liver *igf-1* was sensitive to fasting but not correlated with grow rate. On the other hand, muscle *igf-1* was less sensitive to fasting but showed a moderate, positive correlation with growth rate. This is consistent with the finding in a hybrid striped bass (*Morone chrysops* x *M. saxatilis*) where muscle *igf-1* was positively correlated with growth rate ($r^2 = 0.23$) (Picha et al., 2014). However, it should be noted that the same authors reported that liver *igf-1* also correlated with growth rate ($r^2 = 0.48$) in the hybrid striped bass (Picha et al., 2008b). Although there is species difference in relationships of liver and muscle *igf-1* with growth rate, our finding shows that muscle *igf-1* is a better indicator of growth rate than liver *igf-1* in juvenile chum salmon in SW.

In addition to muscle *igf-1*, *igfbp-1a* and *-1b* may also be alternatives of serum IGF-I. We reported that circulating protein and hepatic mRNA levels of IGFBP-1a and -1b were negatively correlated with growth rate in post-smolt masu salmon in FW (Kawaguchi et al.,

2013). In accord with the previous study, hepatic *igfbp-1a* and *-1b* had negative relationships with growth rate in juvenile chum salmon. Their coefficients of determination were moderate being not as strong as that of serum IGF-I. Small serum volume prevented us to measure serum levels of IGFBP-1a and -1b in the present study. Although their protein and hepatic mRNA levels are expected to be similar as is the case in masu salmon (Kawaguchi et al., 2013), it needs to be confirmed in future research. Muscle *igfbp-1a* responded to fasting but showed no significant relationship with growth rate, indicating that the liver is the tissue to be analyzed when *igfbp-1a* is used as a growth index.

RNA/DNA ratio has been used as a biochemical index of growth in fish and marine organisms (Chícharo and Chícharo, 2008). The ratio is believed to reflect the degree of protein synthesis (RNA) per unit cell (DNA) and thus growth. Muscle RNA/DNA has been shown to be well correlated with growth rate in Atlantic salmon smolts (Maclean and Caldarone, 2008). The present study analyzed liver and muscle RNA/DNA ratio to compare its utility with tissue *igf-1* and *igfbp-1s* as an alternative of plasma IGF-I. As results, in juvenile chum salmon liver and muscle RNA/DNA ratio showed low or no correlation with growth rate, making it difficult to use as an alternative of serum IGF-I in this species.

In conclusion, our findings suggest that in chum salmon IGF-I and IGFBP-1s adjust growth under fasting condition by responding systemically as well as locally. The present study also shows that serum IGF-I is superior as a growth index for juvenile chum salmon. However, when juveniles are too small to collect blood, liver *igfbp-1a* and *-1b* and muscle *igf-1* are alternatives of IGF-I.

Acknowledgment

We thank staffs of the Kitami Salmon Enhancement Programs Association for providing juvenile chum salmon. This work was supported by the Suhara Memorial Foundation and Hokusui Foundation.

References

- Amaral, I.P.G., Johnston, I.A., 2011. Insulin-like growth factor (IGF) signalling and genome-wide transcriptional regulation in fast muscle of zebrafish following a single-satiating meal. *J. Exp. Biol.* 214, 2125-2139.
- Amaral, I.P.G., Johnston, I.A., 2012. Experimental selection for body size at age modifies early life-history traits and muscle gene expression in adult zebrafish. *J. Exp. Biol.* 215, 3895-3904.
- Bax, N.J., 1983. Early marine mortality of marked juvenile chum salmon (*Oncorhynchus keta*) released into Hood Canal, Puget Sound, Washington, in 1980. *Can. J. Fish. Aquat. Sci.* 40, 426-435.
- Beamish, R.J., Mahnken, C., 2001. A critical size and period hypothesis to explain natural regulation of salmon abundance and the linkage to climate and climate change. *Prog. Oceanogr.* 49, 423-437.
- Beckman, B.R., 2011. Perspectives on concordant and discordant relations between insulin-like growth factor 1 (IGF1) and growth in fishes. *Gen. Comp. Endocrinol.* 170, 233-252.
- Beckman, B.R., Shimizu, M., Gadberry, B.A., Cooper, K.A., 2004. Response of the somatotrophic axis of juvenile coho salmon to alterations in plane of nutrition with an analysis of the relationships among growth rate and circulating IGF-I and 41 kDa IGFBP. *Gen. Comp. Endocrinol.* 135, 334-344.
- Breves, J.P., Tipsmark, C.K., Stough, B.A., Seale, A.P., Flack, B.R., Moorman, B.P., Lerner, D.T., Grau, E.G., 2014. Nutritional status and growth hormone regulate insulin-like growth factor binding protein (*igfbp*) transcripts in Mozambique tilapia. *Gen. Comp. Endocrinol.* 207, 66-73.
- Chícharo, M.A., Chícharo, L., 2008. RNA : DNA ratio and other nucleic acid derived indices in marine ecology. *Int. J. Mol. Sci.* 9, 1453-1471.
- Daughaday, W.H., Rotwein, P., 1989. Insulin-like growth factors I and II. Peptide, messenger ribonucleic acid and gene structures, serum, and tissue concentrations. *Endocr. Rev.* 10, 68-91.
- Daza, D.O., Sundstrom, G., Bergqvist, C.A., Duan, C.M., Larhammar, D., 2011. Evolution of the insulin-like growth factor binding protein (IGFBP) family. *Endocrinology* 152, 2278-2289.
- Delgadín, T.H., Sirkin, D.I.P., Di Yorio, M.P., Arranz, S.E., Vissio, P.G., 2015. GH, IGF-I and

501 GH receptors mRNA expression in response to growth impairment following a food
 502 deprivation period in individually housed cichlid fish *Cichlasoma dimerus*. Fish Physiol.
 503 Biochem. 41, 51-60.

504 Fiess, J.C., Kunkel-Patterson, A., Mathias, L., Riley, L.G., Yancey, P.H., Hirano, T., Grau, E.G.,
 505 2007. Effects of environmental salinity and temperature on osmoregulatory ability,
 506 organic osmolytes, and plasma hormone profiles in the Mozambique tilapia
 507 (*Oreochromis mossambicus*). Comp. Biochem. Physiol. A 146, 252-264.

508 Firth, S.M., Baxter, R.C., 2002. Cellular actions of the insulin-like growth factor binding
 509 proteins. Endocr. Rev. 23, 824-854.

510 Fox, B.K., Breves, J.P., Davis, L.K., Pierce, A.L., Hirano, T., Grau, E.G., 2010. Tissue-specific
 511 regulation of the growth hormone/insulin-like growth factor axis during fasting and
 512 re-feeding: Importance of muscle expression of IGF-I and IGF-II mRNA in the tilapia.
 513 Gen. Comp. Endocrinol. 166, 573-580.

514 Fukada, H., Murashita, K., Furutani, T., Masumoto, T., 2012. Yellowtail insulin-like growth
 515 factor 1: molecular cloning and response to various nutritional conditions. Domest.
 516 Anim. Endocrinol. 42, 220-229.

517 Fukuwaka, M., Suzuki, T., 2002. Early sea mortality of mark-recaptured juvenile chum salmon
 518 in open coastal waters. J. Fish Biol. 60, 3-12.

519 Gabillard, J.C., Kamangar, B.B., Montserrat, N., 2006. Coordinated regulation of the GH/IGF
 520 system genes during refeeding in rainbow trout (*Oncorhynchus mykiss*). J. Endocrinol.
 521 191, 15-24.

522 Grémare, A., Vétion, G. 1994. Comparison of several spectrofluorimetric methods for
 523 measuring RNA and DNA concentrations in the deposit-feeding bivalve *Abra ovata*.
 524 Comp. Biochem. Physiol. 107B, 297-308.

525 Hasegawa, S., Hirano, T., Ogasawara, T., Iwata, M., Akiyama, T., Arai, S. 1987.
 526 Osmoregulatory ability of chum salmon, *Oncorhynchus keta*, reared in fresh water for
 527 prolonged periods. Fish Physiol. Biochem. 4, 101-110.

528 Iwata, M., Hirano, T., Hasegawa, S. 1982. Behavior and plasma sodium regulation of chum
 529 salmon fry during ransition into seawater. Aquaculture 28, 133-142.

530 Iwata, M., Kinoshita, K., Moriyama, S., Kurosawa, T., Iguma, K., Chiba, H., Ojima, D.,
 531 Yoshinaga, T., Arai, T., 2012. Chum salmon fry grow faster in seawater, exhibit greater
 532 activity of the GH/IGF axis, higher Na⁺, K⁺-ATPase activity, and greater gill chloride
 533 cell development. Aquaculture 362, 101-108.

534 Jones, J.I., Clemmons, D.R., 1995. Insulin-like growth factors and their binding proteins:
535 biological actions. *Endocr. Rev.* 16, 3-34.

536 Jürss, K., Bittorf, T., Vokler, T., Wacke, R., 1987. Effects of temperature, food deprivation and
537 salinity on growth, RNA/DNA ratio and certain enzyme activities in rainbow trout
538 (*Salmo gairdneri* Richardson). *Comp. Biochem. Physiol. B* 87, 241-253.

539 Kajimura, S., Duan, C., 2007. Insulin-like growth factor-binding protein-1: an evolutionarily
540 conserved fine tuner of insulin-like growth factor action under catabolic and stressful
541 conditions. *J. Fish Biol.* 71, 309-325.

542 Kamei, H., Lu, L., Jiao, S., Li, Y., Gyrup, C., Laursen, L.S., Oxvig, C., Zhou, J., Duan, C., 2008.
543 Duplication and diversification of the hypoxia-inducible IGFBP-1 gene in zebrafish.
544 *PLoS One* 3, e3091.

545 Kaneko, N., Taniyama, N., Inatani, Y., Nagano, Y., Fujiwara, M., Torao, M., Miyakoshi, Y.,
546 Shimizu, M., 2015. Circulating insulin-like growth factor I in juvenile chum salmon:
547 relationship with growth rate and changes during downstream and coastal migration in
548 northeastern Hokkaido, Japan. *Fish Physiol. Biochem.* 41, 991-1003.

549 Kawaguchi, K., Kaneko, N., Fukuda, M., Nakano, Y., Kimura, S., Hara, A., Shimizu, M., 2013.
550 Responses of insulin-like growth factor (IGF)-I and two IGF-binding protein-1 subtypes
551 to fasting and re-feeding, and their relationships with individual growth rates in yearling
552 masu salmon (*Oncorhynchus masou*). *Comp. Biochem. Physiol. A.* 165, 191-198.

553 Kelley, K.M., Price, T.D., Galima, M.M., Sak, K., Reyes, J.A., Zepeda, O., Hagstrom, R.,
554 Truong, T.A., Lowe, C.G., 2006. Insulin-like growth factor-binding proteins (IGFBPs)
555 in fish: beacons for (disrupted) growth endocrine physiology. in: Reinecke, M., Zaccone,
556 G., Kapoor, B.G. (Eds.), *Fish Endocrinology*. Science Publishers, Enfield, New
557 Hampshire, pp. 167-195.

558 Kojima, H., Iwata, M., Kurokawa, T. 1993. Development and temporal decrease in seawater
559 adaptability during early growth in chum salmon, *Oncorhynchus keta*. *Aquaculture* 118,
560 141-150.

561 Lee, P.D., Conover, C.A., Powell, D.R., 1993. Regulation and function of insulin-like growth
562 factor-binding protein-1. *Proc. Soc. Exp. Biol. Med.* 204, 4-29.

563 Le Roith, D., Bondy, C., Yakar, S., Liu, J.L., Butler, A., 2001. The somatomedin hypothesis:
564 2001. *Endocr. Rev.* 22, 53-74.

565 MacLean, S.A., Caldarone, E.M., 2008. Estimating recent growth rates of Atlantic salmon
566 smolts using RNA-DNA ratios from nonlethally sampled tissues. *Trans. Am. Fish. Soc.*

567 137, 1279-1284.

568 Macqueen, D.J., de la Serrana, D.G., Johnston, I.A., 2013. Evolution of ancient functions in the
569 vertebrate insulin-like growth factor system uncovered by study of duplicated salmonid
570 fish genomes. *Mol. Biol. Evol.* 30, 1060-1076.

571 Miyakoshi, Y., Nagata, M., Kitada, S., Kaeriyama, M., 2013. Historical and current hatchery
572 programs and management of chum salmon in Hokkaido, northern Japan. *Rev. Fish. Sci.*
573 21, 469-479.

574 Ohlsson, C., Mohan, S., Sjogren, K., Tivesten, A., Isgaard, J., Isaksson, O., Jansson, J.O.,
575 Svensson, J., 2009. The role of liver-derived insulin-like growth factor-I. *Endocr. Rev.*
576 30, 494-535.

577 Pedroso, F.L., Fukada, H., Masumoto, T., 2009. Molecular characterization, tissue distribution
578 patterns and nutritional regulation of IGFBP-1, -2, -3 and -5 in yellowtail, *Seriola*
579 *quinqueradiata*. *Gen. Comp. Endocrinol* 161, 344-353.

580 Peterson, B.C., Waldbieser, G.C., 2009. Effects of fasting on IGF-I, IGF-II, and IGF-binding
581 protein mRNA concentrations in channel catfish (*Ictalurus punctatus*). *Domest. Anim.*
582 *Endocrinol.* 37, 74-83.

583 Picha, M.E., Turano, M.J., Beckman, B.R., Borski, R.J., 2008a. Endocrine biomarkers of
584 growth and applications to aquaculture: A minireview of growth hormone, insulin-like
585 growth factor (IGF)-I, and IGF-Binding proteins as potential growth indicators in fish.
586 *N. Am. J. Aquaculture* 70, 196-211.

587 Picha, M.E., Turano, M.J., Tipsmark, C.K., Borski, R.J. 2008b. Regulation of endocrine and
588 paracrine sources of Igfs and Gh receptor during compensatory growth in hybrid striped
589 bass (*Morone chrysops* X *Morone saxatilis*). *J. Endocrinol.* 199, 81-94.

590 Picha, M.E., Biga, P.R., Galt, N., McGinty, A.S., Gross, K., Hedgpeth, V.S., Siopes, T.D., Borski,
591 R.J., 2014. Overcompensation of circulating and local insulin-like growth factor-1
592 during catch-up growth in hybrid striped bass (*Morone chrysops* X *Morone saxatilis*)
593 following temperature and feeding manipulations. *Aquaculture* 428, 174-183.

594 Pierce, A.L., Shimizu, M., Beckman, B.R., Baker, D.M., Dickhoff, W.W., 2005. Time course of
595 the GH/IGF axis response to fasting and increased ration in chinook salmon
596 (*Oncorhynchus tshawytscha*). *Gen. Comp. Endocrinol.* 140, 192-202.

597 Poppinga, J., Kittilson, J., McCormick, S.D., Sheridan, M.A., 2007. Effects of somatostatin on
598 the growth hormone-insulin-like growth factor axis and seawater adaptation of rainbow
599 trout (*Oncorhynchus mykiss*). *Aquaculture* 273, 312-319.

600 Quabius ES, Balm PHM, Bonga SEW (1997) Interrenal stress responsiveness of tilapia
 601 (*Oreochromis mossambicus*) is impaired by dietary exposure to PCB 126. Gen. Comp.
 602 Endocrinol. 108, 472-482.

603 Rajaram, S., Baylink, D.J., Mohan, S., 1997. Insulin-like growth factor-binding proteins in
 604 serum and other biological fluids: regulation and functions. Endocr. Rev. 18, 801-831.

605 Reinecke, M., 2010. Influences of the environment on the endocrine and paracrine fish growth
 606 hormone-insulin-like growth factor-I system. J. Fish Biol. 76, 1233-1254.

607 Sakamoto, T., Hirano, T., 1993. Expression of insulin-like growth factor I gene in
 608 osmoregulatory organs during seawater adaptation of the salmonid fish: possible mode
 609 of osmoregulatory action of growth hormone. Proc. Natl. Acad. Sci. USA. 90,
 610 1912-1916.

611 Sharif, S., Shoaie, A., Amiri, B.M., Farahmand, H., 2015. The effects of brackish water on
 612 growth hormone/insulin-like growth factor-1 gene expression of the Caspian trout,
 613 *Salmo trutta caspius* (Kessler, 1877), during the early stage of smoltification. J. World
 614 Aquaculture Soc. 46, 201-209.

615 Shepherd, B.S., Drennon, K., Johnson, J., Nichols, J.W., Playle, R.C., Singer, T.D., Vijayan,
 616 M.M., 2005. Salinity acclimation affects the somatotrophic axis in rainbow trout. Am. J.
 617 Physiol. Regul. Integr. Comp. Physiol. 288, R1385-1395.

618 Shimizu M, Swanson P, Fukada H, Hara A, Dickhoff WW (2000) Comparison of extraction
 619 methods and assay validation for salmon insulin-like growth factor-I using
 620 commercially available components. Gen. Comp. Endocrinol. 119, 26-36.

621 Shimizu, M., Kishimoto, K., Yamaguchi, T., Nakano, Y., Hara, A., Dickhoff, W.W., 2011.
 622 Circulating salmon 28- and 22-kDa insulin-like growth factor binding proteins
 623 (IGFBPs) are co-orthologs of IGFBP-1. Gen. Comp. Endocrinol. 174, 97-106.

624 Small BC, Peterson BC (2005) Establishment of a time-resolved fluoroimmunoassay for
 625 measuring plasma insulin-like growth factor I (IGF-I) in fish: effect of fasting on
 626 plasma concentrations and tissue mRNA expression of IGF-I and growth hormone (GH)
 627 in channel catfish (*Ictalurus punctatus*). Domest. Anim. Endocrinol. 28, 202-215.

628 Stefansson, S.O., Imsland, A.K., Handeland, S.O., 2009. Food-deprivation, compensatory
 629 growth and hydro-mineral balance in Atlantic salmon (*Salmo salar*) post-smolts in sea
 630 water. Aquaculture 290, 243-249.

631 Wertheimer, A.C., Thrower, F.P., 2007. Mortality rates of chum salmon during their early marine
 632 residency. in: Grimes, C.B., Brodeur, R.D., Haldorson, L.J., McKinnell, S.M. (Eds.),

633 The ecology of juvenile salmon in the northeast Pacific Ocena: regional comparisons.
634 American Fisheries Society, Bethesda, MD, pp. 233-247.
635 Wood, A.W., Duan, C., Bern, H.A., 2005. Insulin-like growth factor signaling in fish. Int. Rev.
636 Cytol. 243, 215-285.
637

Table 1
Fork length (FL), body weight (BW) and condition factor (K) of juvenile chum salmon used for seawater transfer and fasting experiments in May

Parameter	Treatment	Time after SW transfer		Time after fasting	
		0 week	1 week	1 week (2 wk)	2 week (3 wk)
FL	FW-Fed	4.58±0.08 ^B	4.91±0.14 ^A	5.21±0.12 ^b	5.70±0.16 ^a
	FW-Fasted			4.95±0.10 ^{bc}	4.95±0.06 ^{bc}
	SW-Fed		4.78±0.04 ^{AB}	4.96±0.08 ^{bc}	5.71±0.05 ^a
	SW-Fasted			4.88±0.07 ^c	4.85±0.10 ^c
BW	FW-Fed	0.64±0.03 ^B	0.73±0.04 ^A	0.95±0.05 ^b	1.45±0.15 ^a
	FW-Fasted			0.74±0.04 ^{cd}	0.75±0.04 ^{cd}
	SW-Fed		0.68±0.02 ^{AB}	0.88±0.04 ^{bc}	1.14±0.06 ^a
	SW-Fasted			0.70±0.03 ^{cd}	0.65±0.04 ^d
K	FW-Fed	0.67±0.04 ^A	0.62±0.03 ^A	0.67±0.03 ^{bc}	0.77±0.02 ^a
	FW-Fasted			0.60±0.01 ^d	0.62±0.03 ^{cd}
	SW-Fed		0.62±0.01 ^A	0.72±0.02 ^{ab}	0.76±0.03 ^a
	SW-Fasted			0.60±0.01 ^d	0.57±0.02 ^d

Values are expressed as mean ±SE (n=8). Numbers in parentheses are weeks after seawater transfer. Within a parameter, groups sharing the same letter are not significantly different. Effects of SW transfer, and SW transfer x fasting were analyzed separately as shown by upper and lower case letters.

Table 2
Correlation coefficients (r) between endocrine/biochemical parameters and growth parameters

	FL	BW	K	SGR-FL	SGR-BW
Liver R/D	-	-	0.65	-	0.50
Muscle R/D	-	-	-	-	-
Serum IGF-I	-	0.65	0.46	0.88	0.92
Liver <i>igf-1</i>	-	-	-	-	-
Muscle <i>igf-1</i>	-	-	0.56	0.44	0.65
Liver <i>igfbp-1a</i>	-	-0.42	-0.71	-	-0.63
Muscle <i>igfbp-1a</i>	-	-	-0.53	-	-
Liver <i>igfbp-1b</i>	-	-0.54	-0.60	-0.48	-0.59

FL: fork length; BW: body weight; K: condition factor; SGR: specific growth rate; (-): not significant.

Figure legends

Fig. 1. Effects of SW transfer on gill NKA activity in juvenile chum salmon. Values are expressed as means $\pm$ SE (n=8). Groups sharing the same letter are not significantly different.

Fig. 2. Effects of SW transfer on serum IGF-I levels in juvenile chum salmon. Values are expressed as means $\pm$ SE (n=6-8).

Fig. 3. Effects of SW transfer on liver (a,c) and muscle (b,d) RNA/DNA ratio (a,b) and *igf-I* mRNA levels(c,d) in juvenile chum salmon. Values are expressed as means $\pm$ SE (n=6-8). Groups sharing the same letter or those without letter are not significantly different.

Fig. 4. Effects of SW transfer on liver (a,c) and muscle (b) *igfbp-1a* (a,b) and *igfbp-1b* (c) mRNA levels in juvenile chum salmon. Values are expressed as means $\pm$ SE (n=6-8). Groups sharing the same letter or those without letter are not significantly different.

Fig. 5. Effects of salinity and fasting on gill NKA activity in juvenile chum salmon. Values are expressed as means $\pm$ SE (n=8). Asterisks indicate overall effects. Groups sharing the same letter are not significantly different.

Fig. 6. Effects of salinity and fasting on serum IGF-I levels in juvenile chum salmon. Values are expressed as means $\pm$ SE (n=7-8). An asterisk indicates an overall effect. Groups sharing the same letter are not significantly different.

Fig. 7. Effects of salinity and fasting on liver (a,c) and muscle (b,d) RNA/DNA ratio (a,b) and *igf-I* mRNA levels (c,d) in juvenile chum salmon. Values are expressed as means $\pm$ SE (n=6-8). Asterisks and crosses indicate overall effects and interactions, respectively. Groups sharing the same letter are not significantly different.

Fig. 8. Effects of salinity and fasting on liver (a,c) and muscle (b) *igfbp-1a* (a,b) and *igfbp-1b* (c) mRNA levels in juvenile chum salmon. Values are expressed as means $\pm$ SE (n=7-8). Asterisks and crosses indicate overall effects and interactions, respectively. Groups sharing the same letter are not significantly different.

671

672 Fig. 9. Relationships with SGR-BW of biochemical and endocrine parameters. (a, d-f): liver, (b,
673 e,g): muscle, (c) serum. (a,b) RNA/DNA ratio, (c) IGF-I, (d,e) *igf-1*, (f,g): *igfbp-1a*, (h) *igfbp-1b*.
674 Regression lines were drawn when the relationships were significant. r^2 : coefficient of
675 determination. Data on muscle RNA/DNA ratio and serum IGF-I were from Kaneko et al.
676 (2015).

Supplemental Table 1

Growth parameters of juvenile chum salmon from fasting experiment in SW

Parameter	Treatment	Day 0	Day 10
FL	Fed	5.80 ± 0.17	5.80 ± 0.09
	Fasted		5.80 ± 0.14
BW	Fed	1.37 ± 0.12	1.37 ± 0.06
	Fasted		1.16 ± 0.10
K	Fed	0.69 ± 0.02^a	0.70 ± 0.01^a
	Fasted		0.59 ± 0.02^b
SGR-FL	Fed	-	0.51 ± 0.10^a
	Fasted		0.03 ± 0.04^b
SGR-BW	Fed	-	1.46 ± 0.29^a
	Fasted		-2.07 ± 0.18^b

FL: fork length; BW: body weight; K: condition factor. Values are expressed as mean $\pm$ SE (initial n=8, Fed n=14, Fasted n=10). (-): not significant. For each parameter, groups sharing the same letter are not significantly different.