



Title	Commutation properties of the partial isometries associated with anticommuting self-adjoint operators
Author(s)	Arai, Asao
Citation	Hokkaido University Preprint Series in Mathematics, 121, 2-25
Issue Date	1991-08
DOI	10.14943/83266
Doc URL	http://hdl.handle.net/2115/68868
Type	bulletin (article)
File Information	pre121.pdf



[Instructions for use](#)

**Commutation Properties of the Partial
Isometries Associated with
Anticommuting Self-adjoint Operators**

Asao Arai

Series #121. August 1991

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 94: Y. Okabe, A. Inoue, On the exponential decay of the correlation functions for KMO-Langevin equations, 13 pages. 1990.
- # 95: T. Sano, Y. Watatani, Angles between two subfactors, 62 pages. 1990.
- # 96: S. Ninomiya, The Fourier-Sato transformation of pure sheaves, 22 pages. 1990.
- # 97: Y. Okabe, A. Inoue, The theory of KM_2O -Langevin equations and its applications to data analysis (II): Causal analysis (1), 51 pages. 1990.
- # 98: J. Lawrynowicz, S. Koshi and O. Suzuki, Dualities generated by the generalised Hurwitz problem and variation of the Yang-Mills field, 17 pages. 1991.
- # 99: R. Agemi, K. Kubota and H. Takamura, On certain integral equations related to nonlinear wave equations, 52 pages. 1991.
- # 100: S. Izumiya, Geometric singularities for Hamilton-Jacobi equation, 13 pages. 1991.
- # 101: S. Izumiya, Legendrian singularities and first order differential equations, 16 pages. 1991.
- # 102: A. Munemasa, Y. Watatani, Orthogonal pairs of $*$ -subalgebras and association schemes, 11 pages. 1991.
- # 103: A. Arai, O. Ogurisu, Meromorphic $N = 2$ Wess-Zumino supersymmetric quantum mechanics, 27 pages. 1991.
- # 104: H. Takamura, Global existence of classical solutions to nonlinear wave equations with spherical symmetry for small data with noncompact support in three space dimensions, 14 pages. 1991.
- # 105: R. Agemi, Blow-up of solutions to nonlinear wave equations in two space dimensions, 11 pages. 1991.
- # 106: T. Nakazi, Extremal problems in H^p , 13 pages. 1991.
- # 107: T. Nakazi, ρ -dilations and hypo-Dirichlet algebras, 15 pages. 1991.
- # 108: A. Arai, An abstract sum formula and its applications to special functions, 25 pages. 1991.
- # 109: Y.-G. Chen, Y. Giga and S. Goto, Analysis toward snow crystal growth, 18 pages. 1991.
- # 110: T. Hibi, M. Wakayama, A q -analogue of Capelli's identity for $GL(2)$, 7 pages. 1991.
- # 111: T. Nishimori, A qualitative theory of similarity pseudogroups and an analogy of Sacksteder's theorem, 13 pages. 1991.
- # 112: K. Matsuda, An analogy of the theorem of Hector and Duminy, 10 pages. 1991.
- # 113: S. Takahashi, On a regularity criterion up to the boundary for weak solutions of the Navier-Stokes equations, 23 pages. 1991.
- # 114: T. Nakazi, Sum of two inner functions and exposed points in H^1 , 18 pages. 1991.
- # 115: A. Arai, De Rham operators, Laplacians, and Dirac operators on topological vector spaces, 27 pages. 1991.
- # 116: T. Nishimori, A note on the classification of non-singular flows with transverse similarity structures, 17 pages. 1991.
- # 117: T. Hibi, A lower bound theorem for Ehrhart polynomials of convex polytopes, 6 pages. 1991.
- # 118: R. Agemi, H. Takamura, The lifespan of classical solutions to nonlinear wave equations in two space dimensions, 30 pages. 1991.
- # 119: S. Altschuler, S. Angenent and Y. Giga, Generalized motion by mean curvature for surfaces of rotation, 15 pages. 1991.
- # 120: T. Nakazi, Invariant subspaces in the bidisc and commutators, 20 pages. 1991.

**Commutation Properties of the Partial Isometries Associated with
Anticommuting Self-adjoint Operators**

ASAO ARAI

Department of Mathematics, Hokkaido University, Sapporo 060, Japan

Abstract. It is proven that, for every pair $\{A, B\}$ of anticommuting self-adjoint operators, iAB is essentially self-adjoint on a suitable domain and its closure $C(A, B)$ anticommutes with A and B . For every self-adjoint operator S , a partial isometry U_S is defined by the polar decomposition $S = U_S|S|$. Let P_S be the orthogonal projection onto $(\text{Ker } S)^\perp$. The commutation properties of the operators $U_A, U_B, U_{C(A, B)}, P_A, P_B$, and $P_A P_B$ are investigated. These operators multiplied by some constants satisfy a set of commutation relations, which may be regarded as an extension of that satisfied by the standard basis of the Lie algebra $su(2, \mathbb{C})$ of the special unitary group $SU(2)$. It is shown that there exists a Lie algebra \mathfrak{M} associated with those operators and that, if A and B are injective, then \mathfrak{M} gives a completely reducible representation of $su(2, \mathbb{C})$ with the highest weight of each irreducible component being $1/2$. Moreover, the "diagonalization" of $A + B$ is given.

I. INTRODUCTION

Recently S. Pedersen [2] has presented several equivalent definitions of anticommutativity of (unbounded) self-adjoint operators and discussed some fundamental properties of anticommuting self-adjoint operators (cf. also [4]).

In this paper we further develop theory of anticommuting self-adjoint operators. The outline is as follows. In Section II, after summarizing some fundamental facts on anticommuting self-adjoint operators, we consider the product iAB of two anticommuting self-adjoint operators A and B in a Hilbert space \mathcal{H} . We show that it is essentially self-adjoint on a suitable domain and its closure $C(A, B)$ anticommutes with A and B (Theorems 2.3, 2.5, and 2.6). This means that for every pair $\{A, B\}$ of anticommuting self-adjoint operators, we have a family $\{A, B, C(A, B)\}$ of mutually anticommuting self-adjoint operators.

Section III is concerned with the partial isometries appearing in the polar decompositions of A, B , and $C(A, B)$. It is known [1, p.358] that the polar decomposition of a self-adjoint operator A with spectral family $\{E_A(\lambda)\}$ is of the form

$$A = U_A |A|, \tag{1.1}$$

with

$$U_A = 1 - E_A(0) - E_A(-0), \tag{1.2}$$

which is a partial isometry with initial space, $(\text{Ker } A)^\perp$, and final space, $\overline{\text{Ran } A}$. We call U_A the *partial isometry associated with* A . Let P_A be the orthogonal projection onto $(\text{Ker } A)^\perp$. We prove that the family $\mathfrak{A} = \{U_A, U_B, U_{C(A, B)}, P_A, P_B, P_A P_B\}$ of bounded self-adjoint operators satisfies some interesting commutation relations (Theorem 3.5), which may be regarded as an extension of those satisfied by the standard basis of the Lie algebra $su(2, \mathbb{C})$ of the special unitary group $SU(2)$. In deriving these commutation relations, a formula for U_A plays an important role (Lemma 3.1). We see that there exists a Lie algebra \mathfrak{M} associated with \mathfrak{A} (Theorem 3.6). In particular, if A and B are injective, then \mathfrak{M} gives a completely reducible representation of $su(2, \mathbb{C})$ with the highest weight of each irreducible component being $1/2$ (Theorems 3.7 and 3.8).

In the last section, we consider the sum $A + B$ in the case where B is injective. The Hilbert space \mathcal{H} has an orthogonal decomposition into two closed subspaces which diagonalizes B (Theorem 4.1). We prove by an explicit construction that there exists a unitary transformation diagonalizing $A + B$ with respect to the decomposition (Theorem 4.4).

II. PRODUCTS OF ANTICOMMUTING SELF-ADJOINT OPERATORS

We first recall a definition of the anticommutativity of self-adjoint operators [2](cf. also [4]): Two self-adjoint operators A and B in a Hilbert space \mathcal{H} are said to *anticommute* if $\exp(itA)B \subset B \exp(-itA)$ for all $t \in \mathbb{R}$. This definition of the anticommutativity is symmetric in A and B (see [2]). For the reader's convenience, we summarize as a lemma some known facts on anticommuting self-adjoint operators.

LEMMA 2.1 (VASILESCU [4], PEDERSEN [2]). *Let A and B be anticommuting self-adjoint operators. Then the following (i)-(vii) hold:*

- (i) $U_B A \subset -A U_B$ and $U_A B \subset -B U_A$.
- (ii) $U_B |A| \subset |A| U_B$ and $U_A |B| \subset |B| U_A$.
- (iii) $|A|$ and $|B|$ commute.
- (iv) $U_A U_B = -U_B U_A$.
- (v) A and $|B|$ commute and B and $|A|$ commute.
- (vi) $D(A) \cap D(B) \cap D(AB) = D(A) \cap D(B) \cap D(BA)$ and

$$(AB + BA)f = 0, \quad f \in D(A) \cap D(B) \cap D(AB).$$

- (vii) $A + B$ is self-adjoint.

Let A and B be anticommuting self-adjoint operators. Then, by Lemma 2.1(iii), we can define a two dimensional spectral measure E by

$$E = E_{|A|} \otimes E_{|B|}. \tag{2.1}$$

Let

$$\mathcal{D} = \bigcup_{a,b>0} \text{Ran } E([-a, a] \times [-b, b]). \tag{2.2}$$

Then \mathcal{D} is dense.

LEMMA 2.2. *The spectral measure E commutes with U_A and U_B , respectively. In particular, U_A and U_B leave \mathcal{D} invariant.*

PROOF: It follows from (1.2) that U_A commutes with $|A|$ and hence with $E_{|A|}$. Lemma 2.1(ii) implies that U_A commutes with $E_{|B|}$. Therefore U_A commutes with E . Similarly one can prove that U_B commutes with E . ■

For two anticommuting self-adjoint operators A and B , we define

$$C_0(A, B) = iAB \quad (2.3)$$

with $D(C_0(A, B)) = D(A) \cap D(B) \cap D(AB)$. It follows from Lemma 2.1(vi) that

$$D(C_0(A, B)) = D(C_0(B, A)), \quad (2.4)$$

and

$$[C_0(A, B) + C_0(B, A)]f = 0, \quad f \in D(C_0(A, B)). \quad (2.5)$$

THEOREM 2.3.

- (i) *For all $k \in \mathbb{N}$, $C_0(A, B)^k$ is essentially self-adjoint on \mathcal{D} .*
- (ii) *Let $C(A, B)$ be the closure of $C_0(A, B)$. Then*

$$C(A, B) = -C(B, A). \quad (2.6)$$

PROOF: (i) The Hermiteness of $C_0(A, B)^k$ follows from (2.4) and (2.5). We show that \mathcal{D} is an invariant set of analytic vectors for $C_0(A, B)^k$. Then the assertion follows from a variant of Nelson's analytic vector theorem [3, X.6, Corollary 2]. Let

$$L = C_0(A, B)^k.$$

We have by (1.1)

$$AB = U_A|A|U_B|B|,$$

which, together with Lemma 2.2, implies that $\mathcal{D} \subset D(AB)$ and AB leaves \mathcal{D} invariant. Hence L leaves \mathcal{D} invariant and

$$L^n f = (U_A U_B)^{n+k} (|A||B|)^{n+k} f$$

for all $f \in \mathcal{D}$. Since $\|U_A\|, \|U_B\| \leq 1$, we have

$$\|L^n f\| \leq a^{n+k} b^{n+k} \|f\|$$

for $f \in \text{Ran } E([-a, a] \times [-b, b]) \in \mathcal{D}$. This implies that f is an analytic vector for L .

(ii) This follows from part(i) with $k = 1$ and (2.5). ■

It is interesting to find other cores for the self-adjoint operator $C(A, B)$. For this purpose, we need a lemma.

LEMMA 2.4. *Let A and B be anticommuting self-adjoint operators. Then, for all $m, n \in \mathbb{N} \cup \{0\}$, $A^{2m} + B^{2n}$ is self-adjoint and nonnegative. Moreover, the subspace \mathcal{D} given by (2.2) is a core for $A^{2m} + B^{2n}$.*

PROOF: The self-adjointness of $A^2 + B^2$ was proved in [4]. Let

$$T_{m,n} = A^{2m} + B^{2n}.$$

Then we have

$$T_{m,n} = |A|^{2m} + |B|^{2n}.$$

It follows from Lemma 2.1(iii) and the operator calculus for commuting self-adjoint operators that $|A|^{2m} + |B|^{2n}$ is self-adjoint. Note that $D(|A|^{2m}) \cap D(|B|^{2n}) = D(A^{2m}) \cap D(B^{2n})$. Thus $T_{m,n}$ is self-adjoint. The nonnegativity of $T_{m,n}$ is obvious.

We have $\mathcal{D} \subset D(A^{2m} + B^{2n})$. Let $f \in D(A^{2m} + B^{2n})$ and define

$$f_k = E([-k, k]) \times [-k, k] f, \quad k \in \mathbb{N}.$$

Then, using the operator calculus with respect to the spectral measure E given by (2.3), one can easily show that, for all $k, f_k \in \mathcal{D}$, and

$$f_k \rightarrow f, \quad T_{m,n} f_k \rightarrow T_{m,n} f,$$

as $k \rightarrow \infty$. Hence \mathcal{D} is a core for $T_{m,n}$. ■

THEOREM 2.5. *Let A and B be anticommuting self-adjoint operators. Then $C(A, B)$ is essentially self-adjoint on every core of $A^2 + B^2$.*

PROOF: By Lemma 2.4, the operator

$$N = A^2 + B^2 + 1$$

is self-adjoint with $N \geq 1$ and is essentially self-adjoint on \mathcal{D} . Since A^2 and B^2 are commuting nonnegative self-adjoint operators, it follows that

$$\|A^2 f\|^2 + \|B^2 f\|^2 \leq \|Nf\|^2, \quad f \in D(N) = D(A^2) \cap D(B^2).$$

We denote by $\langle \cdot, \cdot \rangle$ the inner product of the Hilbert space \mathcal{H} . For all $f \in \mathcal{D}$,

$$\begin{aligned} \|C(A, B)f\|^2 &= \langle A^2 f, B^2 f \rangle \\ &\leq \|A^2 f\| \|B^2 f\| \\ &\leq \frac{1}{2}(\|A^2 f\|^2 + \|B^2 f\|^2) \\ &\leq \frac{1}{2}\|Nf\|^2. \end{aligned}$$

Moreover, one can see that

$$\langle C(A, B)f, Ng \rangle - \langle Nf, C(A, B)g \rangle = 0, \quad f, g \in \mathcal{D}.$$

Therefore we can apply a variant of Nelson's commutator theorem [3, Theorem X.37] to obtain the desired result. ■

THEOREM 2.6. *The operator $C(A, B)$ anticommutes with A, B , and $A + B$.*

PROOF: In the same way as in the proof of Theorem 2.3, we can show that \mathcal{D} is a set of analytic vectors for A . We already know that $C(A, B)\mathcal{D} \subset \mathcal{D}$. Hence, for all $f \in \mathcal{D}$ and all $t \in \mathbb{R}$,

$$e^{itA}C(A, B)f = \lim_{M \rightarrow \infty} g_M$$

with

$$g_M = \sum_{m=0}^M \frac{(it)^m}{m!} A^m C(A, B)f.$$

It follows from (2.6) that

$$A^m C(A, B)f = (-1)^m C(A, B)A^m f.$$

Hence we can write

$$g_M = C(A, B)h_M$$

with

$$h_M = \sum_{m=0}^M \frac{(-it)^m}{m!} A^m f.$$

We have

$$h_M \rightarrow e^{-itA} f \quad (M \rightarrow \infty)$$

and

$$C(A, B)h_M \rightarrow e^{itA} C(A, B)f \quad (M \rightarrow \infty).$$

Since $C(A, B)$ is closed, we conclude that $\exp(-itA)f \in D(C(A, B))$ and

$$C(A, B)e^{-itA} f = e^{itA} C(A, B)f. \quad (2.7)$$

By Theorem 2.3(i), \mathcal{D} is a core for $C(A, B)$. Hence (2.7) implies that

$$e^{itA} C(A, B) \subset C(A, B)e^{-itA},$$

i.e., $C(A, B)$ and A anticommute. Similarly we can prove the anticommutativity of $C(A, B)$ and B . The anticommutativity of $C(A, B)$ with $A + B$ follows from that of $C(A, B)$ with A and B . ■

III. COMMUTATION RELATIONS OF THE PARTIAL ISOMETRIES ASSOCIATED WITH ANTICOMMUTING SELF-ADJOINT OPERATORS

Let A and B be anticommuting self-adjoint operators in a Hilbert space \mathcal{H} . In this section we compute commutation relations of the partial isometries U_A, U_B , and $U_{C(A,B)}$ and show that there exists a Lie algebra containing these partial isometries. If A and B are injective, then this algebra gives a completely reducible (infinite dimensional) representation of $su(2, \mathbb{C})$ with the highest weight of each irreducible component being $1/2$.

A key tool to compute products of the partial isometries associated with self-adjoint operators is the following formula.

LEMMA 3.1. *Let A be a self-adjoint operator. Then*

$$U_A = s - \lim_{\epsilon \rightarrow +0} A(A^2 + \epsilon)^{-1/2} \quad (3.1)$$

PROOF: For $f \in \text{Ker } A$, (3.1) is obvious, since $\text{Ker } A = \text{Ker } U_A$. Let $f \in (\text{Ker } A)^\perp$ and $\epsilon > 0$. Then

$$\| |A|(A^2 + \epsilon)^{-1/2} f - f \|^2 = \int u_\epsilon(\lambda) d\|E_A(\lambda)f\|^2,$$

where

$$u_\epsilon(\lambda) = \left| \frac{|\lambda|}{\sqrt{|\lambda|^2 + \epsilon}} - 1 \right|^2.$$

Obviously we have

$$0 \leq u_\epsilon(\lambda) \leq 4$$

and

$$\lim_{\epsilon \rightarrow +0} u_\epsilon(\lambda) = 0, \quad \lambda \in \mathbb{R} \setminus \{0\}.$$

In the present case, the spectral measure $\|E_A(\cdot)f\|^2$ has no mass at $\lambda = 0$. Hence, by the dominated convergence theorem, we obtain

$$\lim_{\epsilon \rightarrow +0} |A|(A^2 + \epsilon)^{-1/2} f = f.$$

Since U_A is bounded, it follows that

$$\lim_{\epsilon \rightarrow +0} U_A |A| (A^2 + \epsilon)^{-1/2} f = U_A f,$$

which, together with (1.1), implies that

$$\lim_{\epsilon \rightarrow +0} A (A^2 + \epsilon)^{-1/2} f = U_A f.$$

Thus (3.1) follows. ■

Remark. The partial isometry U_A has another representation given by

$$U_A = \operatorname{sgn}(A) P_A,$$

where $\operatorname{sgn}(\lambda) = \lambda/|\lambda|$. This follows from operator calculus as above.

Let A and B be anticommuting self-adjoint operators and $\epsilon > 0$. We define

$$T_\epsilon(A, B) = \int (\lambda^2 + \epsilon)^{1/2} (\mu^2 + \epsilon)^{1/2} dE(\lambda, \mu), \quad (3.2)$$

where E is the spectral measure given by (2.1). The operator $T_\epsilon(A, B)$ is self-adjoint with

$$T_\epsilon(A, B) \geq \epsilon.$$

The operator calculus with respect to E gives

$$(A^2 + \epsilon)^{1/2} (B^2 + \epsilon)^{1/2} \subset T_\epsilon(A, B)$$

and

$$T_\epsilon(A, B)^{-1} = (A^2 + \epsilon)^{-1/2} (B^2 + \epsilon)^{-1/2}. \quad (3.3)$$

It follows that $|A||B|T_\epsilon(A, B)^{-1}$ is bounded.

We have

$$C(A, B)^2 \upharpoonright \mathcal{D} = (A^2 B^2) \upharpoonright \mathcal{D} = (B^2 A^2) \upharpoonright \mathcal{D} \subset \int \lambda^2 \mu^2 dE(\lambda, \mu).$$

By Theorem 2.3, the operator on the left hand side is essentially self-adjoint. Hence we obtain

$$C(A, B)^2 = \int \lambda^2 \mu^2 dE(\lambda, \mu), \quad (3.4)$$

which implies that $|A||B|(C(A, B)^2 + \epsilon)^{-1/2}$ is bounded.

LEMMA 3.2. We have

$$s - \lim_{\epsilon \rightarrow +0} \left\{ |A||B|T_\epsilon(A, B)^{-1} - |A||B|(C(A, B)^2 + \epsilon)^{-1/2} \right\} = 0. \quad (3.5)$$

PROOF: We have for all $f \in \mathcal{H}$

$$\| \{ |A||B|T_\epsilon(A, B)^{-1} - |A||B|(C(A, B)^2 + \epsilon)^{-1/2} \} f \|^2 = \int_{\lambda, \mu \geq 0} w_\epsilon(\lambda, \mu) d\|E(\lambda, \mu)f\|^2,$$

where

$$w_\epsilon(\lambda, \mu) = \left| \frac{\lambda\mu}{\sqrt{(\lambda^2 + \epsilon)(\mu^2 + \epsilon)}} - \frac{\lambda\mu}{\sqrt{\lambda^2\mu^2 + \epsilon}} \right|^2.$$

It is easy to see that

$$|w_\epsilon(\lambda, \mu)| \leq 4$$

and

$$\lim_{\epsilon \rightarrow +0} w_\epsilon(\lambda, \mu) = 0, \quad \lambda, \mu \geq 0.$$

Hence, by the dominated convergence theorem, we obtain (3.5). ■

For a self-adjoint operator A , we denote by P_A the orthogonal projection onto $(\text{Ker } A)^\perp$.

LEMMA 3.3. Let A and B be anticommuting self-adjoint operators. Then :

- (i) P_A and P_B commute.
- (ii) P_A and U_B commute and P_B and U_A commute.

PROOF: Since $\text{Ker } A = \text{Ker } |A|$, it follows that

$$P_A = P_{|A|},$$

Note that $P_{|A|} = E_{|A|}((0, \infty))$. Since $E_{|A|}$ and $E_{|B|}$ commute, part(i) follows.

The proof of Lemma 2.2 gives that U_B commutes with $E_{|A|}$ and hence with $P_{|A|}$. Thus U_B commutes with P_A . It follows from symmetry in A and B that P_B and U_A commute. ■

THEOREM 3.4. *The following formulae hold:*

$$U_A U_B = -i U_{C(A,B)}. \quad (3.6)$$

$$U_{C(A,B)} U_A = -i P_A U_B = -i U_B P_A \quad (3.7)$$

$$U_{C(A,B)} U_B = i P_B U_A = i U_A P_B. \quad (3.8)$$

PROOF: It follows from Lemma 2.1(v) that

$$(A^2 + \epsilon)^{-1/2} B \subset B(A^2 + \epsilon)^{-1/2}.$$

Using this fact and formula (3.1), we have

$$\begin{aligned} U_A U_B &= s - \lim_{\epsilon \rightarrow +0} A(A^2 + \epsilon)^{-1/2} B(B^2 + \epsilon)^{-1/2} \\ &= s - \lim_{\epsilon} A B T_{\epsilon}(A, B)^{-1} \\ &= s - \lim_{\epsilon} U_A U_B |A| |B| T_{\epsilon}(A, B)^{-1}. \end{aligned}$$

It follows from (3.5) and the boundedness of $U_A U_B$ that

$$s - \lim_{\epsilon \rightarrow +0} \left\{ U_A U_B |A| |B| T_{\epsilon}(A, B)^{-1} - U_A U_B |A| |B| (C(A, B)^2 + \epsilon)^{-1/2} \right\} = 0.$$

Hence

$$U_A U_B = s - \lim_{\epsilon \rightarrow +0} U_A U_B |A| |B| (C(A, B)^2 + \epsilon)^{-1/2},$$

which implies that for all $f \in \mathcal{D}$ and $g \in \mathcal{H}$,

$$\begin{aligned} \langle f, U_A U_B g \rangle &= s - \lim_{\epsilon \rightarrow +0} \langle |B| |A| U_B U_A f, (C(A, B)^2 + \epsilon)^{-1/2} g \rangle \\ &= s - \lim_{\epsilon \rightarrow +0} \langle i C(A, B) f, (C(A, B)^2 + \epsilon)^{-1/2} g \rangle \\ &= s - \lim_{\epsilon \rightarrow +0} \langle f, (-i) C(A, B) (C(A, B)^2 + \epsilon)^{-1/2} g \rangle \\ &= \langle f, -i U_{C(A,B)} g \rangle. \end{aligned}$$

Thus (3.6) follows.

Let $f \in \mathcal{D}$. Then $A(A^2 + \epsilon)^{-1/2}f \in \mathcal{D}$ and hence

$$\begin{aligned} C(A, B)A(A^2 + \epsilon)^{-1/2}f &= iABA(A^2 + \epsilon)^{-1/2}f \\ &= -iBA^2(A^2 + \epsilon)^{-1/2}f \\ &= -iU_B|B|A^2(A^2 + \epsilon)^{-1/2}f. \end{aligned}$$

Using the commutativity of U_B with $|C(A, B)|$, which follows from Theorem 2.6 and Lemma 2.1(ii), we have

$$\begin{aligned} U_{C(A, B)}U_A f &= \lim_{\epsilon \rightarrow +0} C(A, B)(C(A, B)^2 + \epsilon)^{-1/2}A(A^2 + \epsilon)^{-1/2}f \\ &= \lim_{\epsilon \rightarrow +0} (C(A, B)^2 + \epsilon)^{-1/2}C(A, B)A(A^2 + \epsilon)^{-1/2}f \\ &= -iU_B \lim_{\epsilon \rightarrow +0} f_\epsilon, \end{aligned}$$

where

$$f_\epsilon = (C(A, B)^2 + \epsilon)^{-1/2}|B|A^2(A^2 + \epsilon)^{-1/2}f.$$

The operator calculus with respect to the spectral measure E gives

$$f_\epsilon = \int_{\lambda, \mu \geq 0} u_\epsilon(\lambda, \mu) dE(\lambda, \mu) f,$$

where

$$u_\epsilon(\lambda, \mu) = \frac{\lambda^2 \mu}{\sqrt{(\lambda^2 \mu^2 + \epsilon)(\lambda^2 + \epsilon)}}.$$

We have

$$|u_\epsilon(\lambda, \mu)| \leq 1, \quad \lambda, \mu \geq 0,$$

and

$$\lim_{\epsilon \rightarrow +0} u_\epsilon(\lambda, \mu) = 1, \quad \lambda, \mu > 0.$$

Let $f \in (\text{Ker } A)^\perp \cap (\text{Ker } B)^\perp$. Then, for all Borel sets $M \subset \mathbb{R}$,

$$\|E(\{0\} \times M)f\|^2 = \|E(M \times \{0\})f\|^2 = 0.$$

Theorefore, by the dominated convergence theorem, we obtain

$$\lim_{\epsilon \rightarrow +0} f_\epsilon = f.$$

Thus

$$U_{C(A,B)}U_A f = -iU_B f.$$

It is obvious that, for $f \in \text{Ker } A \cup \text{Ker } B$, $f_\epsilon = 0$ and hence $U_{C(A,B)}U_A f = 0$. For all $f \in \mathcal{D}$, we have

$$f = P_A P_B f + P_A(1 - P_B)f + (1 - P_A)P_B f + (1 - P_A)(1 - P_B)f.$$

Note that $P_A P_B f \in (\text{Ker } A)^\perp \cap (\text{Ker } B)^\perp$ and

$$P_A(1 - P_B) \in \text{Ker } B, (1 - P_A)P_B \in \text{Ker } A, (1 - P_A)(1 - P_B) \in \text{Ker } A \cap \text{Ker } B.$$

Hence it follows that, for all $f \in \mathcal{D}$,

$$U_{C(A,B)}U_A f = -iU_B P_A P_B f = -iP_A U_B P_B f.$$

Moreover, we have $U_B P_B f = U_B f$, because $\text{Ker } B = \text{Ker } U_B$. Hence we obtain

$$U_{C(A,B)}U_A f = -iP_A U_B f = -iU_B P_A f.$$

Since \mathcal{D} is dense, this equation extends to all $f \in \mathcal{H}$. Thus (3.7) follows.

Formula (3.8) follows from (3.7) with A (resp. B) replaced by B (resp. A) and the fact $U_{C(A,B)} = -U_{C(B,A)}$. ■

Let A and B be anticommuting self-adjoint operators and define

$$X_1 = i\frac{U_A}{2}, X_2 = i\frac{U_B}{2}, X_3 = i\frac{U_{C(A,B)}}{2}, \quad (3.9)$$

$$Y_1 = 1, Y_2 = P_B, Y_3 = P_A, Y_4 = P_A P_B. \quad (3.10)$$

For bounded linear operators X, Y on \mathcal{H} , we define

$$[X, Y] = XY - YX. \quad (3.11)$$

THEOREM 3.5. *The following commutation relations hold:*

$$[X_j, X_k] = \sum_{\ell=1}^3 \epsilon_{j k \ell} X_\ell Y_j, \quad , j, k = 1, 2, 3, \quad (3.12)$$

$$[X_j, Y_m] = [Y_m, Y_n] = 0, \quad j = 1, 2, 3, m, n = 1, 2, 3, 4, \quad (3.13)$$

where $\epsilon_{j k \ell}$ is the Levi-Civita symbol with $\epsilon_{123} = 1$.

PROOF: This follows from Lemma 3.3 , Theorem 3.4 and the fact that $U_A U_B = -U_B U_A$ for any two anticommuting self-adjoint operators A and B (Lemma 2.1(iv)). ■

The vector space of all bounded linear operators on \mathcal{H} is a Lie algebra with the Lie bracket $[\cdot, \cdot]$ given by (3.11). We denote it by $\mathfrak{L}(\mathcal{H})$.

THEOREM 3.6. *Let $\mathfrak{M} \subset \mathfrak{L}(\mathcal{H})$ be a subspace spanned by $X_k Y_m, k = 1, 2, 3, m = 1, 2, 3, 4$. Then \mathfrak{M} is a Lie subalgebra of $\mathfrak{L}(\mathcal{H})$.*

PROOF: We need only to show that $[\mathfrak{M}, \mathfrak{M}] \subset \mathfrak{M}$. By Theorem 3.5, we have for $j, k = 1, 2, 3$, and $m, n = 1, 2, 3, 4$

$$[X_j Y_m, X_k Y_n] = [X_j, X_k] Y_m Y_n = \sum_{\ell=1}^3 \epsilon_{j k \ell} X_\ell Y_j Y_m Y_n.$$

It follows from (3.10) that there exists a number $\alpha(j, m, n) \in \{1, 2, 3, 4\}$ such that

$$Y_j Y_m Y_n = Y_{\alpha(j, m, n)}.$$

Hence $[X_j Y_m, X_k Y_n] \in \mathfrak{M}$. Thus \mathfrak{M} is a Lie algebra. ■

As is well-known, the Lie algebra $\mathfrak{su}(2, \mathbb{C})$ of the special unitary group $SU(2)$ is the set of 2×2 complex skew-Hermitian matrices of trace zero and has a basis $\{e_j\}_{j=1}^3$ which satisfy the commutation relations

$$[e_j, e_k] = \sum_{\ell=1}^3 \epsilon_{j k \ell} e_\ell, \quad j, k = 1, 2, 3. \quad (3.14)$$

We define a linear map $\rho : su(2, \mathbb{C}) \rightarrow \mathfrak{L}(\mathcal{H})$ by

$$\rho\left(\sum_{j=1}^3 \alpha_j e_j\right) = \sum_{j=1}^3 \alpha_j X_j, \quad \alpha_j \in \mathbb{C}, j = 1, 2, 3. \quad (3.15)$$

THEOREM 3.7. *Suppose that A and B are injective. Then ρ is a faithful representation of the $su(2, \mathbb{C})$ with $\text{Ran } \rho = \mathfrak{M}$.*

PROOF: Under the present assumption, we have $P_A = P_B = 1$. Hence $Y_m = 1$, $m = 1, 2, 3, 4$. Thus (3.12) and (3.13) reduce to

$$[X_j, X_k] = \sum_{l=1}^3 \epsilon_{jkl} X_l, \quad j, k = 1, 2, 3. \quad (3.16)$$

Therefore ρ is a representation of $su(2, \mathbb{C})$. The map ρ is injective, since $X_j, j = 1, 2, 3$ are linear independent. Hence ρ is faithful. ■

If the Hilbert space \mathcal{H} is infinite dimensional, then ρ is an infinite dimensional representation of $su(2, \mathbb{C})$. We analyze the structure of this representation in detail. We prove the following theorem.

THEOREM 3.8. *The representation ρ of $su(2, \mathbb{C})$ is completely reducible and the highest weight of each irreducible component of it is $1/2$.*

To prove this theorem, we prepare some lemmas. We denote by $\sigma(S)$ the spectrum of operator S .

LEMMA 3.9. *Let A and B be anticommuting self-adjoint operators and injective. Then :*

- (i) U_A, U_B and $U_{C(A,B)}$ are unitary operators.
- (ii)

$$\sigma(U_A) = \sigma(U_B) = \sigma(U_{C(A,B)}) = \{-1, 1\}. \quad (3.17)$$

PROOF: (i) In general, if a self-adjoint operator is injective, then the associated partial isometry is unitary. This easily follows from the definition of the polar decomposition. Hence the assertion about U_A and U_B follows. If $f \in \text{Ker } U_{C(A,B)} = \text{Ker } C(A,B)$, then we have from (3.16)

$$(U_A U_B - U_B U_A)f = 0.$$

Since $U_A U_B = -U_B U_A$, we obtain

$$U_A U_B f = 0,$$

which means that $U_B f \in \text{Ker } U_A = \{0\}$, i.e., $U_B f = 0$. Hence $f = 0$. Thus $U_{C(A,B)}$ is injective and hence unitary.

(ii) Since U_A is self-adjoint and unitary by part (i), either 1 or -1 is contained in $\sigma(U_A)$. Let $1 \in \sigma(U_A)$ and $U_A f = f$ ($f \neq 0$). Then

$$U_B U_A f = U_B f.$$

The left hand side is equal to $-U_A U_B f$. Since $U_B f \neq 0$, it follows that $U_B f$ is an eigenvector of U_A with eigenvalue -1 . Hence $-1 \in \sigma(U_A)$. Similarly, one can show that, if $-1 \in \sigma(U_A)$, then $1 \in \sigma(U_A)$. Thus (3.17) follows. ■

Let A and B be anticommuting, injective self-adjoint operators. The Weyl canonical basis of the representation ρ is given by the triple $\{H, E_+, E_-\}$ defined by

$$E_+ = \frac{1}{2}(iX_1 - X_2) = -\frac{1}{4}(U_A + iU_B),$$

$$E_- = iX_1 + X_2 = \frac{1}{2}(-U_A + iU_B)$$

$$H = iX_3 = -\frac{1}{2}U_{C(A,B)}.$$

These operators satisfy the following commutation relations:

$$[H, E_+] = E_+, [H, E_-] = -E_-, [E_+, E_-] = H. \quad (3.18)$$

Note that

$$\begin{aligned} E_{\pm}^2 &= 0, H^2 = \frac{1}{4}, \\ E_+^* &= \frac{1}{2}E_-. \end{aligned} \quad (3.19)$$

Moreover, we have from (3.17)

$$\sigma(H) = \left\{ \pm \frac{1}{2} \right\}. \quad (3.20)$$

Let

$$\begin{aligned} \mathcal{K}_{\pm} &= \{f \in \mathcal{H} \mid U_{C(A,B)}f = \mp f\} \\ &= \{f \in \mathcal{H} \mid Hf = \pm \frac{1}{2}f\} \end{aligned} \quad (3.21)$$

Then we have the orthogonal decomposition

$$\mathcal{H} = \mathcal{K}_+ \oplus \mathcal{K}_-. \quad (3.22)$$

LEMMA 3.10. *The operator E_- (resp. $2E_+$) is a unitary operator from \mathcal{K}_+ (resp. \mathcal{K}_-) onto \mathcal{K}_- (resp. \mathcal{K}_+).*

PROOF: Let $f, g \in \mathcal{K}_+$. Then we have from (3.19) and the last commutation relation in (3.18)

$$\begin{aligned} \langle E_-f, E_-g \rangle &= 2 \langle f, E_+E_-g \rangle = 2 \langle f, Hg \rangle + 2 \langle f, E_-E_+g \rangle \\ &= \langle f, g \rangle + 2 \langle f, E_-E_+g \rangle. \end{aligned}$$

The first commutation relation in (3.18) implies that $HE_+g = (3/2)E_+g$. But, by (3.20), $3/2$ cannot be an eigenvalue of H . Hence E_+g must be zero. Thus we obtain

$$\langle E_-f, E_-g \rangle = \langle f, g \rangle,$$

i.e., E_- is an isometry from \mathcal{K}_+ to \mathcal{K}_- . Similarly we can show that $2E_+$ is an isometry from \mathcal{K}_- to \mathcal{K}_+ . To prove the surjectivity of E_- , let $h \in \mathcal{K}_-$ and $f = 2E_+h$. As in the

case of E_+g , we can show that $E_-h = 0$. Using this fact and (3.18), we see that $f \in \mathcal{K}_+$ and

$$E_-f = h$$

Hence E_- is surjective. Thus E_- is unitary from \mathcal{K}_+ to \mathcal{K}_- . Similarly we can prove that $2E_+$ is unitary from \mathcal{K}_- to \mathcal{K}_+ . ■

PROOF OF THEOREM 3.8: Let $\{f_n\}_{n=1}^{\infty}$ be the complete orthonormal system (C.O.N.S) of \mathcal{K}_+ . Then, by Lemma 3.10, $\{E_-f_n\}_{n=1}^{\infty}$ is a C.N.O.S of \mathcal{K}_- . Therefore $\{f_n, E_-f_n\}_{n=1}^{\infty}$ forms a C.N.O.S of \mathcal{H} . Thus we obtain the orthogonal decomposition

$$\mathcal{H} = \bigoplus_{n=1}^{\infty} \mathcal{H}_n$$

with

$$\mathcal{H}_n = \{\alpha f_n + \beta E_-f_n \mid \alpha, \beta \in \mathbb{C}\}.$$

Note that H, E_{\pm} leave \mathcal{H}_n invariant and ρ is irreducible in \mathcal{H}_n with the highest weight $1/2$. Thus the desired result follows. ■

IV. DIAGONALIZATION OF THE SUM OF TWO ANTICOMMUTING SELF-ADJOINT OPERATORS

In [2, Corollary 3.3] it was shown that, for all injective anticommuting self-adjoint operators A and B in a Hilbert space \mathcal{H} , there exists a decomposition

$$\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_- = \begin{pmatrix} \mathcal{H}_+ \\ \mathcal{H}_- \end{pmatrix} \quad (4.1)$$

into orthogonal closed subspaces \mathcal{H}_\pm such that

$$A = \begin{pmatrix} 0 & L_A a^* \\ a L_A & 0 \end{pmatrix},$$

and

$$B = \begin{pmatrix} B_+ & 0 \\ 0 & -a B_+ a^* \end{pmatrix},$$

where a is an isometric isomorphism mapping from \mathcal{H}_+ onto \mathcal{H}_- and B_+ and L_A are injective nonnegative self-adjoint commuting operators on \mathcal{H}_+ (Remark: our A (resp. B) is B (resp. A) in [2, Corollary 3.3]). This theorem can be generalized to the case where A is not necessarily injective.

THEOREM 4.1. *Let A and B be anticommuting self-adjoint operators and B be injective. Then there exists a decomposition (4.1) such that*

$$A = \begin{pmatrix} 0 & a^* L_A^{(-)} \\ a L_A^{(+)} & 0 \end{pmatrix}, \quad (4.2)$$

and

$$B = \begin{pmatrix} B_+ & 0 \\ 0 & -B_- \end{pmatrix}, \quad (4.3)$$

where a is a partial isometry from \mathcal{H}_+ to \mathcal{H}_- , B_+ (resp. B_-) and $L_A^{(+)}$ (resp. $L_A^{(-)}$) are nonnegative self-adjoint commuting operators in \mathcal{H}_+ (resp. \mathcal{H}_-), and B_- is a nonnegative self-adjoint operator in \mathcal{H}_- with $a B_+ \subset -B_- a$.

The proof of this theorem is similar to that of [2, Corollary 3.3]. Hence we omit it.

We consider the sum

$$T = A + B \quad (4.4)$$

of anticommuting self-adjoint operators A and B . The purpose of the rest of this section is to show that T can be diagonalized with respect to the decomposition given in Theorem 4.1. Let A and B be anticommuting self-adjoint operators and B be injective. Then the operator

$$\Lambda = \arctan(|A||B|^{-1}), \quad (4.5)$$

defined by the operator calculus with respect to the spectral measure E given by (2.1), is a bounded self-adjoint operator.

LEMMA 4.2. *Let Λ be given by (4.5), Then*

$$\sin \Lambda = |A|(A^2 + B^2)^{-1/2}, \quad (4.6)$$

$$\cos \Lambda = |B|(A^2 + B^2)^{-1/2}. \quad (4.7)$$

Moreover,

$$[X_j, \Lambda] = [Y_m, \Lambda] = 0, \quad j = 1, 2, 3, m = 1, 2, 3, 4, \quad (4.8)$$

where X_j and Y_m are defined by (3.9) and (3.10).

PROOF: Since $\sigma(\Lambda) \subset [0, \pi/2]$, we have

$$\sin \Lambda = \tan \Lambda (1 + (\tan \Lambda)^2)^{-1/2},$$

$$\cos \Lambda = (1 + (\tan \Lambda)^2)^{-1/2}.$$

It follows from operator calculus that

$$|A|(A^2 + B^2)^{-1/2} \subset \tan \Lambda (1 + (\tan \Lambda)^2)^{-1/2}.$$

It is easy to see that the operator on the left hand side is a bounded self-adjoint. Hence (4.6) follows. Similarly we can prove (4.7).

The commutation relations (4.8) follow from the commutativity of E with $U_A, U_B, U_{C(A,B)}, P_A,$ and $P_B (= 1)$. ■

Since $-iX_3$ and Λ are commuting bounded self-adjoint operators, $-iX_3\Lambda$ is bounded and self-adjoint. Hence the operator

$$V = e^{X_3\Lambda} \quad (4.9)$$

is unitary.

LEMMA 4.3. Let Λ be given by (4.5). Then

$$VX_1V^{-1} = (1 - P_A)X_1 + P_A(X_1 \cos \Lambda + X_2 \sin \Lambda), \quad (4.10)$$

$$VX_2V^{-1} = (1 - P_A)X_2 + P_A(-X_1 \sin \Lambda + X_2 \cos \Lambda), \quad (4.11)$$

PROOF: Since $X_3\Lambda$ and X_1 are bounded, we have

$$VX_1V^{-1} = \sum_{n=0}^{\infty} \frac{1}{n!} (\text{ad}X_3\Lambda)^n X_1,$$

where

$$(\text{ad}X)^0 Y = Y, \quad (\text{ad}X)^n Y = [X, (\text{ad}X)^{n-1} Y], \quad n \geq 1.$$

It is easy to check that

$$(\text{ad}X_3\Lambda)^{2n-1} X_1 = (-1)^{n-1} \Lambda^{2n-1} X_2 P_A,$$

$$(\text{ad}X_3\Lambda)^{2n} X_1 = (-1)^n \Lambda^{2n} X_1 P_A, \quad n \geq 1.$$

Thus (4.10) follows. Similarly, using the formulae

$$(\text{ad}X_3\Lambda)^{2n-1} X_2 = (-1)^n \Lambda^{2n-1} X_1 P_A,$$

$$(\text{ad}X_3\Lambda)^{2n} X_2 = (-1)^n \Lambda^{2n} X_2 P_A,$$

we can prove (4.11). ■

THEOREM 4.4. *Let A and B be anticommuting self-adjoint operators and B be injective. Then*

$$VTV^{-1} = P_A U_B (A^2 + B^2)^{1/2} + (1 - P_A)B. \quad (4.12)$$

In particular, if A is also injective, then

$$VTV^{-1} = U_B (A^2 + B^2)^{1/2}. \quad (4.13)$$

PROOF: we can write

$$T = -2i(X_1|A| + X_2|B|).$$

Since V commutes with $|A|$ and $|B|$, it follows that

$$V|A|V^{-1} = |A|, \quad V|B|V^{-1} = |B|.$$

Hence, for all $f \in D(A) \cap D(B)$, we have

$$VTV^{-1}f = -2i(VX_1V^{-1}|A|f + VX_2V^{-1}|B|f).$$

Putting (4.10) and (4.11) into the right hand side and using the fact $(1 - P_A)|A| = 0$, we obtain (4.12). If A is injective, then $P_A = 1$. Hence (4.12) gives (4.13). ■

One can easily see that the right hand sides of (4.12) and (4.13) are diagonal with respect to the decomposition given in Theorem 4.1.

REFERENCES

1. T. Kato, "Perturbation Theory for Linear Operators," 2nd Ed., Springer-Verlag, Berlin, Heidelberg, 1976.
2. S. Pedersen, *Anticommuting selfadjoint operators*, J.Funct.Anal. 89 (1990), 428-443.
3. M. Reed and B. Simon, "Methods of Modern Mathematical Physics II: Fourier Analysis, Self-adjointness," Academic Press, New York, 1975.
4. F.H. Vasilescu, *Anticommuting self-adjoint operators*, Rev.Roumaine Math.Pures Appl. 28 (1983), 77-91.