



Title	Perestroikas of optical wave fronts and graphlike Legendrian unfoldings
Author(s)	Izumiya,S.
Citation	Hokkaido University Preprint Series in Mathematics, 156, 1-13
Issue Date	1992-06
DOI	10.14943/83300
Doc URL	http://hdl.handle.net/2115/68902
Type	bulletin (article)
File Information	pre156.pdf



[Instructions for use](#)

**PERESTROIKAS OF OPTICAL WAVE
FRONTS AND GRAPHLIKE
LEGENDRIAN UNFOLDINGS**

Shyūichi Izumiya

Series #156. June 1992

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 127: G. Ishikawa, S. Izumiya and K. Watanabe, Vector fields near a generic submanifold, 9 pages. 1991.
- # 128: A. Arai, I. Mitoma, Comparison and nuclearity of spaces of differential forms on topological vector spaces, 27 pages. 1991.
- # 129: K. Kubota, Existence of a global solution to a semi-linear wave equation with initial data of non-compact support in low space dimensions, 53 pages. 1991.
- # 130: S. Altschuler, S. Angenent and Y. Giga, Mean curvature flow through singularities for surfaces of rotation, 62 pages. 1991.
- # 131: M. Giga, Y. Giga and H. Sohr, L^p estimates for the Stokes system, 13 pages. 1991.
- # 132: Y. Okabe, T. Ootsuka, Applications of the theory of KM_2O -Langevin equations to the non-linear prediction problem for the one-dimensional strictly stationary time series, 27 pages. 1992.
- # 133: Y. Okabe, Applications of the theory of KM_2O -Langevin equations to the linear prediction problem for the multi-dimensional weakly stationary time series, 22 pages. 1992.
- # 134: P. Aviles, Y. Giga and N. Komuro, Duality formulas and variational integrals, 22 pages. 1992.
- # 135: S. Izumiya, The Clairaut type equation, 6 pages. 1992.
- # 136: S. Izumiya, Singular solutions of first order differential equations, 6 pages. 1992.
- # 137: S. Izumiya, W.L. Marar, The Euler characteristic of a generic wave front in a 3-manifold, 6 pages. 1992.
- # 138: S. Izumiya, W.L. Marar, The Euler characteristic of the image of a stable mapping from a closed n -manifold to a $(2n - 1)$ -manifold, 5 pages. 1992.
- # 139: Y. Giga, Z. Yoshida, A bound for the pressure integral in a plasma equilibrium, 20 pages. 1992.
- # 140: S. Izumiya, What is the Clairaut equation ?, 13 pages. 1992.
- # 141: H. Takamura, Weighted deformation theorem for normal currents, 27 pages. 1992.
- # 142: T. Morimoto, Geometric structures on filtered manifolds, 104 pages. 1992.
- # 143: G. Ishikawa, T. Ohmoto, Local invariants of singular surfaces in an almost complex four-manifold, 9 pages. 1992.
- # 144: K. Kubota, K. Mochizuki, On small data scattering for 2-dimensional semilinear wave equations, 22 pages. 1992.
- # 145: T. Nakazi, K. Takahashi, Hyponormal Toeplitz operators and extremal problems of Hardy spaces, 30 pages. 1992.
- # 146: N. Hayashi, T. Ozawa, Remarks on nonlinear Schrödinger equations in one space dimension, 10 pages. 1992.
- # 147: M. Sato, Interface evolution with Neumann boundary condition, 16 pages. 1992.
- # 148: Y. Okabe, Langevin equations and causal analysis, 49 pages. 1992.
- # 149: Y. Giga, S. Takahashi, On global weak solutions of the nonstationary two-phase Stokes flow, 25 pages. 1992.
- # 150: G. Ishikawa, Determinacy of envelope of the osculating hyperplanes to a curve, 9 pages. 1992.
- # 151: G. Ishikawa, Developable of a curve and determinacy relative to osculation-type, 15 pages. 1992.
- # 152: H. Kubo, Global existence of solutions of semilinear wave equations with data of non compact support in odd space dimensions, 25 pages. 1992.
- # 153: Y. Watatani, Lattices of intermediate subfactors, 33 pages. 1992.
- # 154: T. Ozawa, On critical cases of Sobolev inequalities, 11 pages. 1992.
- # 155: M. Ohnuma, M. Sato, Singular degenerate parabolic equations with applications to geometric evolutions, 20 pages. 1992.

PERESTROIKAS* OF OPTICAL WAVE FRONTS AND GRAPHLIKE LEGENDRIAN UNFOLDINGS

Shyūichi IZUMIYA

Department of Mathematics
Hokkaido University

0. Introduction

It is well known that parallels of given smooth hypersurface in \mathbb{R}^n have Legendrian singularities ([1,3]). If we are concerned the way in which these parallels change as we alter the distance, it will be appeared perestroikas of wave front sets. This problem has been considered by several people ([2,3,4,5]). Roughly speaking, it has been shown that generic perestroikas of the singularities of parallels of smooth hypersurfaces are generic perestroikas of Legendrian singularities. In [15] Zakalyukin proved that generic perestroikas of Legendrian singularities are stable perestroikas of Legendrian singularities in the case $n \leq 5$ and classified these perestroikas. In this paper we shall consider the realization problem of the perestroikas of Legendrian singularities. In relation to this problem, Arnol'd ([2], p40) mentioned "One may find in the literature the statement that the local perestroikas of the wave fronts generated by the general Legendre mapping over space-time and of the equidistants (i.e. parallel) of the smooth hypersurface are same. It seems this has never been correctly proved." However, he has corrected his mistakes in his another book ([3], p60) "Indeed, the non-trivial perestroikas A_1 and A_2 change the number of connected components of the Legendrian manifold. Hence they cannot occur as perestroikas of equidistant hypersurfaces." This fact has already known by Bill Bruce in 1983 ([4]). Then we have the following natural question.

Question. What sort of a class of one-parameter families of Legendrian immersion germs is the correct class to describe perestroikas of parallels of hypersurfaces?

Here we shall give a candidate of this class which we call *graphlike Legendrian unfoldings*. Roughly speaking, a graphlike Legendrian unfolding is a Legendrian submanifold germ with submersive generating function. The general properties of graphlike Legendrian unfoldings will be studied in §2.

On the other hand, it should be natural to consider this problem in the framework of Hamiltonian systems which also contains the situation of parallels of hypersurfaces in a Riemannian manifold (cf. [5]).

All maps considered here are class C^∞ unless stated otherwise.

* The author uses this word in honor of Professor V. I. Arnol'd.

Acknowledgment. A part of this work has been done during the author's stay in the University of Liverpool mainly supported by Anglo-Japanese Scientific Exchange programme (The Royal Society and JSPS). The author would like to thank Professor C.T.C. Wall and his colleagues in the Department of Pure Mathematics for their kind encouragement and hospitality. The author is also grateful to Professor G. Ishikawa and Dr. T. Ohmoto for their useful discussions and advises.

1. Optical wave fronts

In this section we shall introduce the notion of optical wave fronts which is a generalization of parallels of hypersurfaces. We need some notions and results on Legendrian singularity theory. For this we refer [1,2,3,12,14].

Let $J^1(\mathbb{R}^n, \mathbb{R})$ be the 1-jet bundle of functions of n -variables. Since we only consider the local situation, the 1-jet bundle $J^1(\mathbb{R}^n, \mathbb{R})$ may be consider as \mathbb{R}^{2n+1} and the canonical coordinate is given by $(x_1, \dots, x_n, t, p_1, \dots, p_n)$. Then the canonical 1-form is given by $\theta = dt - \sum_{i=1}^n p_i dx_i$. Let $T^*\mathbb{R}^n$ be the cotangent bundle over \mathbb{R}^n whose canonical coordinate is given by $(x_1, \dots, x_n, p_1, \dots, p_n)$. We also have the canonical 1-form $\alpha = \sum_{i=1}^n p_i dx_i$ on $T^*\mathbb{R}^n$. There are natural projections. Namely,

$$\begin{aligned} \pi : J^1(\mathbb{R}^n, \mathbb{R}) &\rightarrow \mathbb{R}^n \times \mathbb{R} ; \pi(x, t, p) = (x, t), \\ \tilde{\pi} : T^*\mathbb{R}^n &\rightarrow \mathbb{R}^n ; \tilde{\pi}(x, p) = x, \\ \Pi : J^1(\mathbb{R}^n, \mathbb{R}) &\rightarrow T^*\mathbb{R}^n ; \Pi(x, t, p) = (x, p), \\ \pi_2 : J^1(\mathbb{R}^n, \mathbb{R}) &\rightarrow \mathbb{R} ; \pi_2(x, t, p) = t. \end{aligned}$$

Let $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ be a function germ. We call it a *Hamiltonian function germ*. We say that H is *starshaped* if it satisfies $\sum_{i=1}^n p_i \frac{\partial H}{\partial p_i}(x, p) \neq 0$ at any $(x, p) \in H^{-1}(c)$. We remark that if H is starshaped then $H^{-1}(c)$ is transverse to fibres of $\tilde{\pi}$. It is easy to prove the following proposition.

Proposition 1.1. *If H is starshaped, then $\alpha|_{H^{-1}(c)}$ gives a contact structure on $H^{-1}(c)$ and $\tilde{\pi}|_{H^{-1}(c)} : H^{-1}(c) \rightarrow \mathbb{R}^n$ is a Legendrian fibration.*

Let $(\ell, (x_0, p_0)) \subset (T^*\mathbb{R}^n, (x_0, p_0))$ be a submanifold germ. We say that ℓ is an *optical Legendrian submanifold relative to H* if $\dim \ell = n - 1$, $\alpha|_{\ell} = 0$ and $\ell \subset H^{-1}(c)$. We call the image $\tilde{\pi}(\ell)$ an *optical wave front relative to H* .

By this proposition, if ℓ is an optical Legendrian submanifold relative to H , then ℓ is a Legendrian submanifold of $H^{-1}(c)$ and $\tilde{\pi}(\ell)$ is a wave front set in the ordinary sense (see [1,2,3,14]). We now give examples of optical wave front sets.

Example 1.2. Let $g = \sum_{ij} g_{ij}(x) dx_i dx_j$ be a Riemannian metric on \mathbb{R}^n . We define a Hamiltonian function H on $T^*\mathbb{R}^n$ by

$$H(x_1, \dots, x_n, p_1, \dots, p_n) = \sum_{ij} g_{ij}(x) p_i p_j.$$

Then we can show that $\sum_{k=1}^n p_k \frac{\partial H}{\partial p_k} = 2 \sum_{ik} g_{ik}(x) p_i p_k = 2$ on $H^{-1}(1)$.

For a Hamiltonian function germ $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$, we define a function germ $\tilde{H} : (J^1(\mathbb{R}^n, \mathbb{R}), (x_0, t_0, p_0)) \rightarrow (\mathbb{R}, c)$ by $\tilde{H}(x, t, p) = H(x, p)$. We call \tilde{H} an induced contact Hamiltonian function germ from H . By the definition, we have the Hamiltonian vector field :

$$X_H = - \sum_{i=1}^n \frac{\partial H}{\partial p_i} \frac{\partial}{\partial x_i} + \sum_{j=1}^n \frac{\partial H}{\partial x_j} \frac{\partial}{\partial p_j} \text{ on } T^*\mathbb{R}^n.$$

For an isotropic $(n-1)$ -dimensional submanifold germ $(\ell, (x_0, p_0)) \subset (T^*\mathbb{R}^n, (x_0, p_0))$ (i.e. $\alpha|_\ell = 0$), we say that ℓ is non-characteristic relative to H at (x_0, p_0) if $X_{H, (x_0, p_0)} \notin T_{(x_0, p_0)}\ell$. Then we have the following lemma.

Lemma 1.3. *If $\sum_{i=1}^n p_i \frac{\partial H}{\partial p_i} \neq 0$ at $(x_0, p_0) \in \ell$, then ℓ is non-characteristic relative to H at (x_0, p_0) .*

Proof. If $X_{H, (x_0, p_0)} \in T_{(x_0, p_0)}\ell$, then $0 = \alpha(T_{(x_0, p_0)}\ell) = \alpha(X_{H, (x_0, p_0)}) = - \sum_{i=1}^n p_i \frac{\partial H}{\partial p_i}$ at (x_0, p_0) . This contradicts to the assumption.

By this lemma, if ℓ is an optical Legendrian submanifold relative to a starshaped Hamiltonian function H , then ℓ is automatically non-characteristic at any points relative to H . Then we can apply characteristic method to solve Cauchy problem of the first order partial differential equation $H = c$ with the given initial condition ℓ (see [12]). We consider the embedding $i : \mathcal{O} \subset H^{-1}(c)$, where $\mathcal{O} \subset \ell \times (-\varepsilon, \varepsilon)$ is some open neighbourhood of $\ell \times 0$, $\varepsilon > 0$; here $i(u, t) = T_t(u)$, $u \in \ell$, $(u, t) \in \mathcal{O}$, and T_t is an one-parameter group of translations along X_H . Then we have a family of optical wave fronts $\tilde{\pi}(\ell_t)$.

Example 1.4. We now consider the Hamiltonian function $H = p_1^2 + \dots + p_n^2$. Let $F : (\mathbb{R}^n, x_0) \rightarrow (\mathbb{R}, 0)$ be a submersion germ. We define an $(n-1)$ -dimensional submanifold of $T^*\mathbb{R}^n$ by

$$\ell_F = \{(x, p) | p = \frac{1}{(\sum_{i=1}^n (F_{x_i})^2)^{\frac{1}{2}}} (F_{x_1}, \dots, F_{x_n}) \text{ and } F(x) = 0\},$$

where $F_{x_i} = \frac{\partial F}{\partial x_i}$.

By the definition, $\ell \subset H^{-1}(1)$ and $\alpha|_\ell = 0$. Parallels of hypersurface $F^{-1}(0)$ in \mathbb{R}^n are $\tilde{\pi}(\ell_t)$ which are constructed by the above argument with the initial submanifold ℓ_F .

Our purpose is to construct a framework to describe the perestroika of singularities of $\tilde{\pi}(\ell_t)$. Then we consider what the time t is. We can construct a Legendrian submanifold $\mathcal{L} \subset J^1(\mathbb{R}^n, \mathbb{R})$ associated with the Lagrangian submanifold $L \subset T^*\mathbb{R}^n$ by the following way : We fix a point $(x_0, p_0) \in L$. Then for every point $(x, p) \in L$ and every path $\gamma = \gamma(t)$, $\gamma(0) = (x_0, p_0)$, $\gamma(1) = (x, p)$, we define a function

$$S(\gamma, (x, p)) = \int_\gamma \alpha|_L.$$

Then

$$\mathcal{L} = \bigcup_{([\gamma], (x, p))} \{(x, S(x, p), p)\},$$

where $[\gamma]$ is the homotopy class represented by the path γ joining (x_0, p_0) and (x, p) . Since we consider the local situation, then \mathcal{L} does not depend on the choice of γ . For any $(x, p) \in \ell_t \subset L$, we put $(x', p') = T_{-1}(x, p) \in \ell$. Then there exists a path γ in ℓ such that $\gamma(0) = (x_0, p_0)$ and $\gamma(1) = (x', p')$. We denote $\phi(t) = (x(t), p(t))$ which is defined by $\phi(\tau) = T_\tau(x', p')$. Since $\alpha|_\ell = 0$, then we have

$$\begin{aligned} S(x(t), p(t)) &= \int_{\phi \cdot \gamma} \alpha|_L = \int_0^t \sum_{i=1}^n p_i(\tau) \frac{dx_i}{dt}(\tau) d\tau \\ &= \int_0^t \sum_{i=1}^n p_i(\tau) \frac{\partial H}{\partial p_i}(x(\tau), p(\tau)) d\tau. \end{aligned}$$

Since H is starshaped, then we have $\frac{dS(x(t), p(t))}{dt}|_{t=0} \neq 0$. This means that $\pi_2|_{\mathcal{L}} : \mathcal{L} \rightarrow \mathbb{R}$ is a submersion germ. According to this fact, we will introduce the notion of graphlike Legendrian unfoldings in the next section.

2. Graphlike Legendrian unfoldings

In which we shall introduce the notion of graphlike Legendrian unfoldings and study some fundamental properties. In ([10,11]) we defined the notion of Legendrian unfoldings which is very useful in the theory of first order partial differential equations. However it is still too general for our purpose.

Let $(\mathcal{L}, (x_0, t_0, p_0)) \subset J^1(\mathbb{R}^n, \mathbb{R})$ be a Legendrian submanifold germ (i.e. $\dim \mathcal{L} = n$ and $\theta|_{\mathcal{L}} = 0$), we say that \mathcal{L} is a *graphlike Legendrian unfolding* if $\pi_2|_{\mathcal{L}}$ is a submersion germ. We remark that $\mathcal{L}_{t_0} = \mathcal{L} \cap \{t = t_0\}$ is an $(n-1)$ -dimensional submanifold and $\Pi(\mathcal{L}_{t_0}) = \ell_{t_0}$ is an isotropic submanifold in $(T^*\mathbb{R}^n, (x_0, p_0))$. Here we call $\tilde{\pi}(\ell_{t_0})$ a *wave front* of ℓ_{t_0} . In order to study perestroikas of wave fronts of graphlike Legendrian unfoldings, we now introduce the following equivalence relation among graphlike Legendrian unfoldings. Let $(\mathcal{L}_i, (x_i, t_i, p_i)) \subset J^1(\mathbb{R}^n, \mathbb{R})$ ($i = 0, 1$) be graphlike Legendrian unfoldings. We say that $(\mathcal{L}_0, (x_0, t_0, p_0))$ and $(\mathcal{L}_1, (x_1, t_1, p_1))$ are *P-Legendrian equivalent* if there exists a contact diffeomorphism

$$\Phi : (J^1(\mathbb{R}^n, \mathbb{R}), (x_0, t_0, p_0)) \rightarrow (J^1(\mathbb{R}^n, \mathbb{R}), (x_1, t_1, p_1))$$

of the form

$$\Phi(x, t, p) = (\phi_1(x, t), \phi_2(t), \phi_3(x, t, p))$$

such that $\Phi(\mathcal{L}_0) = \mathcal{L}_1$.

This equivalence relation preserves the diffeomorphic type of perestroikas of wave fronts of graphlike Legendrian unfoldings. We can define the notion of the stability with respect to P-Legendrian equivalence exactly the same way as that of the ordinary Legendrian stability (see [1,2,3,14]).

By the Arnol'd-Zakalyukin's theory of Legendrian singularities, we can construct generating families of graphlike Legendrian unfoldings. Let $F : (\mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^k, 0) \rightarrow (\mathbb{R}, 0)$ be a function germ such that $\frac{\partial F}{\partial t} \neq 0$ and $(d_2 F, F)|_{\{0\} \times \mathbb{R}^n \times \mathbb{R}^k}$ is non-singular,

where $d_2F(x, q) = (\frac{\partial F}{\partial q_1}(x, q), \dots, \frac{\partial F}{\partial q_k}(x, q))$. Then (d_2F, F) is a submersion germ, so that $C_*(F) = (d_2F, F)^{-1}(0)$ is an n -dimensional smooth submanifold. We now define a map germ

$$\Phi_F : (C_*(F), 0) \rightarrow J^1(\mathbb{R}^n, \mathbb{R})$$

by

$$\Phi_F(t, x, q) = (x, t, \frac{\partial F}{\partial t}, \dots, \frac{\partial F}{\partial t}).$$

It is easy to show that $\Phi_F^*\theta = 0$ and $\pi_2 \circ \Phi_F$ is a submersion germ, so that Φ_F is a graphlike Legendrian unfolding. By the theory of Arnol'd-Zakalyukin, we can show the following :

Proposition 2.1. *All graphlike Legendrian unfolding germs are constructed by the above method.*

We call F a *generating family* of Φ_F . We also consider an equivalence relation among generating families of graphlike Legendrian unfoldings. Let

$$F_i : (\mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^k, 0) \rightarrow (\mathbb{R}, 0) \quad (i = 0, 1)$$

be generating families of Φ_{F_i} . We say that F_0 and F_1 are *t-P-K-equivalent* if there exists a diffeomorphism germ

$$\Phi : (\mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^k, 0) \rightarrow (\mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^k, 0)$$

of the form

$$\Phi(t, x, q) = (\phi_1(t), \phi_2(t, x), \phi_3(t, x, q))$$

such that

$$\langle F_1 \circ \Phi \rangle_{\mathcal{E}(t, x, q)} = \langle F_0 \rangle_{\mathcal{E}(t, x, q)},$$

where $\langle F_0 \rangle_{\mathcal{E}(t, x, q)}$ denotes the ideal generated by F_0 in the ring of function germs of (t, x, q) -variables at the origin. The definition of *the stable t-P-K-equivalence* is given by the usual way (see [1,14]).

For a generating family F of Φ_F , we put

$$T_e(\text{P-K})(f) = \langle \frac{\partial f}{\partial q_1}, \dots, \frac{\partial f}{\partial q_k}, f \rangle_{\mathcal{E}(x, q)} + \langle \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \rangle_{\mathcal{E}_x},$$

where $f = F|_{0 \times \mathbb{R}^n \times \mathbb{R}^k}$. We also say that F is a *P-K-versal deformation* of f if

$$\mathcal{E}_{(x, q)} = \langle \frac{\partial F}{\partial t} |_{0 \times \mathbb{R}^n \times \mathbb{R}^k} \rangle_{\mathbb{R}} + T_e(\text{P-K})(f).$$

Then we have the following proposition whose proof is just like as that of the ordinary theory of Legendrian singularities ([1,14]).

Proposition 2.2. (1) Let F_i ($i = 0, 1$) be generating families of Φ_{F_i} . Then Φ_{F_0} and Φ_{F_1} are P -Legendrian equivalent if and only if F_0 and F_1 are stably t - P - \mathcal{K} -equivalent.
(2) Let F be a generating family of Φ_F , then Φ_F is stable with respect to the P -Legendrian equivalence if and only if F is a P - \mathcal{K} -versal deformation of f .

We can classify generic graphlike Legendrian unfoldings by the P -Legendrian equivalence for $n \leq 5$ by the aid of classifications of one parameter perestroikas of wave fronts. In [15] Zakalyukin has given a generic classification of function germs $F : (\mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^k, 0) \rightarrow (\mathbb{R}, 0)$ by the stable t - P - \mathcal{K} -equivalence. Since the set of function germs $F(t, x, q)$ which satisfy $\frac{\partial F}{\partial t} \neq 0$ is an open subset, then such a function germ is stably t - P - \mathcal{K} -equivalent to one of germs in the following list :

$$({}^0A_r) \quad q_1^{r+1} \pm q_2^2 + \sum_{i=1}^r x_i q_1^{i-1} + t \quad (1 \leq r \leq n)$$

$$({}^0D_r) \quad q_1^2 q_2 \pm q_2^{r-1} + \sum_{i=2}^r x_i q_2^{i-2} + x_1 q_1 + t \quad (4 \leq r \leq n)$$

$$({}^1A_r) \quad q_1^{r+1} \pm q_2^2 + q_1^{r-1} (t \pm x_r^2 \pm \dots \pm x_n^2) + \sum_{i=1}^{r-1} x_i q_1^{i-1} + t \quad (2 \leq r \leq n+1)$$

$$({}^1D_r) \quad q_1^2 q_2 \pm q_2^{r-1} + q_2^{r-2} (t \pm x_r^2 \pm \dots \pm x_n^2) + \sum_{i=2}^{r-1} x_i q_2^{i-2} + x_1 q_1 + t \quad (4 \leq r \leq n+1)$$

$$({}^1E_6) \quad q_1^3 + q_2^4 + (q_1 q_2^2 + 1)t + x_5 q_1 q_2 + x_4 q_2^2 + x_3 q_1 + x_2 q_2 + x_1.$$

Since $(d_2 F, F)|_0 \times \mathbb{R}^n \times \mathbb{R}^k$ for germs 1A_1 and 1A_2 are not submersive, then these germs can be removed from the list of generating families of graphlike Legendrian unfoldings. Thus we have the following :

Theorem 2.3. For $n \leq 5$, the generic graphlike Legendrian unfolding is P -Legendrian equivalent to one of germs of the following type :

$${}^0A_r \quad (1 \leq r \leq n), \quad {}^0D_r \quad (4 \leq r \leq n), \quad {}^1A_r \quad (3 \leq r \leq n), \quad {}^1D_r \quad (4 \leq r \leq n), \quad {}^1E_6.$$

We remark that we can choose a generating family of the graphlike Legendrian unfolding Φ_F of the form $F(t, x, q) = f(x, q) - t$ by the implicit function theorem. In this case the function $f(x, q)$ should satisfy that $(d_2 f, f)$ is non-singular, so that it gives a generating family of the initial isotropic submanifold $\Phi_F|_{t=0}$.

3. Optical graphlike Legendrian unfoldings

In this section we return to study perestroikas of optical wave fronts. For our purpose, we now introduce the following : Let $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ be a starshaped Hamiltonian function germ and $(\mathcal{L}, (x_0, t_0, p_0)) \subset J^1(\mathbb{R}^n, \mathbb{R})$ be a graphlike Legendrian unfolding. We say that \mathcal{L} is *optical relative to H* if $\mathcal{L} \subset \tilde{H}^{-1}(c)$, where \tilde{H} is the induced contact Hamiltonian function germ from H . Then we have the following theorem

Theorem 3.1. (1) Let $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ be a starshaped Hamiltonian function germ and $(\ell, (x_0, p_0)) \subset T^*\mathbb{R}^n$ be an optical Legendrian submanifold relative to H . Then there exists a graphlike Legendrian unfolding $(\mathcal{L}, (x_0, 0, p_0))$ which is optical relative to H such that $\tilde{\pi}(\mathcal{L} \cap \{t = 0\}) = \ell$.

(2) Let $(\mathcal{L}, (x_0, t_0, p_0)) \subset J^1(\mathbb{R}^n, \mathbb{R})$ be a graphlike Legendrian unfolding. Then there exists a starshaped Hamiltonian function germ $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ such that \mathcal{L} is optical relative to H .

In order to prove this theorem, we need the following quite useful theorem in symplectic geometry.

Theorem 3.2. (Kostant-Sternberg [8]) Let (P, ω) be a symplectic manifold, L a Lagrangian submanifold and β a smooth 1-form on P with $\beta|_L = 0$ and $d\beta = \omega$. Then there exists a tubular neighbourhood V of L in P , and a unique vector bundle isomorphism $K : V \rightarrow (T^*L, \alpha_L)$ such that K is the identity on L and $K^*\alpha_L = \beta$. Here, α_L is the canonical 1-form on T^*L .

Proof. (1) This assertion has been already proved in §1.

(2) Since \mathcal{L} is a graphlike Legendrian unfolding, then we can choose a local parametrization $\phi : (\mathbb{R}^n, (u_0, t_0)) \rightarrow (\mathcal{L}, (x_0, t_0, p_0))$ of \mathcal{L} of the form $\phi(u, t) = (x(u, t), t, p(u, t))$, where $u = (u_1, \dots, u_{n-1})$. Furthermore, we have $\phi^*\theta = 0$, so that it is equivalent to

$$(a) \quad \sum_{i=1}^n p_i(u, t) \frac{\partial x_i}{\partial u_j}(u, t) = 0 \quad j = 1, \dots, n-1,$$

$$(b) \quad \sum_{i=1}^n p_i(u, t) \frac{\partial x_i}{\partial t}(u, t) = 1.$$

We now consider a vector field

$$X = \left(\frac{\partial x}{\partial t}(u, t), 1, \frac{\partial p}{\partial t}(u, t) \right)$$

on \mathcal{L} , so that we have

$$\tilde{\pi}_*X = \left(\frac{\partial x}{\partial t}(u, t), \frac{\partial p}{\partial t}(u, t) \right) \in TL,$$

where $L = \tilde{\pi}(\mathcal{L})$. Since L is a Lagrangian submanifold germ in $(T^*\mathbb{R}^n, (x_0, p_0))$, then there exist a decomposition $I \cup J = \{1, \dots, n\}$ and a function germ $S(x_I, p_J)$ such that

$$L = \left\{ (x_I, x_J, p_I, p_J) \mid x_J = -\frac{\partial S}{\partial p_J}, p_I = \frac{\partial S}{\partial x_I} \right\},$$

(for detail see [1,14]). If we define an 1-form β on $T^*\mathbb{R}^n$ by $\beta = dS - p_I dx_I + x_J dp_J$, then we have $\beta|_L = 0$. Thus we can use β as the 1-form in Theorem 3.2. It follows that there exists a tubular neighbourhood V of L in $T^*\mathbb{R}^n$ and a unique vector bundle isomorphism $K : V \rightarrow (T^*L, \alpha_L)$ such that $k|_L = id_L$ and $K^*\alpha_L = \beta$.

Since $\tilde{\pi}_*X$ is a vector field on L , then we can lift this vector field $X_{H'}$ on T^*L , where $H' : (T^*L, (x_0, p_0)) \rightarrow \mathbb{R}$ is a Hamiltonian function germ. If we adopt the canonical local coordinate (\bar{x}, \bar{p}) of T^*L around (x_0, p_0) , where \bar{x} is a local coordinate of L , then L is the submanifold defined by $\bar{p}_i = 0$ ($i = 1, \dots, n$). We now define a Hamiltonian function germ $H'' : (T^*L, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ by $H''(\bar{x}, \bar{p}) = H'(\bar{x}, \bar{p}) - H'(\bar{x}, 0) + c$. By the definition, we have $H''(L) = c$. The Hamiltonian vector field $X_{H''}$ on T^*L has the following component :

$$\begin{cases} \frac{\partial \bar{p}_i}{\partial t} = \frac{\partial H''}{\partial \bar{x}_i} \\ \frac{\partial \bar{x}_i}{\partial t} = -\frac{\partial H''}{\partial \bar{p}_i}, \end{cases}$$

$i = 1, \dots, n$.

On the other hand, we have $\frac{\partial H''}{\partial \bar{p}_i} = \frac{\partial H'}{\partial \bar{p}_i}$ and

$$\frac{\partial H''}{\partial \bar{x}_i}(\bar{x}, 0) = \frac{\partial H'}{\partial \bar{x}_i}(\bar{x}, 0) - \frac{\partial H'}{\partial \bar{x}_i}(\bar{x}, 0) = 0 \text{ on } L,$$

so that $X_{H''}|_L = X_{H'}|_L = \tilde{\pi}_*X$.

Finally, we define a Hamiltonian function germ $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ by $H = H'' \circ K$. Since $K|_L = id_L$, then the Hamiltonian vector field X_H is an extension of $\tilde{\pi}_*X$ on $T^*\mathbb{R}^n$ and $H(L) = c$. It follows that we have relations

$$\begin{cases} \frac{\partial x_i}{\partial t}(u, t) = \frac{\partial H}{\partial p_i}(x(u, t), p(u, t)), \\ \frac{\partial p_i}{\partial t}(u, t) = -\frac{\partial H}{\partial x_i}(x(u, t), p(u, t)) \end{cases}$$

on L . By the equality (b), we have

$$1 = \sum_{i=1}^n p_i(u, t) \frac{\partial x_i}{\partial t}(u, t) = \sum_{i=1}^n p_i(u, t) \frac{\partial H}{\partial p_i}(x(u, t), p(u, t)).$$

Thus, H is starshaped near (x_0, p_0) . This completes the proof

This theorem guarantees that the class of graphlike Legendrian unfoldings supplies the correct class which describes perestroikas of optical wave fronts. However, for our first question, we must deal with in the case when the Hamiltonian function germ H is fixed.

4. Optical graphlike Legendrian unfoldings relative to a fixed Hamiltonian

In this section we consider the perestroikas of optical wave fronts relative to a fixed Hamiltonian function H . Let $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}^n, 0)$ be a starshaped Hamiltonian function germ. Since $\tilde{\pi} : H^{-1}(c) \rightarrow \mathbb{R}^n$ is a Legendrian fibration, then there exists a

local contact diffeomorphism $K : PT^*\mathbb{R}^n \rightarrow H^{-1}(c)$ and a diffeomorphism $h : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $\tilde{\pi} \circ K = h \circ p$, where $p : PT^*\mathbb{R}^n \rightarrow \mathbb{R}^n$, is the projective cotangent bundle (see [1,2,3,14]). If ℓ is a Legendrian submanifold germ in $PT^*\mathbb{R}^n$, then $K(\ell)$ is also a Legendrian submanifold germ in $H^{-1}(c)$. Of course, ℓ and $K(\ell)$ are Legendrian equivalent, so that any wave front can be realized by an optical one. However, for the perestroikas of wave fronts, the situation is different from the above. Consider the following example :

Example 4.1. Let $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ be a Hamiltonian function germ defined by $H(x, p) = p_n$, where $(x_0, p_0) = (0, \dots, 0, 1)$. Let $F(t, x, q) = f(x, q) - t$ be a generating family of the graphlike Legendrian unfolding Φ_F . Suppose that $\Phi_F|_{t=0}$ is an optical Legendrian submanifold relative to H . Then we have

$$H(x, \frac{\partial f}{\partial x}(x, q)) \equiv 1 \pmod{\langle f, \frac{\partial f}{\partial q} \rangle_{\mathcal{E}(x, q)}}.$$

Because $H(x, \frac{\partial f}{\partial x}(x, q)) = \frac{\partial f}{\partial x_n}(x, q)$, we have

$$-1 \in \langle \frac{\partial f}{\partial x_n} \rangle_{\mathcal{E}_x} + \langle f, \frac{\partial f}{\partial x} \rangle_{\mathcal{E}(x, q)} \subset T_e(p\mathcal{K})(f).$$

It follows that $F(t, x, q)$ is not a $P\mathcal{K}$ -versal deformation of $f(x, q)$ if $P\mathcal{K}\text{-cod}(f) \geq 1$. Thus, for example, types of 1A_r and 1D_r in Theorem 2.3 cannot be realized by an optical one relative to H .

Hence, we should assume a kind of non-degeneracy condition on a Hamiltonian function germ. We say that a Hamiltonian function germ $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ is *non-degenerated* if

(1) H is starshaped.

(2) The quadratic form $\sum_{1 \leq i, j \leq n} \frac{\partial^2 H}{\partial p_i \partial p_j}(x_0, p_0) \xi_i \xi_j$ does not vanish on the null space of $(\frac{\partial H}{\partial p_1}(x_0, p_0), \dots, \frac{\partial H}{\partial p_n}(x_0, p_0))$.

The second condition of the above definition is given by Duistermaat (Proposition 5.2.1 in [7]) in order to study optical Lagrangian submanifolds. It is easy to check that the Hamiltonian function in Example 1.2 satisfies the above conditions. Then we have the following theorem which gives an almost complete answer to our first question.

Theorem 4.2. Let $H : (T^*\mathbb{R}^n, (x_0, p_0)) \rightarrow (\mathbb{R}, c)$ be a non-degenerated Hamiltonian function and $(\mathcal{L}, (x_1, t_1, p_1)) \subset J^1(\mathbb{R}^n, \mathbb{R})$ be a P -Legendrian stable graphlike Legendrian unfolding. Then there exists an optical graphlike Legendrian unfolding $(\mathcal{L}', (x_0, t_0, p_0))$ relative to H such that \mathcal{L} and \mathcal{L}' are P -Legendrian equivalent.

Proof. Without loss of generality, we may assume that $(x_0, t_0) = (0, 0)$. Let

$$G : (\mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^k, 0) \rightarrow (\mathbb{R}, 0)$$

be a generating family of the graphlike Legendrian unfolding \mathcal{L} . Since $(d_2G, G)|_{0 \times \mathbb{R}^n \times \mathbb{R}^k}$ is non-singular, then the set

$$\phi_g = \{((x_1, \dots, x_n), [\frac{\partial g}{\partial x_1}(x, q); \dots; \frac{\partial g}{\partial x_n}(x, q)]) | d_2g(x, q) = g(x, q) = 0\}$$

is a Legendrian submanifold in $PT^*\mathbb{R}^n$, where $g = G|_{0 \times \mathbb{R}^n \times \mathbb{R}^k}$. Then $\ell_0 = K(\psi_g)$ is an optical Legendrian submanifold relative to H which is Legendrian equivalent to ψ_g , where K is the contact diffeomorphism like as the previous arguments. It follows from Theorem 3.1 that there exists an optical graphlike Legendrian unfolding \mathcal{L}' relative to H such that $\Pi(\mathcal{L}'|_{t=0}) = \ell_0$.

We now choose a generating family $F(t, x, q) = f(x, q) - t$ of \mathcal{L}' . by the definition, $\mathcal{L}'|_{t=0}$ is given by

$$\Phi_F|_{t=0} = \{(x, 0, \frac{\partial f}{\partial x_1}(x, q), \dots, \frac{\partial f}{\partial x_n}(x, q) | d_2 f(x, q) = f(x, q) = 0\},$$

so that f and g are stably P- \mathcal{K} -equivalent by the Arnol'd-Zakalyukin's theory ([1,2,3,14]). Thus we may assume that f and g are P- \mathcal{K} -equivalent.

If P- \mathcal{K} -cod(f) = 0, then P- \mathcal{K} -cod(g) = 0, so that $f(x, q)$ is already a P- \mathcal{K} -versal deformation of itself. Hence, $F(t, x, q)$ is also a P- \mathcal{K} -versal deformation of $f(x, q)$. It follows from the uniqueness theorem of P- \mathcal{K} -versal deformations (see [6,8]) that $F(t, x, q)$ and $G(t, x, q)$ are t-P- \mathcal{K} -equivalent. Then $\mathcal{L}' = \Phi_F$ and $\mathcal{L} = \Phi_G$ are P-Legendrian equivalent.

Thus, we now assume that P- \mathcal{K} -cod(g) = 1, so that P- \mathcal{K} -cod(f) = 1. If $-1 = \frac{\partial F}{\partial t}|_{t=0} \notin T_e(\text{P-}\mathcal{K})(f)$, then we can get the required assertion by the uniqueness of the P- \mathcal{K} -versal deformation like as the previous case.

Suppose that $1 \in T_e(\text{P-}\mathcal{K})(f)$ for any generating family of \mathcal{L}' of the form $F(t, x, q) = f(x, q) - t$. Since $H(\ell_0) = c$, then we have a relation

$$H(x, \frac{\partial f}{\partial x}(x, q)) \equiv c \pmod{\langle f, \frac{\partial f}{\partial q} \rangle_{\mathcal{E}_{(x,q)}}},$$

so that

$$H(0, \frac{\partial f}{\partial x}(0, q)) \equiv c \pmod{\langle f_0, \frac{\partial f_0}{\partial q} \rangle_{\mathcal{E}_q}},$$

where $f_0 = f|_{x=0}$. We may assume that $f_0 \in \mathfrak{M}_k^3$. We now consider the Taylor polynomial of $H(x, p)$ of degree 2 at (x, p_0) with respect to $p = (p_1, \dots, p_n)$ -variables as follows :

$$\begin{aligned} H(x, p) &= H(x, p_0) + \sum_{i=1}^n \frac{\partial H}{\partial p_i}(x, p_0)(p_i - p_{0,i}) \\ &\quad + \frac{1}{2} \sum_{i,j} \frac{\partial^2 H}{\partial p_i \partial p_j}(x, p_0)(p_i - p_{0,i})(p_j - p_{0,j}) + \text{higher term.} \end{aligned}$$

Since $H(0, p_0) = c$, then we have

$$\begin{aligned} H(0, \frac{\partial f}{\partial x}(0, q)) &= c + \sum_{i=1}^n \frac{\partial H}{\partial p_i}(0, \frac{\partial f}{\partial x}(0, q))(\frac{\partial f}{\partial x_i}(0, q) - p_{0,i}) \\ &\quad + \frac{1}{2} \sum_{i,j} \frac{\partial^2 H}{\partial p_i \partial p_j}(0, \frac{\partial f}{\partial x}(0, q))(\frac{\partial f}{\partial x_i}(0, q) - p_{0,i})(\frac{\partial f}{\partial x_j}(0, q) - p_{0,j}) + \text{higher term.} \end{aligned}$$

Because $H(0, \frac{\partial f}{\partial x}(0, q)) - c \in \langle \frac{\partial f_0}{\partial q}, f_0 \rangle_{\mathcal{E}_q} \subset \mathfrak{M}_k^2$, the right hand side of the above equality is a function of $q = (q_1, \dots, q_k)$ -variables whose order is at least two. Since $F(t, x, q) = f(x, q) - t$ is a generating family of the graphlike Legendrian unfolding \mathcal{L}' , then

$$\text{rank} \left(\frac{\partial^2 f}{\partial q_i \partial x_j}(0) \right) = k.$$

Then we may assume that $(\frac{\partial^2 f}{\partial q_i \partial x_j}(0) = \delta_{ij}$ for $i, j = 1, \dots, k$ and $\frac{\partial f}{\partial x_\ell}(0, q) \in \mathfrak{M}_k^2$ for $\ell = k+1, \dots, n$, then $\frac{\partial f}{\partial x_i}(0, q) = q_i + \psi(q)$, where $\psi(q) \in \mathfrak{M}_k^2$, so that

$$\sum_{i=1}^k \frac{\partial H}{\partial p_i}(0, p_0) q_i = 0.$$

Since H is non-degenerated, then

$$\sum_{i=1}^k \frac{\partial H}{\partial p_i \partial p_j}(0, p_0) q_i q_j \neq 0.$$

On the other hand, we have

$$1 \in \langle \frac{\partial f}{\partial x_1}(0, q), \dots, \frac{\partial f}{\partial x_n}(0, q) \rangle_{\mathbb{R}} + \langle f_0, \frac{\partial f_0}{\partial q} \rangle_{\mathcal{E}_q}$$

by the assumption. It follows that

$$\begin{aligned} & \frac{1}{2} \sum_{1 \leq i, j \leq n} \frac{\partial^2 H}{\partial p_i \partial p_j}(0, p_0) \left(\frac{\partial f}{\partial x_i}(0, q) - p_{0,i} \right) \left(\frac{\partial f}{\partial x_j}(0, q) - p_{0,j} \right) \\ & \equiv H(0, \frac{\partial f}{\partial x}(0, q)) - c - \sum_{i=1}^n \frac{\partial H}{\partial p_i}(0, p_0) \left(\frac{\partial f}{\partial x_i}(0, q) - p_{0,i} \right) \text{ mod } \mathfrak{M}_k^3 \\ & = H(0, \frac{\partial f}{\partial x}(0, q)) - c - \sum_{i=1}^n \frac{\partial H}{\partial p_i}(0, p_0) \frac{\partial f}{\partial x_i}(0, q) + \sum_{i=1}^n p_{0,i} \frac{\partial H}{\partial p_i}(0, p_0) \\ & \in \langle \frac{\partial f}{\partial x_1}(0, q), \dots, \frac{\partial f}{\partial x_n}(0, q) \rangle_{\mathbb{R}} + \langle f_0, \frac{\partial f_0}{\partial q} \rangle_{\mathcal{E}_q} \text{ mod } \mathfrak{M}_k^3. \end{aligned}$$

Let $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear isomorphism, then we have

$$\frac{\partial f(Ax, q)}{\partial x_i} = \sum_{j=1}^n A_{ij} \frac{\partial f}{\partial x_j}(Ax, q).$$

Since

$$\begin{aligned} & \sum_{1 \leq i, j \leq n} \frac{\partial^2 H}{\partial p_i \partial p_j}(0, p_0) \left(\frac{\partial f}{\partial x_i}(0, q) - p_{0,i} \right) \left(\frac{\partial f}{\partial x_j}(0, q) - p_{0,j} \right) \\ & \equiv \sum_{1 \leq i, j \leq k} \frac{\partial^2 H}{\partial p_i \partial p_j}(0, p_0) q_i q_j \text{ mod } \mathfrak{M}_k^3 \end{aligned}$$

and the vector space

$$\left\langle \frac{\partial f}{\partial x_1}(0, q), \dots, \frac{\partial f}{\partial x_n}(0, q) \right\rangle_{\mathbb{R}} + \left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q} \bmod \mathfrak{M}_k^3$$

is invariant under the action of linear isomorphisms $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$, then every quadratic forms of $q = (q_1, \dots, q_k)$ -variables are contained in the above vector space. If there exists a quadratic form of q -variables such that it is contained in $\left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q} \bmod \mathfrak{M}_k^3$, then every quadratic forms are contained in it by the same reason as that of the above. In this case, since the vector space $\left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q} \bmod \mathfrak{M}_k^3$ has dimension k , then k should be 1 and f_0 is an A_2 -type function germ. It follows from Theorem 2.3 that $F(t, x, q)$ is 0A_2 -type, so that this case is contained in the case of $\text{P-}\mathcal{K}\text{-cod}(f) = 0$.

We may assume that every quadratic forms of q -variables are contained in

$$\left\langle \frac{\partial f}{\partial x_1}(0, q), \dots, \frac{\partial f}{\partial x_n}(0, q) \right\rangle_{\mathbb{R}} \bmod \mathfrak{M}_k^3.$$

Since $\mathcal{K}\text{-cod}(f_0)$ is finite (for the definition of \mathcal{K} -finiteness, see [13]), then there exists $r \in \mathbb{N}$ such that $\mathfrak{M}_k^r \subset \left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q}$. By the same arguments as those of the previous paragraph, we can assert that every monomials of q -variables of degree 3 are contained in the vector space

$$\left\langle \frac{\partial f}{\partial x_1}(0, q), \dots, \frac{\partial f}{\partial x_n}(0, q) \right\rangle_{\mathbb{R}} + \left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q} \bmod \mathfrak{M}_k^4.$$

If there exists a monomial of degree 3 which is contained in $\left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q} \bmod \mathfrak{M}_k^4$, then k should be 1 and f_0 is an A_3 -type function germ. It follows that

$$\dim_{\mathbb{R}} \left\langle \frac{\partial f}{\partial x_1}(0, q), \dots, \frac{\partial f}{\partial x_n}(0, q) \right\rangle_{\mathbb{R}} \geq 3 \bmod \mathfrak{M}_k^4.$$

By Theorem 2.3, $F(t, x, q)$ should be 0A_3 -type, then this case is contained in the case of $\text{P-}\mathcal{K}\text{-cod}(f) = 0$.

Since we can remove the assumption $n \leq 5$ for ${}^0A_\ell$ or ${}^1A_\ell$ -type germs in Theorem 2.3 (cf. Theorem 2.2 in [15]), we can continue this procedure up to degree $r - 1$. Eventually, it remains the case that every polynomial of degree $r - 1$ are contained in the vector space

$$\left\langle \frac{\partial f}{\partial x_1}(0, q), \dots, \frac{\partial f}{\partial x_n}(0, q) \right\rangle_{\mathbb{R}} + \left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q} \bmod \mathfrak{M}_k^r.$$

Since $\mathfrak{M}_k^r \subset \left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q}$, then we have

$$\mathcal{E}_q = \left\langle \frac{\partial f}{\partial x_1}(0, q), \dots, \frac{\partial f}{\partial x_n}(0, q) \right\rangle_{\mathbb{R}} + \left\langle f_0, \frac{\partial f_0}{\partial q} \right\rangle_{\mathcal{E}_q}.$$

It follows that

$$\mathcal{E}_{(x, q)} = T_e(\text{P-}\mathcal{K})(f) + \mathfrak{M}_k \mathcal{E}_{(x, q)},$$

so that we have

$$\mathcal{E}_{(x, q)} = T_e(\text{P-}\mathcal{K})(f)$$

by the Malgrange preparation theorem. This contradicts to the fact that $\text{P-}\mathcal{K}\text{-cod}(f) = 1$. This completes the proof.

Remark. The above proof is motivated by the proof of Proposition 5.2.1 in [7]. In fact, Theorem 4.2 is invented by the attempt to understand the proof of the proposition.

References

1. V. I. Arnol'd, S. M. Gusein-Zade and A. N. Varchenko, *Singularities of Differentiable Maps vol. I*, Birkhäuser, 1986.
2. V. I. Arnol'd, *Contact geometry and wave propagation*, Monographie de l'Enseignement. Math. 34 (1989).
3. V. I. Arnol'd, *Singularities of Caustics and Wave fronts, Mathematics and Its Applications (Soviet Series)*, vol. 62, Kluwer Academic Publisher, 1990.
4. J. W. Bruce, *Wavefronts and parallels in Euclidean space*, Math. Proc. Cambridge Phil. Soc. 93 (1983), 323-333.
5. J. W. Bruce and D. J. Hurley, *Focal sets in certain Riemannian manifolds*, Proc. Edinburgh Math. Soc. 27 (1984), 209-214.
6. J. Damon, *The unfolding and determinacy theorems for subgroups of \mathcal{A} and \mathcal{K}* , Memoirs of Amer. Math. Soc. 50-306 (1984).
7. J. J. Duistermaat, *Oscillatory integrals, Lagrange immersions and Unfolding of singularities*, Commu. pure and applied Math. 27 (1974), 207-281.
8. M. Golubitsky and V. Guillemin, *Contact equivalence for Lagrange manifolds*, Adv. In Math. 15 (1975), 375-387.
9. S. Izumiya, *Generic bifurcations of varieties*, Manuscripta Math. 46 (1984), 137-164.
10. S. Izumiya, *The theory of Legendrian unfoldings and first order differential equations*, to appear in Proceedings of the Royal Society of Edinburgh.
11. S. Izumiya, *Completely integrable holonomic systems of first order differential equations*, preprint.
12. V. V. Lychagin, *Local classification of non-linear first order partial differential equations*, Russian Math. Surveys 30 (1975), 105-175.
13. C. T. C. Wall, *Finite determinacy of smooth map germs*, Bull. London Math. Soc. 14 (1981), 481-539.
14. V. M. Zakalyukin, *Lagrangian and Legendrian singularities*, Funct. Anal. Appl. 10 (1976), 23-31.
15. V. M. Zakalyukin, *Reconstructions of fronts and caustics depending on a parameter and versality of mappings*, Journal of Soviet Math. 27 (1983), 2713-2735.

Department of Mathematics, Faculty of Science, Hokkaido University, Sapporo 060, JAPAN
E-mail address: izumiya@math.hokudai.ac.jp